

## A Weierstrass Theorem for Real Banach Spaces

VASILE I. ISTRĂȚESCU

*Institut de Mathématique, Calea Griviței, 21, Bucarest 12, Romania*

*Communicated by R. C. Buck*

Received March 25, 1975

Let  $X$  be a real separable Banach space and  $K$  be a compact subset;  $C(K)$  denotes the family of continuous functions from  $K$  into  $X$  together with the uniform norm topology

$$\|f - g\| = \max_{x \in K} \|f(x) - g(x)\|.$$

In some recent papers [3, 4], Prenter has proved that if  $X$  is a Hilbert space or a separable Banach space with “Property M” then the Weierstrass theorem holds in  $C(K)$ . It is the purpose of the present note to give a general theorem without any restriction on  $X$ . Our method is entirely different from Prenter’s.

### 1. PRELIMINARIES

Let  $X$  be a Banach space and  $k \geq 1$ .  $X^k$  denotes the Banach space

$$\underbrace{X \times \cdots \times X}_{k \text{ times}}.$$

DEFINITION 1.1. A  $k$ -linear operator  $T$  on  $X$  is a function on  $X^k$  into  $X$  which is linear in each of its arguments separately.

A 0-linear operator  $L_0$  on  $X$  is a constant function on  $X$  into  $X$  and we shall identify a 0-linear operator  $L_0$  with its range.

The norm of a  $k$ -linear operator  $T$  is the number

$$\|T\| = \sup \|T(x_1, \dots, x_k)\| / \|x_1\| \|x_2\| \cdots \|x_k\|.$$

If  $T$  is a  $k$ -linear operator and  $S$  is a  $q$ -linear operator then we can define in a natural way the products  $TS$  and  $ST$  which are obviously  $(k + q)$ -linear operators. By a polynomial we mean any function of the form

$$P_n(x) = L_0x + \cdots + L_nx^n,$$

where  $L_i$  are  $i$ -linear operators. The number  $n$  is called the degree,  $n = \deg P_n$ .

Some explanation of notation is in order. For any  $k \geq 1$  and  $T$  a  $k$ -linear operator on  $X$  we note

$$T(x, x, \dots, x) = Tx^k.$$

It is obvious that the set of polynomials forms an algebra (generally non-commutative) with properties:

1.  $\deg(P_n + P_m) \leq \max\{\deg P_n, \deg P_m\}$ ,
2.  $\deg(P_n \cdot P_m) \leq n + m$ .

For details see any book of functional analysis.

## 2

We consider now a Banach space  $X$ ; we let  $K$  be a compact subset in  $X$  and introduce

- (1)  $C(K, X) = \{f, f: K \rightarrow X \text{ continuously}\}$ ,
- (2)  $P(K, X) = \{p, p \text{ is a continuous polynomial}\}$ ,

and if  $Y \subset R$  (the reals)

- (3)  $C(K, Y) = \{f, f: K \rightarrow Y \text{ continuously}\}$ ,
- (4)  $P(K, Y) = \{f, f \text{ is a continuous polynomial from } K \text{ to } Y\}$ . Since all spaces in (1)–(3) are clearly defined, we give some details for (4). By real polynomial we mean any function of the form

$$p_n(x) = a_0 + a_1x + \dots + a_nx^n,$$

where  $a_0$  is a real constant and

$$a_i: \underbrace{X \times \dots \times X}_i \text{ times} \rightarrow Y \quad (Y \text{ the reals}),$$

with  $a_i(x, \dots, x) = a_i x^i$ . It is clear that  $P(K, Y)$  forms an algebra. Our first basic result gives a relation between  $C(K, Y)$  and  $P(K, Y)$ .

**THEOREM 2.1.** *The algebra  $P(K, R)$  is dense in the uniform norm topology in  $C(K, R)$ .*

*Proof.* Our proof depends upon a basic idea used by de Branges [1]. This consists of remarking that the support of some measures has special proper-

ties with respect to some subspaces and using this for commutative subalgebras of the algebra of all continuous functions on a compact Hausdorff space.

We need the following notion.

**DEFINITION 2.1.** A level set for a family of functions  $S \subset C(K, R)$  is a subset  $K_0 \subset K$  such that  $S/K_0$  contains only constant functions. It is obvious that any point of  $K$  is a level set, and using Zorn's lemma we see that any point is contained in a maximal level set; the same is true for any level set.

**LEMMA 2.2.** *Let  $F$  be a continuous linear functional on  $C(K, R)$  such that*

(1)  *$F$  is an extreme point of the set of all linear functionals  $\lambda$  on  $C(K, R)$  such that  $\|\lambda\| \leq 1$  and  $\lambda$  annihilates  $P(K, R)$ ,*

(2)  *$\mu$  is the measure determining  $F$  so that  $F(f) = \int_K f(s) d\mu(s)$ . Then the support of  $\mu$ ,  $\text{supp } \mu$ , is a level set of  $P(K, R)$ .*

*Proof.* Suppose that is not so. We find an element of  $P(K, R)$ , say  $g_0$ , such that it is not constant on  $\text{supp } \mu$  (which is equal to  $\text{supp } \|\mu\|$ ,  $\|\mu\|$  the total variation of  $\mu$ ). We may suppose without loss of generality that

$$0 \leq g_0 \leq 1.$$

We can consider also the following measures

$$\mu_1 = g_0 \mu, \quad \mu_2 = (1 - g_0) \mu;$$

it is easy to see that

$$\|\mu_1\| + \|\mu_2\| = 1,$$

and thus

$$\mu_1^* = \mu_1 / \|\mu_1\|, \quad \mu_2^* = \mu_2 / \|\mu_2\|$$

have the property

$$\mu = \|\mu_1\| \mu_1^* + \|\mu_2\| \mu_2^*,$$

which represents a contradiction, since  $\mu$  is a measure representing an extreme point. The lemma is proved.

*Remark.* The lemma is valid also for the closure of  $P(K, R)$ ,  $\overline{P(K, R)}$  and the proof is the same.

The proof of Theorem 2.1 is now very simple. Indeed the Hahn–Banach theorem implies that any level set of  $P(K, R)$  contains only one point. By the Krein–Milman and Hahn–Banach theorems combined with the Riesz–Kakutani representation theorem there exists a nonzero functional annihila-

ting  $P(K, R)$ , and it is an extreme point. By the lemma, the support of the corresponding measure reduces to a point, and then our functional is of the form

$$F(f) = f(t_0),$$

for some  $t_0 \in K$ , which is an impossibility since no  $f \in P(K, R)$  vanishes at  $t_0$ . This completes the proof of the theorem.

**DEFINITION 2.3.** A function  $f: K \rightarrow R$  is called normal if it is continuous and  $f(x) \in [0, 1]$ .

The following is well known [2].

**THEOREM 2.4.** For any covering  $V_1, \dots, V_m$ , with open sets, of  $K$ , there exist normal functions  $h_1, \dots, h_m$  such that

- (1)  $h_i(t) = 0$  in the exterior of  $V_i$ ,
- (2)  $\sum_{i=1}^m h_i(t) = 1$ .

We can prove the Weierstrass theorem.

**THEOREM 2.5.** The space  $P(K, X)$  is dense in  $C(K, X)^*$ .

*Proof.* Let  $f \in C(K, X)$  and  $\epsilon > 0$ . For each  $t \in K$ , let

$$V_t = \{s, \|f(t) - f(s)\| < \epsilon\},$$

which is open. Thus  $\{V_t\}$  is an open covering of  $K$  and there exists a finite covering of  $K$ , say,  $V_{t_1}, \dots, V_{t_m}$ . From Theorem 2.4 we find the normal functions  $h_1, \dots, h_m$  and we can construct the function

$$F^*(t) = \sum_1^m f(t_i) h_i(t).$$

It is easy to see that

$$\|f(t) - F^*(t)\| < \epsilon,$$

and, since each  $h_i$  can be approximated by elements of  $P(K, R)$ , we find easily that  $f$  can be approximated by elements of  $P(K, X)$ .

In a paper which is now in preparation we extend these constructions to complex Banach spaces proving in full generality a Weierstrass theorem.

#### ACKNOWLEDGMENT

The author is grateful to Professor R. C. Buck for some useful suggestions.

## REFERENCES

1. L. DE BRANGES, The Stone–Weierstrass theorem, *Proc. Amer. Math. Soc.* **10** (1959), 822–824.
2. G. E. SHILOV, O koltzah funktsii s ravnomernoï skhodimostiū, *Ukrainian Math. J.* (1951), t.III, Nr. 4.
3. P. M. PRENTER, A Weierstrass theorem for real separable Hilbert spaces, *J. Approximation Theory* **3** (1970), 341–351.
4. P. M. PRENTER, A Weierstrass theorem for normed linear spaces, *Bull. Amer. Math. Soc.* **4** (1969) 860–862, and MRC Summary Report #957, Jan., 1969.
5. V. I. ISTRĂȚESCU, A Weierstrass theorem for complex Banach spaces, in preparation.