

# Uniqueness of Solutions to a Two-Dimensional Mean Problem

I. BOROSH AND C. K. CHUI

*Department of Mathematics, Texas A & M University, College Station, Texas 77843*

*Communicated by Oved Shisha*

Received May 11, 1976

Let  $0 < r_m < 1$ ,  $r_m^m \leq \rho$  for all large  $m$ , and let  $w_n = e^{i2\pi/n}$ ,  $n = 1, 2, \dots$ . For a function  $f(z) = \sum a_n z^n$ , holomorphic in the open unit disk  $U$ , let  $s_n(f) = (1/n) \sum_{k=1}^n f(r_n w_n^k)$ , the  $n$ th arithmetic mean of  $f$  over the circle  $|z| = r_n$ . We prove that if  $\rho < 1$  and  $a_n = O(n^{-\alpha_1})$  for  $\alpha_1 = 1.728\dots$ , then  $f$  is uniquely determined by the two-dimensional means  $s_n(f)$ ,  $n = 1, 2, \dots$ . We also prove that for each  $\rho$ ,  $0 < \rho < 1$ , there is a nontrivial  $f$ , holomorphic in  $U$ , such that  $s_n(f) = 0$  for  $n = 1, 2, \dots$  with  $r_n = \rho^{1/n}$ .

## 1. INTRODUCTION AND RESULTS

Let  $U$  denote the open unit disk  $|z| < 1$  in the complex plane and  $H$  the space of functions holomorphic in  $U$ . Let  $0 < r_n < 1$ ,  $n = 1, 2, \dots$ , and consider the means

$$s_n(f) = \frac{1}{n} \sum_{k=1}^n f(r_n e^{i2\pi k/n}) \tag{1.1}$$

of  $f \in H$  on the concentric circles  $|z| = r_n$ . In this note, we study the problem of uniqueness of  $f$  when  $s_n(f)$ ,  $n = 1, 2, \dots$ , are given. This problem was posed in [2] and discussed in [1]. It was proved, in particular, that if  $r_n^n \leq \rho$ ,  $\rho \leq \frac{1}{2}$ , for all  $n$ , and  $f(z) = \sum a_n z^n$  with  $\sum |a_n| < \infty$ , then  $f$  is uniquely determined by the sequence  $s_n(f)$ ,  $n = 1, 2, \dots$ . The condition  $\rho \leq \frac{1}{2}$  was a technical one. Here, by using a different method we prove a uniqueness result for any  $\rho$ ,  $0 < \rho < 1$ . Of course a "smoothness" condition on  $f$  is required. In fact, we also obtain a negative result for each  $\rho$ ,  $0 < \rho < 1$ . We state our main results in the following theorems.

**THEOREM 1.** *Let  $0 < r_n < 1$  with  $r_n^n \leq \rho < 1$  for all large  $n$ . Then there exists an  $\alpha_\rho$ ,  $1 \leq \alpha_\rho < \alpha_1 = 1.728\dots$ , such that any function  $f(z) = \sum_{n=0}^\infty a_n z^n$ , satisfying  $a_n = O(n^{-\alpha_\rho})$  and  $s_n(f) = 0$  for  $n = 1, 2, \dots$ , must be identically zero.*

In the above theorem,  $\alpha_\rho$  is uniquely determined by  $\phi(\rho, \alpha_\rho) = 2\rho$ , where

$$\phi(z, s) = \sum_{n=1}^{\infty} \frac{z^n}{n^s} \tag{1.2}$$

is the so-called polylogarithm function (cf. [3, 4]). In particular,  $\alpha_1$  satisfies  $\zeta(\alpha_1) = 2$  where  $\zeta$  is the Riemann zeta-function. Calculation gives  $\alpha_1 = 1.728\dots$ . As a simple consequence of the above theorem, we have

**COROLLARY 1.** *Let  $0 < r_n < 1$  with  $r_n^n \leq 0.79$  for all large  $n$ . Then any function  $f(z) = \sum_{n=0}^{\infty} a_n z^n$ , satisfying  $a_n = o(n^{-1})$  and  $s_n(f) = 0$  for  $n = 1, 2, \dots$ , must be the zero function.*

On the other hand, we have the following negative result.

**THEOREM 2.** *Let  $0 < \rho < 1$  be given and  $r_n = \rho^{1/n}$ . Then there exists a complex number  $\beta = \beta(\rho)$  such that the function  $f(z) = \phi(z, \beta)$  satisfies  $s_n(f) = 0$  for all  $n, n = 1, 2, \dots$*

## 2. PROOF OF THE MAIN RESULTS

We first prove the following:

**LEMMA.** *Let  $C = (c_{i,j}), i, j = 1, 2, \dots$ , be an upper triangular matrix with nonzero diagonal elements. Suppose that for some  $\alpha \geq 0$ ,*

$$\sum_{j=k+1}^{\infty} |c_{k,j}| j^{-\alpha} \leq k^{-\alpha} |c_{k,k}|, \quad k = 1, 2, \dots \tag{2.1}$$

*Then for every  $b = (b_1, b_2, \dots)^T$  satisfying  $b_n = o(n^{-\alpha})$  and  $Cb = 0, b = 0$ .*

*Proof.* Suppose  $b \neq 0$ . Since  $n^\alpha b_n \rightarrow 0$ , there exists a  $k$  such that  $k^\alpha |b_k|$  is maximum. We pick the largest such  $k$ , so that  $k^\alpha |b_k| > n^\alpha |b_n|$  for all  $n > k$ . Hence, since  $Cb = 0$  and  $c_{k,k} \neq 0$ , we have

$$|c_{k,k}| |b_k| \leq \sum_{j=k+1}^{\infty} |c_{k,j}| |b_j| < k^\alpha |b_k| \sum_{j=k+1}^{\infty} |c_{k,j}| j^{-\alpha}.$$

This is a contradiction to (2.1).

We can now prove Theorem 1. Without loss of generality, we assume that  $r_n^n \leq \rho$  for all  $n$ . Let  $f(z) = \sum a_n z^n$  be in  $H$  satisfying  $a_n = o(n^{-\alpha\rho})$  and  $n = 1, 2, \dots$ . As in [1], it can easily be shown that  $f(0) = a_0 = 0$ , and

$$s_n(f) = \sum_{k=1}^{\infty} r_k^{nk} a_{nk}. \tag{2.2}$$

Hence,  $a = (a_1, a_2, \dots)^T$  satisfies the equation  $Ca = 0$  with  $C = (c_{k,j})$  and

$$\begin{aligned} c_{k,j} &= 0 && \text{if } k \nmid j \\ &= r_k^i && \text{if } k \mid j. \end{aligned}$$

Since  $\phi(x, \alpha_\rho)/x$  is monotone increasing in  $x$ , and  $r_k^k \leq \rho$  for all  $k$ , we have  $\phi(r_k^k, \alpha_\rho)/r_k^k \leq \phi(\rho, \alpha_\rho)/\rho = 2$ . Thus, we have

$$\sum_{t=2}^{\infty} r_k^{kt} t^{-\alpha_\rho} \leq r_k^k$$

or

$$\sum_{t=2}^{\infty} r_k^{kt} (kt)^{-\alpha_\rho} \leq k^{-\alpha_\rho} r_k^k,$$

which is (2.1) with  $\alpha = \alpha_\rho$ . Hence,  $a = 0$ , or  $f \equiv 0$ , by the above lemma.

To prove Corollary 1, we observe that  $\phi(x, 1)/x = -[\log(1 - x)]/x = 2$  for  $x = 0.79\dots$ . Hence, as above, if  $r_n^n \leq 0.79$ , then  $\phi(r_n^n, 1) < 2r_n^n$ , which gives (2.1) with  $\alpha = 1$ .

To prove Theorem 2, we set  $r_n = \rho^{1/n}$  where  $\rho$  is any given positive number less than 1. For this fixed  $\rho$ , it was shown in [3] that the polylogarithm function  $\phi(\rho, s)$  has many complex zeros. Let  $\beta = \beta(\rho)$  be one of them, and define  $f(z) = \phi(z, \beta)$ . We have

$$s_n(f) = \sum_{t=1}^{\infty} \frac{\rho^t}{(nt)^\beta} = \frac{1}{n^\beta} \phi(\rho, \beta) = 0$$

for all  $n = 1, 2, \dots$

We remark that from [3],  $\text{Re } \beta(\rho) < 1$  and we can choose  $\beta(\rho)$  such that  $\beta(\rho) \rightarrow 1$  as  $\rho \rightarrow 1^-$ .

REFERENCES

1. G. R. BLAKLEY, I. BOROSH, AND C. K. CHUI, A two-dimensional mean problem, *J. Approximation Theory* **22** (1978), 11–26.
2. C. K. CHUI AND C. H. CHING, Approximation of functions from their means, in “Approximation Theory” (G. G. Lorentz, Ed.), pp. 307–311, Academic Press, New York, 1973.
3. B. FORNBERG AND K. S. KÖLBIG, Complex zeros of the Jonquiere or polylogarithm function, *Math. Comp.* **29** (1975), 582–599.
4. C. TRUESDELL, On a function which occurs in the theory of the structure of polymers, *Ann. of Math.* **40** (1945), 144–157.