

Note

A Multidimensional Generalization of Slater's Inequality

JOSIP E. PEČARIĆ

*Department of Mathematics and Physics, Faculty of Civil Engineering,
University of Beograd, 11000 Beograd, Yugoslavia*

Communicated by Oved Shisha

Received February 22, 1983; revised September 18, 1984

1

Slater [1] proved the following companion to Jensen's inequality for convex functions:

Suppose that f is convex and nondecreasing (nonincreasing) on (a, b) . Then for $x_1, \dots, x_n \in (a, b)$, $p_1, \dots, p_n \geq 0$, $P_n = p_1 + \dots + p_n > 0$, and $p_1 f'_+(x_1) + \dots + p_n f'_+(x_n) \neq 0$, we have

$$\frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \leq f \left(\frac{\sum_{i=1}^n p_i x_i f'_+(x_i)}{\sum_{i=1}^n p_i f'_+(x_i)} \right). \quad (1)$$

An integral analog of this result is also valid. Both results remain true if at any occurrence of $f'_+(x)$ we write instead any value in the interval $[f'_-(x), f'_+(x)]$.

The following simple generalization of this result was given in [2]:

Suppose that f is a convex function on (a, b) . If for $x_1, \dots, x_n \in (a, b)$, $p_1, \dots, p_n \geq 0$, and $p_1 f'_+(x_1) + \dots + p_n f'_+(x_n) \neq 0$, we have

$$\frac{\sum_{i=1}^n p_i x_i f'_+(x_i)}{\sum_{i=1}^n p_i f'_+(x_i)} \in (a, b),$$

then (1) is valid.

Note that a similar companion inequality to Jensen-Steffensen's inequality was also given in [2]. Some other inequalities, complementary to Jensen's inequality for convex functions, are given in [3 and 4] (see also [5 and 6]).

All these results hold for convex functions of one variable. However, we shall show that a generalization of Slater's inequality to convex functions of several variables is also valid.

2

If $x, y \in R^m$, say, $x = (x_1, \dots, x_m)$, $y = (y_1, \dots, y_m)$, then $\langle x, y \rangle = x_1 y_1 + \dots + x_m y_m$. We shall say that a real function f is convex on an open set I ($I \subseteq R^m$) if the following inequality holds: $f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y)$, $\forall x, y \in I$ and $\forall \lambda \in [0, 1]$.

THEOREM. Let $f: I \rightarrow R$ ($I \subseteq R^m$) be a convex function, and let $x_1, \dots, x_n \in I$, $p_1, \dots, p_n \geq 0$, $P_n > 0$. If $A \in I$ exists such that

$$\left\langle A, \sum_{k=1}^n p_k f'_+(x_k) \right\rangle \geq \sum_{k=1}^n p_k \langle x_k, f'_+(x_k) \rangle, \quad (2)$$

where $f'_+(x) = (f'_{1+}(x), \dots, f'_{m+}(x))$ and f'_{1+}, \dots, f'_{m+} are right partial derivatives of f , then

$$\frac{1}{P_n} \sum_{i=1}^n p_i f(x_i) \leq f(A). \quad (3)$$

Proof. If f is convex on I , then

$$f(A) \geq f(x_k) + \langle A - x_k, f'_+(x_k) \rangle,$$

i.e.,

$$f(A) \geq f(x_k) + \langle A, f'_+(x_k) \rangle - \langle x_k, f'_+(x_k) \rangle$$

for $k = 1, \dots, n$.

Multiply the k th inequality by p_k and add the inequalities thus obtained; we obtain

$$\begin{aligned} f(A) P_n &\geq \sum_{k=1}^n p_k f(x_k) + \left\langle A, \sum_{k=1}^n p_k f'_+(x_k) \right\rangle - \sum_{k=1}^n p_k \langle x_k, f'_+(x_k) \rangle \\ &\geq \sum_{k=1}^n p_k f(x_k) \end{aligned}$$

since (2) holds.

COROLLARY. Let f , $x_1 = (x_{11}, \dots, x_{1m}), \dots, x_n = (x_{n1}, \dots, x_{nm})$, p_1, \dots, p_n satisfy the conditions of the theorem. If f is also nondecreasing (nonin-

creasing) in each of its m variables and if $p_1 f'_{j+}(x_1) + \cdots + p_n f'_{j+}(x_n) \neq 0$ ($j = 1, \dots, m$), then (3) is valid if

$$A = (A_1, \dots, A_m) \\ = \left(\frac{\sum_{k=1}^n p_k x_{k1} f'_{1+}(x_k)}{\sum_{k=1}^n p_k f'_{1+}(x_k)}, \dots, \frac{\sum_{k=1}^n p_k x_{km} f'_{m+}(x_k)}{\sum_{k=1}^n p_k f'_{m+}(x_k)} \right).$$

Proof. Observe that $A \in I$ since A_j is a convex combination of x_{1j}, \dots, x_{mj} . Since

$$\begin{aligned} \left\langle A, \sum_{k=1}^n p_k f'_+(x_k) \right\rangle &= \sum_{k=1}^n p_k \langle A, f'_+(x_k) \rangle = \sum_{k=1}^n p_k \sum_{j=1}^m A_j f'_{j+}(x_k) \\ &= \sum_{j=1}^m A_j \sum_{k=1}^n p_k f'_{j+}(x_k) = \sum_{j=1}^m \sum_{k=1}^n p_k x_{kj} f'_{j+}(x_k) \\ &= \sum_{k=1}^n p_k \sum_{j=1}^m x_{kj} f'_{j+}(x_k) = \sum_{k=1}^n p_k \langle x_k, f'_+(x_k) \rangle, \end{aligned}$$

the theorem implies the corollary.

Remark. One can prove the integral analogs of the above results (i.e., generalizations of inequality (4) of [1]).

REFERENCES

1. M. L. SLATER, A companion inequality to Jensen's inequality, *J. Approx. Theory* **32** (1981), 160–166.
2. J. E. PEČARIĆ, A companion to Jensen–Steffensen's inequality, *J. Approx. Theory* **44**, 1985.
3. P. LAH AND M. RIBARIĆ, Converse of Jensen's inequality for convex functions, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* **412–460** (1973), 201–205.
4. D. S. MITRINOVIĆ AND P. M. VASIĆ, The centroid method in inequalities, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* **498–541** (1975), 3–16.
5. P. M. VASIĆ AND J. E. PEČARIĆ, On the Jensen inequality, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* **634–677** (1979), 50–54.
6. P. R. BEESACK, On inequalities complementary to Jensen's, *Canad. J. Math.* **35** (1983), 324–338.