

An Analogue of a Problem of
P. Erdős and E. Feldheim on
 L_p Convergence of Interpolatory Processes

A. K. VARMA

Department of Mathematics, University of Florida, Gainesville, Florida 32611, U.S.A.

AND

J. PRASAD

*Department of Mathematics and Computer Science,
California State University, Los Angeles, California 90032, U.S.A.*

Communicated by Paul Erdős

Received March 12, 1987

DEDICATED TO PROFESSOR R. BOJANIC

For the Lagrange interpolation polynomial $L_n(f, x)$ of degree $\leq n-1$, defined by

$$L_n(f, x_{kn}) = f(x_{kn}), \quad k = 1, 2, \dots, n, \quad (1.1)$$

of a given function f (based on n distinct points $-1 \leq x_{nn} < x_{n-1,n} < \dots < x_{1n} \leq 1$), P. Erdős and E. Feldheim [6] proved the following

THEOREM A. *If $f \in C[-1, 1]$ and x_{kn} , $k = 1, 2, \dots, n$, are the zeros of $T_n(x)$, the n th Chebyshev polynomial, then for any fixed $p > 0$,*

$$\lim_{n \rightarrow \infty} \int_{-1}^1 |L_n[f, x] - f(x)|^p (1-x^2)^{-1/2} dx = 0. \quad (1.2)$$

For a more detailed study of this kind of work we refer the reader to P. Erdős and P. Turán [7], R. Askey [1], P. G. Nevai [11], P. Vertesi [20], and A. K. Varma and P. Vertesi [19]. The corresponding study of mean convergence of the Hermite–Fejér interpolation process was recently initiated by Nevai and Vertesi [12]. In Ref. [12] weighted L_p convergence of Hermite–Fejér interpolation based on the zeros of a generalized Jacobi polynomial was investigated. The main result of Ref. [12] gives necessary and sufficient conditions for such convergence for all continuous functions.

They mentioned that the main reason for the lack of the general theory appears to be the complicated structure of explicit representation for the Hermite-Fejér interpolating polynomial.

The object of this paper is to consider the problem of degree of approximation (in the L_p norm for any fixed $p > 0$) of a given continuous function by various interpolatory (Hermite) processes based on the Tchebycheff nodes. Now we turn to describe these results.

Let

$$x_k \equiv x_{kn} = \cos \frac{(2k-1)\pi}{2n}, \quad k = 1, 2, \dots, n, \quad (1.3)$$

be the zeros of $T_n(x) = \cos n\theta$, $\cos \theta = x$, the n th degree Tchebycheff polynomial of the first kind. In this case the well-known Hermite-Fejér interpolation polynomial is given by [8], [10]

$$H_n(f, x) = \sum_{k=1}^n f(x_k) h_k(x), \quad (1.4)$$

where

$$h_k(x) = (1 - xx_k) \left(\frac{T_n(x)}{n(x - x_k)} \right)^2 \geq 0, \quad \sum_{k=1}^n h_k(x) \equiv 1. \quad (1.5)$$

Next, we define a polynomial $Q_n(f, x)$ of degree $\leq 2n + 1$ by

$$\begin{aligned} Q_n(f, x) = & H_n(f, x) + (f(1) - H_n(f, 1)) \left(\frac{1+x}{2} \frac{T_n(x)}{T_n(1)} \right)^2 \\ & + (f(-1) - H_n(f, -1)) \left(\frac{1-x}{2} \frac{T_n(x)}{T_n(-1)} \right)^2. \end{aligned} \quad (1.6)$$

It is known [3] that

$$\begin{aligned} Q_n(f, x_k) &= f(x_k), & Q'_n(f, x_k) &= 0, & k &= 1, 2, \dots, n, \\ Q_n(f, \pm 1) &= f(\pm 1). \end{aligned} \quad (1.7)$$

Concerning $H_n(f, x)$ and $Q_n(f, x)$ we shall prove the following

THEOREM 1. *Let $f \in C[-1, 1]$ and let $H_n(f, x)$ be as defined by (1.4). Then for any fixed $p > 0$ we have*

$$\left(\int_{-1}^1 |H_n(f, x) - f(x)|^p (1-x^2)^{-1/2} dx \right)^{1/p} \leq c_1 w_f(1/n), \quad (1.8)$$

and

$$\left(\int_{-1}^1 |Q_n(f, x) - f(x)|^p (1-x^2)^{-1/2} dx \right)^{1/p} \leq c_2 w_f(1/n), \tag{1.9}$$

where c_1, c_2 are positive absolute constants and $w_f(\delta)$ is the modulus of continuity of f .

Remark. It is well known that for $f \in \text{Lip } 1$ the error function $\|f(x) - H_n(f, x)\|_{L^x}$ is of order $O(\log n/n)$ and this is best possible (see [2], [14], and [16]). On the other hand we can conclude from (1.8) that the corresponding estimate in L_p norm is $O(1/n)$.

If $f \in C^1[-1, 1]$ then it is known that a Hermite interpolation polynomial $H_n^*(f, x)$ of degree $\leq 2n - 1$ which satisfies the conditions

$$H_n^*(f, x_k) = f(x_k), \quad \frac{d}{dx} [H_n^*(f, x)]_{x_k} = f'(x_k), \quad k = 1, 2, \dots, n. \tag{1.10}$$

is given by

$$H_n^*(f, x) = \sum_{k=1}^n f(x_k) h_k(x) + \sum_{k=1}^n f'(x_k) \sigma_k(x), \tag{1.11}$$

where $h_k(x)$ is as in (1.5), the x_k 's are as in (1.3), and $\sigma_k(x)$ is given by

$$\sigma_k(x) = (x - x_k) l_k^2(x), \quad l_k(x) = \frac{T_n(x)}{(x - x_k) T_n'(x_k)}. \tag{1.12}$$

For the polynomial $H_n^*(f, x)$ we shall prove the following:

THEOREM 2. *Let $f(x)$ be defined and have a continuous derivative $f'(x)$ on $[-1, 1]$. Then for the Hermite interpolation polynomial $H_n^*(f, x)$ corresponding to the Tchebycheff abscissas of the first kind*

$$\left(\int_{-1}^1 |f(x) - H_n^*(f, x)|^2 (1-x^2)^{-1/2} dx \right)^{1/2} \leq c_3 n^{-1} E_{2n-2}(f'), \tag{1.13}$$

where $E_{2n-2}(f')$ is the best approximation to $f'(x)$ by polynomials of degree at most $2n - 2$ and c_3 is a positive absolute constant.

If we change the nodes of interpolation to the zeros of Tchebycheff polynomials of the second kind,

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta}, \quad x = \cos \theta, \tag{1.14}$$

then it is known that for many interpolation processes (this includes Lagrange as well as Hermite-Fejér interpolation) convergence behaviour is very poor with respect to the nodes of (1.14) especially near 1 or -1 . The situation is not improved for the Lagrange interpolation on these nodes even in problems of mean convergence. E. Feldheim [9] proved that for the same abscissas it is not true that ($r \geq 1$)

$$\lim_{n \rightarrow \infty} \int_{-1}^1 |f(x) - L_n(f, x)|^{2r} dx = 0. \quad (1.15)$$

In fact the superior limit of the integrals in question may be $+\infty$ if $f(x)$ is a properly chosen continuous function. But the situation changes if we consider $Q_n^*(f, x)$, the so-called quasi-Hermite-Fejér interpolation polynomial of degree $\leq 2n+1$ based on the extended Tchebycheff nodes of the second kind. It is given by

$$Q_n^*(f, x) = \left(\frac{1+x}{2} f(1) + \frac{1-x}{2} f(-1) \right) \frac{U_n^2(x)}{(n+1)^2} + \sum_{k=1}^n f(t_k) (1-x^2)(1-xt_k) \left(\frac{U_n(x)}{(n+1)(x-t_k)} \right)^2, \quad (1.16)$$

where the t_k 's are the zeros of $U_n(x)$.

In this case, P. Szasz [15] proved that $\lim_{n \rightarrow \infty} Q_n^*(f, x) = f(x)$ uniformly on $[-1, 1]$ provided $f \in C[-1, 1]$. Later Saxena and Mathur [21] proved that if $f \in C[-1, 1]$ we have

$$|Q_n^*(f, x) - f(x)| \leq \frac{c}{n} \sum_{k=1}^n w_f \left(\frac{\sqrt{1-x^2}}{k} + \frac{1}{k^2} \right). \quad (1.17)$$

Next let us denote by $R_n(f, x)$ the Hermite-Fejér interpolation polynomial of degree $\leq 2n+3$ satisfying the conditions

$$\begin{aligned} R_n(f, t_k) &= f(t_k), & R'_n(f, t_k) &= 0, & k &= 1, 2, \dots, n \\ R_n(f, \pm 1) &= f(\pm 1), & R'_n(f, \pm 1) &= 0, \end{aligned} \quad (1.18)$$

where the t_k 's are the zeros $U_n(x)$ given by (1.14). Concerning $R_n(f, x)$, the following pointwise estimate was obtained by Bojanic, Prasad, and Saxena [3]. It is given by ($-1 \leq x \leq 1$)

$$|R_n(f, x) - f(x)| \leq \frac{c_4}{n} \sum_{k=1}^n w \left(\frac{\sqrt{1-x^2}}{k} + \frac{1}{k^2} \right) + \frac{c_5}{n^2}, \quad (1.19)$$

where c_4, c_5 are positive absolute constants.

Concerning $Q_n^*(f, x)$ and $R_n(f, x)$ we shall prove the following:

THEOREM 3. *Let $f \in C[-1, 1]$ and let $Q_n^*(f, x)$ be the quasi-Hermite-Fejér interpolation polynomial of degree $\leq 2n + 1$ as stated in (1.16). Then for any fixed $p > 0$, we have*

$$\left(\int_{-1}^1 |Q_n^*(f, x) - f(x)|^p (1 - x^2)^{-1.2} dx \right)^{1/p} \leq c_6 w_f(1/n). \quad (1.20)$$

Also, we have

$$\left(\int_{-1}^1 |R_n(f, x) - f(x)|^p (1 - x^2)^{-1.2} dx \right)^{1/p} \leq c_7 w_f(1/n), \quad (1.21)$$

where $R_n(f, x)$ is defined by (1.18).

2. PRELIMINARIES

Here we state some known results which we shall need later on. If $l_k(x)$ is given by (1.12) then for $-1 \leq x \leq 1$ it is known that

$$l_k^2(x) \leq 2, \quad \sum_{k=1}^n l_k^2(x) \leq 2. \quad (2.1)$$

Also, from (1.4) it follows that for $-1 \leq x \leq 1$

$$\sum_{k=1}^n h_k(x) = 1, \quad h_k(x) \geq 0, \quad k = 1, 2, \dots, n. \quad (2.2)$$

Next, if

$$\lambda_k(x) = \frac{U_n(x)}{(x - t_k) U_n'(t_k)} = \frac{(-1)^{k+1} (1 - t_k^2) U_n(x)}{(n + 1)(x - t_k)}, \quad k = 1, 2, \dots, n, \quad (2.3)$$

then due to Erdős [5], Varma and Vertesi [19], and Varma [18] we have for $-1 \leq x \leq 1$ respectively

$$|\lambda_k(x)| \leq c_8, \quad \frac{(1 - x^2)^{1.2}}{(1 - t_k^2)^{1/2}} |\lambda_k(x)| \leq \sqrt{2} \quad (2.4)$$

$$\sum_{k=1}^n \frac{(1 - x^2)^2}{(1 - t_k^2)^2} \lambda_k^2(x) \leq 2, \quad \sum_{k=1}^n \frac{1 - x^2}{1 - t_k^2} \lambda_k^2(x) \leq 2, \quad (2.5)$$

where the t_k 's are given by

$$t_k = \cos \frac{k\pi}{n+1}, \quad k = 1, 2, \dots, n. \quad (2.6)$$

Further, from a known theorem of S. A. Teljakovskii [17] it follows that there exists a polynomial $P_n(x)$ of degree $\leq n$ such that for all x , $-1 \leq x \leq 1$,

$$|f(x) - P_n(x)| \leq c_{11} w_f \left(\frac{\sqrt{1-x^2}}{n} \right) \quad (2.7)$$

and

$$(1-x^2)^{1/2} |P'_n(x)| \leq c_{12} n w_f(1/n). \quad (2.8)$$

Also for $-1 \leq x \leq 1$, we know that

$$|T_n(x)| \leq 1. \quad (2.9)$$

3. SOME LEMMAS

In this section we state and prove several lemmas which will be needed later on.

LEMMA 3.1. *If $l_k(x)$ is as in (1.12) then for $k = 1, 2, \dots, n$,*

$$\begin{aligned} \int_{-1}^1 l_k(x) l_j(x) (1-x^2)^{-1/2} dx &= \frac{\pi}{n}, & k=j \\ &= 0, & k \neq j. \end{aligned} \quad (3.1)$$

Proof. For the proof we refer the reader to Erdős and Turán [7].

LEMMA 3.2. *If $\sigma_k(x)$ is given by (1.12) then*

$$\begin{aligned} \int_{-1}^1 \sigma_k(x) \sigma_j(x) (1-x^2)^{-1/2} dx &= 0, & k \neq j \\ &= \frac{\pi}{2n^3} (1-x_k^2), & k=j. \end{aligned} \quad (3.2)$$

Proof. From (1.12), (3.1), and using the orthogonal property of Tchebycheff polynomials we have

$$\begin{aligned} & \int_{-1}^1 \sigma_k(x) \sigma_j(x) (1-x^2)^{-1/2} dx \\ &= \int_{-1}^1 (x-x_k) l_k^2(x) (x-x_j) l_j^2(x) (1-x^2)^{-1/2} dx \\ &= \frac{1}{T'_n(x_k) T'_n(x_j)} \int_{-1}^1 \frac{1+T_{2n}(x)}{2} l_k(x) l_j(x) (1-x^2)^{-1/2} dx \\ &= \frac{1}{2T'_n(x_k) T'_n(x_j)} \int_{-1}^1 l_k(x) l_j(x) (1-x^2)^{-1/2} dx \\ &= 0, \quad k \neq j. \end{aligned}$$

Similarly, if $k = j$ then due to (3.1) and

$$|T'_n(x_k)| = \frac{n}{(1-x_k^2)^{1/2}}, \quad k = 1, 2, \dots, n, \tag{3.3}$$

it follows that

$$\int_{-1}^1 \sigma_k^2(x) \frac{1}{\sqrt{1-x^2}} dx = \frac{\pi}{2n(T'_n(x_k))^2} = \frac{\pi}{2n^3} (1-x_k^2). \tag{3.4}$$

This proves Lemma 3.2.

LEMMA 3.3. *If $\sigma_k(x)$ is given by (1.12) then*

$$\int_{-1}^1 \left[\sum_{k=1}^n P'_n(x_k) \sigma_k(x) \right]^2 (1-x^2)^{-1/2} dx \leq c_{13} [w_f(1/n)]^2, \tag{3.5}$$

where $P_n(x)$ is the polynomial for which (2.7) and (2.8) are valid.

Proof. We have

$$\begin{aligned} & \int_{-1}^1 \left[\sum_{k=1}^n P'_n(x_k) \sigma_k(x) \right]^2 (1-x^2)^{-1/2} dx \\ &= \sum_{j=1}^n \sum_{k=1}^n P'_n(x_j) P'_n(x_k) \int_{-1}^1 \sigma_k(x) \sigma_j(x) (1-x^2)^{-1/2} dx. \end{aligned}$$

So from (3.2) and (2.8) we have

$$\begin{aligned} & \int_{-1}^1 \left[\sum_{k=1}^n P'_n(x_k) \sigma_k(x) \right]^2 (1-x^2)^{-1/2} dx \\ &= \sum_{k=1}^n (P'_n(x_k))^2 \frac{\pi}{2n^3} (1-x_k^2) \leq \frac{\pi}{2n^3} c_{12}^2 n^2 \sum_{k=1}^n (w_f(1/n))^2 \\ &= c_{13} (w_f(1/n))^2, \end{aligned}$$

which yields (3.5). This proves Lemma 3.3.

In the work of Erdős and Feldheim [6] the following result played an important role. Let $l_{v_n}(x) \equiv l_v(x)$, $v=1, 2, \dots, n$, be the fundamental polynomial of Lagrange interpolation based on the zeros of $T_n(x)$. Then

$$\int_{-1}^1 l_{v_1}(x) l_{v_2}(x) \cdots l_{v_{2k}}(x) (1-x^2)^{-1/2} dx = 0,$$

where v_1, v_2, \dots, v_{2k} are distinct integers between 1 and n . For the L_p convergence of a quasi-Hermite-Fejér interpolation process based on the nodes $(1-x^2) U_n(x)$ the corresponding result is given by the following lemma:

LEMMA 3.4. *Let v_1, v_2, \dots, v_{2k} be distinct integers between 1 and n . Then we have*

$$\int_{-1}^1 \chi_{v_1}(x) \chi_{v_2}(x) \cdots \chi_{v_{2k}}(x) (1-x^2)^{-1/2} dx = 0, \tag{3.6}$$

where

$$\chi_v(x) = \frac{\sqrt{1-x^2}}{\sqrt{1-t_v^2}} \lambda_v(x) = \frac{\sqrt{1-x^2}}{\sqrt{1-t_v^2}} \frac{U_n(x)}{(t-t_v) U'_n(t_v)}. \tag{3.7}$$

Proof. From the earlier result [19, page 72] it follows that

$$(1-x^2)^k (U_n(x))^{2k-1} = \sum_{i=n}^{(2k-1)(n+1)+1} \mu_i \cos i\theta. \tag{3.8}$$

Since $t_{v_1}, \dots, t_{v_{2k}}$ are distinct it follows that $U_n(x)/(x-t_{v_1}) \cdots (x-t_{v_{2k}})$ is indeed a polynomial of degree $\leq n-2k$. Next we also note that

$$\chi_{v_1}(x) \cdots \chi_{v_{2k}}(x) = \alpha(v_1, v_2, \dots, v_{2k}) \frac{U_n(x)(1-x^2)^k (U_n(x))^{2k-1}}{(x-t_{v_1}) \cdots (x-t_{v_{2k}})}. \tag{3.9}$$

Therefore on using the orthogonality of a Tchebycheff polynomial of the first kind it follows that

$$\int_{-1}^1 \chi_{v_1}(x) \cdots \chi_{v_{2k}}(x)(1-x^2)^{-1/2} dx$$

$$\alpha(v_1, v_2 \cdots v_{2k}) \int_{-1}^1 q_{n-2k}(x) \sum_{i=n}^{(2k-1)(n+1)+1} \mu_i T_i(x)(1-x^2)^{-1/2} dx = 0.$$

Let us introduce the linear operator

$$L_n^*[f, x] = \sum_{k=1}^n f(t_k) \chi_k(x), \tag{3.10}$$

where $\chi_k(x)$ is defined by (3.7) and the t_k 's are the zeros of $U_n(x)$ given by (2.6). On using (3.6) and some simple computation we obtain

$$\int_{-1}^1 \chi_j(x) \chi_k(x)(1-x^2)^{-1/2} dx = 0, \quad j \neq k$$

$$= \frac{\pi}{n+1}, \quad j = k. \tag{3.11}$$

On using (3.11) we obtain

$$\int_{-1}^1 (L_n^*[f, x])^2(1-x^2)^{-1/2} dx = \frac{\pi}{n+1} \sum_{k=1}^n f^2(t_k) \leq \pi \max_{-1 \leq x \leq 1} |f^2(x)|. \tag{3.12}$$

Also, from (3.12) and the Cauchy-Schwarz inequality for integrals we also have

$$\int_{-1}^1 |L_n^*[f, x]| (1-x^2)^{-1/2} dx \leq \pi \max_{-1 \leq x \leq 1} |f(x)|. \tag{3.13}$$

4. PROOF OF THE THEOREMS

For the proof of Theorem 1 and Theorem 3 we follow the method of Erdős and Feldheim [6]. It is enough to prove the theorems for even values of p only. In the case of the proof of Theorem 3 we limit for the case $p=4$. For arbitrary fixed even p the proof is similar.

Proof of Theorem 1. Let $P_n(x)$ be the polynomial of degree $\leq n$ for which (2.7) and (2.8) are valid. Due to (1.4) we have

$$H_n(f, x) - f(x) = H_n(f - P_n, x) + H_n(P_n, x) - P_n(x) + P_n(x) - f(x). \quad (4.1)$$

On using (2.2) and (2.7) it follows that

$$\begin{aligned} |H_n(f - P_n, x)| &= \left| \sum_{k=1}^n (f(x_k) - P_n(x_k)) h_k(x) \right| \\ &\leq w_f(1/n) \sum_{k=1}^n h_k(x) \leq w_f(1/n). \end{aligned} \quad (4.2)$$

Thus to complete the proof of (1.8) we must prove that

$$\left(\int_{-1}^1 |H_n(P_n, x) - P_n(x)|^{2p} (1-x^2)^{-1/2} dx \right)^{1/2p} \leq c_{14} w_f(1/n). \quad (4.3)$$

From the uniqueness of Hermite interpolation we have

$$\begin{aligned} P_n(x) - H_n(P_n, x) &= \sum_{k=1}^n P'_n(x_k) \sigma_k(x) = \sum_{k=1}^n P'_n(x_k) (x - x_k) l_k^2(x) \\ &= \frac{1}{n} \sum_{k=1}^n P'_n(x_k) (-1)^{k-1} (1-x_k^2)^{1/2} l_k(x) T_n(x) \\ &= T_n(x) \sum_{k=1}^n g_0(x_k) l_k(x) \equiv T_n(x) L_n[g_0, x], \end{aligned}$$

where

$$g_0(x) = \frac{(1-x^2) U_{n-1}(x)}{n} P'_n(x).$$

Now, on using Theorem A of Erdős and Feldheim [6] and $|g_0(x)| \leq c_{15} w_f(1/n)$ we have

$$\begin{aligned} &\left(\int_{-1}^1 |P_n(x) - H_n(P_n, x)|^{2p} (1-x^2)^{-1/2} dx \right)^{1/2p} \\ &\leq c_{16} \left(\int_{-1}^1 |L_n[g_0, x]|^{2p} (1-x^2)^{-1/2} dx \right)^{1/2p} \\ &\leq c_{17} \max_{-1 \leq x \leq 1} |g_0(x)| \\ &\leq c_{18} w_f(1/n). \end{aligned}$$

This proves (4.3) and at the same time inequality (1.8). Proof of the inequality of (1.9) is a direct consequence of (1.8), (1.6), and the known estimate [3] (see formula (1.4)) $|H_n(f, \pm 1) - f(\pm 1)| \leq c_{19} w_f(1/n)$.

This proves Theorem 1. Next we turn to prove Theorem 2.

Proof of Theorem 2. One can easily see that for $-1 \leq x \leq 1$

$$H_n^*(f, x) - f(x) = H_n^*(f, x) - H_n^*(S_{2n-1}, x) + S_{2n-1}(x) - f(x), \tag{4.4}$$

where $S_{2n-1}(x)$ is the polynomial of best approximation of $f(x)$ and $H_n^*(f, x)$ is given by (1.11). From (4.4) it follows that

$$\begin{aligned} & \int_{-1}^1 [H_n^*(f, x) - f(x)]^2 (1-x^2)^{-1/2} dx \\ & \leq 2 \int_{-1}^1 [H_n^*(f - S_{2n-1}, x)]^2 (1-x^2)^{-1/2} dx \\ & \quad + 2 \int_{-1}^1 [S_{2n-1}(x) - f(x)]^2 (1-x^2)^{-1/2} dx \equiv \Delta_1 + \Delta_2. \end{aligned} \tag{4.5}$$

From the definition of $S_{2n-1}(x)$ we have for $-1 \leq x \leq 1$,

$$|S_{2n-1}(x) - f(x)| \leq E_{2n-1}(f), \tag{4.6}$$

where $E_{2n-1}(f)$ is the best approximation of $f(x)$. Consequently due to (4.6) we get

$$\Delta_2 \leq 2(E_{2n-1}(f))^2 \int_{-1}^1 (1-x^2)^{-1/2} dx = 2\pi E_{2n-1}^2(f). \tag{4.7}$$

Next, we turn to estimate Δ_1 . We have

$$\begin{aligned} \Delta_1 & \leq 4 \int_{-1}^1 \left[\sum_{k=1}^n |f(x_k) - S_{2n-1}(x_k)| h_k(x) \right]^2 (1-x^2)^{-1/2} dx \\ & \quad + 4 \int_{-1}^1 \left[\sum_{k=1}^n (f'(x_k) - S'_{2n-1}(x_k)) \sigma_k(x) \right]^2 (1-x^2)^{-1/2} dx \\ & = \Delta_3 + \Delta_4. \end{aligned} \tag{4.8}$$

Now, from (4.6) and (2.2) it follows that

$$\begin{aligned} \Delta_3 & \leq 4E_{2n-1}^2(f) \int_{-1}^1 \left[\sum_{k=1}^n h_k(x) \right]^2 (1-x^2)^{-1/2} dx \\ & = 4\pi E_{2n-1}^2(f). \end{aligned} \tag{4.9}$$

Further, on using Lemma 3.2 we have

$$\begin{aligned}
 A_4 &= 4 \int_{-1}^1 \sum_{k=1}^n (f'(x_k) - S'_{2n-1}(x_k))^2 \sigma_k^2(x) (1-x^2)^{-1/2} dx \\
 &= \frac{2\pi}{n^3} \sum_{k=1}^n (f'(x_k) - S'_{2n-1}(x_k))^2 (1-x_k^2).
 \end{aligned}
 \tag{4.10}$$

Next, on using a theorem of J. Czipser and G. Freud [4] and Corollary 1.44 of T. J. Rivlin [13, p. 23], we get

$$(1-x_k^2)^{1/2} |f'(x_k) - S'_{2n-1}(x_k)| \leq 40E_{2n-2}(f').
 \tag{4.11}$$

Consequently from (4.10), (4.11) it follows that

$$A_4 \leq c_{20} n^{-2} E_{2n-2}^2(f').
 \tag{4.12}$$

Thus, from (4.8), (4.9), and (4.12) we obtain

$$\begin{aligned}
 A_1 &\leq 4\pi E_{2n-1}^2(f) + c_{21} n^{-2} E_{2n-2}^2(f') \\
 &\leq c_{22} n^{-2} E_{2n-2}^2(f').
 \end{aligned}
 \tag{4.13}$$

Due to Rivlin [13, p. 23], we have

$$E_{2n-1}(f) \leq \frac{6}{2n-1} E_{2n-2}(f').
 \tag{4.14}$$

On using (4.13), (4.7), and (4.14) we have (1.13). This completes the proof of Theorem 2 as well.

Proof of Theorem 3. First we need to show that for $f \in C[-1, 1]$,

$$\int_{-1}^1 (Q_n^*(f, x) - f(x))^4 (1-x^2)^{-1/2} dx \leq c_{23} (w_f(1/n))^4.
 \tag{4.15}$$

Since

$$Q_n^*(f, x) - f(x) = Q_n^*(f - P_n, x) + Q_n^*(P_n, x) - P_n(x) + P_n(x) - f(x),
 \tag{4.16}$$

where $P_n(x)$ is the polynomial which satisfies (2.7) and (2.8), in view of (2.7) and since $Q_n^*(f, x)$ is such that $|f| \leq c_{24}$ implies $\|Q_n^*(f, x)\| \leq c_{24}$ we have

$$\|Q_n^*(f - P_n, x)\| \leq \|f - P_n\| \leq c_{25} w_f(1/n).
 \tag{4.17}$$

Next, we consider

$$\mu = \int_{-1}^1 (P_n(x) - Q_n^*(P_n, x))^4 (1-x^2)^{-1/2} dx. \tag{4.18}$$

Further, one can easily see that

$$P_n(x) - Q_n^*(P_n, x) = (1-x^2)^{1/2} U_n(x) \sum_{k=1}^n g(t_k) \chi_k(x), \tag{4.19}$$

where

$$g(x) = -\frac{T_{n+1}(x) P_n'(x) (1-x^2)^{1/2}}{n+1} \tag{4.20}$$

and

$$\chi_k(x) = \frac{(1-x^2)^{1/2} \lambda_k(x)}{(1-t_k^2)^{1/2}}. \tag{4.21}$$

Also it is well known that for $-1 \leq x \leq 1$

$$(1-x^2)^{1/2} (U_n(x)) \leq 1. \tag{4.22}$$

Consequently from (4.19) and (4.22) it follows that

$$\mu \leq \int_{-1}^1 \left(\sum_{k=1}^n g(t_k) \chi_k(x) \right)^4 (1-x^2)^{-1/2} dx \equiv \mu_1. \tag{4.23}$$

We also note (see (2.8) and (2.4)) that for $-1 \leq x \leq 1$

$$|\chi_k(x)| \leq \sqrt{2}, \quad |g(x)| \leq c_{30} w_f(1/n). \tag{4.24}$$

Now, we may write

$$\mu \leq \mu_1 = \mu_{11} + \mu_{12} + \mu_{13} + \mu_{14} + \mu_{15}, \tag{4.25}$$

where on using (2.5) and (2.4)

$$\begin{aligned} \mu_{11} &= \int_{-1}^1 \sum_{k=1}^n g^4(t_k) \chi_k^4(x) (1-x^2)^{-1/2} dx \\ &\leq c_{31} (w_f(1/n))^4 \int_{-1}^1 \sum_{k=1}^n \chi_k^4(x) (1-x^2)^{-1/2} dx \leq c_{32} (w_f(1/n))^4. \end{aligned} \tag{4.26}$$

Next, we note that

$$\begin{aligned}
 \mu_{12} &\equiv \int_{-1}^1 \sum_{k \neq j} g^2(t_k) g^2(t_j) \chi_k^2(x) \chi_j^2(x) (1-x^2)^{-1/2} dx \\
 &\leq \int_{-1}^1 \sum_{k=1}^n \sum_{j=1}^n g^2(t_k) g^2(t_j) \chi_k^2(x) \chi_j^2(x) (1-x^2)^{-1/2} dx \\
 &\leq c_{33} (w_f(1/n))^4 \int_{-1}^1 \left(\sum_{k=1}^n \chi_k^2(x) \right) \left(\sum_{j=1}^n \chi_j^2(x) \right) (1-x^2)^{-1/2} dx \\
 &\leq 4c_{33} (w_f(1/n))^4 \int_{-1}^1 (1-x^2)^{-1/2} dx = 4c_{33} \pi (w_f(1/n))^4. \quad (4.27)
 \end{aligned}$$

Next, we observe that (Lemma 3.4)

$$\begin{aligned}
 \mu_{13} &\equiv \int_{-1}^1 \sum_{k \neq j \neq i \neq m} g(t_k) g(t_j) g(t_i) g(t_m) \\
 &\quad \times \chi_k(x) \chi_j(x) \chi_i(x) \chi_m(x) (1-x^2)^{-1/2} dx \\
 &= 0. \quad (4.28)
 \end{aligned}$$

Next, we observe that

$$\begin{aligned}
 \mu_{14} &\equiv \int_{-1}^1 \sum_{k \neq j} g^3(t_k) g(t_j) \chi_k^3(x) \chi_j(x) (1-x^2)^{-1/2} dx \\
 &= \int_{-1}^1 \left[\left(\sum_{k=1}^n g^3(t_k) \chi_k^3(x) \right) \left(\sum_{k=1}^n g(t_k) \chi_k(x) \right) \right. \\
 &\quad \left. - \sum_{k=1}^n g^4(t_k) \chi_k^4(x) \right] (1-x^2)^{-1/2} dx. \quad (4.29)
 \end{aligned}$$

Hence, on account of (2.4), (2.5), (4.24), (4.26), (3.10), and (3.13)

$$\begin{aligned}
 |\mu_{14}| &\leq c_{34} (w_f(1/n))^3 \int_{-1}^1 \left| \sum_{k=1}^n g(t_k) \chi_k(x) \right| \frac{dx}{(1-x^2)^{1/2}} \\
 &\quad + c_{35} (w_f(1/n))^4 \\
 &\leq c_{34} (w_f(1/n))^3 \pi \max_{-1 \leq x \leq 1} |g(x)| + c_{35} (w_f(1/n))^4 \\
 &\leq c_{34} (w_f(1/n))^3 \pi c_{30} w_f(1/n) + c_{35} (w_f(1/n))^4 \\
 &\leq c_{36} (w_f(1/n))^4. \quad (4.30)
 \end{aligned}$$

Finally, we consider

$$\begin{aligned} \mu_{15} &= \int_{-1}^1 \sum_{k \neq j \neq i} \sum g^2(t_k) \chi_k^2(x) g(t_j) \chi_j(x) g(t_i) \chi_i(x) (1-x^2)^{-1,2} dx \\ &= \int_{-1}^1 \sum_{k=1}^n g^2(t_k) \chi_k^2(x) \\ &\quad \times \left(\sum_{k=1}^n g(t_k) \chi_k(x) \right)^2 (1-x^2)^{-1,2} dx - \mu_{14} - \mu_{12}. \end{aligned} \quad (4.31)$$

Therefore, on using (4.27), (4.30), and Lemma 3.4 we have

$$\begin{aligned} |\mu_{15}| &\leq c_{37}(w_f(1/n))^2 \int_{-1}^1 \left(\sum_{k=1}^n g(t_k) \chi_k(x) \right)^2 (1-x^2)^{-1,2} dx \\ &\quad + c_{38}(w_f(1/n))^4 \\ &\leq c_{37}(w_f(1/n))^2 \int_{-1}^1 \sum_{k=1}^n g^2(t_k) \chi_k^2(x) (1-x^2)^{-1,2} dx \\ &\quad + c_{38}(w_f(1/n))^4 \\ &\leq c_{39}(w_f(1/n))^4 \int_{-1}^1 \sum_{k=1}^n \chi_k^2(x) (1-x^2)^{-1,2} dx + c_{38}(w_f(1/n))^4 \\ &\leq c_{40}(w_f(1/n))^4. \end{aligned} \quad (4.32)$$

On combining (4.32), (4.30), (4.27), (4.28), (4.26), and (4.25) we obtain

$$\mu \leq c_{41}(w_f(1/n))^4. \quad (4.33)$$

Now, on using (4.16), (4.17), (4.18), and (4.33) we obtain (1.20). Proof of (1.21) is a simple consequence of [3]

$$|R_n(f, \pm 1) - f(\pm 1)| \leq c_{42} w_f(1/n),$$

the representation given in the work of Bojanic, Prasad, and Saxena [3], and inequality (1.20).

REFERENCES

1. R. ASKEY, Mean convergence of orthogonal series and Lagrange interpolation, *Acta Math. Acad. Sci. Hungar.* **23** (1972), 79-85.
2. R. BOJANIC, A note on the precision of interpolation by Hermite-Fejér interpolation, in "Proceedings of the Conference on Constructive Theory of Functions" pp. 69-76, Akad. Kiadó, Budapest, 1972.

3. R. BOJANIC, J. PRASAD, AND R. B. SAXENA, An upper bound for the rate of convergence of the Hermite-Fejér process on the extended Chebyshev nodes of the Second kind, *J. Approx. Theory* **26** (1969), 195–203.
4. J. CZIPSZER AND G. FREUD, Sur l'approximation d'une fonction periodique et de ses dérivées successives par un polynome trigonometrique et par ses dérivées successives, *Acta Math.* **99** (1958), 33–51.
5. P. ERDÖS, On the maximum of the fundamental functions of the ultraspherical polynomials, *Ann. of Math. (2)* **45** (1944), 335–339.
6. P. ERDÖS AND E. FELDHEIM, Sur le mode de convergence pour l'interpolation de Lagrange, *C. R. Acad. Sci. Paris Sér. I Math.* **203** (1936), 913–915.
7. P. ERDÖS AND P. TURÁN, On interpolation, I, *Ann. of Math. (2)* **38** (1937), 142–155.
8. L. FEJÉR, Über interpolation, *Göttinger Nachr.* (1916), 66–91.
9. E. FELDHEIM, Sur le mode de convergence dans l'interpolation de Lagrange, *Soviet Math. Dokl.* **14** (1937), 327–331.
10. L. FEJÉR, Lagrangesche interpolation und zugehörigen konjugierten punkte, *Math. Ann.* **106** (1932), 1–55.
11. P. G. NEVAI, Mean convergence of Lagrange interpolation, I, *J. Approx. Theory* **18** (1976), 363–377.
12. P. NEVAI AND P. VERTESI, Mean convergence of Hermite-Fejér Interpolation, *J. Math. Anal. Appl.* **105** (1906), 26–58.
13. T. J. RIVLIN, Introduction to the approximation of functions, Ginn (Blaisdell), Boston, 1969.
14. O. SHISHA AND B. MOND, The rapidly of convergence of the Hermite-Fejér approximation to functions of one and several variables, *Proc. Amer. Math. Soc.* **16** (1965), 1269–1276.
15. P. SZASZ, On quasi-Hermite-Fejér interpolation, *Acta Math. Acad. Sci. Hungar.* **10** (1959), 413–439.
16. G. SZEGÖ, Orthogonal polynomials, in "Amer. Math. Soc. Coll. Publ. No. XXIII," New York, 1959.
17. S. A. TELJAKOVSKII, Two theorems on the approximation of functions by algebraic polynomials, *Math. Sbornik* **70**, No. 2 (1966), 252–265.
18. A. K. VARMA, On a problem of P. Turán on lacunary interpolation, *Canad. Math. Bull.* **10**, No. 4 (1967), 531–557.
19. A. K. VARMA AND P. VERTESI, Some Erdős-Feldheim type theorems on mean convergence of Lagrange interpolation, *J. Math. Anal. Appl.* **91** (1983), 68–79.
20. P. VERTESI, Note on mean convergence of Lagrange parabolas, *J. Approx. Theory* **28** (1980), 30–35.
21. K. K. MATHUR AND R. B. SAXENA, On the convergence of quasi-Hermite-Fejér interpolation, *Pac. J. Math.* **20** (1967), 245–259.