

Generalized Jacobi Weights, Christoffel Functions, and Zeros of Orthogonal Polynomials*

TAMÁS ERDÉLYI AND PAUL NEVAI

*Department of Mathematics, The Ohio State University,
P. O. Box 3341, Columbus, Ohio 43210-0341*

Communicated by V. Totik

Received January 7, 1991

We study generalized Jacobi weight functions in terms of their (generalized) degree. We obtain sharp lower and upper bounds for the corresponding Christoffel functions, and for the distance of the consecutive zeros of the corresponding orthogonal polynomials. The novelty of our results is that our constants depend only on the degree of the weight function but not on the weight itself. © 1992 Academic Press, Inc.

1. INTRODUCTION AND NOTATION

The sets of all real-valued algebraic polynomials and trigonometric polynomials of degree at most n will be denoted by Π_n and H_n , respectively. The function

$$w(z) = |\omega| \prod_{j=1}^k |z - z_j|^{r_j}, \quad \omega \neq 0, z_j \in \mathbb{C}, z \in \mathbb{C}, r_j > 0, \quad (1.1)$$

will be called a *generalized Jacobi weight function* of degree

$$N = \sum_{j=1}^k r_j \quad (1.2)$$

with positive exponents. The set of all generalized Jacobi weight functions of degree at most N with positive exponents will be denoted by $|\text{GCAP}|_N$. This notation comes from the fact that the function w defined by (1.1) is

* This material is based upon work supported by the National Science Foundation under Grant DMS-8814488, and by Nato under Grant CRG.870806.

the modulus of a generalized complex algebraic polynomial of degree N defined by (1.2). In the trigonometric case, the function

$$w(z) = |\omega| \prod_{j=1}^k \left| \sin \frac{z - z_j}{2} \right|^{r_j}, \quad \omega \neq 0, z_j \in \mathbb{C}, z \in \mathbb{C}, r_j > 0, \quad (1.3)$$

will be called a *generalized trigonometric Jacobi weight function* of degree

$$N = \frac{1}{2} \sum_{j=1}^k r_j \quad (1.4)$$

with positive exponents. The set of all generalized trigonometric Jacobi weight functions of degree at most N with positive exponents will be denoted by $|\text{GCTP}|_N$. The function w defined by (1.3) is the modulus of a generalized complex trigonometric polynomial of degree N defined by (1.4) which explains our notation.

Let α be a nonnegative, finite Borel measure in $[-1, 1]$. Given $0 < p < \infty$, we define the Christoffel function $\lambda_n(\alpha)$ for $n = 1, 2, \dots$, by

$$\lambda_n(\alpha, p, z) = \min_{Q \in \Pi_{n-1}} \int_{-1}^1 \frac{|Q(t)|^p}{|Q(z)|^p} d\alpha(t), \quad z \in \mathbb{C}, \quad (1.5)$$

and the generalized Christoffel function $\lambda_n^*(\alpha)$ for real $n \geq 1$ by

$$\lambda_n^*(\alpha, p, z) = \inf_{f \in |\text{GCAP}|_{n-1}} \int_{-1}^1 \frac{f^p(t)}{f^p(z)} d\alpha(t), \quad z \in \mathbb{C}. \quad (1.6)$$

For $M > 0$, we also introduce the functions

$$A_M(x) = \max\{M^{-1}\sqrt{1-x^2}, M^{-2}\}, \quad -1 \leq x \leq 1, \quad (1.7)$$

and

$$w_M(x) = \int_{|t-x| \leq A_M(x)} w(t) dt, \quad -1 \leq x \leq 1, \quad (1.8)$$

where we assume that $w(t)$ is defined if $|t-x| \leq A_M(x)$.

For $g(\geq 0) \in L^1(0, 2\pi)$, the Szegő function $D(g)$ is defined by

$$D(g, z) = \exp\left(\frac{1}{4\pi} \int_0^{2\pi} \log g(\theta) \frac{1 + ze^{-i\theta}}{1 - ze^{-i\theta}} d\theta\right), \quad |z| < 1. \quad (1.9)$$

The boundary value $D(g, e^{it})$ can be defined as the non-tangential limit of $D(g, z)$; this exists for almost every real t . It is important to note that $D(g) \in H^2(|z| \leq 1)$, and $|D(g, e^{it})|^2 = |g(t)|$ holds for almost every real t . Other properties of the Szegő function may be found in, e.g., [6, Chap. V].

In Sections 2 and 3, we give sharp lower and upper bounds for Christoffel functions and for generalized Christoffel functions in $[-1, 1]$ corresponding to generalized Jacobi weight functions. These results are proved in Sections 6 and 7. In Section 4, we give sharp lower and upper bounds for the distance of consecutive zeros of orthogonal polynomials corresponding to generalized Jacobi weight functions with positive exponents. These results are proved in Section 8. The properties of generalized Jacobi weight functions with positive exponents were the subject of study in a series of recent papers [1–5]. A number of these results which are used in this paper are summarized in Section 5.

The novelty of the results in the present paper is that the constants depend only on the degree of the weight function but not on the weight itself. For *fixed* generalized Jacobi weights, all our principal results have already been known (see [9, Theorem 6.3.28, p. 120, and Theorem 9.22, p. 166] for the results corresponding to Theorems 2.1, 2.2, 3.1, and 4.1, whereas the “fixed” version of Theorem 3.2 was essentially proved in [7, formula (13), p. 150]).

2. UPPER BOUNDS FOR $\lambda_n(\alpha, p, x)$ FOR $-1 \leq x \leq 1$

THEOREM 2.1. *Given $0 < p < \infty$, $0 \leq \Gamma < \infty$, and $n = 1, 2, \dots$, let $M = 1 + p(n - 1)/(\Gamma + p + 1)$. Then there exists an absolute constant $c_1 > 0$ such that*

$$\lambda_n(\alpha, p, x) \leq c_1^{\Gamma + p + 1} w_M(x), \quad -1 \leq x \leq 1,$$

for every measure α such that $d\alpha = w dt$ with a $w \in |\text{GCAP}|_\Gamma$.

THEOREM 2.2. *Let $0 < p < \infty$, $0 \leq \Gamma < \infty$, and $n = 1, 2, \dots$. Let $w = w^{(T)}/w^{(B)}$, where $w^{(T)}$ and $w^{(B)}$ belong to $|\text{GCAP}|_\Gamma$, and let $d\alpha = w dt$. Let d denote the number of different zeros of $w^{(B)}$. Then there exists an absolute constant $c_2 > 0$ such that*

$$\lambda_n(\alpha, p, x) \leq c_2^{\Gamma + pd + p + 1} w_M(x), \quad -1 \leq x \leq 1,$$

with

$$M = \frac{p(n - 1 - d) - \Gamma}{2\Gamma + 4 + p}$$

whenever $M \geq 1$, and

$$\lambda_n^*(\alpha, p, x) \leq c_2^{\Gamma + 1} w_M(x), \quad -1 \leq x \leq 1,$$

with

$$M = \frac{p(n-1) - \Gamma}{2\Gamma + 4}$$

whenever $M > 0$.

3. LOWER BOUNDS FOR $\lambda_n^*(\alpha, p, x)$ FOR $-1 \leq x \leq 1$

Since $\lambda_n^*(\alpha, p) \leq \lambda_n(\alpha, p)$, we will give lower bounds for $\lambda_n^*(\alpha, p)$ instead of $\lambda_n(\alpha, p)$. Our first theorem deals with the case $d\alpha(t) = w(t) dt$, where $w \in |\text{GCAP}|_\Gamma$, and the second one gives a lower bound in a more general case, when the weight function w satisfies the Szegő condition of logarithmic integrability.

THEOREM 3.1. *Given $0 < p < \infty$, $0 \leq \Gamma < \infty$, and $1 \leq n < \infty$, let $M = 1 + p(n-1)/(\Gamma + p + 1)$. Then there exists an absolute constant $c_3 > 0$ such that*

$$\lambda_n^*(\alpha, p, x) \geq c_3^{\Gamma + p + 1} w_M(x), \quad -1 \leq x \leq 1,$$

for every measure α such that $d\alpha = w dt$ with $w \in |\text{GCAP}|_\Gamma$.

THEOREM 3.2. *Let w be a nonnegative, integrable weight function in $[-1, 1]$ such that $\log(w(\cos \cdot)) \in L^1(-\pi, \pi)$. Let $0 < p < \infty$, and let $d\alpha = w dt$. Then there exists an absolute constant $c_4 > 0$ such that*

$$\lambda_n^*(\alpha, p, x) \geq c_4 \Delta_{p(n-1)+1}(x) |D(w(\cos \cdot), re^{i\theta})|^2, \quad -1 \leq x \leq 1,$$

where $r = (p(n-1) + 1)/(p(n-1) + 3)$, $x = \cos \theta$, and $D(g)$ is the Szegő function defined by (1.9).

4. ZEROS OF ORTHOGONAL POLYNOMIALS CORRESPONDING TO GENERALIZED JACOBI WEIGHTS WITH POSITIVE EXPONENTS

We will use the standard notations. Let α be a nonnegative, finite Borel measure with $\text{supp}(\alpha) \subset [-1, 1]$, and let $\{p_n(\alpha)\}_{n=0}^\infty$ denote the corresponding orthonormal polynomials. In addition, $\{x_{j,n}(\alpha)\}_{j=1}^n$ denote the zeros of $p_n(\alpha)$ in decreasing order, $x_{0,n} = 1$, $x_{n+1,n} = -1$, and $x_{j,n} = \cos \theta_{j,n}$ for $j = 1, 2, \dots, n$. The main goal of this paper is to establish sharp lower and upper bounds for the distance of consecutive zeros of orthogonal polynomials associated with generalized Jacobi weight functions with positive

exponents. The novelty of these estimates lies in the fact that our constants depend only on the degree of the weight function.

THEOREM 4.1. *Let $0 \leq \Gamma < \infty$ and let $dx = w dt$, where $w \in |\text{GCAP}|_\Gamma$. Then there exist two absolute constants $c_5 > 0$ and $c_6 > 0$ such that the zeros of the corresponding orthogonal polynomials satisfy*

$$\theta_{j,n} - \theta_{j-1,n} \leq \frac{c_5^{\Gamma+1}}{n}, \quad j = 1, 2, \dots, n+1, \tag{4.1}$$

and

$$\theta_{j,n} - \theta_{j-1,n} \geq \frac{c_6^{\Gamma+1}}{n}, \quad j = 2, 3, \dots, n, \tag{4.2}$$

for $n = 1, 2, \dots$.

We believe that (4.2) actually holds for $j = 1$ and $j = n + 1$ as well, but, alas, our method does not seem to work in these cases. The latter would generalize [8, Theorem 3, p. 367; 9, Theorem 9.22, p. 166].

5. AUXILIARY RESULTS

The extremal properties of generalized Jacobi weight functions with positive exponents were studied in a number of recent papers. In this section we list those theorems which will be used to prove our new results in Sections 2–4. We will use a pointwise Remez-type inequality for which we introduce the class

$$|\text{GCAP}|_N(s) = \{f \in |\text{GCAP}|_N : m(\{x \in [-1, 1] : f(x) \leq 1\}) \geq 2 - s\},$$

$$0 < s < 2,$$

where $m(A)$ denotes the Lebesgue measure of a measurable set $A \subset \mathbb{R}$. The following theorem was proved in [1, Theorem 4].

THEOREM 5.1. *There exists an absolute constant $c_7 > 0$ such that, given $0 \leq N < \infty$ and $0 < s \leq 1$, the inequality*

$$f(x) \leq \exp \left(c_7 N \min \left\{ \frac{s}{\sqrt{1-x^2}}, \sqrt{s} \right\} \right), \quad -1 \leq x \leq 1,$$

holds for every $f \in |\text{GCAP}|_N(s)$.

We will need Nikol'skii-type inequalities for the classes $|\text{GCAP}|_N$ and $|\text{GCTP}|_N$ as well (see Section 1 for the notations). For $0 \leq N < \infty$ and $f \in |\text{GCAP}|_N$ or for $n = 0, 1, 2, \dots$, and $f \in \Pi_n$, let

$$\|f\|_\infty = \max_{-1 \leq t \leq 1} |f(t)| \quad \text{and} \quad \|f\|_p = \left(\int_{-1}^1 |f(t)|^p dt \right)^{1/p}, \quad 0 < p < \infty,$$

whereas for $0 \leq N < \infty$ and $f \in |\text{GCTP}|_N$ or for $n = 0, 1, 2, \dots$, and $f \in H_n$, let

$$\|f\|_\infty = \max_{-\pi \leq \theta \leq \pi} |f(\theta)| \quad \text{and} \quad \|f\|_p = \left(\int_{-\pi}^{\pi} |f(\theta)|^p d\theta \right)^{1/p}, \quad 0 < p < \infty.$$

The following two inequalities were proved in [5, Theorems 5 and 6].

THEOREM 5.2. *Given $0 \leq N < \infty$, there exists an absolute constant $c_8 > 0$ such that*

$$\|f\|_p \leq [c_8(1 + qN)]^{2/q - 2/p} \|f\|_q, \quad 0 < q < p \leq \infty,$$

holds for every $f \in |\text{GCAP}|_N$. For instance, $c_8 = e^2(2\pi)^{-1}$ is a suitable choice.

THEOREM 5.3. *Given $0 \leq N < \infty$, there exists an absolute constant $c_9 > 0$ such that*

$$\|f\|_p \leq [c_9(1 + qN)]^{1/q - 1/p} \|f\|_q, \quad 0 < q < p \leq \infty,$$

holds for every $f \in |\text{GCTP}|_N$. For instance, $c_9 = e(4\pi)^{-1}$ is a suitable choice.

If $f \in |\text{GCAP}|_N$, where $0 \leq N < \infty$, then $g(\theta) = |f(\cos \theta) \sin \theta| \in |\text{GCTP}|_{N+1}$, and, thus, applying Theorem 5.3 to the function g with $q = 1$ and $p = \infty$, we obtain

$$f(x) \leq 2c_9 \frac{N+2}{\sqrt{1-x^2}} \int_{-1}^1 f(t) dt, \quad -1 \leq x \leq 1,$$

for every $f \in |\text{GCAP}|_N$. This, together with Theorem 5.2, gives

THEOREM 5.4. *Given $0 \leq N < \infty$, there exists an absolute constant $c_{10} > 0$ such that*

$$f(x) \leq c_{10} (A_{N+2}(x))^{-1} \int_{-1}^1 f(t) dt, \quad -1 \leq x \leq 1,$$

holds for every $f \in |\text{GCAP}|_N$. For instance, $c_{10} = e^2(2\pi)^{-1}$ is a suitable choice.

Using a linear transformation, we easily deduce from Theorem 5.2 that

$$w_M(x) \leq 2\Delta_M(x) \max_{|t-x| \leq \Delta_m(x)} w(t) \leq 2c_8^2(\Gamma+1)^2 w_M(x), \quad -1 \leq x \leq 1, \tag{5.1}$$

holds for every $w \in |\text{GCAP}|_\Gamma$, where w_M is defined by (1.8).

To prove (4.2) in Theorem 4.1, we will need weighted Markov and Bernstein type inequalities. The following results are particular cases of [4, Theorems 1 and 3].

THEOREM 5.5. *Given $n = 0, 1, \dots$, and $0 \leq \Gamma < \infty$, there exists an absolute constant $c_{11} > 0$ such that*

$$\|Q'w\|_\infty \leq c_{11}(n + \Gamma)^2 \|Qw\|_\infty$$

holds for every algebraic polynomial $Q \in \Pi_n$ and weight $w \in |\text{GCAP}|_\Gamma$.

THEOREM 5.6. *Given $n = 0, 1, \dots$, and $0 \leq \Gamma < \infty$, there exists an absolute constant $c_{12} > 0$ such that*

$$\sqrt{1-x^2} |Q'(x) w(x)| \leq c_{12}(\Gamma+1)(n + \Gamma) \|Qw\|_\infty, \quad -1 < x < 1,$$

holds for every $Q \in \Pi_n$ and $w \in |\text{GCAP}|_\Gamma$.

With the notation of (1.7), and combining Theorems 5.5 and 5.6, we obtain the inequality

$$\Delta_n(x) |Q'(x) w(x)| \leq c_{13}(\Gamma+1)^2 \|Qw\|_\infty, \quad -1 \leq x \leq 1, \tag{5.2}$$

for every $Q \in \Pi_n$ and $w \in |\text{GCAP}|_\Gamma$, where $n = 1, 2, \dots, 0 \leq \Gamma < \infty$, and $c_{13} > 0$ is an absolute constant.

We think that the factor $(\Gamma+1)$ in Theorems 5.6 can perhaps be dropped.

6. PROOF OF THEOREMS 2.1 AND 2.2

To prove Theorem 2.1 we need the following

LEMMA 6.1. *Given $a > 0, b \in \mathbb{R}$, and $0 \leq \Gamma < \infty$, the inequality*

$$w(y) \leq 2^\Gamma a^{-\Gamma} |y - b|^\Gamma \max_{b-a \leq t \leq b+a} w(t), \quad y \in \mathbb{R} \setminus (b - a, b + a),$$

holds for every $w \in |\text{GCAP}|_\Gamma$.

Proof of Lemma 6.1. Without loss of generality we may assume that $a = 1$ and $b = 0$; the general case follows from this case by a linear substitution. Let T_m denote the Chebyshev polynomial of degree m , and let $-1 < x_{1,m} < x_{2,m} < \dots < x_{m,m} < 1$ denote the zeros of T_m , where $x_{j,m} = -x_{m-j,m}$ for $j = 1, 2, \dots, m$. Given $m = 1, 2, \dots$, a well-known inequality of Bernstein yields

$$\frac{|Q(y)|}{\|Q\|_\infty} \leq \frac{|T_m(y)|}{\|T_m\|_\infty} = 2^{m-1} \prod_{j=1}^m (y - x_{j,m}) \leq 2^m |y|^m, \quad y \in \mathbb{R} \setminus (-1, 1), \tag{6.1}$$

for every $Q \in \Pi_m$. Now let

$$w(z) = |\omega| \prod_{j=1}^k |z - z_j|^{r_j} \in |\text{GCAP}|_r. \tag{6.2}$$

If each exponent r_j is a positive rational number, say, $r_j = q_j/q$ for $j = 1, 2, \dots, k$, with some positive integers q_j and q , then applying (6.1) to the polynomial

$$Q(z) = \omega^{2q} \prod_{j=1}^k (z - z_j)^{q_j} (z - \bar{z}_j)^{q_j} \in \Pi_{2q\Gamma},$$

and taking the $2q$ th root of its modulus, we obtain the inequality in the lemma. The case of positive real exponents r_j in (6.2) can be reduced to that of rational exponents by approximation. ■

Proof of Theorem 2.1. Let v denote the Chebyshev weight, that is, $v(x) = (1/\pi)(1 - x^2)^{-1/2}$, and let

$$T_m(x) = \cos(m\theta) \quad \text{and} \quad U_m(x) = \frac{\sin((m+1)\theta)}{\sin \theta}, \quad -1 \leq x \leq 1, x = \cos \theta,$$

be the Chebyshev polynomials of first and second kinds, respectively. We define the corresponding reproducing kernel function $K_m(v)$ by

$$K_m(v, x, t) = 1 + 2 \sum_{j=1}^m T_j(x) T_j(t). \tag{6.3}$$

It is easy to see that

$$K_m(v, x, x) = \frac{2m+1 + U_{2m}(x)}{2},$$

and, thus, by a straightforward calculation,

$$c_{14}(m + 1) \leq K_m(v, x, x) < 2m + 1, \quad -1 \leq x \leq 1, \tag{6.4}$$

where $c_{14} > 0$ is an absolute constant. Therefore,

$$\left| \frac{K_m(v, x, t)}{K_m(v, x, x)} \right| \leq \frac{2m + 1}{c_{14}(m + 1)} \leq \frac{2}{c_{14}}, \quad -1 \leq x, t \leq 1. \tag{6.5}$$

In addition, we also have¹

$$|K_m(v, x, t)| \leq c_{15} \min \left\{ m, 1 + \frac{\sqrt{1-x^2} + \sqrt{2-t^2}}{|x-t|} \right\}, \tag{6.6}$$

$-1 \leq x, t \leq 1, m = 1, 2, \dots$

with an absolute constant $c_{15} > 0$. To show (6.6), we may assume without loss of generality that $0 \leq x \leq 1$, that is, $x = \cos \gamma$ with $0 \leq \gamma \leq \pi/2$. By the Christoffel–Darboux formula,

$$K_m(v, x, t) = \frac{T_m(x) T_{m-1}(t) - T_m(t) T_{m-1}(x)}{x - t}. \tag{6.7}$$

First, by (6.3),

$$|K_m(v, x, t)| \leq 2m + 1, \quad -1 \leq t \leq 1. \tag{6.8}$$

Second, if $t = \cos \theta$ with $0 \leq \theta \leq 2\pi/3$, then

$$\begin{aligned} & |T_m(x) T_{m-1}(t) - T_m(t) T_{m-1}(x)| \\ &= |T_{m-1}(t)(T_m(x) - T_{m-1}(x)) - T_{m-1}(x)(T_m(t) - T_{m-1}(t))| \\ &\leq |\cos(m\gamma) - \cos((m-1)\gamma)| + |\cos(m\theta) - \cos((m-1)\theta)| \\ &\leq 2|\sin((2m-1)\gamma/2) \sin(\gamma/2)| + 2|\sin((2m-1)\theta/2) \sin(\theta/2)| \\ &\leq \left| \frac{\sin \gamma}{\cos(\gamma/2)} \right| + \left| \frac{\sin \theta}{\cos(\theta/2)} \right| \leq 2(\sqrt{1-x^2} + \sqrt{1-t^2}). \end{aligned} \tag{6.9}$$

Thus, inequality (6.6) follows from (6.8) and (6.9).

Now, given $0 < p < \infty$, $0 \leq \Gamma < \infty$, and $n = 1, 2, \dots$, let²

$$s = \left[\frac{2\Gamma + 4}{p} \right] + 1 \quad \text{and} \quad m = \left[\frac{n-1}{s} \right]. \tag{6.10}$$

¹ See [9, Lemma 6.3.8, p. 108] where the constant 1 was accidentally left out.

² Here $[\cdot]$ denotes the integer part.

Given $x \in [-1, 1]$, define the polynomial $Q = Q_x \in \Pi_{n-1}$ by

$$Q(t) = \left(\frac{K_m(v, x, t)}{K_m(v, x, x)} \right)^s. \quad (6.11)$$

If $m=0$, then $p(n-1) \leq 2\Gamma+4$ and $Q(t) \equiv 1$, so we can obtain the inequality in Theorem 2.1 immediately as an application of (5.1) and Lemma 6.1. More specifically,

$$\begin{aligned} \lambda_n(\alpha, p, x) &\leq \int_{-1}^1 w(t) dt \leq 2 \max_{-1 \leq t \leq 1} w(t) \\ &\leq 2 \left(\frac{4}{\Delta_M(x)} \right)^{\Gamma} \max_{|t-x| \leq \Delta_M(x)} w(t) \\ &\leq 2 \left(\frac{4}{\Delta_M(x)} \right)^{\Gamma} \Delta_m^{-1}(x) c_8^2(\Gamma+1)^2 w_M(x) \\ &\leq \frac{1}{2} \left[2 \left(1 + \frac{p(n-1)}{\Gamma+p+1} \right) \right]^{2\Gamma+2} c_8^2(\Gamma+1)^2 w_M(x) \\ &\leq \frac{1}{2} \left[2 \left(1 + \frac{2\Gamma+4}{\Gamma+p+1} \right) \right]^{2\Gamma+2} c_8^2(\Gamma+1)^2 \omega_M(x) \leq c_{16}^{\Gamma+1} w_M(x), \\ &\quad -1 \leq x \leq 1, \end{aligned}$$

where $c_{16} > 0$ is an absolute constant.

Thus, in what follows, we may assume $m \geq 1$. We may also assume without loss of generality that $0 \leq x \leq 1$. For the sake of brevity, we introduce the intervals

$$I_m(x) = [x - \Delta_m(x), x + \Delta_m(x)], \quad -1 \leq x \leq 1.$$

It follows from (6.5), (6.10), and (6.11) that the inequality

$$\int_{I_m(x) \cap [-1, 1]} |Q(t)|^p w(t) dt \leq c_{17}^{2\Gamma+4+p} \Delta_m(x) \max_{t \in I_m(x)} w(t) \quad (6.12)$$

holds with an absolute constant $c_{17} > 0$. By Lemma 6.1,

$$w(t) \leq \left[\frac{2|t-x|}{\Delta_m(x)} \right]^{\Gamma} \max_{y \in I_m(x)} w(y), \quad t \notin I_m(x), \quad (6.13)$$

for every $w \in |\text{GCAP}|_{\Gamma}$.

Using (6.11), (6.10), (6.6), (6.4), and elementary estimates, it can be shown that the inequalities

$$|Q(t)|^p \leq c_{18}^{2\Gamma+4+p} \left(\frac{\sqrt{1-x^2}}{m|x-t|} \right)^{\Gamma+2}, \quad t \in [2x-1, 1] \setminus I_m(x), \quad (6.14)$$

$$|Q(t)|^p \leq c_{18}^{2\Gamma+4+p} (m\sqrt{1-t^2})^{-2\Gamma-4}, \quad t \in [-0.5, 2x-1] \setminus I_m(x), x \geq 0.5, \quad (6.15)$$

and

$$|Q(t)|^p \leq c_{18}^{2\Gamma+4+p} m^{-2\Gamma-4}, \quad x, t \in [-1, 1], |t-x| \geq 0.5, \quad (6.16)$$

hold with some absolute constant $c_{18} > 0$, and we can assume that $c_{18} \geq 4$. For instance, to prove (6.14), we notice that $sp > \Gamma + 2$, and $1 + (\sqrt{1-x^2} + \sqrt{1-t^2})|x-t|^{-1}$ on the right-hand side of (6.6) can be estimated by $5\sqrt{1-x^2}|x-t|^{-1}$ for $t \in [2x-1, 1] \setminus I_m(x)$, whereas for (6.15), we use $1-x \leq 1-t$ and $0 \leq x-t \leq 1-t \leq 2(x-t)$ for $t \in [-0.5, 2x-1]$ and $x \geq 0.5$.

Using (6.13) and (6.14) we obtain

$$\begin{aligned} & \int_{[2x-1, 1] \setminus I_m(x)} |Q(t)|^p w(t) dt \\ & \leq 2^\Gamma c_{18}^{2\Gamma+4+p} \left(\frac{\sqrt{1-x^2}}{m \Delta_m(x)} \right)^\Gamma \Delta_m^2(x) \\ & \quad \times \max_{y \in I_m(x)} w(y) \int_{[2x-1, 1] \setminus I_m(x)} |t-x|^{-2} dt \\ & \leq c_{18}^{3\Gamma+5+p} \Delta_m(x) \max_{y \in I_m(x)} w(y). \end{aligned} \quad (6.17)$$

In addition, by (6.13), (6.15), and by $0 \leq x-t \leq 1-t$,

$$\begin{aligned} & \int_{[-0.5, 2x-1] \setminus I_m(x)} |Q(t)|^p w(t) dt \\ & \leq 2^\Gamma c_{18}^{2\Gamma+4+p} m^{-2\Gamma-4} \max_{y \in I_m(x)} w(y) \\ & \quad \times \int_{[-0.5, 2x-1] \setminus I_m(x)} \left[\frac{(1-t^2)}{\Delta_m(x)} \right]^\Gamma (1-t^2)^{-\Gamma-2} dt \\ & \leq c_{18}^{3\Gamma+4+p} m^{-4} \max_{y \in I_m(x)} w(y) \int_{[-0.5, 2x-1] \setminus I_m(x)} (1-t^2)^{-2} dt \\ & \leq c_{18}^{3\Gamma+5+p} m^{-4} \Delta_m^{-1}(x) \max_{y \in I_m(x)} w(y) \leq c_{18}^{3\Gamma+5+p} \Delta_m(x) \max_{y \in I_m(x)} w(y) \end{aligned} \quad (6.18)$$

for $0.5 \leq x \leq 1$. Finally, from (6.13) and (6.16) we obtain

$$\begin{aligned} & \int_{[-1, 1] \setminus [x-0.5, x+0.5]} |Q(t)|^p w(t) dt \\ & \leq 4^\Gamma c_{18}^{2\Gamma+4+p} m^{-4} (m^2 \Delta_m(x))^{-\Gamma} \max_{y \in I_m(x)} w(y) \\ & \leq c_{18}^{3\Gamma+4+p} \Delta_m(x) \max_{y \in I_m(x)} w(y) \end{aligned} \tag{6.19}$$

by a generous estimate, where we used the fact that $|x - t| \leq 2$ in (6.13). Since $Q \in \Pi_{n-1}$ and $Q(x) = 1$, the extremal property (1.5) and the inequalities (6.12) and (6.17)–(6.19) yield

$$\lambda_n(\alpha, p, x) \leq \int_{-1}^1 |Q(t)|^p w(t) dt \leq c_{19}^{\Gamma+p+1} \Delta_m(x) \max_{y \in I_m(x)} w(y) \tag{6.20}$$

with an absolute constant $c_{19} \geq 4$. By (6.10),

$$\frac{p(n-1)}{2\Gamma+4+p} \leq m \leq \frac{p(n-1)}{2\Gamma+4}. \tag{6.21}$$

Therefore, Lemma 6.1, inequalities (6.21), the facts that $m \geq 1$ and $M = 1 + p(n-1)/(\Gamma + p + 1)$, and inequality (5.1) imply

$$\Delta_m(x) \max_{y \in I_m(x)} w(y) \leq c_{20}^{\Gamma+p+1} \Delta_M(x) \max_{y \in I_M(x)} w(y) \leq c_{21}^{\Gamma+p+1} w_M(x)$$

with some appropriately chosen absolute constants $c_{20} > 0$ and $c_{21} > 0$, which together with (6.20) yields Theorem 2.1. ■

Proof of Theorem 2.2. We will prove the appropriate upper bound for λ_n only; the corresponding estimate for λ_n^* is analogous, and it requires only minor modifications of the following proof. Let

$$w^{(B)}(z) = |\omega| \prod_{j=1}^d |z - z_j|^{r_j} \in |\text{GCAP}|_\Gamma, \quad z_i \neq z_j \text{ if } i \neq j. \tag{6.22}$$

Set³

$$q_j = \left\lceil \frac{r_j}{p} \right\rceil + 1, \quad q = \sum_{j=1}^k q_j \leq \frac{\Gamma + pd}{p}, \tag{6.23}$$

$$s = \left\lceil \frac{2\Gamma + 4}{p} \right\rceil + 1, \quad m = \left\lceil \frac{n-1-q}{s} \right\rceil \geq 1, \tag{6.24}$$

$$P(\cdot) = |\omega| \prod_{j=1}^d (\cdot - z_j)^{q_j} \in \Pi_q, \tag{6.25}$$

³ Here $\lceil \cdot \rceil$ denotes the integer part.

and

$$\tilde{P}_x(\cdot) = |\omega| \prod_{j=1}^d (\cdot - \tilde{z}_j)^{q_j} \in \Pi_q, \quad -1 \leq x \leq 1, \tag{6.26}$$

where

$$\tilde{z}_j = \begin{cases} z_j, & \text{if } |z_j - x| \geq \frac{1}{2} \Delta_m(x) \\ x - \Delta_m(x), & \text{if } |z_j - x| < \frac{1}{2} \Delta_m(x). \end{cases} \tag{6.27}$$

Let

$$Q(\cdot) = \left(\frac{K_m(v, x, \cdot)}{K_m(v, x, x)} \right)^s \tilde{P}_x(\cdot) \in \Pi_{n-1}. \tag{6.28}$$

Obviously,

$$|Q(x)|^p = |\tilde{P}_x(x)|^p. \tag{6.29}$$

Introducing the sets

$$J_1 = [x - \Delta_m(x), x + \Delta_m(x)] \cap [-1, 1] \quad \text{and} \quad J_2 = [-1, 1] \setminus J_1, \tag{6.30}$$

replacing w by $W = |\tilde{P}_x|^p w \in |\text{GCAP}|_{\Gamma+pd}$ in the proof of Theorem 2.1, and using (6.22)–(6.30) and (5.1), we obtain

$$\begin{aligned} \int_{J_2} |Q(t)|^p w(t) dt &= \int_{J_2} \left| \frac{K_m(v, x, t)}{K_m(v, x, x)} \right|^{sp} |\tilde{P}_x(t)|^p w(t) dt \\ &\leq 4^{\Gamma+pd} \int_{J_2} \left| \frac{K_m(v, x, t)}{K_m(v, x, x)} \right|^{sp} |P(t)|^p w(t) dt \\ &\leq 4^{\Gamma+pd} c_{22}^{\Gamma+pd+p+1} \int_{|t-x| \leq \Delta_m(x)} |P(t)|^p w(t) dt \end{aligned} \tag{6.31}$$

with an absolute constant $c_{22} > 0$ chosen in such a way that $c_{19}^{\Gamma+pd+p+1} c_8^2 (\Gamma+pd+1)^2 \leq c_{22}^{\Gamma+pd+p+1}$. Here we used that $|\tilde{P}_x(t)| \leq 4^{\Gamma+pd} |P(t)|$ for every $t \in J_2$ which follows from (6.22)–(6.27). Observe that (6.22)–(6.27) and (6.30) imply $|P(t)|^p \leq 2^{\Gamma+pd} |\tilde{P}_x(x)|^p$ for every $t \in J_2$. This, together with (6.31), yields

$$\begin{aligned} \int_{J_2} |Q(t)|^p w(t) dt &\leq 8^{\Gamma+pd} c_{22}^{\Gamma+pd+p+1} |\tilde{P}_x(x)|^p \int_{|t-x| \leq \Delta_m(x)} w(t) dt \\ &\leq c_{23}^{\Gamma+pd+p+1} |\tilde{P}_x(x)|^p \int_{|t-x| \leq \Delta_m(x)} w(t) dt \end{aligned} \tag{6.32}$$

with an absolute constant $c_{23} > 0$. In addition, (6.22)–(6.27) and (6.30)

imply $|\tilde{P}_x(t)|^p \leq 3^{\Gamma+pd} |\tilde{P}_x(x)|^p$ for every $t \in J_1$, so that by (6.28), (6.23), (6.24), and (6.5)

$$\begin{aligned} \int_{J_1} |Q(t)|^p w(t) dt &\leq c_{24}^{2\Gamma+4+p} \int_{J_1} |\tilde{P}_x(t)|^p w(t) dt \\ &\leq c_{24}^{2\Gamma+4+p} 3^{\Gamma+pd} |\tilde{P}_x(x)|^p \int_{J_1} w(t) dt \\ &\leq c_{25}^{\Gamma+pd+p+1} |\tilde{P}_x(x)|^p \int_{J_1} w(t) dt, \end{aligned} \tag{6.33}$$

where $c_{24} > 0$ and $c_{25} > 0$ are absolute constants. Therefore, the extremal property (1.5) and formulas (6.32), (6.33), (6.30), (6.28), (6.23), and (6.24) yield

$$\begin{aligned} \lambda_n(\alpha, p, x) &\leq |\tilde{P}_x(x)|^{-p} \int_{-1}^1 |Q(t)|^p w(t) dt \leq c_{26}^{\Gamma+pd+p+1} w_M(x) \\ &\leq c_{26}^{\Gamma+pd+p+1} w_M(x), \end{aligned}$$

where $c_{26} > 0$ is an absolute constant and $M = (p(n-1-d) - \Gamma)/(2\Gamma+4+p)$, and, hence, the theorem has completely been proved. ■

7. PROOF OF THEOREMS 3.1 AND 3.2

Proof of Theorem 3.1. Let $x \in [-1, 1]$ be fixed, $N = \Gamma + p(n-1)$, and $M = 1 + p(n-1)/(\Gamma + p + 1)$, and let⁴

$$I_{M,\eta}(x) = [x - \eta \Delta_M(x), x + \eta \Delta_M(x)], \quad 0 < \eta \leq 1.$$

Let the weight $w \in |\text{GCAP}|_\Gamma$ be of the form

$$w(z) = |\omega| \prod_{j=1}^k |z - z_j|^{r_j}, \quad \omega \neq 0, z_j \in \mathbb{C}, z \in \mathbb{C}, r_j > 0, \sum_{j=1}^k r_j \leq \Gamma. \tag{7.1}$$

In what follows, $A_M(x)$ denotes the open disk in the complex plane centered at x with radius $0.125 \cdot \Delta_M(x)$. Define \tilde{z}_j and \tilde{w} by

$$\tilde{z}_j = \begin{cases} z_j, & \text{if } z_j \in \mathbb{C} \setminus A_M(x) \\ x - 0.125 \cdot \text{sgn}(x) \Delta_M(x), & \text{if } z_j \in A_M(x), \end{cases} \tag{7.2}$$

and

$$\tilde{w}(\cdot) = |\omega| \prod_{j=1}^k |\cdot - \tilde{z}_j|^{r_j} \in |\text{GCAP}|_\Gamma, \tag{7.3}$$

⁴ See formula (1.7) for the definition of $\Delta_M(x)$.

respectively. In what follows, we assume that $Q \in |\text{GCAP}|_{n-1}$. Observe that $Q \in |\text{GCAP}|_{n-1}$ and $\tilde{w} \in |\text{GCAP}|_\Gamma$ imply $|Q|^\rho \tilde{w} \in |\text{GCAP}|_N$ with $N = \Gamma + p(n-1)$, and, therefore, using Theorem 5.1 with $0 < s = \Delta_M(x) < 1$, where $M = 1 + p(n-1)/(\Gamma + p + 1)$, we obtain

$$m(\{t \in [-1, 1]: |Q(t)|^\rho \tilde{w}(t) \geq \exp(-c_7(\Gamma + p + 1))|Q(x)|^\rho \tilde{w}(x)\}) \geq \Delta_M(x).$$

Hence, there is a set

$$E \subset [-1, 1] \setminus [x - 0.25 \cdot \Delta_M(x), x + 0.25 \cdot \Delta_M(x)] \tag{7.4}$$

such that

$$\Delta_M(x) \leq 2m(E) \tag{7.5}$$

and

$$|Q(t)|^\rho \tilde{w}(t) \geq \exp(-c_7(\Gamma + p + 1))|Q(x)|^\rho \tilde{w}(x), \quad t \in E. \tag{7.6}$$

Observe that (7.1), (7.2), (7.3), and (7.4) yield

$$\tilde{w}(x) \geq (2/3)^\Gamma \max_{t \in I_{m,0.0625}(x)} w(t) \tag{7.7}$$

and

$$w(t) \geq 3^{-\Gamma} \tilde{w}(t), \quad t \in E. \tag{7.8}$$

From inequalities (7.5), (7.6), and (7.7) we obtain

$$\begin{aligned} & \frac{2 \exp(c_7(\Gamma + p + 1))}{\Delta_M(x)} \int_E |Q(t)|^\rho \tilde{w}(t) dt \\ & \geq |Q(x)|^\rho \tilde{w}(x) \geq |Q(x)|^\rho (2/3)^\Gamma \max_{t \in I_{m,0.0625}(x)} w(t). \end{aligned}$$

We can use this estimate combined with (7.8) and Lemma 6.1 to obtain the existence of an absolute constant $c_{27} > 0$ such that

$$\begin{aligned} & \int_{-1}^1 |Q(t)|^\rho w(t) dt \\ & \geq \int_E |Q(t)|^\rho w(t) dt \geq 3^{-\Gamma} \int_E |Q(t)|^\rho \tilde{w}(t) dt \\ & \geq 3^{-\Gamma} \exp(-c_7(\Gamma + 1)) 2^{-1} \Delta_M(x) (2/3)^\Gamma |Q(x)|^\rho \max_{t \in I_{m,0.0625}(x)} w(t) \\ & \geq c_{27}^{\Gamma+p+1} \Delta_M(x) |Q(x)|^\rho \max_{|t-x| \leq \Delta_M(x)} w(t) \geq c_{27}^{\Gamma+p+1} |Q(x)|^\rho w_M(x). \end{aligned} \tag{7.9}$$

Since (7.9) holds for every $Q \in |\text{GCAP}|_{n-1}$, Theorem 3.1 has completely been proved in view of the extremal property (1.6) of the generalized Christoffel function λ^* . ■

Proof of Theorem 3.2. In what follows we will assume that $0 < p < \infty$ and $x \in [-1, 1]$. We start with the inequality

$$|P_n(z)|^p |g(rz)|^2 \leq \frac{(2+pn)e}{8\pi} \int_{-\pi}^{\pi} |P_n(e^{i\theta})|^p |g(e^{i\theta})|^2 d\theta, \quad (7.10)$$

$$|z| = 1, \quad r = \frac{pn}{pn+2},$$

which holds for every $g \in H^2(|z| \leq 1)$ and for every complex algebraic polynomial P_n of degree at most $n = 1, 2, \dots$ (cf. [7, Theorem 6, p. 148]). A simple calculation shows that for every real trigonometric polynomial R_n of degree at most n there is an algebraic polynomial $P_{2n} \in \Pi_{2n}$ such that $R_n^2(\theta) = |P_{2n}(e^{i\theta})|^2$. Therefore, (7.10) yields

$$|R_n(\gamma)|^p |g(re^{i\gamma})|^2 \leq \frac{(1+pn)e}{4\pi} \int_{-\pi}^{\pi} |R_n(\theta)|^p |g(e^{i\theta})|^2 d\theta, \quad (7.11)$$

$$\gamma \in \mathbb{R}, \quad r = \frac{pn}{pn+2},$$

holds for every real trigonometric polynomial R_n and for every $g \in H^2(|z| \leq 1)$. First, we extend (7.11) to every $f \in |\text{GCTP}|_N$, writing N in place of n . To this end, first assume f can be written as

$$f(z) = |\omega| \prod_{j=1}^k \left| \sin \frac{z-z_j}{2} \right|^{r_j}, \quad \omega \neq 0, z_j \in \mathbb{C}, z \in \mathbb{C}, \sum_{j=1}^k r_j \leq 2N,$$

with some rational exponents $r_j > 0$. If $r_j = q_j/q$ with some positive integers q_j and q for $1 \leq j \leq k$, then, applying (7.11) with $p/(2q)$ instead of p to the trigonometric polynomial R_n of degree at most $2qN$, where

$$R_n(z) = |\omega|^{2q} \prod_{j=1}^k \left(\sin \frac{z-z_j}{2} \sin \frac{z-\bar{z}_j}{2} \right)^{q_j},$$

we obtain

$$|f(\gamma)|^p |g(re^{i\gamma})|^2 \leq \frac{(1+pN)e}{4\pi} \int_{-\pi}^{\pi} |f(\theta)|^p |g(e^{i\theta})|^2 d\theta, \quad (7.12)$$

$$\gamma \in \mathbb{R}, \quad r = \frac{pN}{pN+2},$$

for every $f \in |\text{GCTP}|_N$ and for every $g \in H^2(|z| \leq 1)$. Now let $1 \leq n < \infty$, $0 < p < \infty$, and let $Q \in |\text{GCAP}|_{n-1}$. Applying (7.12) with f , where $f(\theta) = |Q(\cos \theta)| |\sin \theta|^{1/p} \in |\text{GCTP}|_{n-1+1/p}$, and choosing $g = D(w(\cos \cdot)) \in H^2(|z| \leq 1)$, we obtain

$$\begin{aligned} & |Q(\cos \gamma)|^p |\sin \gamma| |D(w(\cos \theta), re^{i\gamma})|^2 \\ & \leq \frac{(2 + p(n-1))e}{4\pi} \int_{-\pi}^{\pi} |Q_n(\cos \theta)|^p |\sin \theta| |D(w(\cos \cdot), e^{i\theta})|^2 d\theta, \\ & r = \frac{p(n-1) + 1}{p(n-1) + 3}, \end{aligned} \tag{7.13}$$

for every $Q \in |\text{GCAP}|_{n-1}$. Since $|D(w(\cos \cdot), e^{i\theta})|^2 = w(\cos \theta)$ for almost every real θ , we can use the change of variables $t = \cos \theta$ and $x = \cos \gamma$ to obtain from (7.13) the inequality

$$\begin{aligned} & |Q(x)|^p \sqrt{1-x^2} \\ & \leq |D(w(\cos \cdot), re^{i\theta})|^{-2} \frac{(2 + p(n-1)) 2e}{4\pi} \int_{-1}^1 |Q(t)|^p w(t) dt, \\ & r = \frac{p(n-1) + 1}{p(n-1) + 3}, \end{aligned}$$

for every $Q \in |\text{GCAP}|_{n-1}$. To finish the proof, we refer to our Schur-type inequality [5, Theorem 7] which states that given $0 < N < \infty$ and $0 < p < \infty$,

$$\max_{-1 \leq t \leq 1} |Q(t)|^p \leq e(1 + pN) \max_{-1 \leq t \leq 1} (\sqrt{1-t^2} |Q(t)|^p)$$

for every $Q \in |\text{GCAP}|_N$. Now Theorem 3.2 follows from the last two inequalities, and from the definition of $\Delta_M(x)$ in (1.7). ■

8. PROOF OF THEOREM 4.1

Proof of Theorem 4.1. Given $n = 1, 2, \dots$ and $0 \leq \Gamma < \infty$, let $M = 1 + 2(n-1)/(\Gamma + 3)$. First we prove (4.1). It is sufficient to prove the existence of a positive constant c_{28} such that

$$x_{m-1,n} - x_{m,n} \leq c_{28}^{\Gamma+1} (\Delta_M(x_{m-1,n}) + \Delta_M(x_{m,n})), \quad m = 1, 2, \dots, n+1, \tag{8.1}$$

since (8.1) implies (4.1) by a straightforward calculation (cf. (1.7)). Assume that there is an integer m such that $1 \leq m \leq n+1$ and

$$x_{m-1,n} - x_{m,n} > \Delta_M(x_{m-1,n}) + \Delta_M(x_{m,n}), \tag{8.2}$$

since, otherwise, we already have the desired upper bound for this particular value of m . We introduce the Christoffel numbers

$$\lambda_{m,n} = \lambda_n(\alpha, 2, x_{m,n}), \quad m = 0, 1, \dots, n+1. \quad (8.3)$$

By the Markov–Stieltjes inequality [6, formula (5.4), p. 29] we have

$$\int_{x_{m,n}}^{x_{m-1,n}} w(t) dt \leq \lambda_{m-1,n} + \lambda_{m,n}, \quad m = 1, 2, \dots, n+1. \quad (8.4)$$

Since $w \in |\text{GCAP}|_\Gamma$, Theorem 2.1 yields

$$\begin{aligned} \lambda_{m,n} &\leq c_1^{\Gamma+3} w_M(x_{m,n}) = c_1^{\Gamma+3} \int_{|t-x_{m,n}| \leq \Delta_M(x_{m,n})} w(t) dt \\ &\leq c_1^{\Gamma+3} \Delta_M(x_{m,n}) \max_{|t-x_{m,n}| \leq \Delta_M(x_{m,n})} w(t), \quad m = 0, 1, \dots, n+1. \end{aligned} \quad (8.5)$$

Observe that assumption (8.2) and Lemma 6.1 imply

$$\max_{|t-x_{m,n}| \leq \Delta_M(x_{m,n})} w(t) \leq 6^\Gamma \max_{x_{m,n} \leq t \leq x_{m-1,n}} w(t), \quad m = 1, 2, \dots, n+1, \quad (8.6)$$

and

$$\max_{|t-x_{m-1,n}| \leq \Delta_M(x_{m-1,n})} w(t) \leq 6^\Gamma \max_{x_{m,n} \leq t \leq x_{m-1,n}} w(t), \quad m = 1, 2, \dots, n+1. \quad (8.7)$$

Now inequalities (8.4)–(8.7) and Theorem 5.2 yield

$$\begin{aligned} &(x_{m-1,n} - x_{m,n}) \max_{x_{m,n} \leq t \leq x_{m-1,n}} w(t) \\ &\leq 2c_3^2(\Gamma+1)^2 \int_{x_{m,n}}^{x_{m-1,n}} w(t) dt \leq 2c_3^2(\Gamma+1)^2(\lambda_{m-1,n} + \lambda_{m,n}) \\ &\leq 2c_3^2(\Gamma+1)^2 c_1^{\Gamma+3} 6^\Gamma (\Delta_M(x_{m-1,n}) + \Delta_M(x_{m,n})) \max_{x_{m,n} \leq t \leq x_{m-1,n}} w(t), \end{aligned}$$

for $m = 1, 2, \dots, n+1$, and, thus, (8.1) follows. Hence, (4.1) has been proved.

The proof of (4.2) is somewhat more complicated; it is based on a method of Erdős and Turán (cf. [6, pp. 111–112; 8, p. 369; 9, pp. 164–165]). Given $n = 1, 2, \dots$, and $0 \leq \Gamma < \infty$, let $M = 1 + 2(n-1)/(\Gamma+3)$, and let

$$I_{M,n}(x) = [x - \eta \Delta_M(x), x + \eta \Delta_M(x)], \quad 0 < \eta \leq 1.$$

Let $\{l_{m,n}\}_{m=1}^n$ denote the fundamental polynomials of Lagrange interpolation defined by

$$l_{m,n} \in \Pi_{n-1} \quad \text{and} \quad l_{m,n}(x_{j,n}) = \delta_{j,m}, \quad j, m = 1, 2, \dots, n, \quad (8.8)$$

where $\delta_{j,m}$ is the Kronecker-delta symbol. The identity

$$\frac{1}{\lambda_n(\alpha, 2, t)} = \sum_{j=1}^n \frac{l_{j,n}^2(t)}{\lambda_{j,n}}, \quad (8.9)$$

where $\lambda_{j,n}$ is given by (8.3), is well known (cf. [6, formula (4.7), p. 25]). From (8.9), Theorems 2.1 and 3.1, Lemma 6.1, and (5.1) we obtain

$$\begin{aligned} l_{m,n}^2(t) &= \frac{l_{m,n}^2(t)}{\lambda_{m,n}} \lambda_{m,n} \leq \sum_{j=1}^n \frac{l_{j,n}^2(t)}{\lambda_{j,n}} \lambda_{m,n} = \frac{\lambda_{m,n}}{\lambda_n(\alpha, 2, t)} \\ &\leq c_{29}^{r+3} \frac{\Delta_M(x_{m,n}) \max_{|y-x_{m,n}| \leq \Delta_M(x_{m,n})} w(y)}{\Delta_M(t) \max_{|y-t| \leq \Delta_M(t)} w(y)} \leq c_{30}^{r+1}, \quad t \in I_{m,1} \cap [-1, 1], \end{aligned} \quad (8.10)$$

with some appropriate absolute constants $c_{29} > 0$ and $c_{30} > 0$. Let $A_n(x)$ denote the open disk centered at x with radius $0.5 \cdot \Delta_n(x)$. Given

$$w(\cdot) = \prod_{j=1}^k |\cdot - z_j|^{r_j}, \quad \sum_{j=1}^k r_j \leq r, \quad (8.11)$$

we define

$$\tilde{z}_j = \begin{cases} z_j, & \text{if } z_j \in \mathbb{C} \setminus A_n(x_{m,n}) \\ x_{m,n} - 0.5 \cdot \text{sgn}(x_{m,n}) \Delta_M(x_{m,n}), & \text{if } z_j \in A_n(x_{m,n}), \end{cases} \quad (8.12)$$

and

$$\tilde{w}(\cdot) = \prod_{j=1}^k |\cdot - \tilde{z}_j|^{r_j}. \quad (8.13)$$

We will estimate

$$\int_{-1}^1 l_{m,n}^2(t) \tilde{w}(t) dt.$$

To this end we introduce the sets

$$J_1 = I_{M,1}(x_{m,n}) \cap [-1, 1] \quad \text{and} \quad J_2 = [-1, 1] \setminus J_1. \quad (8.14)$$

It follows from (8.10)

$$\begin{aligned} \int_{J_1} l_{m,n}^2(t) \tilde{w}(t) dt &\leq c_{30}^{\Gamma+1} \Delta_M(x_{m,n}) \max_{|t-x_{m,n}| \leq \Delta_M(x_{m,n})} \tilde{w}(t) \\ &\leq c_{30}^{\Gamma+1} \Delta_M(x_{m,n}) \cdot 8^\Gamma \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t). \end{aligned} \quad (8.15)$$

Observe that (8.11)–(8.14) imply $\tilde{w}(t) \leq 3^\Gamma w(t)$ for $t \in J_2$ and $w(t) \leq 3^\Gamma \tilde{w}(t)$ for $t \in I_{M,0.25}(x_{m,n})$. Therefore, by the Gauss–Jacobi quadrature formula (cf. [6, formula (3.9), p.23]), Theorem 2.1 and Lemma 6.1 yield

$$\begin{aligned} \int_{J_2} l_{m,n}^2(t) \tilde{w}(t) dt &\leq 3^\Gamma \int_{J_2} l_{m,n}^2(t) w(t) dt \leq 3^\Gamma \int_{-1}^1 l_{m,n}^2(t) w(t) dt = 3^\Gamma \lambda_{m,n} \\ &= 3^\Gamma \lambda_n(\alpha, 2, x_{m,n}) \leq 3^\Gamma c_1^{\Gamma+3} \Delta_M(x_{m,n}) \max_{|t-x_{m,n}| \leq \Delta_M(x_{m,n})} w(t) \\ &\leq 3^\Gamma c_1^{\Gamma+3} \Delta_M(x_{m,n}) \cdot 8^\Gamma \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} w(t) \\ &\leq 3^\Gamma c_1^{\Gamma+3} \Delta_M(x_{m,n}) \cdot 8^\Gamma \cdot 3^\Gamma \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t). \end{aligned} \quad (8.16)$$

Summarizing (8.15) and (8.16), we get

$$\int_{-1}^1 l_{m,n}^2(t) \tilde{w}(t) dt \leq c_{31}^{\Gamma+1} \Delta_M(x_{m,n}) \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t) \quad (8.17)$$

with an appropriate absolute constant $c_{31} > 0$. When proving (4.2), we can assume without loss of generality that $x_{m,n} \leq 0$. We may also assume that $x_{m-1,n} \leq 0.25$; otherwise, there is nothing to prove. Now Theorem 5.4, (8.17), and the inequalities $x_{m,n} \leq 0$ and $x_{m-1,n} \leq 0.25$ imply

$$\begin{aligned} l_{m,n}^2(x) \tilde{w}(x) &\leq c_{10} (\Delta_{2n+\Gamma}(x))^{-1} \int_{-1}^1 l_{m,n}^2(t) \tilde{w}(t) dt \\ &\leq c_{32}^{\Gamma+1} (\Delta_{2n+\Gamma}(x))^{-1} \Delta_M(x_{m,n}) \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t) \\ &\leq c_{33}^{\Gamma+1} \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t), \\ &|x| \leq |x_{m,n}| + 0.5 \cdot (1 - |x_{m,n}|), \end{aligned} \quad (8.18)$$

with appropriate absolute constants $c_{32} > 0$ and $c_{33} > 0$. Assume that

$$x_{m-1,n} - x_{m,n} \leq 0.25 \cdot \Delta_M(x_{m,n}), \tag{8.19}$$

since, otherwise, we already have the desired lower bound for this particular value of m . In view of (8.11)–(8.13) and (8.19), we have

$$\tilde{w}(x_{m,n}) \leq (1.5)^F \tilde{w}(\xi), \quad \forall \xi \in [x_{m,n}, x_{m-1,n}], \tag{8.20}$$

and

$$\max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t) \leq (1.5)^F \tilde{w}(x_{m,n}). \tag{8.21}$$

Finally, using (8.8), the *Mean Value Theorem*, inequality (5.2) transformed linearly to the interval

$$[-y, y] \equiv [-|x_{m,n}| - 0.5 \cdot (1 - |x_{m,n}|), |x_{m,n}| + 0.5 \cdot (1 - |x_{m,n}|)],$$

the assumptions $x_{m,n} \leq 0$ and $x_{m-1,n} \leq 0.25$, and inequalities (8.18), (8.20), and (8.21), we can find a point $\xi \in (x_{m,n}, x_{m-1,n})$ such that

$$\begin{aligned} \tilde{w}(x_{m,n}) &= |I_{m,n}^2(x_{m-1,n}) - I_{m,n}^2(x_{m,n})| \tilde{w}(x_{m,n}) \\ &= (x_{m-1,n} - x_{m,n}) |(I_{m,n}^2)'(\xi)| \tilde{w}(x_{m,n}) \\ &\leq (x_{m-1,n} - x_{m,n}) (1.5)^F |(I_{m,n}^2)'(\xi)| \tilde{w}(\xi) \\ &\leq (x_{m-1,n} - x_{m,n}) c_{34} (\Gamma + 1)^2 y^{-1} \Delta_n^{-1}(\xi/y) \max_{|x| \leq |x_{m,n}| + 0.5 \cdot (1 - |x_{m,n}|)} \tilde{w}(x) \\ &\leq (x_{m-1,n} - x_{m,n}) c_{35}^{\Gamma+1} \Delta_M^{-1}(x_{m,n}) c_{16}^{\Gamma+1} \max_{|t-x_{m,n}| \leq 0.25 \cdot \Delta_M(x_{m,n})} \tilde{w}(t) \\ &\leq (x_{m-1,n} - x_{m,n}) c_{36}^{\Gamma+1} \Delta_M^{-1}(x_{m,n}) (1.5)^F \tilde{w}(x_{m,n}) \end{aligned}$$

with appropriate absolute constants $c_{34} > 0$, $c_{35} > 0$, and $c_{36} > 0$, and this, together with (1.7), gives the lower bound in (4.2). Thus, the theorem has completely been proved. ■

REFERENCES

1. T. ERDÉLYI, Remez-type inequalities on the size of generalized polynomials, *J. London Math. Soc.*, to appear.
2. T. ERDÉLYI, Markov and Bernstein type inequalities for generalized non-negative polynomials, *Canad. J. Math.* **43** (1991), 495–505.
3. T. ERDÉLYI, Nikol'skii-type inequalities for generalized polynomials and zeros of orthogonal polynomials, *J. Approx. Theory* **66** (1991), 80–92.

4. T. ERDÉLYI, Weighted Markov and Bernstein type inequalities for generalized non-negative polynomials, *J. Approx. Theory* **68** (1992), 283–305.
5. T. ERDÉLYI, A. MÁTÉ, AND P. NEVAI, Inequalities for generalized non-negative polynomials, *Constr. Approx.*, to appear.
6. G. FREUD, “Orthogonal Polynomials,” Pergamon, Oxford, 1971.
7. A. MÁTÉ AND P. NEVAI, Bernstein’s inequality in L^p for $0 < p < 1$ and $(C, 1)$ bounds for orthogonal polynomials, *Ann. of Math.* **111** (1980), 145–154.
8. P. NEVAI, Mean convergence of Lagrange interpolation, I, *J. Approx. Theory* **18** (1976), 363–377.
9. P. NEVAI, “Orthogonal Polynomials,” Mem. Amer. Math. Soc., Vol. 213, Amer. Math. Soc., Providence, RI, 1979.