

## Lieb–Thirring Inequalities for Jacobi Matrices

Dirk Hundertmark and Barry Simon<sup>1,2</sup>

*Department of Mathematics 253-37, California Institute of Technology, Pasadena,  
California 91125, U.S.A.*

E-mail: dirkh@caltech.edu, bsimon@caltech.edu

*Communicated by Walter Van Assche*

Received November 30, 2001; accepted in revised form April 3, 2002

For a Jacobi matrix  $J$  on  $\ell^2(\mathbb{Z}_+)$  with  $Ju(n) = a_{n-1}u(n-1) + b_nu(n) + a_nu(n+1)$ , we prove that

$$\sum_{|E|>2} (E^2 - 4)^{1/2} \leq \sum_n |b_n| + 4 \sum_n |a_n - 1|.$$

We also prove bounds on higher moments and some related results in higher dimension. © 2002 Elsevier Science (USA)

### 1. INTRODUCTION

Let  $J$  be a Jacobi matrix, that is, a tridiagonal matrix

$$J = \begin{pmatrix} b_1 & a_1 & 0 & 0 & \dots \\ a_1 & b_2 & a_2 & 0 & \dots \\ 0 & a_2 & b_3 & a_3 & \dots \\ 0 & 0 & a_3 & b_4 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

viewed as an operator on  $\ell^2(\mathbb{Z}_+)$  via

$$(Ju)(n) = a_{n-1}u(n-1) + b_nu(n) + a_nu(n+1). \quad (1.1)$$

Here  $a_n > 0$  and  $b_n \in \mathbb{R}$ . We will sometimes denote the variables in  $J$  explicitly by writing  $J(\{a_n\}_{n \geq 1}, \{b_n\}_{n \geq 1})$ . We are interested in perturbations

<sup>1</sup>To whom correspondence should be addressed.

<sup>2</sup>Supported in part by NSF Grant DMS-9707661.

of the special case  $a_n \equiv 1$ ,  $b_n = 0$ , called  $J_0$ , the free Jacobi matrix and, in particular, the case where  $J - J_0$  is compact, viz.,  $a_n \rightarrow 1$ ,  $b_n \rightarrow 0$  as  $n \rightarrow \infty$ . Then  $\sigma_{\text{ess}}(J) = \sigma_{\text{ess}}(J_0) = [-2, 2]$  and  $J$  has simple eigenvalues  $\{E_n^\pm\}_{n=1}^{N_\pm}$  with ( $N_+$  or  $N_-$  or both may be infinite)

$$E_1^+ > E_2^+ > \dots > 2 > -2 > \dots > E_2^- > E_1^-. \quad (1.2)$$

One of our main goals in this paper is to prove the following bound

**THEOREM 1.**

$$\sum_{n=1}^{N_+} [(E_n^+)^2 - 4]^{1/2} + \sum_{n=1}^{N_-} [(E_n^-)^2 - 4]^{1/2} \leq \sum_n |b_n| + 4 \sum_n |a_n - 1|. \quad (1.3)$$

As we will see, the constants 1 in front of the  $b$  sum and 4 in the  $a_n - 1$  sum are both optimal. Equation (1.3) is optimal in another regime, namely, large coupling for  $b$ . Specifically, let  $J_\lambda$  be defined with  $a_n = a_n^{(0)}$  and  $b_n = \lambda b_n^{(0)}$ . Let  $\tilde{b}_n^\pm$  be a reordering of the  $b_n$ 's with  $\pm \tilde{b}_n^\pm > 0$  so  $\tilde{b}_1^+ \geq \tilde{b}_2^+ \geq \dots \geq 0$  and  $\tilde{b}_1^- \leq \tilde{b}_2^- \leq \dots \leq 0$ . Then it is not hard to see that

$$\lim_{\lambda \rightarrow \infty} \lambda^{-1} E_n^\pm(J_\lambda) = \tilde{b}_n^\pm, \quad (1.4)$$

which shows that the ratio of the two sides of (1.3) goes to 1 as  $\lambda \rightarrow \infty$  for any  $b_n$  with  $\sum |b_n| < \infty$ .

Since

$$\begin{aligned} (E_n^\pm)^2 - 4 &= |E_n^\pm \mp 2| |E_n^\pm \pm 2| \\ &\geq 4 |E_n^\pm \mp 2|, \end{aligned}$$

(1.3) implies that

$$\sum_n (|E_n^+ - 2|^{1/2} + |E_n^- + 2|^{1/2}) \leq \frac{1}{2} \left( \sum_n |b_n| + 4 \sum_n |a_n - 1| \right). \quad (1.5)$$

More generally, we will prove

**THEOREM 2.**

$$\sum_n (|E_n^+ - 2|^p + |E_n^- + 2|^p) \leq c_p \left[ \sum_n |b_n|^{p+1/2} + 4 \sum_n |a_n - 1|^{p+1/2} \right] \quad (1.6)$$

for any  $p \geq \frac{1}{2}$  where

$$c_p = \frac{1}{2} 3^{p-1/2} \frac{\Gamma(p+1) \Gamma(2)}{\Gamma(p+\frac{3}{2}) \Gamma(\frac{3}{2})}.$$

As for sums of moments for  $p < \frac{1}{2}$ , we will prove

**THEOREM 3.** *Let  $0 \leq p < \frac{1}{2}$ . Let  $\|\cdot\|$  be any translation invariant norm on pairs of sequences  $\{a_n\}_{n=0}^\infty, \{b_n\}_{n=0}^\infty$ . For any  $\varepsilon > 0$ , there exists a Jacobi matrix with  $a_n = 1, b_n = 0$  for  $n$  large so that  $\|(a, b)\| \leq \varepsilon$  but  $\sum_n |E_n^+ - 2|^p + |E_n^- - 2|^p \geq \varepsilon^{-1}$ .*

As (1.4) shows, (1.5) and (1.6) are poor as  $\lambda \rightarrow \infty$ , since the left side grows like  $\lambda^p$  and the right side as  $\lambda^{p+1/2}$ . It is better to use

$$\begin{aligned} (E_n^\pm)^2 - 4 &= |E_n^\pm - 2| |E_n^\pm \pm 2| \\ &\geq |E_n^\pm - 2|^2 \end{aligned}$$

and (1.3) to obtain

$$\sum_n (|E_n^+ - 2| + |E_n^- + 2|) \leq \sum_n |b_n| + 4 \sum_n |a_n - 1| \tag{1.7}$$

and the related.

**THEOREM 4.**

$$\sum_n (|E_n^+ - 2|^p + |E_n^- + 2|^p) \leq \sum_n ((b_n^+ + 2|a_n - 1|)^p + (b_n^- + 2|a_n - 1|)^p) \tag{1.8}$$

for any  $p \geq \frac{1}{2}$ .

As (1.4) shows, the ratio of the two sides of (1.8) is 1 as  $\lambda \rightarrow \infty$ .

We got interested in this problem because Killip–Simon [14] needed a bound like Theorem 1 to prove a conjecture of Nevai [20, 21] that if the right-hand side of (1.3) is finite, then a condition of Szegő holds. They and we expected bounds like (1.3) to hold because of the analogous results for Schrödinger operators.

Nevai’s conjecture says that if  $\sum_n |b_n| + \sum_n |a_n - 1| < \infty$ , then, with  $m$ , the  $m$ -function defined by

$$m(E) = (J - E)_{11}^{-1},$$

we have

$$\int_{-2}^2 \log \operatorname{Im} m(E + i0) \frac{dE}{\sqrt{4 - E^2}} > -\infty. \tag{1.9}$$

Killip–Simon [14] use a sum rule of Case [4, 5] that

$$Z(m) = \sum_n \log|a_n| + \sum \log|\beta_j|, \tag{1.10}$$

where  $\beta_j$  is defined by  $|\beta_j| > 1$  and  $\beta_j + \beta_j^{-1}$  are the listing of the eigenvalues of  $J$  outside  $[-2, 2]$ . In (1.10),  $Z(m)$  is defined by

$$Z(m) = \frac{1}{2\pi} \int_{-2}^2 \log \left( \frac{\sqrt{4 - E^2}}{\operatorname{Im} m(E + i0)} \right) \frac{dE}{\sqrt{4 - E^2}}. \tag{1.11}$$

Equation (1.10) is only proven initially for  $J$  with  $J - J_0$  finite rank. (Or, in any event, not initially for all  $J$ 's with  $J - J_0$  trace class. Eventually, using our bounds here and the theory of Nevanlinna functions, Killip–Simon [14] do prove (1.10) for trace class  $J - J_0$ .) Killip–Simon show  $Z(m)$  is lower semicontinuous as a trace class  $J$  is approximated by cutoff  $J$ 's with  $J - J_0$  finite rank. Thus to prove  $Z(m) < \infty$  (i.e., that (1.9) holds), they need to control the right-hand side of (1.10). Since  $\sum |a_n - 1| < \infty$ , the  $\sum_n \log|a_n|$  is absolutely convergent. Since  $|\beta_j| \sim 1 + (|E_j| - 2)^{1/2}$  for  $E_j$  close to 2, (1.5) implies that  $\sum \log|\beta_j|$  is uniformly bounded.

Theorem 1 should also be interesting in connection with some recent results of Peherstorfer–Yuditskii [22], who focus on the finiteness of the left-hand side of (1.3).

Bounds for Schrödinger operator eigenvalues of the form

$$\sum_{n=1}^{\infty} |E_n|^p \leq L_{p,v} \int_{\mathbb{R}^v} |V(x)|^{p+v/2} d^v x, \tag{1.12}$$

where  $E_n$  are the negative eigenvalues of  $-\Delta + V$  on  $L^2(\mathbb{R}^v)$ , go back 25 years to the work of Lieb and Thirring [18, 19], who used the case  $p = 1, v = 3$  in their celebrated proof of the stability of matter. They proved (1.12) for  $p > 0, v \geq 2$ , and  $p > \frac{1}{2}, v = 1$ , and shortly thereafter, Cwikel [7], Lieb [17], and Rozenblum [24] proved (1.12) in case  $p = 0, v \geq 3$ . It is easy to see (e.g., [15, pp. 156–157; 26]) that it is false in case  $p = 0, v = 2$ .

For many years, the case  $p = \frac{1}{2}, v = 1$  was open. Only in 1996 did Weidl [27] establish this result for  $p = \frac{1}{2}, v = 1$ . For  $v = 1$ , Lieb–Thirring [19] conjectured the optimal value of  $L_{p,v}$  for all  $p \geq \frac{1}{2}$ . They proved their conjecture when  $v = 1$  for  $p = \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \dots$ , and subsequently, Aizenman–Lieb [1] for all  $p \geq \frac{3}{2}$ . Shortly after Weidl's work, Hundertmark *et al.* [12] found a

new proof which yielded the optimal constant  $L_{1/2,1}$ . A partially alternate proof of a part of the argument in [12] can be found in Hundertmark *et al.* [11].

Unlike the discrete case, the continuum theory has a scaling symmetry: taking  $V(x) \rightarrow \lambda^2 V(\lambda x)$  yields  $E_n \rightarrow \lambda^2 E_n$  since there is a unitary operator that implements  $x \rightarrow \lambda x$ . This forces the power  $|E|^p$  on the right-hand side of (1.12) given the scaling behavior of  $d^v x$ . Thus, the same power properly captures large and small  $E$ 's. In the discrete case, this is not so, which is why we have two bounds (1.5) and (1.8). As noted, (1.8) is good for large coupling, but (1.5) is better for small  $E$ 's. In particular, if  $b_n \sim n^{-\alpha}$  (with  $\alpha > 1$ ) for  $n$  large, (1.8) only implies  $\sum |E_n^+ - 2|^p < \infty$  for  $p > \alpha^{-1}$  which (1.5) implies is true for  $p > \alpha^{-1} - \frac{1}{2}$ .

Of course, the best extended estimate would involve powers of  $(E^2 - 4)^{1/2}$  but both the Aizenman–Lieb [1] method to increase powers and the Laptev–Weidl [16] method to increase dimension seem to require powers of  $\text{dist}(E, \sigma_{\text{ess}}(J))$ . However, one can save a little bit of the structure; see the remark at the end of Section 5.

We note one interesting feature of (1.3) vis-à-vis the continuum bound. The continuum  $p = \frac{1}{2}$  bound has an optimal constant, but is off by a factor of 2 in the large coupling limit. For (1.3), as we noted above, the optimal bound for small coupling is also exact in the large coupling limit.

In Section 2, we will prove Theorem 1 when  $a_n \equiv 1$  by closely following [12] and then obtain Theorems 2 and 4 when  $a_n \equiv 1$  by the now standard argument of Aizenman and Lieb [1]. In Section 3, we make a simple but useful observation that allows one to obtain estimates for eigenvalues for arbitrary Jacobi matrices from the estimates for the special case. Section 4 contains some examples and some counterexamples, and proves Theorem 3. Section 5 uses ideas of Laptev–Weidl [16] to prove bounds for the higher-dimensional case. In Appendix A, we show how the ideas in this paper provide a simple proof of a strengthening of the Bargmann-type bound of Geronimo [8, 9].

This paper is aimed towards two rather different audiences: the Schrödinger operator community and the orthogonal polynomial community, who have rather different toolkits. For that reason, we include some material (such as that at the start of Section 2) that one group or the other may regard as elementary.

## 2. BOUNDS FOR DISCRETE SCHRÖDINGER OPERATORS

In this section, we prove Theorems 1, 2, and 4 when all  $a_n = 1$ . We begin with some general preliminaries. Given any self-adjoint operator  $A$ , bounded

from above, we define

$$E_j^+ = \inf_{\varphi_1 \dots \varphi_{j-1}} \sup_{\substack{\psi : \psi \perp \varphi_j \\ \psi \in D, \|\psi\|=1}} \langle \psi, A\psi \rangle. \quad (2.1)$$

Similarly, if  $A$  is bounded below,

$$E_j^- = \sup_{\varphi_1 \dots \varphi_j} \inf_{\substack{\psi : \psi \perp \varphi_j \\ \psi \in D(A), \|\psi\|=1}} \langle \psi, A\psi \rangle. \quad (2.2)$$

We will use  $E_j^\pm(A)$  if the dependence on  $A$  is important. From the definitions,

$$A \leq B \Rightarrow E_j^\pm(A) \leq E_j^\pm(B) \quad (2.3)$$

and

$$E_1^- \leq E_2^- \leq \dots \leq E_2^+ \leq E_1^+. \quad (2.4)$$

The min–max principle [23, Theorem XIII.1] asserts that

- (i)  $E_\infty^\pm = \lim E_j^\pm$  has  $E_\infty^+(A) = \sup \sigma_{\text{ess}}(A)$ ,  $E_\infty^-(A) = \inf \sigma_{\text{ess}}(A)$ .
- (ii) If  $A$  has  $N^+$  (resp.  $N^-$ ) eigenvalues counting multiplicity in the interval  $(E_\infty^+, \infty)$  (resp.  $(-\infty, E_\infty^-)$ ), these eigenvalues are precisely  $E_1^\pm, E_2^\pm, \dots, E_{N_\pm}^\pm$  and  $E_j^\pm = E_\infty^\pm$  if  $j > N_\pm$ .

Next, note from the definition that if  $A_m \rightarrow A$  in norm, then we have convergence of the corresponding eigenvalues since  $|E_j^\pm(A) - E_j^\pm(B)| \leq \|A - B\|$ . It follows if  $f$  is an arbitrary continuous nonnegative function, then

$$\begin{aligned} \sum_{j=1}^k f(E_j^\pm(A)) &= \lim_{m \rightarrow \infty} \sum_{j=1}^k f(E_j^\pm(A_m)) \\ &\leq \liminf_{m \rightarrow \infty} \sum_{j=1}^{\infty} f(E_j^\pm(A_m)), \end{aligned}$$

so taking  $k \rightarrow \infty$ ,

$$\sum_{j=1}^{\infty} (f(E_j^+(A)) + f(E_j^-(A))) \leq \liminf_{m \rightarrow \infty} \sum_{j=1}^{\infty} (f(E_j^+(A_m)) + f(E_j^-(A_m))). \quad (2.5)$$

Equation (2.5) and the min–max principle imply

**PROPOSITION 2.1.** *To prove (1.3)–(1.6), it suffices to prove the special case where only finitely many  $a_n$ 's differ from 1 and finitely many  $b_n$ 's differ from 0.*

Next, we want to note the impact of restriction. Let  $A$  be a bounded self-adjoint operator on  $\mathcal{H}$ . Let  $P$  be an orthogonal projection. By  $A_P$ , we mean  $PAP$  restricted as an operator on  $P\mathcal{H} = \text{Ran } P$ . In (2.1)/(2.2), changing from  $A$  to  $A_P$  adds the condition  $\psi \in \text{Ran } P$  and it decreases sups and increases infs. Thus,

PROPOSITION 2.2.

$$E_j^+(A_P) \leq E_j^+(A), \quad E_j^-(A_P) \geq E_j^-(A). \tag{2.6}$$

We have two applications of (2.6) in mind. First, given two two-sided sequences  $\{a_n\}_{n=-\infty}^\infty, \{b_n\}_{n=-\infty}^\infty$ , define the whole-line operator  $W$  on  $\ell^2(\mathbb{R})$  by

$$(Wu)(n) = a_{n-1}u(n-1) + b_nu(n) + a_nu(n+1). \tag{2.7}$$

Thus, if  $P$  is the projection of  $\ell^2(\mathbb{Z})$  to  $\ell^2(\mathbb{Z}_+) \subset \ell^2(\mathbb{Z})$ ,  $W_P = J$  where  $J$  is built from the projected sequences  $\{a_n\}_{n=1}^\infty$  and  $\{b_n\}_{n=1}^\infty$ . As a result, (2.6) implies

PROPOSITION 2.3. *To prove (1.3)–(1.6), it suffices to prove the analogous result for the whole-line operators.*

One might think that the results are much harder for whole-line operators. After all, it can be shown that if  $\tilde{b}$  has compact support, then  $J(a_n \equiv 1, b_n = \lambda\tilde{b}_n)$  has no spectrum outside  $[-2, 2]$  if  $\lambda$  is small, but  $W(a_n \equiv 1, b_n = \lambda\tilde{b}_n)$  always has eigenvalues outside  $[-2, 2]$  if  $\lambda \neq 0, \tilde{b} \neq 0$ . That is why there is a Bargmann bound for  $J$  but not for  $W$ . However, it is *not* harder because (1.3)–(1.6) have translation invariant quantities for their right side. Let  $P_n$  be the projection onto  $\ell^2(m \in \mathbb{Z}, m \geq n)$ . One can see that as  $n \rightarrow -\infty$ ,  $E_j^\pm(W_{P_n}) \rightarrow E_j^\pm(W)$  so (1.3)–(1.6) for the Jacobi case actually implies it for the whole-line case.

The second application of (2.6) is to the study of the following objects that will play a role below:

$$S_n^\pm(A) = \sum_{j=1}^n E_j^\pm(A). \tag{2.8}$$

PROPOSITION 2.4. *Let  $A$  be a self-adjoint operator.*

- (i)  $S_n^+(A) = \sup\{\text{Tr}(AP) \mid P^* = P, P^2 = P, \text{Tr}(P) = n\}$ .
- (ii)  $S_n^-(A) = \inf\{\text{Tr}(AP) \mid P^* = P, P^2 = P, \text{Tr}(P) = n\}$ .
- (iii)  $A \mapsto S_n^+(A)$  is convex;  $A \mapsto S_n^-(A)$  is concave.

*Remark.* One can see that if  $E_\infty^+ \geq 0$ , then in (i)  $P^2 = P$  can be replaced by  $0 \leq P \leq 1$  which is how it is often written.

*Proof.*

(i) By (2.6),  $S_n^+(A_P) \leq S_n^+(A)$ . But since  $\text{Ran } P$  has dimension  $n$ ,  $S_n^+(A_P) = \text{Tr}(A_P) = \text{Tr}(AP)$ . Thus,

$$S_n^+(A) \geq \sup\{\text{Tr}(AP) \mid P^* = P, P^2 = P, \text{Tr}(P) = n\}.$$

Next, pick  $\varphi_1, \dots, \varphi_n$  as follows. If  $n \leq N^+(A)$ , take  $\varphi_1, \dots, \varphi_n$  to be the eigenfunctions of  $A$  with eigenvalues  $E_1^+, \dots, E_n^+$ . If  $n > N^+$ , pick  $\varphi_1, \dots, \varphi_{N^+}$  to be the eigenfunctions for  $A$  with eigenvalues  $E_1^+, \dots, E_{N^+}^+$  and  $\varphi_{N^++1}, \dots, \varphi_n$  to be arbitrary orthonormal vectors in  $\text{Ran}(P_{[E_\infty^+ - \varepsilon, E_\infty^+]}(A))$ , the range of the spectral projection which is infinite-dimensional when  $N^+ < \infty$  since  $E_\infty^+ = \sup \sigma_{\text{ess}}(A)$ . Let  $P$  be the projection onto the span of  $\varphi_1, \dots, \varphi_n$ . Then,

$$\begin{aligned} \text{Tr}(AP) &= \sum_{j=1}^n (\varphi_j, A\varphi_j) \\ &\geq S_n^+(A) - \varepsilon[n - \min(n, N^+)]. \end{aligned}$$

Since  $\varepsilon$  is arbitrary,

$$S_n^+(A) \leq \sup\{\text{Tr}(AP) \mid P^* = P, P^2 = P, \text{Tr}(P) = n\}.$$

(ii) The same proof as (i).

(iii)  $S_n^\pm$  are the sup and inf of linear functions, so convex and concave, respectively. ■

As a final general preliminary, we note the Birman–Schwinger principle: Let  $A$  be a self-adjoint operator which is bounded above with  $\alpha = \sup \sigma(A)$ . Let  $B$  be a positive relatively form compact, that is,

$$K_\beta \equiv B^{1/2}(\beta - A)^{-1}B^{1/2} \tag{2.9}$$

is compact for one and hence for all  $\beta > \alpha$ .  $K_\beta$  is called the Birman–Schwinger operator.

**PROPOSITION 2.5** (The Birman–Schwinger Principle, Birman [3] and Schwinger [25]). *Let  $\lambda > 0$ .  $\beta > \alpha$  is an eigenvalue of  $A + \lambda B$  if and only if*



$K_\beta$  has eigenvalue  $\lambda^{-1}$ . We have for  $j \leq N^+(A + \lambda B)$ ,

$$E_j^+(K_{E_j^+(A+\lambda B)}) = \lambda^{-1}. \tag{2.10}$$

*Remark.* The point of (2.10) is that the index  $j$  is the same in both  $E_j^+$ 's.

*Proof.* For simplicity, we suppose  $A$  and  $B$  are bounded operators, which is true in the applications we will make. If  $(A + \lambda B)\varphi = \beta\varphi$ , then  $B^{1/2}(\beta - A)^{-1}B^{1/2}(B^{1/2}\varphi) = \lambda^{-1}B^{1/2}\varphi$  and  $B^{1/2}\varphi \neq 0$  since if not, we must have  $A\varphi = \beta\varphi$ , which is impossible since  $\beta > \sup \sigma(A)$ . Conversely, if  $K_\beta\psi = \lambda^{-1}\psi$  and  $\varphi = (\beta - A)^{-1}B^{1/2}\psi$  ( $\neq 0$  since  $\lambda^{-1} \neq 0$ ), we have  $(A + \lambda B)\varphi = \beta\varphi$ . Thus the first expression is true.

Next, note that  $\|K_\beta\| \rightarrow 0$  as  $\beta \rightarrow \infty$  by compactness. Its eigenvalues are continuous, and so by eigenvalue perturbation theory [13, 23], real analytic. If  $e(\beta)$  is a positive eigenvalue of  $K_\beta$  with  $K_\beta\varphi = e\varphi$  and  $\|\varphi\| = 1$ , then by eigenvalue perturbation theory (the Feynman–Hellmann theorem),

$$\frac{de}{d\beta} = \left\langle \varphi, \frac{\partial K_\beta}{\partial \beta} \varphi \right\rangle = -\|(\beta - A)^{-1}B^{1/2}\varphi\|^2 < 0,$$

so  $e$  is strictly monotone. Thus, if  $e(\beta)$  is the  $j$ th eigenvalue of  $K_\beta$  and  $e(\beta_0) > \lambda^{-1}$ , there is exactly one  $\beta > \beta_0$  with  $e(\beta) = \lambda^{-1}$ , so

$$\#\{j \mid E_j^+(K_{\beta_0}) \geq \lambda^{-1}\} = \#\{\beta > \beta_0 \mid E_j^+(K_\beta) = \lambda^{-1}\}$$

(counting multiplicity) from which (2.10) follows. ■

With the general preliminaries out of the way, we compute the Birman–Schwinger operator for  $A = W_0$  and a diagonal (i.e.,  $a_n \equiv 1$ ) perturbation.

**PROPOSITION 2.6.** *Let  $W_0$  be the whole-line matrix with  $a_n \equiv 1$ ,  $b_n \equiv 0$ . Let  $\beta > 2 = \sup \sigma(W_0)$ . Then  $(\beta - W_0)^{-1}$  has matrix elements*

$$[(\beta - W_0)^{-1}]_{nm} = (\mu^{-1} - \mu)^{-1} \mu^{|n-m|}, \tag{2.11}$$

where  $\mu$  is related to  $\beta$  by

$$\beta = \mu + \mu^{-1}, \quad \mu < 1. \tag{2.12}$$

*Remark.* Of course,  $\mu = \frac{1}{2}(\beta - \sqrt{\beta^2 - 4})$  and  $\mu^{-1} = \frac{1}{2}(\beta + \sqrt{\beta^2 - 4})$  so  $\mu^{-1} - \mu = \sqrt{\beta^2 - 4}$ . This is why  $\sqrt{E^2 - 4}$  enters in Theorem 1.

*Proof.* This is a standard calculation. Looking for solutions of

$$\varphi(n - 1) + \varphi(n + 1) = \beta\varphi(n), \tag{2.13}$$

one tries  $\varphi(n) = \zeta^n$  and finds  $\zeta + \zeta^{-1} = \beta$ , so the solutions are  $\zeta = \mu$  and  $\mu^{-1}$ . Let

$$\varphi_{\pm}(n) = \mu^{\pm n}.$$

Both solve (2.13) if  $\mu$  obeys (2.12). Since  $\mu < 1$ ,  $\varphi_+$  is  $\ell^2$  at  $+\infty$ ,  $\varphi_-$  at  $-\infty$ , so the right-hand side of (2.11) which has the form  $(\mu^{-1} - \mu)^{-1} \varphi_-(\min(n, m)) \varphi_+(\max(n, m)) \equiv G_n(m)$  is  $\ell^2$  in  $m$  for each  $n$  with  $((W_0 - \beta)G_n)(m) = 0$  if  $m \neq n$ . By a direct computation (essentially  $\mu^{-1} - \mu$  is the Wronskian of  $\varphi_+$  and  $\varphi_-$ ),  $(\beta_0 - W)G_n = \delta_n$ , that is,  $G_n(m) = [(\beta_0 - W)^{-1} \delta_n](m)$ , proving (2.11). ■

*Remark.* Alternatively, one can use Fourier analysis to compute the inverse.

Because of (2.11), the following operator will enter in our discussion,  $\{b_n\}_{n \in \mathbb{Z}}$  is a positive sequence of finite support,

$$(L_{\mu})_{nm} = b_n^{1/2} \mu^{|n-m|} b_m^{1/2}. \quad (2.14)$$

Recall the definition (2.8) of  $S_m(\cdot)$ . The crucial lemma is

PROPOSITION 2.7. *Let  $0 < \mu < \eta \leq 1$ . Then for any  $n$ ,*

$$S_n^+(L_{\mu}) \leq S_n^+(L_{\eta}). \quad (2.15)$$

*Remark.*

(1) Since  $\text{Tr}(L_{\mu})$  is constant, individual eigenvalues cannot all be monotone.

(2) This is a special case of the warm-up to the proof of Lemma 4 in [12]. Our proof is close to the proof there, except where [12] uses eigenvalue perturbation at  $\mu_j = 0$ , we use symmetry.

*Proof.* Given a bounded positive sequence  $\{\mu_n\}_{n=-\infty}^{\infty}$ , we define

$$(L_{\{\mu_n\}})_{k\ell} = \begin{cases} b_k^{1/2} b_{\ell}^{1/2} \prod_{j=k}^{\ell-1} \mu_j & \text{if } k \leq \ell, \\ (L_{\{\mu_n\}})_{\ell k} & \text{if } k > \ell, \end{cases}$$

so  $L_{\mu}$  is  $L_{\{\mu_n\}}$  when all  $\mu_n = \mu$ . Thus, (2.15) follows if we show  $S_n^+(L_{\{\mu_n\}})$  is monotone in  $\mu_n \in [0, \infty)$  when  $\{\mu_j\}_{j \neq n}$  are held fixed. Let  $f(\mu)$  be this function when  $\mu_n$  takes the value  $\mu$ .  $L_{\{\mu_j\}_{j \neq n}, \mu_n = \mu}$  is affine in  $\mu$  for each matrix element is either constant or a multiple of  $\mu$ . More precisely, in matrix

notation we have

$$L_{\{\mu_j\}_{j \neq n}, \mu_n = \mu} = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} + \mu \begin{pmatrix} 0 & C \\ C^\dagger & 0 \end{pmatrix},$$

where  $A$ ,  $B$ , and  $C$  depend only on  $\{\mu_j\}_{j \neq n}$ . So by Proposition 2.4(iii),  $f(\mu)$  is a convex function of  $\mu$ .

On the other hand, if  $U$  is the diagonal matrix,

$$(U\varphi)(\ell) = \begin{cases} -\varphi(\ell) & \ell \leq n, \\ \varphi(\ell) & \ell \geq n + 1, \end{cases}$$

or, as a block matrix,  $U = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ , then  $UL_{\{\mu\}}U^{-1} = L_{\{\tilde{\mu}\}}$  where

$$\tilde{\mu}_\ell = \begin{cases} \mu_\ell & \text{if } \ell \neq n, \\ -\mu_\ell & \text{if } \ell = n, \end{cases}$$

that is, we have

$$U \begin{pmatrix} A & \mu C \\ \mu C^\dagger & B \end{pmatrix} U^{-1} = \begin{pmatrix} A & -\mu C \\ -\mu C^\dagger & B \end{pmatrix} = L_{\{\mu_j\}_{j \neq n}, \mu_n = -\mu}.$$

Since  $E_j^+$ , and so  $S_j^+$ , are invariant under unitary transformations, we see  $f(-\mu) = f(\mu)$ . An even convex function is monotone increasing on  $[0, \infty)$ , so  $S_n^+(L_{\{\mu_n\}})$  is monotone in each  $\mu_n$  in the region  $\mu_n \geq 0$ . ■

We are now ready to prove what is essentially Theorem 1 in case  $a_n \equiv 1$ .

**THEOREM 2.8.** *Let  $W_0$  be the free whole-line Schrödinger operator and  $B$  a positive finite-rank diagonal matrix. Let  $W = W_0 + B$ . Then*

$$\sum_{j=1}^{N_+(W)} \sqrt{E_j^+(W)^2 - 4} \leq \text{Tr}(B). \tag{2.16}$$

*Proof.* (Following Hundertmark *et al.* [12]). Since  $B$  is finite rank, we know that  $N_j^+(W) < \infty$ . Define  $\mu_j$  by  $\mu_j^{-1} + \mu_j = E_j^+$  with  $\mu_j < 1$ . By (2.9) and the remark after Proposition 2.6,

$$K_{E_j^+} = ((E_j^+)^2 - 4)^{-1/2} L_{\mu_j} \tag{2.17}$$

with  $L_\mu$  given by (2.14). By (2.10),

$$E_j^+(K_{E_j^+}) = 1. \tag{2.18}$$

Since for  $a > 0$ ,  $E_j^+(aA) = aE_j^+(A)$ , (2.17) and (2.18) imply

$$\sqrt{E_j^+(W)^2 - 4} = E_j^+(L_{\mu_j}). \quad (2.19)$$

Thus,

$$\sum_{j=1}^{N^+(W)} \sqrt{E_j^+(W)^2 - 4} = E_1^+(L_{\mu_1}) + E_2^+(L_{\mu_2}) + \cdots + E_{N^+}^+(L_{\mu_{N^+}}). \quad (2.20)$$

But, by (2.15) and  $\mu_1 < \mu_2 < \cdots < \mu_{N^+} < 1$ ,

$$\begin{aligned} E_1^+(L_{\mu_1}) + E_2^+(L_{\mu_2}) &= S_1^+(L_{\mu_1}) + E_2^+(L_{\mu_2}) \\ &\leq S_1^+(L_{\mu_2}) + E_2^+(L_{\mu_2}) \\ &= S_2^+(L_{\mu_2}) \\ &\leq S_2^+(L_{\mu_3}), \end{aligned}$$

so by induction,

$$\sum_{j=1}^k E_j^+(L_{\mu_j}) \leq S_k^+(L_{\mu_k}) \leq S_k^+(L_{\mu_{k+1}})$$

and thus (2.20) implies

$$\sum_{j=1}^{N^+(W)} \sqrt{E_j^+(W)^2 - 4} \leq \sum_{j=1}^{N^+} E_j^+(L_{\mu_{=1}}) = \text{Tr}(B),$$

since  $L_{\mu=1}$  is the rank one operator  $b_n^{1/2} b_m^{1/2}$  with a single nonzero eigenvalue equal to  $\text{Tr}(L_{\mu=1}) = \text{Tr}(B)$ . ■

*Remark.* The proof shows the inequality is strict if  $E_1^+(L_{\mu})$  is strictly monotone. Thus, the inequality is strict if  $\text{rank}(B) \geq 2$ .

There is a standard argument of Aizenman–Lieb [1] which we can use to go from a  $(\frac{1}{2}, 1)$  bound (power of  $E - 2$ , power of  $b$ ) to a general  $(p, p + \frac{1}{2})$  for any  $p \geq \frac{1}{2}$ :

**THEOREM 2.9.** *Under the hypothesis of Theorem 2.8, we have, for any  $p \geq \frac{1}{2}$ ,*

$$\sum_{j=1}^{N_+(W)} |E_j^+(W) - 2|^p \leq \frac{1}{2} \frac{\Gamma(p+1) \Gamma(2)}{\Gamma(p + \frac{3}{2}) \Gamma(\frac{3}{2})} \text{Tr}(|B|^{p+1/2}). \quad (2.21)$$

*Proof.* Note first that since  $b_n \leq (b_n)_+ \equiv \max(0, b_n)$ , if positivity of  $B$  is dropped, we still have that

$$\sum_{j=1}^{N_+(W)} |E_j^+(W) - 2|^{1/2} \leq \frac{1}{2} \sum_n (b_n)_+ \tag{2.22}$$

by using (2.5),  $W_0 + B \leq W_0 + B_+$  and  $\sqrt{E^2 - 4} \leq 2|E - 2|^{1/2}$ .

Let  $r > 0$ . Then

$$(E_j^+(W) - 2 - r)_+ = (E_j^+(W - r\mathbf{1}) - 2)_+,$$

so (2.22) implies

$$\sum_{j=1}^{N_+(W)} |E_j^+(W) - 2 - r|_+^{1/2} \leq \frac{1}{2} \sum_n (b_n - r)_+. \tag{2.23}$$

Now the well-known integral for  $\alpha < p$ ,

$$\frac{\Gamma(p + 1)}{\Gamma(p - \alpha)\Gamma(\alpha + 1)} \int_0^1 (1 - x)^\alpha x^{p-\alpha-1} dx = 1$$

with scaling implies for any  $\alpha < p$ :

$$a_+^p = C_{p,\alpha} \int_0^\infty (a - r)_+^\alpha r^{p-\alpha-1} dr, \tag{2.24}$$

where  $C_{p,\alpha} = \Gamma(p + 1)/\Gamma(p - \alpha)\Gamma(\alpha + 1)$ . Equations (2.23) and (2.24) immediately imply that

$$\sum_{j=1}^{N_+(W)} |E_j^+(W) - 2|^p \leq \frac{1}{2} \frac{C_{p,1/2}}{C_{p+1/2,1}} \sum_n (b_n)_+^{p+1/2}, \tag{2.25}$$

which implies (2.21). ■

Similarly, we have

**THEOREM 2.10.** *Under the hypothesis of Theorem 2.8, we have for any  $p \geq 1$ ,*

$$\sum_{j=1}^{N_+(W)} |E_j(W) - 2|^p \leq \text{Tr}(|B|^p). \tag{2.26}$$

*Proof.* Since  $E \geq 2$  implies

$$\begin{aligned} E^2 - 4 &= (E - 2)^2(E + 2) \\ &\geq (E - 2)^2, \end{aligned}$$

(2.16) implies (2.26) for  $p = 1$ . The result for general  $p \geq 1$  follows as above, where above we get a factor of  $C_{p,1/2}/C_{p-1/2,1}$ , here we get  $C_{p,1}/C_{p,1} = 1$ . ■

So far, we have proven a bound on  $E_j^+$ , but they immediately imply bounds on  $E_j^-$ . One can prove that by analogy, but it is even easier to use the unitary map

$$(Vu)(n) = (-1)^n u(n),$$

which has

$$VW(\{a_n\}, \{b_n\})V^{-1} = W(\{-a_n\}, \{b_n\}) = -W(\{a_n\}, \{-b_n\}),$$

so

$$E_j^-(W(\{a_n\}, \{b_n\})) = -E_j^+(W(\{a_n\}, \{-b_n\})). \quad (2.27)$$

Thus, for example,

$$\begin{aligned} \sum_{j=1}^{N_r} |E_j^-(W)^2 - 4|^{1/2} &\leq \sum_n (-b_n)_+ \\ &= \sum_n (b_n)_-, \end{aligned} \quad (2.28)$$

where  $x_- = (-x)_+ = -\min(0, x)$ , so  $|x| = x_+ + x_-$  and we obtain (1.3) for the case  $a_n \equiv 1$ .

### 3. BOUNDS FOR JACOBI MATRICES

The following elementary observation lets us pass from bounds in case  $a_n \equiv 1$  to the general case. Note that

$$\begin{pmatrix} -|a_n - 1| & 1 \\ 1 & -|a_n - 1| \end{pmatrix} \leq \begin{pmatrix} 0 & a_n \\ a_n & 0 \end{pmatrix} \leq \begin{pmatrix} |a_n - 1| & 1 \\ 1 & |a_n - 1| \end{pmatrix}$$

for any  $a_n$  real since for any  $x$  in  $\mathbb{R}$ ,  $\begin{pmatrix} |x| & x \\ x & |x| \end{pmatrix} \geq 0$  since it has determinant 0 and trace  $2|x| \geq 0$ . This immediately implies by repeated use at each pair of

indices

$$W(\{a_n \equiv 1\}, \{b_n^-\}) \leq W(\{a_n\}, \{b_n\}) \leq W(\{a_n \equiv 1\}, \{b_n^+\}), \quad (3.1)$$

where

$$b_n^\pm = b_n \pm (|a_{n-1} - 1| + |a_n - 1|). \quad (3.2)$$

Equations (3.1) and (2.5) immediately imply

**THEOREM 3.1.** *Let  $f$  be monotone increasing on  $(0, \infty)$  and even. Then*

$$f(E_j^\pm(W(\{a_n\}, \{b_n\}))) \leq f(E_j^\pm(W(\{a_n \equiv 1\}, \{b_n^\pm\}))), \quad (3.3)$$

where  $b_n^\pm$  is given by (3.2).

With this, we can now prove our three main theorems:

*Proof of Theorem 1.* By (3.3), (2.23), and (2.28),

$$\begin{aligned} & \sum_n ([(E_n^+)^2 - 4]^{1/2} + [(E_n^-)^2 - 4]^{1/2}) \\ & \leq \sum_n ([b_n + |a_{n-1} - 1| + |a_n - 1|]_+ + [b_n - |a_{n-1} - 1| - |a_n - 1|]_-) \\ & \leq \sum_n ([b_n]_+ + [b_n]_-) + 4 \sum_n |a_n - 1|. \end{aligned} \quad (3.4)$$

In obtaining (3.4), we used  $[x + y]_+ \leq x_+ + y_+$  and  $[x + y]_- \leq x_- + y_-$  and that a given  $|a_n - 1|$  occurs in four terms with  $[b_n]_\pm$  and  $[b_{n+1}]_\pm$ . ■

*Proof of Theorem 2.* By (3.3), (2.21), and (2.27),

$$\begin{aligned} \sum_n (|E_n^+ - 2|^p + \sum_n |E_n^- + 2|^p) & \leq d_p \sum_n ([b_n + |a_{n-1} - 1| + |a_n - 1|]_+^{p+1/2} \\ & \quad + [b_n - |a_{n-1} - 1| - |a_n - 1|]_-^{p+1/2}), \end{aligned} \quad (3.5)$$

where

$$d_p = \frac{1}{2} \frac{\Gamma(p+1) \Gamma(2)}{\Gamma(p + \frac{3}{2}) \Gamma(\frac{3}{2})}.$$

Now for any  $q \geq 1$  ( $q$  will be  $p + \frac{1}{2}$ ),  $x^q$  is convex, so

$$\begin{aligned} (\alpha + \beta + \gamma)^q &= 3^q \left( \frac{\alpha}{3} + \frac{\beta}{3} + \frac{\gamma}{3} \right)^q \\ &\leq 3^{q-1} [\alpha^q + \beta^q + \gamma^q], \end{aligned}$$

from which (1.6) holds if we note that  $c_p = 3^{(p+1/2)-1} d_p$ . ■

*Proof of Theorem 4.* As stated, (1.8) is an immediate consequence of Theorem 2.10 and (3.1).

We kept this bound in form (1.8) to get an exact result as  $\lambda \rightarrow \infty$ . We could use the same method of proof of Theorem 2 to get

$$\sum_n |E_n^+ - 2|^p + \sum_n |E_n^- + 2|^p \leq 3^{p-1} \left[ \sum_n |b_n|^p + 4 \sum_n |a_n - 1|^p \right]. \quad \blacksquare$$

We believe it could be true that (1.3) holds with  $|a_n - 1|$  replaced by  $(a_n - 1)_+$  and, in particular, we know that (1.6) and (1.8) hold when  $p \geq 1$  if  $|a_n - 1|$  is replaced by  $(a_n - 1)_+$ . To see the latter, we note that—by a convexity plus evenness argument much like that in the proof of Proposition 2.7— $\sum_{j=1}^k E_j^+(W(\{a_n\}, \{b_n\}))$  is monotone in  $a_n$  in the region  $a_n \geq 0$ . Thus for  $p = 1$ , (1.6) and (1.8) hold with  $(a_n - 1)_+$  for we move those  $a$ 's with  $a_n > 1$  to the diagonal as we did in (3.1), and use the monotonicity just noted to move  $a_n$ 's in  $(0, 1)$  up to 1. Once one has the result for  $p = 1$ , it follows for  $p \geq 1$  by the Aizenman–Lieb argument.

The fact that in (1.6) for  $p \geq 1$  and in the Bargmann bound of Appendix A, one can take  $(a_n - 1)_+$  leads us to conjecture (1.3) holds with  $(a_n - 1)_+$  rather than  $|a_n - 1|$ .

#### 4. EXAMPLES

**EXAMPLE 4.1.**  $W$  has all  $a_n = 1$ , all  $n$ , and all  $b_n = 0$  for  $n \neq 0$ . If  $b_0 \equiv b > 0$ , then there is an eigenvalue at energy  $E = \mu + \mu^{-1}$  with  $\mu < 1$  and eigenfunction  $\varphi_n = \mu^{|n|}$ . To have the eigenfunction fit at  $n = 0$ , we need

$$2\mu + b = E$$

or

$$b = \mu^{-1} - \mu = \sqrt{E^2 - 4}.$$

This example has equality in (1.3) for all values of  $b > 0$  (and also  $b < 0$  it turns out) and shows one cannot decrease the value 1 in front on  $\sum |b_n|$ .



EXAMPLE 4.2.  $W$  has all  $b_n = 0$ , all  $n$ , and all  $a_n = 1$ ,  $n \neq 0$ . If  $a_0 \equiv a > 1$ , there is an eigenvalue at energy  $E = \mu + \mu^{-1}$  with  $0 < \mu < 1$ . Then  $\varphi_n = \mu^{-n}$  for  $n \leq 0$  and  $\varphi_n = \mu^{n-1}$  for  $n \geq 1$  since  $\varphi$  must be symmetric around  $n = \frac{1}{2}$ . The eigenfunction condition at 0 reads

$$\mu + a = \mu + \mu^{-1}$$

or  $a = \mu^{-1}$ . Thus,

$$a - a^{-1} = \sqrt{E^2 - 4}.$$

There is a second eigenvalue at energy  $-E$  (there has to be by the symmetry (2.27)). Thus,

$$\begin{aligned} \text{LHS of (1.3)} &= 2(a - a^{-1}) \\ &= 2(1 + a^{-1})(a - 1). \end{aligned}$$

The two sides of (1.3) are not equal for any  $a$ , but the ratio goes to 1 as  $a \downarrow 1$  since  $2(1 + a^{-1}) \uparrow 4$ . Thus, the 4 in front of the  $|a - 1|$  cannot be made smaller. However, both this example and the discussion in Appendix A suggest it might be possible to replace  $|a - 1|$  by  $(a - 1)_+$ .

As noted above, the best constant for the  $W$  case is the same as for the  $J$  case.

EXAMPLE 4.3. (Proof of Theorem 3). Shift to the Jacobi case. Take an example with  $a_n \equiv 1$  and  $b_n = 0$ , except for  $n = m, 2m, \dots, Nm$  where  $b_n = \beta$ . As  $m \rightarrow \infty$ , there are  $n$  eigenvalues above 2 which all approach the solution of  $\sqrt{E^2 - 4} = \beta$ . So long as  $\beta < 1$ ,  $|E_n - 2| \geq \frac{1}{6}\beta^2$ , so

$$\sum_n |E_n - 2|^p \geq N \left( \frac{\beta^2}{6} \right)^p. \quad (4.1)$$

In the translation invariant norm  $\|\cdot\|$ , let  $\alpha = \|(a_n \equiv 1, b_1 = 1, b_n = 0 \text{ for } n \neq 1)\|$ . Then for the  $(a, b)$  of this  $\beta, N, m$  example,

$$\|(a, b)\| \leq N\alpha\beta. \quad (4.2)$$

Let  $N_0(\varepsilon)$ ,  $\beta_0(\varepsilon)$  solve

$$\begin{aligned} N \left( \frac{\beta}{6} \right)^p &= 2\varepsilon^{-1}, \\ N\alpha\beta &= \frac{\varepsilon}{2}, \end{aligned}$$

so

$$\beta = c_1 \varepsilon^{2/1-2p} \rightarrow 0,$$

$$N = c_2 \varepsilon^{-(1+2p)/(1-2p)} \rightarrow \infty,$$

since  $p < \frac{1}{2}$ . Increase  $N$  slightly to be an integer. Thus,

$$\sum_n |E_n - 2|^p \geq \varepsilon^{-1}, \quad \|(a, b)\| \leq \varepsilon,$$

proving Theorem 3.

## 5. BOUNDS IN HIGHER DIMENSION

In this section, we want to use the ideas of Laptev–Weidl [6] to prove bounds on operators on  $\ell^2(\mathbb{Z}^v)$ . We begin with the discrete Schrödinger operator case. Let  $H_0$  be defined on  $\ell^2(\mathbb{Z}^v)$  by

$$(H_0 u)(n) = \sum_{|m-n|=1} u(m)$$

and

$$(Vu)(n) = V(n)u(n).$$

LEMMA 5.1. *Let  $W_0$  act on  $\ell^2(\mathbb{Z}; X)$  where  $X$  is a Hilbert space, and let  $B(n) : X \rightarrow X$  be self-adjoint and trace class with  $\sum_n \text{Tr}(|B(n)|) < \infty$ . Then,*

$$\sum_j (E_j^\pm(W_0 + B)^2 - 4)^{1/2} \leq \text{Tr}_X(B^\pm), \quad (5.1)$$

where  $B^\pm(n) = \max(\pm B(n), 0)$  is defined via the functional calculus.

*Proof.* Suppose  $B(n) \geq 0$ . As with (2.14), define  $L_\mu : \ell^2(\mathbb{Z}; X) \rightarrow \ell^2(\mathbb{Z}; X)$  by

$$(L_\mu)_{mn} = B_n^{1/2} \mu^{|n-m|} B_m^{1/2}.$$

As with Proposition 2.7,  $0 < \mu < \eta \leq 1$  implies

$$S_n^+(L_\mu) \leq S_n^+(L_\eta)$$

and then the proof of (2.16) extends. ■

**THEOREM 5.2.** *If  $V \in L^p(\mathbb{Z}^v; X)$  for  $p \geq 1$  where  $X$  is a Hilbert space, that is,  $V(x): X \rightarrow X$  is a symmetric compact operator such that  $\sum_{x \in \mathbb{Z}^v} \text{Tr}_X |V(x)|^p < \infty$ , then*

$$\sum_j |E_j^+(H_0 + V) - 2v|^p + \sum_j |E_j^-(H_0 + V) + 2v|^p \leq \sum_{x \in \mathbb{Z}^v} \text{Tr}_X |V(x)|^p. \quad (5.2)$$

*Proof.* By the Aizenman–Lieb idea, (2.24), it suffices to prove this for  $p = 1$ . As usual, we can suppose  $V \geq 0$  and prove the result for  $E_j^+$ . Write

$$H_0 = H_{0,1} + H_{0,\{2,\dots,v\}},$$

where  $H_{0,1}$  involves neighbors in one direction and  $H_{0,\{2,\dots,v\}}$  neighbors in the other directions. Note that

$$\begin{aligned} & (H_{0,1} + H_{0,\{2,\dots,v\}} + V - 2v)_+ \\ & \leq (H_{0,1} + (H_{0,\{2,\dots,v\}} + V - 2(v-1))_+ - 2)_+ \end{aligned} \quad (5.3)$$

and thus

$$\begin{aligned} \sum_j |E_j^+(H_0 + V) - 2v| &= \text{Tr}_{\ell^2(\mathbb{Z}^v; X)}((H_0 + V - 2v)_+) \\ &\leq \text{Tr}_{\ell^2(\mathbb{Z}; \ell^2(\mathbb{Z}^{v-1}; X))}((H_{0,1} + (H_{0,\{2,\dots,v\}} + V - 2(v-1))_+ - 2)_+) \\ &\leq \sum_{n_1} \text{Tr}_{\ell^2(\mathbb{Z}^{v-1}; X)}((H_{0,\{2,\dots,v\}} + V(n_1, \cdot) - 2v + 2)_+) \end{aligned}$$

by (5.1) and  $(E^2 - 4)^{1/2} \geq (|E| - 2)$ . An inductive argument completes the proof. ■

For the other moment result, it will be convenient to phrase things in terms of the classical constants,

$$\begin{aligned} L_{p,v}^{cl} &= (2\pi)^{-v/2} \int_{|k| \leq 1} |k|^{2p} d^v k \\ &= 2^{-v} \pi^{-v/2} \frac{\Gamma(p+1)}{\Gamma(p+1 + \frac{v}{2})}. \end{aligned} \quad (5.4)$$

These constants have several important features. First, the argument that led to (2.25) says that if

$$\sum_{j=1}^{N_+} |E_j^+ - 2|^p \leq \alpha L_{p,v}^{cl} \sum_n |b_n|^{p+v/2} \quad (5.5)$$

for some  $p = p_0$ , it holds for all  $p > p_0$ . Second,

$$L_{p=1/2, v=1} = 2^{-1}\pi^{-1/2} \left[ \frac{\frac{1}{2}\sqrt{\pi}}{1} \right] = \frac{1}{4},$$

so the consequence of (5.1) and  $(E^2 - 4)^{1/2} \geq 2(|E| + 2)^{1/2}$  is that (5.5) holds for  $v = 1$ ,  $p = \frac{1}{2}$ , and  $\alpha = 2$ .

Finally, we note that from (5.4) and Fubini, we have

$$L_{p,v}^{c\ell} = \prod_{j=0}^{v-1} L_{p+j/2,1}^{c\ell}. \tag{5.6}$$

**THEOREM 5.3.** *Let  $V \in L^{p+v/2}(\mathbb{Z}^v; X)$  for  $p \geq 1$ . Then*

$$\begin{aligned} & \sum_j |E_j^+(H_0 + V) - 2v|^p + \sum_j |E_j^-(H_0 + V) + 2v|^p \\ & \leq 2^v L_{p,v}^{c\ell} \sum_{x \in \mathbb{Z}^v} \text{Tr}_X |V(x)|^{p+v/2}. \end{aligned} \tag{5.7}$$

*Proof.* We exploit (5.3), but use (5.5) for  $\alpha = 2$ ,  $v = 1$ ,  $p \geq \frac{1}{2}$  at each stage of the induction. We then get (5.7) with a constant

$$\prod_{j=0}^{v-1} 2L_{p+1/2,1}^{c\ell} = 2^v L_{p,v}^{c\ell}$$

by (5.6). ■

As in the one-dimensional case, Theorem 5.2 is better for large coupling. Indeed, it is exact in the large coupling regime, while Theorem 5.3 gives more information on the eigenvalues very close to  $\pm 2v$  in the regime of slow decay of  $V(n)$  at infinity.

As with the one-dimensional case, we can handle nonconstant off-diagonal terms which approach 1 fast enough at infinity. Let  $B(\mathbb{Z}^v)$  be set of bonds in  $\mathbb{Z}^v$ , that is, the set of unordered pairs  $b = (ij)$  with  $i, j \in \mathbb{Z}^v$ ,  $|i - j| = 1$ . Given  $\{a_b\}_{b \in B(\mathbb{Z}^v)}$ , a nonnegative real number  $a_b$  for each bond  $b = (ij)$ , one can define

$$(H_0(a_b)u)(n) = \sum_{|m-n|=1} a_{(nm)}u(m). \tag{5.8}$$

The analog of (3.3) is then

$$H_0 + V^- \leq H_0(a_b) + V \leq H_0 + V^+, \tag{5.9}$$

where

$$V^\pm(n) = V(n) \pm \sum_{|m-n|=1} |a_{(mn)} - 1|, \quad (5.10)$$

so, for example, we get

$$\sum_j [E_j^+(H_0(a_b) + V) + E_j^-(H_0(a_b) + V)] \leq \sum_n |V(n)| + \sum_b 4|a_b - 1|. \quad (5.11)$$

*Remark.* Since the bound in Theorem 1 is optimal both for large and small couplings, the curious reader might wonder whether it is possible to keep some of its structure also in higher dimension. This is indeed the case. For constant diagonal terms and scalar potential, we have the two bounds

$$\sum_{n=1, \dots, N_+} [(E_n^+)^2 - 4]^{1/2} + \sum_{n=1, \dots, N_-} [(E_n^-)^2 - 4]^{1/2} \leq \sum_{x \in \mathbb{Z}^v} |V(x)|$$

and

$$\sum_{n=1, \dots, N_+} [(E_n^+)^2 - 4]^{1/2} + \sum_{n=1, \dots, N_-} [(E_n^-)^2 - 4]^{1/2} \leq 2^{v-1} L_{1, v-1}^{cl} \sum_{x \in \mathbb{Z}^v} |V(x)|^{1+(v-1)/2}.$$

Simply use the induction in the dimension idea to strip off the first coordinate  $x_1$  and then use either Theorem 5.2 or 5.3 in  $v - 1$  dimension. Of course, the above extension to nonconstant diagonal terms also applies.

## ACKNOWLEDGMENTS

We thank Jeff Geronimo, Fritz Gesztesy, Rowan Killip, and Paul Nevai for useful comments.

## APPENDIX A. THE BARGMANN BOUND

Our goal in this appendix is to prove

**THEOREM A.1.** *Let  $N(\{a\}, \{b\})$  be the number of eigenvalues of  $J(\{a\}, \{b\})$  outside  $[-2, 2]$ . Then*

$$N(\{a\}, \{b\}) \leq \sum_{n=1}^{\infty} (n|b_n| + (4n + 2)(a_n - 1)_+), \quad (A.1)$$

where  $(x)_+ = \max(x, 0)$ .

This is related to a result of Geronimo [8, 9]. We provide a proof here because it is easy from our machinery earlier. One of Geronimo’s proofs of this result, Theorem III.1 in [9], has an error in the argument that allows for  $a_n < 1$ , since his Lemma III.1 is wrong. Earlier papers that show  $N < \infty$  if (A.1) holds include Geronimo–Case [10] and Chihara–Nevai [6].

*Note.* (1) If you translate Geronimo’s result in [8] into our normalization (he has  $J_0$  with  $a \equiv \frac{1}{2}$ , not  $a = 1$ ), then where we have  $(4n + 2)(a_n - 1)_+$ , he has  $(4n + 4)(a_n - 1)_+(a_n + 1)$ , which is weaker in two regards:  $4n + 2 < 4n + 4$  and we have no  $a_n + 1$ . We note that by looking at  $b_n = 0$  and  $a_n = 1$  for  $n \geq 2$ , one finds examples with  $N = 2$  and  $(a_1 - 1)_+$  arbitrarily close to  $\sqrt{2} - 1$  so that constant in front of  $(a_1 - 1)_+$  must be at least  $2(\sqrt{2} + 1)$  and, in particular,  $4n$  does not work.

(2) We actually have separate inequalities for  $N_+$  and  $N_-$ .

*Step 1:*  $a_n \equiv 1$ ;  $b_n \geq 0$ . The proof of Bargmann’s bound [2] given by Birman [3] and Schwinger [25] works in this case. By (2.10) and the monotonicity with  $A = J_0$ ,  $B = J - J_0$ , for  $\beta > 2$ ,

$$\begin{aligned} \# \text{ of eigenvalues of } A + B &\geq \beta \\ &= \# \text{ of } \beta' \geq \beta \text{ so that } K_{\beta'} \text{ has eigenvalue } = 1 \\ &= \# \text{ of eigenvalues of } K_\beta \geq 1 \end{aligned} \tag{A.2}$$

$$\leq \text{Tr}(K_\beta) \tag{A.3}$$

$$\leq \text{Tr}(K_2), \tag{A.4}$$

where (A.2) follows from the fact that  $\|K_\beta\| \downarrow 0$  as  $\beta \rightarrow \infty$  and the strict monotonicity of the eigenvalues of  $K_\beta$  noted in the proof of Proposition 2.5. Equation (A.3) holds since

$$\text{Tr}(K_\beta) = \sum_{E_j^+(K_\beta)} E_j^+ \geq \sum_{E_j^+(K_\beta) \geq 1} E_j^+ \geq (\# \text{ of eigenvalues of } K_\beta \geq 1)$$

since  $K_\beta > 0$ . Equation (A.4) holds since  $K_\beta \leq K_2$ .

The same argument that led to Proposition 2.6 shows that  $[(\beta - J_0)^{-1}]_{nm} = w(\beta)^{-1} \varphi_-^{(\beta)}(\min(n, m)) \varphi_+^{(\beta)}(\max(n, m))$  where  $\varphi_\pm$  solve  $J_0 \varphi = \beta \psi$  with  $\varphi_+ \in L^2$  at infinity,  $\varphi_-(0) = 0$ , and  $w$  is their Wronskian. As  $\beta \downarrow 2$ ,  $\varphi_+(n) \rightarrow 1$ ,  $\varphi_-(n) \rightarrow n$ , and their Wronskian is 1 so

$$(K_2)_{nm} = \min(n, m) b_n^{1/2} b_m^{1/2}$$

and

$$\text{Tr}(K_2) = \sum_{n=1}^{\infty} n b_n,$$

proving (A.1) in this case.

*Step 2:*  $a_n \leq 1$ ;  $b_n \geq 0$ . Let  $J_0(\{a_n\})$  be  $J$  with  $b_n = 0$ . We claim if  $a_n \leq 1$  and  $\beta > 2$ , then

$$(\beta - J_0(\{a_n\}))_{nm}^{-1} \leq (\beta - J_0)_{nm}^{-1}. \tag{A.5}$$

This is a simple maximal principle argument. One first notes that if  $f_n(m) = (\beta - J_0(\{a_n\}))_{nm}^{-1}$ , then  $f_n(m) > 0$  (expand  $(1 - \beta^{-1}J)$  in a geometric series). Next, one notes that

$$\begin{aligned} ((\beta - J_0)f_n)(m) &= \delta_{nm} - (1 - a_{m-1})f_n(m - 1) - (1 - a_m)f_n(m) \\ &\leq \delta_{nm}. \end{aligned}$$

Since  $(\beta - J_0)^{-1}$  also has a positive matrix, applying it preserves pointwise matrix inequalities, so

$$f_n(m) \leq [(\beta - J_0)^{-1} \delta_n]_m = (\beta - J_0)_{nm}^{-1},$$

proving (A.5).

Now (A.5) shows the Birman–Schwinger kernel for  $J_0\{a_n\}$  and  $J(\{a_n, b_n\})$  is dominated (in the sense of inequalities on matrix elements) by this for  $J_0$  and  $J(\{a_n \equiv 1, b_n\})$ , so Step 1 implies

$$\text{Tr}(K_2(\{a_n, b_n\})) \leq \text{Tr}(K_2(\{a_n \equiv 1, b_n\})) = \sum_{j=1}^{\infty} nb_n.$$

Note we do not have an *operator* inequality of the form  $(\beta - J_0(\{a_n\}))^{-1} \leq (\beta - J_0)^{-1}$ , so individual eigenvalues may not have an inequality (this is Geronimo’s error in [9]).

*Step 3:* Adding  $b$ ’s of both signs. Fix  $a_n$  with  $0 < a_n \leq 1$ . Let  $J(\{b_n\})$  be the Jacobi matrix with  $b_n$  along the diagonal and  $N_{\pm}(\{b_n\})$  the number of eigenvalues  $E$  with  $\pm E > 2$ . Since  $J(\{-(b_n)_-\}) \leq J(\{b_n\}) \leq J(\{(b_n)_+\})$ , we have

$$N_{\pm}(\{b_n\}) \leq N_{\pm}(\{\pm(b_n)_{\pm}\}),$$

so by (A.1) for  $b_n \geq 0$  and (2.27), we have (A.1) for the case  $0 \leq a_n \leq 1$ .

*Step 4 (General Case):* Now use the idea at the start of Section 3 but only for  $a_n$ ’s with  $a_n > 1$ . Then (A.1) holds in general since this idea reduces to the case  $a_n \leq 1$ . We use here that

$$2n(a_n - 1)_+ + 2(n + 1)(a_n - 1)_+ = (4n + 2)(a_n - 1)_+.$$

## REFERENCES

1. M. Aizenman and E. H. Lieb, On semi-classical bounds for eigenvalues of Schrödinger operators, *Phys. Lett.* **66A** (1978), 427–429.
2. V. Bargmann, On the number of bound states in a central field of force, *Proc. Nat. Acad. Sci. U.S.A.* **38** (1952), 961–966.
3. M. S. Birman, The spectrum of singular boundary problems, *Mat. Sb. (N.S.)* **55(97)** (1961), 125–174 [in Russian]; Translated in *Amer. Math. Soc. Trans.* **53** (1966), 23–80.
4. K. M. Case, Orthogonal polynomials from the viewpoint of scattering theory, *J. Math. Phys.* **15** (1974), 2166–2174.
5. K. M. Case, Orthogonal polynomials. II, *J. Math. Phys.* **16** (1975), 1435–1440.
6. T. S. Chihara and P. Nevai, Orthogonal polynomials and measures with finitely many point masses, *J. Approx. Theory* **35** (1982), 370–380.
7. M. Cwikel, Weak type estimates for singular values and the number of bound states of Schrödinger operators, *Ann. Math.* **106** (1977), 93–100.
8. J. S. Geronimo, An upper bound on the number of eigenvalues of an infinite dimensional Jacobi matrix, *J. Math. Phys.* **23** (1982), 917–921.
9. J. S. Geronimo, On the spectra of infinite-dimensional Jacobi matrices, *J. Approx. Theory* **53** (1988), 251–265.
10. J. S. Geronimo and K. N. Case, Scattering theory and polynomials orthogonal on the real line, *Trans. Amer. Math. Soc.* **258** (1980), 467–494.
11. D. Hundertmark, A. Laptev, and T. Weidl, New bounds on the Lieb–Thirring constants, *Invent. Math.* **140** (2000), 693–704.
12. D. Hundertmark, E. H. Lieb, and L. E. Thomas, A sharp bound for an eigenvalue moment of the one-dimensional Schrödinger operator, *Adv. Theor. Math. Phys.* **2** (1998), 719–731.
13. T. Kato, “Perturbation Theory for Linear Operators,” Springer-Verlag, New York, 1966.
14. R. Killip and B. Simon, Sum rules for Jacobi matrices and their applications to spectral theory, *Ann. Math.*, to appear.
15. L. D. Landau and E. M. Lifshitz, “Quantum Mechanics. Non-relativistic Theory,” Course of Theoretical Physics, Vol. 3, Pergamon Press, London, 1958.
16. A. Laptev and T. Weidl, Sharp Lieb–Thirring inequalities in high dimensions, *Acta Math.* **184** (2000), 87–111.
17. E. H. Lieb, Bounds on the eigenvalues of the Laplace and Schrödinger operators, *Bull. Amer. Math. Soc.* **82** (1976), 751–753.
18. E. H. Lieb and W. Thirring, Bound for the kinetic energy of fermions which proves the stability of matter, *Phys. Rev. Lett.* **35** (1975), 687–689; Errata **35** (1975), 1116.
19. E. H. Lieb and W. Thirring, Inequalities for the moments of the eigenvalues of the Schrödinger Hamiltonian and their relation to Sobolev inequalities, in “Studies in Mathematical Physics. Essays in Honor of Valentine Bargmann” (E. Lieb, B. Simon, and A. Wightman, Eds.), pp. 269–303, Princeton Univ. Press, Princeton, NJ, 1976.
20. P. Nevai, Orthogonal polynomials, recurrences, Jacobi matrices, and measures, in “Progress in Approximation Theory,” (A.A. Gonchar and E.B. Saff, Eds.), Tampa, FL, 1990, pp. 79–104, Springer Series Computational Mathematics, Vol. 19, Springer, New York, 1992.
21. P. Nevai, Research problems in orthogonal polynomials, in “Approximation Theory VI,” Vol. II, College Station, TX, 1989, pp. 449–489, Academic Press, Boston, 1989.
22. F. Peherstorfer and P. Yuditskii, Asymptotics of orthonormal polynomials in the presence of a denumerable set of mass points, *Proc. Amer. Math. Soc.* **120** (2001), 3213–3220.



23. M. Reed and B. Simon, "Methods of Modern Mathematical Physics, IV. Analysis of Operators," Academic Press, New York, 1978.
24. G. V. Rozenblum, Distribution of the discrete spectrum of singular differential operators, *Dokl. Akad. Nauk. SSSR* **202** (1972), 1012–1015; *Izv. Vyssh. Uchebn. Zaved. Mat.* **1** (1976), 75–86.
25. J. Schwinger, On the bound states for a given potential, *Proc. Nat. Acad. Sci. U.S.A.* **47** (1961), 122–129.
26. B. Simon, On the number of bound states of two-body Schrödinger operators: A review, in "Studies in Mathematical Physics. Essays in Honor of Valentine Bargmann," (E. Lieb, B. Simon, and A. Wightman, Eds.), pp. 305–326, Princeton Univ. Press, Princeton, NJ, 1976.
27. T. Weidl, On the Lieb–Thirring constants  $L_{\gamma,1}$  for  $\gamma \geq 1/2$ , *Comm. Math. Phys.* **178** (1996), 135–146.