



ACADEMIC
PRESS

Available at
WWW.MATHEMATICSWEB.ORG
POWERED BY SCIENCE @ DIRECT®

Journal of Approximation Theory 121 (2003) 357–364

JOURNAL OF
**Approximation
Theory**

<http://www.elsevier.com/locate/jat>

Zeros of orthogonal polynomials on the real line

Sergey A. Denisov and Barry Simon^{*,1}

Mathematics 253-37, California Institute of Technology, Pasadena, CA 91125, USA

Received 9 July 2002; accepted 7 February 2003

Communicated by Leonid Golinskii

Abstract

Let $p_n(x)$ be the orthonormal polynomials associated to a measure $d\mu$ of compact support in \mathbb{R} . If $E \notin \text{supp}(d\mu)$, we show there is a $\delta > 0$ so that for all n , either p_n or p_{n+1} has no zeros in $(E - \delta, E + \delta)$. If E is an isolated point of $\text{supp}(\mu)$, we show there is a δ so that for all n , either p_n or p_{n+1} has at most one zero in $(E - \delta, E + \delta)$. We provide an example where the zeros of p_n are dense in a gap of $\text{supp}(d\mu)$.

© 2003 Elsevier Science (USA). All rights reserved.

1. Introduction

Let $d\mu$ be a measure on \mathbb{R} whose support is not a finite number of points and with $\int |x|^n d\mu(x) < \infty$ for all $n = 0, 1, 2, \dots$. The orthonormal polynomials $p_n(x; d\mu)$ or $p_n(x)$ are determined uniquely by

$$p_n(x) = \gamma_n x^n + \text{lower order}, \quad \gamma_n > 0, \tag{1.1}$$

$$\int p_n(x)p_m(x) d\mu(x) = \delta_{nm}. \tag{1.2}$$

There are $a_n > 0$, $b_n \in \mathbb{R}$ for $n \geq 1$ so that

$$xp_n(x) = a_{n+1}p_{n+1}(x) + b_{n+1}p_n(x) + a_n p_{n-1}(x) \tag{1.3}$$

(many works use a_{n-1} , b_{n-1} , where we use a_n , b_n).

*Corresponding author.

E-mail addresses: denissov@its.caltech.edu (S.A. Denisov), bsimon@caltech.edu (B. Simon).

¹Supported in part by NSF Grants DMS-9707661, DMS-0140592.

In this paper, we will be interested in the zeros of $p_n(x; d\mu)$. The following results are classical (see, e.g., Freud's book [4]):

- (1) The zeros of $p_n(x)$ are real and simple.
- (2) If $(a, b) \cap \text{supp}(d\mu) = \emptyset$, then if $a = -\infty$ or $b = +\infty$, p_n has no zeros in (a, b) and, in any event, (a, b) has at most one zero of $p_n(x)$.
- (3) In the determinate case, if $x_0 \in \text{supp}(d\mu)$ and $\delta > 0$, for all large n , $p_n(x)$ has a zero in $(x_0 - \delta, x_0 + \delta)$.

Define

$$N_n(x_0, \delta) = \# \text{ of zeros of } p_n(x) \text{ in } (x_0 - \delta, x_0 + \delta).$$

Then (1)–(3) immediately imply:

- (i) If x_0 is a non-isolated point of $\text{supp}(d\mu)$, then for any $\delta > 0$, $\lim_{n \rightarrow \infty} N_n(x_0, \delta) = \infty$.
- (ii) If x_0 is an isolated point of $\text{supp}(d\mu)$ and $\delta = \text{dist}(x_0, \text{supp}(d\mu) \setminus \{x_0\})$, then $N_n(x_0, \delta)$ is never more than 2, and for all $\delta > 0$ and n large, $N_n(x_0, \delta) \geq 1$.
- (iii) If $x_0 \notin \text{supp}(d\mu)$ and $\delta = \text{dist}(x_0, \text{supp}(d\mu))$, then $N_n(x_0, \delta)$ is never more than 1.

(i) is fairly complete, but (ii), (iii) leave open how often there is one vs. two points in case (ii) and zero vs. one in case (iii). One might guess that a zero near $x_0 \notin \text{supp}(d\mu)$ and two zeros near an isolated x_0 in $\text{supp}(d\mu)$ are not too common occurrences.

Example. If $d\mu$ is even about $x = 0$, then $p_n(-x) = (-1)^n p_n(x)$. Thus, if n is odd, $p_n(0) = 0$. So if $0 \notin \text{supp}(d\mu)$, we still have $N_n(0, \delta) = 1$ for all small δ and n odd. If zero is an isolated point of $d\mu$, p_n for n even has a zero at x_n near 0, but not equal to 0 (since zeros are simple), so also at $-x_n$, that is, $N_n(0, \delta) = 2$ for δ small and n even. So “not too common” can be as often as 50% of the time. Our goal here is to show this 50% is a maximal value.

It is surprising that there do not seem to be any results on these issues until a recent paper of Ambroladze [1], who proved

Theorem (Ambroladze [1]). *If $\text{supp}(d\mu)$ is bounded and $x_0 \notin \text{supp}(d\mu)$, then for some $\delta > 0$, $\liminf_{n \rightarrow \infty} N_n(x_0, \delta) = 0$.*

Thus, we can use $N_n(x_0, \delta)$ to distinguish when $x_0 \in \text{supp}(d\mu)$. Our goal in this paper is to prove

Theorem 1.1. *Let $d = \text{dist}(x_0, \text{supp}(d\mu)) > 0$. Let $\delta_n = d^2 / (d + \sqrt{2} a_{n+1})$ (where a_n is the recursion coefficient given by (1.3)). Then either p_n or p_{n+1} (or both) has no zeros in*

$(x_0 - \delta_n, x_0 + \delta_n)$. In particular, if $a_\infty = \sup_n a_n < \infty$ and $\delta_\infty = d^2 / (d + \sqrt{2}a_\infty)$, then $(x_0 - \delta_\infty, x_0 + \delta_\infty)$ does not have zeros of p_j for two successive values of j .

Theorem 1.2. Let x_0 be an isolated point of $\text{supp}(d\mu)$. Then there exists a $d_0 > 0$, so that if $\delta_n = d_0^2 / (d_0 + \sqrt{2} a_{n+1})$, then at least one of p_n and p_{n+1} has no zeros or one zero in $(x_0 - \delta_n, x_0 + \delta_n)$. In particular, if $a_\infty = \sup_n a_n < \infty$ and $\delta_\infty = d_0^2 / (d_0 + \sqrt{2}a_\infty)$, then for all large n , either $N_n(x_0, \delta_\infty) = 1$ or $N_{n+1}(x_0, \delta_\infty) = 1$.

We will prove Theorem 1.1 in Section 2 and Theorem 1.2 in Section 3. In Section 4, we present an example of a set of polynomials whose zeros are dense in a gap of the spectrum.

It is a pleasure to thank Leonid Golinskii and Paul Nevai for useful correspondence.

2. Points outside the support of $d\mu$

We arrived at the following lemma by trying to abstract the essence of Ambroladze’s argument [1]; it holds for orthogonal polynomials on the complex plane. Let $d\mu$ be a measure on \mathbb{C} with finite moments and infinite support, and let $p_n(z; d\mu)$ be the orthonormal polynomials. Define the reproducing kernel

$$K_n(z, w) = \sum_{j=0}^n p_j(z)\overline{p_j(w)}, \tag{2.1}$$

so in $L^2(\mathbb{C}, d\mu)$, for any polynomial π of degree n or less,

$$\int K_n(z, w)\pi(w) d\mu(w) = \pi(z). \tag{2.2}$$

Lemma 2.1. Suppose $z_0 \in \mathbb{C}$, $p_j(w) = 0$ for some $j \leq n + 1$. Then

$$|z_0 - w| \geq \frac{|p_j(z_0)|}{K_n(z_0, z_0)^{1/2}} \text{dist}(w, \text{supp}(d\mu)). \tag{2.3}$$

Proof. Let $q(z) = p_j(z)/(z - w)$, which has $\text{deg}(q) \leq n$. Thus, by (2.2), $\langle K(\cdot, z_0), q(\cdot) \rangle = q(z_0)$ so, by the Schwarz inequality,

$$\frac{|p_j(z_0)|}{|z_0 - w|} \leq \|q\| \|K(\cdot, z_0)\|.$$

By (2.2), $\|K(\cdot, z_0)\| = K(z_0, z_0)^{1/2}$ and clearly, $\|q\| \leq \text{dist}(w, \text{supp}(d\mu))^{-1} \|p_j\| = \text{dist}(w, \text{supp}(d\mu))^{-1}$. This yields (2.3). \square

The following only holds in the real case:

Lemma 2.2. For any $x \in \mathbb{R}$ and n ,

$$K_n(x, x) \text{dist}(x, \text{supp}(d\mu))^2 \leq a_{n+1}^2 [p_{n+1}^2(x) + p_n^2(x)]. \tag{2.4}$$

Proof. The Christoffel–Darboux formula [4] says

$$K_n(x, y) = a_{n+1} \left[\frac{p_{n+1}(x)p_n(y) - p_{n+1}(y)p_n(x)}{x - y} \right],$$

so since $\langle p_j, p_k \rangle = \delta_{jk}$,

$$\|(x - \cdot)K_n(x, \cdot)\|^2 = |a_{n+1}|^2 [p_{n+1}^2(x) + p_n^2(x)]. \tag{2.5}$$

Clearly,

$$\|(x - \cdot)K_n(x, \cdot)\|^2 \geq \text{dist}(x, \text{supp}(d\mu))^2 \|K_n(x, \cdot)\|^2 \tag{2.6}$$

and, as above, $\|K_n(x, \cdot)\|^2 = K_n(x, x)$, which yields (2.4). \square

Remark. An alternate way of seeing (2.5) is to let ψ be the trial vector $(p_0(x), \dots, p_n(x), 0, 0, \dots)$ and note that in terms of the standard Jacobi matrix $((J - x)\psi)_j = 0$ unless $j = n, n + 1$, in which case the values are $-a_{n+1}p_{n+1}(x)$ and $a_{n+1}p_n(x)$, (2.6) is then just $\|(J - x)\psi\| \geq \text{dist}(x, \text{supp}(d\mu))\|\psi\|$.

Proof of Theorem 1.1. By (2.4), we have that

$$K_n(x_0, x_0) \text{dist}(x_0, \text{supp}(d\mu))^2 \leq 2a_{n+1}^2 p_{n+1}^2(x_0) \tag{2.7}$$

and/or

$$K_n(x_0, x_0) \text{dist}(x_0, \text{supp}(d\mu))^2 \leq 2a_{n+1}^2 p_n^2(x_0). \tag{2.8}$$

Suppose (2.7) holds. Then, by (2.3), if w is a zero of $p_{n+1}(x)$ and if $d = \text{dist}(x_0, \text{supp}(d\mu))$,

$$\begin{aligned} |x_0 - w| &\geq \frac{1}{\sqrt{2}} \frac{1}{a_{n+1}} d \text{dist}(w, \text{supp}(d\mu)) \\ &\geq \frac{1}{\sqrt{2}} \frac{1}{a_{n+1}} d(d - |w - x_0|) \end{aligned}$$

which leads directly to $|x_0 - w| \geq d^2 / (d + a_{n+1}\sqrt{2})$. \square

Remark. There is also a Christoffel–Darboux result for polynomials on the unit circle $\partial D = \{z \mid |z| = 1\}$ in \mathbb{C} . This leads to the following: If $d\mu$ is a measure on ∂D and $z_0 \in \partial D$ has $d = \text{dist}(z_0, \text{supp}(d\mu)) > 0$, then the circle of radius $d^2 / (2 + d)$ has no zeros of the orthogonal polynomials. Golinskii has pointed out that the theorem of Fejér [3] that the zeros lie in the convex hull of $\text{supp}(d\mu)$ implies there are no zeros in the circle of radius $d^2 / 2$ and this is a stronger result, so we do not provide the details.

3. Isolated points of the support of $d\mu$

To prove Theorem 1.2, we will make use of the second kind polynomials [4,7] associated to $d\mu$ and $\{p_n\}$. This is a second family of polynomials, q_n defined by recursion coefficients, \tilde{a}_n, \tilde{b}_n with

$$\tilde{a}_n = a_{n+1}, \quad \tilde{b}_n = b_{n+1}. \tag{3.1}$$

They have the following two critical properties:

Proposition 3.1. (i) *The zeros of p_{n+1} and q_n interlace. In particular, between any two zeros of p_{n+1} is a zero of q_n .*

(ii) *If x_0 is an isolated point of $d\mu$ and dv is a suitable measure with respect to which the q 's are orthogonal, then $x_0 \notin \text{supp}(dv)$.*

These are well known. (i) follows from the fact that the zeros of p_{n+1} are eigenvalues of the matrix

$$J_{ij}^{(n+1)} = b_i \delta_{ij} + a_i \delta_{ij+1} + a_{i-1} \delta_{ij-1}, \quad 1 \leq i, j \leq n+1$$

and the zeros of q_n are the eigenvalues of

$$\tilde{J}_{ij}^{(n)} = \tilde{b}_i \delta_{ij} + \tilde{a}_i \delta_{ij+1} + \tilde{a}_{i-1} \delta_{ij-1}, \quad 1 \leq i, j \leq n$$

which is the matrix $J_{ij}^{(n+1)}$ with the top row and left column removed. (ii) follows because of the relation that v obeys for all $z \in \mathbb{C} \setminus \mathbb{R}$ (see, e.g., [7]):

$$\int \frac{dv(x)}{x-z} = a_1^{-2} \left[b_1 - z - \left(\int \frac{d\mu(x)}{x-z} \right)^{-1} \right] \tag{3.2}$$

(if the moment problem is indeterminate, this is one possible v). Isolated points of $d\mu$ are poles of $\int d\mu(x)/(x-z)$ so $\int dv(x)/(x-z)$ is regular there.

Proof of Theorem 1.2. Let $d_0 = \text{dist}(x_0, \text{supp}(dv)) > 0$ by (ii) of Proposition 3.1. By Theorem 1.1 and (3.1), either q_{n-1} or q_n has no zeros in $(x_0 - \delta_n, x_0 + \delta_n)$. By the intertwining result (Proposition 3.1(i)), either p_n or p_{n+1} cannot have two zeros in this interval. \square

Remark. Let x_0 be an isolated point of $d\mu$. If $b \in \text{supp}(d\mu)$ is such that $|x_0 - b| = \text{dist}(x_0, \text{supp}(d\mu) \setminus \{x_0\})$ and $\int d\mu(y)/|y-b| = \infty$, then dv has an isolated point in between x_0 and b , and so d_0 may be strictly less than $\text{dist}(x_0, \text{supp}(d\mu) \setminus \{x_0\})$.

4. An example of dense zeros in the gap

Nevai raised the issue of whether as n varies, the single possible zero of p_n in a gap (a, b) of $\text{supp}(d\mu)$ can yield all of (a, b) as limit points, or if the situation of a single

(or finite number of) limit point as in the example in Section 1 is the only possibility. In this section, we describe an explicit bounded Jacobi matrix so that $\text{supp}(d\mu) = [-5, -1] \cup [1, 5]$ but the set $\{x \in (-1, 1) \mid p_n(x) = 0 \text{ for some } n\}$ is dense in $[-1, 1]$.

Note: While this paper was in the refereeing process, we received a paper of Peherstorfer [5], who also constructs examples, very different from ours, of dense zeros in a gap.

Let $\{\beta_j\}_{j=1}^\infty$ be the sequence

$$\beta_1, \beta_2, \dots = 0, -\frac{1}{2}, 0, \frac{1}{2}, -\frac{3}{4}, -\frac{1}{2}, -\frac{1}{4}, 0, \frac{1}{4}, \frac{3}{4}, -\frac{7}{8}, \dots$$

which goes through *all* dyadic rationals in $(-1, 1)$ with denominator 2^k successively for $k = 1, 2, 3, \dots$ with each $j/2^k$ “covered” multiple times. Let L be the Jacobi matrix with

$$a_{2n-1} = 3, \quad a_{2n} = 1, \quad n = 1, 2, \dots, \tag{4.1}$$

$$b_k = \beta_n \quad \text{if } 2n^2 \leq k < 2(n+1)^2, \tag{4.2}$$

$$b_1 = \beta_1.$$

We claim that

- (1) $\text{supp}(d\mu) = [-5, -1] \cup [1, 5]$.
- (2) There is an x_n with $|x_n - \beta_n| \leq 2 \cdot 3^{-2n}$ so that

$$p_{2(n+1)^2-1}(x_n) = 0. \tag{4.3}$$

This provides the claimed example.

Remarks. (1) By adjusting a_1 and a_2 (but keeping $a_{2n+1} = a_1$; $a_{2n} = a_2$), we can replace $[-5, -1] \cup [1, 5]$ by $[-3 - \varepsilon, -1] \cup [1, 3 + \varepsilon]$, but our method seems to require bands larger than the size of the gap.

(2) One can replace (4.2) by $b_k = \beta_n$ for $\ell_n \leq k < \ell_{n+1}$ so long as $\ell_{n+1} - \ell_n \rightarrow \infty$.

(3) We believe that the measure associated to L is purely singular.

To prove the claims, we let L_0 be the Jacobi matrix with a 's given by (4.1) but $b_n = 0$, and L_∞ the period two, doubly infinite matrix on \mathbb{Z} which equals L_0 when restricted to \mathbb{Z}^+ . By the general theory of periodic Schrödinger operators [6], the spectrum of L_∞ is the two bands where $|\Delta(x)| \leq 2$ where Δ is the discriminant, that is, the trace of the two-step transfer matrix. If a_1, a_2 are the two values of a (so $a_1 = 3, a_2 = 1$ in our example), a simple calculation shows that

$$\Delta(x) = \frac{1}{a_1 a_2} (x^2 - (a_1^2 + a_2^2)),$$

so $\Delta(x) = \pm 2$ occurs at $x = \pm |a_1 \pm a_2|$. Thus,

$$\text{spec}(L_\infty) = [-4, -2] \cup [2, 4]. \tag{4.4}$$

The orthonormal polynomials $p_n^{(0)}$ for L_0 at $x = 0$ obey the recursion relation

$$p_{2n+2}^{(0)}(0) = -3p_{2n}^{(0)}(0),$$

so we have

$$p_{2n+1}^{(0)} = 0, \quad p_{2n}^{(0)}(0) = (-3)^n. \tag{4.5}$$

By the general theory of restricting periodic operators to the half-line, $\text{spec}(L_0)$ is $\text{spec}(L_\infty)$ plus a possible single eigenvalue in the gap $(-2, 2)$. Since there is a symmetry, the only possible eigenvalue is at $x = 0$, but (4.5) says that 0 is not an eigenvalue since $\sum_{j=0}^\infty |p_j(0)|^2 = \infty$. Thus, $\text{spec}(L_0) = [-4, -2] \cup [2, 4]$ also. $L - L_0$ is a diagonal matrix, so it is easy to see $\|L - L_0\| = \sup_j |\beta_j| = 1$. Thus, $\text{spec}(L) \subset \bigcup_{x \in [-1, 1]} x + \text{spec}(L_0) = [-5, -1] \cup [1, 5]$. On the other hand, since the b 's are equal to β_j on arbitrary long runs, a Weyl vector argument (see [2, p. 36]) shows that

$$\text{spec}(L) \supset \overline{\bigcup_j \beta_j + \text{spec}(L_0)} = [-5, -1] \cup [1, 5]$$

so claim 1 is proven.

Let $L_{n,F}$ be the $n \times n$ matrix obtained by taking the first n rows and columns of L . Then the zeros of $p_n(x)$ are precisely the eigenvalues of $L_{n,F}$ (see [7, Proposition 5.6]). Let φ_j be the j component vector with $(p_0^{(0)}(0), p_1^{(0)}(0), \dots, p_{j-1}^{(0)}(0))$. Then if j is odd so $p_j^{(0)}(0) = 0$, and we have $L_{0;j,F} \varphi_j = 0$. Thus, if $j = 2(n + 1)^2 - 1$,

$$[(L_{j,F} - \beta_n) \varphi_j]_k = (b_k - \beta_n) \varphi_{j,k}. \tag{4.6}$$

If $2n^2 \leq k \leq 2(n + 1)^2 - 1$, the right-hand side is zero and its absolute value is always less than $2|\varphi_{j,k}|$. Thus,

$$\begin{aligned} \frac{\| (L_{j,F} - \beta_n) \varphi_j \|^2}{\| \varphi_j \|^2} &\leq \frac{4 \sum_{k=0}^{n^2-1} 3^{2k}}{\sum_{k=0}^{(n+1)^2-1} 3^{2k}} \\ &\leq 4 \cdot 3^{-4n} \end{aligned}$$

by (4.5) and a simple estimate. Thus, $L_{j,F}$ has an eigenvalue within $2 \cdot 3^{-2n}$ of β_n , proving claim 2.

This completes the example.

References

- [1] A. Ambroladze, On exceptional sets of asymptotic relations for general orthogonal polynomials, J. Approx. Theory 82 (1995) 257–273.
- [2] H.L. Cycon, R.G. Froese, W. Kirsch, B. Simon, Schrödinger Operators With Application to Quantum Mechanics and Global Geometry, Springer, Berlin, 1987.
- [3] L. Fejér, Über die Lage der Nullstellen von Polynomen, die aus Minimum-forderungen gewisser Art entspringen, Math. Ann. 85 (1922) 41–48.
- [4] G. Freud, Orthogonal Polynomials, Pergamon, Oxford–New York, 1971.
- [5] F. Peherstorfer, Zeros of polynomials orthogonal on several intervals, Internat. Math. Res. Notices (7) (2003) 361–385.

- [6] M. Reed, B. Simon, *Methods of Modern Mathematical Physics, IV: Analysis of Operators*, Academic Press, New York, 1978.
- [7] B. Simon, The classical moment problem as a self-adjoint finite difference operator, *Adv. Math.* 137 (1998) 82–203.