



ELSEVIER

Available at

www.ElsevierMathematics.com

POWERED BY SCIENCE @ DIRECT®

Journal of Approximation Theory 124 (2003) 194–218

JOURNAL OF
Approximation
Theory

<http://www.elsevier.com/locate/jat>

Finding the projection of a point onto the intersection of convex sets via projections onto half-spaces

Lev M. Bregman,^a Yair Censor,^{b,*} Simeon Reich,^c and
Yael Zepkowitz-Malachi^b

^a*The Institute for Industrial Mathematics, 4 Yehuda Hanakhtom Street, Beer-Sheva 84311, Israel*

^b*Department of Mathematics, University of Haifa, Mt. Carmel, Haifa 31905, Israel*

^c*Department of Mathematics, The Technion–Israel Institute of Technology, Haifa 32000, Israel*

Received 9 May 2002; accepted in revised form 1 August 2003

Communicated by Frank Deutsch

Abstract

We present a modification of Dykstra's algorithm which allows us to avoid projections onto general convex sets. Instead, we calculate projections onto either a half-space or onto the intersection of two half-spaces. Convergence of the algorithm is established and special choices of the half-spaces are proposed.

The option to project onto half-spaces instead of general convex sets makes the algorithm more practical. The fact that the half-spaces are quite general enables us to apply the algorithm in a variety of cases and to generalize a number of known projection algorithms.

The problem of projecting a point onto the intersection of closed convex sets receives considerable attention in many areas of mathematics and physics as well as in other fields of science and engineering such as image reconstruction from projections.

In this work we propose a new class of algorithms which allow projection onto certain super half-spaces, i.e., half-spaces which contain the convex sets. Each one of the algorithms that we present gives the user freedom to choose the specific super half-space from a family of such

*Corresponding author.

E-mail addresses: bregman@math.bgu.ac.il (L.M. Bregman), yair@math.haifa.ac.il (Y. Censor), sreich@technion.technion.ac.il (S. Reich), yael@math.haifa.ac.il (Y. Zepkowitz-Malachi).

half-spaces. Since projecting a point onto a half-space is an easy task to perform, the new algorithms may be more useful in practical situations in which the construction of the super half-spaces themselves is not too difficult.

© 2003 Elsevier Inc. All rights reserved.

Keywords: Convex set; Dykstra's algorithm; Half-space; Projection

1. Introduction

The Dykstra algorithm is an iterative procedure which (asymptotically) finds the nearest point projection (also called the orthogonal projection) of any given point onto the intersection of a given family of closed convex sets. It iterates by passing sequentially over the individual sets and projecting onto each one a *deflected version* of the previous iterate. A precise description of the algorithm can be found in [26]. The algorithm was first proposed and analyzed by Dykstra [21] in 1983 for a family of closed convex cones in R^n . Boyle and Dykstra [9] studied the algorithm for general convex sets in Hilbert space. Han rediscovered the algorithm in 1988 [26] and investigated its behavior in R^n by using the duality theory of mathematical programming. Han and Lou [27] proposed a simultaneous version of the algorithm in R^n .

Gaffke and Mathar [25] studied the Dykstra algorithm in Hilbert space from a duality standpoint and showed its relation to the method of component-wise cyclic minimization over a Cartesian product. They also proposed a fully simultaneous Dykstra algorithm. Iusem and De Pierro showed in [32] convergence of the simultaneous Dykstra algorithm in both consistent and inconsistent cases in R^n , using Pierra's [38] formalism, see also [31]. Crombez [17] did a similar analysis in Hilbert space. Combettes [15] included the Dykstra algorithm in his review. Bauschke and Borwein [3] analyzed the algorithm for two sets in Hilbert space and generalized the work of Iusem and De Pierro [32] to this setting. Deutsch and Hundal published a rate of convergence study for the polyhedral case in [20], and in [30] established generalizations to an infinite family of sets and to random, rather than cyclic, order control, see also [19].

Han [26], as well as Iusem and De Pierro [32], showed that for linear inequality constraints and for linear interval inequality constraints (the polyhedral case), the method of Dykstra becomes the Hildreth algorithm, first published in [29] and studied further by D'Esopo [18], and by Lent and Censor [35].

Censor and Reich [13] proposed a synthesis of Dykstra's algorithm with Bregman distances and obtained a new algorithm that solves the best approximation problem with Bregman projections. However, they established convergence of the resulting Dykstra algorithm with Bregman projections only when the constraints are half-spaces. Bauschke and Lewis [8] provided the first proof for general closed convex constraint sets. Their analysis relies on some strong properties of Bregman distances corresponding to Legendre functions which were treated earlier by Bauschke and Borwein [5]. Bauschke and Lewis [8] also discovered the close relationship between

the Dykstra algorithm with Bregman projections and the very general and powerful algorithmic framework of Tseng [41]. Bregman, Censor and Reich show that the Dykstra algorithm with Bregman projections is precisely the nonlinear extension of the Bregman optimization algorithm in [11], see also Dykstra [22]. This recognition goes beyond the fact that the two algorithms coincide in the linear constraint case, as was shown by Censor and Reich [13]. It enables these authors to present a new proof and convergence analysis of Dykstra's algorithm with Bregman projections for almost cyclic control sequences, which rests on Bregman's original work [10]. Their work also offers an intuitive geometric interpretation of the iterative steps.

Other algorithms for finding the projection of a point onto the intersection of convex sets are available. Haugazeau proposed such an algorithm in [28] and his ideas were further abstracted and developed by Bauschke and Combettes [6]. The latter was extended to Bregman projections by Bauschke and Combettes [7]. Earlier, Pierra discussed yet another method, which is, however, related to Haugazeau's work, in [37,38]. Combettes constructed a block-iterative outer approximations method in [16], and Bauschke [2] extended an algorithm of Wittman.

In the present paper we propose an algorithmic scheme which is a modification of Dykstra's algorithm. It allows us to replace projections onto convex sets by projections onto either a half-space or the intersection of two half-spaces. A method, which replaces projections onto convex sets by projections onto either a half-space or the intersection of two half-spaces, was proposed and studied by Iusem and Svaiter [33,34], but in a different way. In our work, when we have to project onto the intersection of two half-spaces, our scheme enables us to choose one of the half-spaces from a family of possible half-spaces and the convergence theorem is true for the whole family of possible half-spaces. This feature allows us to construct many specific algorithms within our general scheme and to obtain the result of Iusem and Svaiter (both the algorithm and the convergence theorem) as a special case by making a specific choice of the half-spaces.

2. δ -Super half-spaces and δ -super hyperplanes: definitions and construction

The orthogonal projection x' of a point x onto a nonempty closed convex set $E \subseteq R^n$ can be viewed the orthogonal projection of x onto the particular hyperplane H which separates x from E and supports E at x' , the closest point to x in E . (For the definitions of a separating hyperplane and a supporting hyperplane consult any book on convex analysis or optimization theory or look, e.g., in [14].)

But, of course, at the time of performing such an orthogonal projection neither the point x' , nor the separating and supporting hyperplane H are available. In view of the simplicity of performing an orthogonal projection onto a hyperplane, it is natural to ask whether in the construction of iterative projection algorithms one could use other separating supporting hyperplanes, instead of that particular hyperplane H through the closest point to x .

Aside from theoretical interest, this approach leads to algorithms that can be used in practice, provided that the computational effort of finding such other hyperplanes

competes favorably with the work involved in performing orthogonal projections directly onto the given sets.

Such an approach was taken by Aharoni et al. [1], where the (δ, η) -algorithm for the convex feasibility problem replaces orthogonal projections onto the convex sets by projections onto separating hyperplanes (see also [14, Algorithm 5.5.1]).

In the present paper we use the rationale behind the (δ, η) -algorithm to deal with another class of mathematical problems, namely, finding the optimal point that minimizes a given objective function over the intersection of given convex sets. We construct a family of half-spaces and hyperplanes with particular properties and replace projections onto convex sets by projections onto a half-space or a hyperplane from this family or onto the intersection of two half-spaces.

In the next subsection we define δ -super half-spaces and δ -super hyperplanes which play an important role throughout this work.

2.1. Definition and construction of δ -super half-spaces and δ -super hyperplanes

Let $E \subseteq R^n$ be a nonempty closed convex set defined by $E := \{x \in R^n \mid e(x) \leq 0\}$, where $e : R^n \rightarrow R$ is a convex function, and let $z \in R^n$ be a given point. For $z \notin E$ we wish to construct a half-space which contains the set E , but does not contain any point of the interior of a ball centered at z with radius $\delta e(z)$, for some fixed δ , $0 < \delta \leq 1$. Such a half-space will be called a δ -super half-space (δ -SHS for short) with respect to the ball described above and E , and its boundary will be called a δ -super hyperplane (δ -SHP for short). If $z \in E$, then the only possible δ -SHS is defined to be R^n (see Fig. 1).

Definition 1 (δ -Super half-space). Given are a point $z \in R^n$, a real number δ , $0 < \delta \leq 1$, and a nonempty closed convex set $E := \{x \in R^n \mid e(x) \leq 0\}$, where $e : R^n \rightarrow R$ is a convex function. For $z \notin E$ define the ball

$$B(z, \delta e(z)) := \{x \in R^n \mid \|x - z\| \leq \delta e(z)\} \tag{1}$$




Fig. 1. Geometric description of a δ -super half-space and a δ -super hyperplane.

and a half-space

$$S_E(z) := \{x \in \mathbb{R}^n \mid \langle t^E(z), x \rangle \leq \theta_E(z)\}, \tag{2}$$

where $t^E(z) \neq 0$ and $\theta_E(z) \in \mathbb{R}$. The set $S_E(z)$ will be called a δ -super half-space with respect to $B(z, \delta e(z))$ and E if and only if the following two conditions hold:

$$y \in B(z, \delta e(z)) \text{ implies } \langle t^E(z), y \rangle \geq \theta_E(z), \tag{3}$$

i.e.,

$$S_E(z) \cap \text{int } B(z, \delta e(z)) = \emptyset \tag{4}$$

and

$$E \subseteq S_E(z). \tag{5}$$

Definition 2 (δ -Super hyperplane). If $S_E(z)$ is a δ -SHS, as in Definition 1, and $z \notin E$, then its bounding hyperplane is called a δ -super hyperplane (δ -SHP for short) with respect to $B(z, \delta e(z))$ and E . If $z \in E$ then the δ -SHP is empty.

Here is an example of the construction of a δ -SHS. This particular example plays an important role in a special case of our new algorithm (see Section 5.1 below).

Example 3. Let E be a convex set, $E := \{x \in \mathbb{R}^n \mid e(x) \leq 0\}$, where $e(x)$ is a convex function. Let $z \notin E$ and denote by $e'(z)$ any subgradient of e at z . Assume that there exists an $M > 0$ such that $\|e'(z)\| \leq M$ for all z in some bounded set $G \subseteq \mathbb{R}^n$. Then

$$H := \{x \in \mathbb{R}^n \mid e(z) + \langle e'(z), x - z \rangle \leq 0\} \tag{6}$$

is a δ -SHS with respect to $B(z, \delta e(z))$ and E for all $z \in G$ and $\delta \leq 1/M$. In order to prove this claim we need to show that (4) and (5) hold. Indeed, let $y \in H$. Then

$$\langle e'(z), y - z \rangle \leq -e(z), \tag{7}$$

by (6). It follows that

$$e(z) \leq |\langle e'(z), y - z \rangle| \leq \|e'(z)\| \|z - y\|. \tag{8}$$

Using the assumption on the boundedness of the subgradients we obtain

$$e(z) \leq M \|z - y\| \tag{9}$$

or

$$e(z) \frac{1}{M} \leq \|z - y\|. \tag{10}$$

The last inequality shows that whenever δ is chosen such that $\delta \leq 1/M$, we obtain $y \notin \text{int } B(z, \delta e(z))$ which implies that $H \cap \text{int } B(z, \delta e(z)) = \emptyset$. Next, we show that $E \subseteq H$. Let $x \in E$. Using the well-known subgradient inequality (see, e.g., [39, p. 214]) we have

$$e(x) - e(z) \geq \langle e'(z), x - z \rangle. \tag{11}$$

Since $x \in E$, $e(x) \leq 0$. Thus

$$-e(z) \geq \langle e'(z), x - z \rangle, \tag{12}$$

which implies that $x \in H$, by (6), and as a result of (7)–(12) we obtain that H is indeed a δ -SHS, as claimed.

The last example is very important because there are many algorithms that use projections onto half-spaces of the form of (6), see, e.g., [23,24,33]. We will need to make use of the following condition.

Condition 4. *For any E , as in Definition 1, and any bounded set $G \subseteq R^n$, there exists a $\delta \in (0, 1)$ such that $z \in G$ and $z \notin E$ imply that the intersection $B(z, \delta e(z)) \cap E$ is empty.*

Remark 5. Condition 4 is necessary and sufficient to enable us to construct a δ -SHS. In our general algorithmic scheme, which will be presented next, we have to assume that this condition holds. For the special cases of the algorithm, that we treat separately, we show how to choose a δ such that Condition 4 actually holds. Example 3 illustrates such a special case. However, formulating a general sufficient condition for Condition 4 to hold for our general algorithmic scheme still eludes us.

3. The δ -SHS algorithm

3.1. The algorithm

We consider the optimization problem

$$\min \{f(x) \mid x \in Q\}, \tag{13}$$

where $Q = \bigcap_{i=1}^m Q_i$, $Q_i := \{x \in R^n \mid q_i(x) \leq 0\}$, and $f, \{q_i\}_{i=1}^m$ are real-valued functions the effective domains of which, $\text{dom } f$ and $\text{dom } q_i$, are subsets of R^n . We make the following assumptions regarding the constraints:

Assumption A1. $q_i(x)$ is convex, $1 \leq i \leq m$.

Assumption A2. $Q \neq \emptyset$.

Assumption A3. $Q \cap (\text{dom } f) \neq \emptyset$ and $Q_i \cap \text{int}(\text{dom } f) \neq \emptyset$, for all $i, 1 \leq i \leq m$. (But $Q \cap \text{int}(\text{dom } f)$ may be empty.)

Assumption A4. The sets $\text{dom } q_i, 1 \leq i \leq m$, are “wide enough” in the sense that all points appearing in the new Algorithm 8, defined below, belong to $\text{int}(\text{dom } q_i), 1 \leq i \leq m$.

We assume that f is a *Bregman function with zone* $S = \text{int}(\text{dom } f)$ (see the definition of $D_f(x, y)$ in, e.g., [12,14, Definition 2.1.1]).

Remark 6. If f is a Bregman function with zone $S = \text{int}(\text{dom} f)$ and if a point $a \in S = \text{int}(\text{dom} f)$ satisfies $\nabla f(a) = 0$, then the problem

$$\min\{f(x) \mid x \in Q \cap (\text{dom} f)\} \tag{14}$$

achieves its minimum at the Bregman projection with respect to f of the point a onto Q . This follows immediately from the definition of the Bregman projection of a point onto Q , and the fact that $\nabla f(a) = 0$, i.e.,

$$\begin{aligned} P_Q^f(a) &= \arg \min\{D_f(x, a) \mid x \in Q \cap (\text{dom} f)\} \\ &= \arg \min\{f(x) - f(a) - \langle \nabla f(a), x - a \rangle \mid x \in Q \cap \bar{S}\} \\ &= \arg \min\{f(x) \mid x \in Q \cap (\text{dom} f)\}. \end{aligned} \tag{15}$$

We add the next three assumptions on f to make sure that the algorithm is well defined:

Assumption B1. The function f is co-finite, which, since it is a Bregman function, implies that the mapping $y = \nabla f(x)$ is a one-to-one mapping of $\text{int}(\text{dom} f)$ onto R^n (see, e.g., [39, Theorem 26.5]).

Assumption B2. The function f is *zone consistent* with respect to any half-space and with respect to the intersection of two half-spaces containing points from $\text{int}(\text{dom} f)$.

Remark 7. Bauschke and Borwein showed in [5, Theorem 3.14] that if f is a Legendre function, then it is zone consistent. Rockafellar [39, Lemma 26.7] gives a characterization of co-finiteness for differentiable convex functions.

Assumption B3. The function f has a global minimum.

The precise description of the new algorithm that we propose for problem (13) is as follows:

Algorithm 8. 1. Data at the beginning of the k th iterative step

- 1.1. *Current approximation* $x^k \in \text{int}(\text{dom} f)$.
- 1.2. m vectors $a_i^k \in R^n$ and m real numbers α_i^k , $1 \leq i \leq m$, such that each pair (a_i^k, α_i^k) defines a half-space L_i^k ,

$$L_i^k := \{x \in R^n \mid \langle a_i^k, x \rangle \leq \alpha_i^k\}, \tag{16}$$

containing Q_i .

2. Initialization

2.1. x^0 is a (global) minimum point of $f(x)$ on R^n , i.e.,

$$\nabla f(x^0) = 0. \tag{17}$$

2.2. Set $a_i^0 = 0$ and $\alpha_i^0 = 0$ for all $1 \leq i \leq m$.

2.3. Choose a real $\delta \in (0, 1]$ such that Condition 4 holds for the bounded set

$$G = \{x \in R^n \mid D_f(y, x) \leq f(y) - f(x^0)\}, \text{ for some } y \in Q \cap \text{dom} f \tag{18}$$

and for each $Q_i, i = 1, 2, \dots, m$. (Note that G is bounded because the partial level sets of the Bregman distance D_f are always bounded, see, e.g., [17 Definition 2.1.1].)

3. Iterative step

3.1. Choose an operating control index $i(k)$ from the almost cyclic control sequence. Recall (see, e.g., [14, Definition 5.1.1]) that an almost cyclic control sequence on $\{1, 2, \dots, m\}$ is a sequence $\{i(k)\}_{k=0}^\infty$ such that $1 \leq i(k) \leq m$ for all $k \geq 0$ and there exists a constant (called the almost cyclicity constant) $T \geq m$ such that, for all $k \geq 0$,

$$\{1, 2, \dots, m\} \subseteq \{i(k+1), i(k+2), \dots, i(k+T)\}. \tag{19}$$

3.2. Calculate $z^k \in R^n$ such that

$$\nabla f(z^k) = \nabla f(x^k) + a_{i(k)}^k. \tag{20}$$

Such a vector $z^k \in \text{int}(\text{dom} f)$ exists because of Assumption B1.

3.3. Set $x^{k+1}, \alpha_{i(k)}^{k+1}, a_{i(k)}^{k+1}$ by one of the following two possible options:

3.3.1. If $q_{i(k)}(x^k) \leq 0$, then let x^{k+1} be the Bregman projection with respect to f of z^k onto the half-space

$$L_{i(k)}^k := \{x \in R^n \mid \langle a_{i(k)}^k, x \rangle \leq \alpha_{i(k)}^k\}. \tag{21}$$

That is,

if $z^k \in L_{i(k)}^k$, then let $x^{k+1} = z^k$ and define $\lambda_k := 0$,

if $z^k \notin L_{i(k)}^k$, then x^{k+1} and λ_k are calculated from the Karush–Kuhn–

Tucker (see, e.g., [36]) conditions

$$\begin{cases} \nabla f(x^{k+1}) = \nabla f(z^k) - \lambda_k a_{i(k)}^k, \\ \lambda_k \geq 0, \\ \langle a_{i(k)}^k, x^{k+1} \rangle \leq \alpha_{i(k)}^k, \\ \lambda_k (\langle a_{i(k)}^k, x^{k+1} \rangle - \alpha_{i(k)}^k) = 0. \end{cases} \tag{22}$$

Next, set

$$\begin{cases} a_{i(k)}^{k+1} = \lambda_k a_{i(k)}^k, \\ \alpha_{i(k)}^{k+1} = \lambda_k \alpha_{i(k)}^k. \end{cases} \tag{23}$$

3.3.2. If $q_{i(k)}(x^k) > 0$, then let x^{k+1} be the Bregman projection with respect to f of z^k onto the intersection of the following two half-spaces: $L_{i(k)}^k$ which was defined by (21) and

$$S_{Q_{i(k)}}(x^k) := \{x \in R^n \mid \langle t^{i(k)}, x \rangle \leq \theta_{i(k)}\}, \tag{24}$$

which is a δ -SHS with respect to the ball $B(x^k, \delta q_{i(k)}(x^k))$ and $Q_{i(k)}$. In other words,

$$\begin{cases} \langle t^{i(k)}, x \rangle \leq \theta_{i(k)} & \text{for all } x \in Q_{i(k)}, \\ \langle t^{i(k)}, x \rangle \geq \theta_{i(k)} & \text{for all } x \in B(x^k, \delta q_{i(k)}(x^k)), \end{cases} \tag{25}$$

where $t^{i(k)} = t^{i(k)}(x^k) \neq 0$ and $\theta_{i(k)} = \theta_{i(k)}(x^k)$ (see Section 2 for the definition and construction of the δ -SHS). Thus, we calculate the vector x^{k+1} , λ_k and μ_k from the Karush–Kuhn–Tucker conditions

$$\begin{cases} \nabla f(x^{k+1}) - \nabla f(z^k) + \lambda_k a_{i(k)}^k + \mu_k t^{i(k)} = 0, \\ \lambda_k \geq 0, \quad \mu_k \geq 0, \\ \langle a_{i(k)}^k, x^{k+1} \rangle \leq \alpha_{i(k)}^k, \\ \langle t^{i(k)}, x^{k+1} \rangle \leq \theta_{i(k)}, \\ \lambda_k (\langle a_{i(k)}^k, x^{k+1} \rangle - \alpha_{i(k)}^k) = 0, \\ \mu_k (\langle t^{i(k)}, x^{k+1} \rangle - \theta_{i(k)}) = 0, \end{cases} \tag{26}$$

and then set

$$\begin{cases} a_{i(k)}^{k+1} = \lambda_k a_{i(k)}^k + \mu_k t^{i(k)}, \\ \alpha_{i(k)}^{k+1} = \lambda_k \alpha_{i(k)}^k + \mu_k \theta_{i(k)}. \end{cases} \tag{27}$$

3.4. For $i \neq i(k)$ do not change a_i^k and α_i^k , i.e., set

$$\begin{aligned} a_i^{k+1} &= a_i^k, \\ \alpha_i^{k+1} &= \alpha_i^k. \end{aligned} \tag{28}$$

Fig. 2 describes geometrically the various possibilities of the iterative step of Algorithm 8, in the following way: Cases (i)–(iii) describe iterative steps in which x^k belongs to the convex set $Q_{i(k)}$ (but the modified point z^k can be in $Q_{i(k)}$, in $L_{i(k)}^k$ and not in $Q_{i(k)}$, and not in $L_{i(k)}^k$, respectively). Case (iv) describes the iterative step in which x^k does not belong to the convex set $Q_{i(k)}$.

Note that the Bregman projection of a point onto a half-space or onto the intersection of two half-spaces exists. According to Assumption B2 the projection belongs to $\text{int}(\text{dom} f)$, thus we have that x^{k+1} , λ_k and μ_k in (22) and (26) exist and $x^{k+1} \in \text{int}(\text{dom} f)$.




Fig. 2. Geometric interpretation of Algorithm 8.

Remark 9. Observe that no relaxation parameters appear in our algorithm. Some of the special cases discussed below employ a sequence $\{\beta_k\}_{k \geq 0}$ of relaxation parameters. Loosely speaking, these parameters overdo or underdo the move prescribed in an iterative step. Relaxation parameters add an extra degree of freedom to the way a method might actually be implemented, and have important consequences for the performance of the method in practice. We, however, do have the flexibility to choose a δ -SHS which will lie closer to x^k at each iterate step. In this way, some underrelaxation, i.e., $\varepsilon \leq \beta_k < 1$ for some arbitrarily small $\varepsilon > 0$, can be actually incorporated in our algorithm.

The three lemmas below will be used to prove the convergence of Algorithm 8.

Lemma 10. For any k , the half-space $L_{i(k)}^{k+1} := \{x \in R^n \mid \langle a_{i(k)}^{k+1}, x \rangle \leq \alpha_{i(k)}^{k+1}\}$, defined by the pair $(a_{i(k)}^{k+1}, \alpha_{i(k)}^{k+1})$, generated by Algorithm 8, contains $Q_{i(k)}$.

Proof. When Step 3.3.1 in Algorithm 8 holds, λ_k is nonnegative (see [14, Lemma 2.2.2]). Notice that there is a sign difference of λ_k between Lemma 2.2.2 in [14] and our lemma, because of a different definition of the Lagrangian function). Thus, by definition (23) of $a_{i(k)}^{k+1}$ and $\alpha_{i(k)}^{k+1}$ we obtain $L_{i(k)}^{k+1} = L_{i(k)}^k$, and $L_{i(k)}^k$ contains Q_i (by Step 1.2 of Algorithm 8). When Step 3.3.2 of Algorithm 8 holds, let $x \in Q_{i(k)}$. We will show that this implies that $x \in L_{i(k)}^{k+1}$. We have

$$\langle a_{i(k)}^{k+1}, x \rangle = \lambda_k \langle a_{i(k)}^k, x \rangle + \mu_k \langle t^{i(k)}, x \rangle \leq \lambda_k \alpha_{i(k)}^k + \mu_k \theta_{i(k)} = \alpha_{i(k)}^{k+1}, \tag{29}$$

where the first equality uses (27), the inequality uses (21) and (24), and the last equality again uses (27). We got that if $x \in Q_{i(k)}$, then $\langle a_{i(k)}^{k+1}, x \rangle \leq a_{i(k)}^{k+1}$, i.e., $x \in L_{i(k)}^{k+1}$. Thus, the lemma is true for Step 3.3.2 too. So Lemma 10 is true in all cases. \square

Lemma 11. *If $\{x^k\}_k$ and $\{a_i^k\}_{k \geq 0}$ are generated by Algorithm 8, then, for all k , we have*

$$\nabla f(x^k) + \sum_{i=1}^m a_i^k = 0. \tag{30}$$

Proof. For $k = 0$ the statement is true by (17) and Step 2.2 in the initialization of Algorithm 8. Now we suppose that it is true for some k and we prove it for $k + 1$. For the proof we will use the fact (see Step 3.4 of Algorithm 8) that

$$\sum_{\substack{i=1 \\ i \neq i(k)}}^m a_i^{k+1} = \sum_{\substack{i=1 \\ i \neq i(k)}}^m a_i^k. \tag{31}$$

When Step 3.3.1 in Algorithm 8 holds, then using (22), (20), the induction hypothesis, (31) and (23) we obtain

$$\begin{aligned} \nabla f(x^{k+1}) &= \nabla f(z^k) - \lambda_k a_{i(k)}^k = \nabla f(x^k) + a_{i(k)}^k - \lambda_k a_{i(k)}^k \\ &= \left(- \sum_{i=1}^m a_i^k \right) + a_{i(k)}^k - \lambda_k a_{i(k)}^k \\ &= \left(- \sum_{\substack{i=1 \\ i \neq i(k)}}^m a_i^{k+1} \right) - a_{i(k)}^k + a_{i(k)}^k - \lambda_k a_{i(k)}^k \\ &= \left(- \sum_{\substack{i=1 \\ i \neq i(k)}}^m a_i^{k+1} \right) - a_{i(k)}^{k+1} = - \sum_{i=1}^m a_i^{k+1}. \end{aligned} \tag{32}$$

When Step 3.3.2 in Algorithm 8 holds, then we use (26), (20), the induction hypothesis, (31) and (27), to obtain

$$\begin{aligned} \nabla f(x^{k+1}) &= \nabla f(z^k) - \lambda_k a_{i(k)}^k - \mu_k t^{i(k)} = \nabla f(x^k) + a_{i(k)}^k - \lambda_k a_{i(k)}^k - \mu_k t^{i(k)} \\ &= \left(- \sum_{i=1}^m a_i^k + a_{i(k)}^k \right) - \lambda_k a_{i(k)}^k - \mu_k t^{i(k)} \end{aligned}$$

$$\begin{aligned}
 &= \left(- \sum_{\substack{i=1 \\ i \neq i(k)}}^m a_i^{k+1} \right) - a_{i(k)}^k + a_{i(k)}^k - \lambda_k a_{i(k)}^k - \mu_k t^{i(k)} \\
 &= \left(- \sum_{\substack{i=1 \\ i \neq i(k)}}^m a_i^{k+1} \right) - a_{i(k)}^{k+1} = - \sum_{i=1}^m a_i^{k+1}.
 \end{aligned} \tag{33}$$

This completes the proof of Lemma 11. \square

Lemma 12. *If $\{x^k\}_{k \geq 0}$, $\{a_i^k\}_{k \geq 0}$ and $\{\alpha_i^k\}_{k \geq 0}$ are generated by Algorithm 8, then, for $k \geq 0$, there exists an integer $r = r(k) \geq 0$ and r vectors*

$$y^j \in \{x^0, x^1, \dots, x^{k-1}\}, \quad 1 \leq j \leq r,$$

such that, for all i , $1 \leq i \leq m$, a_i^k can be represented as a finite linear combination of the normal vectors t^i , generated in Step 3.3.2 of Algorithm 8, at y^j with nonnegative coefficients, that is,

$$a_i^k = \sum_{j=1}^r \gamma_j t^i(y^j), \quad \gamma_j \geq 0, \tag{34}$$

and, for all i , $1 \leq i \leq m$, α_i^k can be represented as

$$\alpha_i^k = \sum_{j=1}^r \gamma_j \theta_i(y^j), \quad \gamma_j \geq 0, \tag{35}$$

where the $\theta_i(y^j)$ s correspond to the $t^i(y^j)$ s generated above.

Proof. For $k = 0$, the left-hand sides of (34) and (35) are zero by Step 2.2 of Algorithm 8. So both statements are true with $\gamma_j = 0, j = 1, 2, \dots, r$. We now assume that the lemma is true for some k and prove it for $k + 1$.

For $i \neq i(k)$ both (34) and (35) hold, because a_i^k and α_i^k do not change, according to Step 3.4 of the algorithm.

For $i = i(k)$, in Step 3.3.1 or in Step 3.3.2 with $\mu_k = 0$, (34) and (35) hold, because, by (23) or (27) and the induction hypothesis,

$$a_{i(k)}^{k+1} = \lambda_k a_{i(k)}^k = \lambda_k \left(\sum_{j=1}^r \gamma_j t^i(y^j) \right) \tag{36}$$

and

$$\alpha_{i(k)}^{k+1} = \lambda_k \alpha_{i(k)}^k = \lambda_k \left(\sum_{j=1}^r \gamma_j \theta_{i(k)}(y^j) \right). \tag{37}$$

In Step 3.3.2 with $\mu_k > 0$, (34) and (35) hold by (27), the induction hypothesis and the fact that a vector x^k is added to the set $\{x^0, x^1, \dots, x^{k-1}\}$, so we have

$t^{i(k)} = t^{i(k)}(y^j)$ for some y^j . Thus we have

$$\alpha_{i(k)}^{k+1} = \lambda_k \alpha_{i(k)}^k + \mu_k t^{i(k)}(x^k) = \lambda_k \left(\sum_{j=1}^r \gamma_j t^{i(k)}(y^j) \right) + \mu_k t^{i(k)}(y^j), \tag{38}$$

$$\alpha_{i(k)}^{k+1} = \lambda_k \alpha_{i(k)}^k + \mu_k \theta_{i(k)}(x^k) = \lambda_k \left(\sum_{j=1}^r \gamma_j \theta_{i(k)}(y^j) \right) + \mu_k \theta_{i(k)}(y^j). \quad \square \tag{39}$$

4. Convergence of Algorithm 8

We have to make now an additional assumption on the functions q_i . Denote by I_1 the subset of $I = \{1, 2, \dots, m\}$ for which the q_i are affine functions.

Assumption A5. There exists a point $\bar{y} \in Q \cap \text{dom} f$ such that $q_i(\bar{y}) < 0$ for all $i \in I_2 := I \setminus I_1$.

4.1. The convergence theorem

Theorem 13. *Let f be a Bregman function, let $q_i, 1 \leq i \leq m$, be convex functions and let Assumptions A1–A5 and B1–B3 hold. Then any sequence $\{x^k\}_{k \geq 0}$, generated by Algorithm 8, converges to the solution of (13).*

Proof. The proof is divided into five steps. In Step 1 we define the sequence $\{\varphi_k\}_{k \geq 0}$ by

$$\varphi_k := f(x^k) + \sum_{i=1}^m (\langle a_i^k, x^k \rangle - \alpha_i^k) \tag{40}$$

and show that it is increasing. Step 2 proves that

$$\lim_{k \rightarrow \infty} D_f(x^{k+1}, x^k) = 0. \tag{41}$$

In Step 3 we show that the sequences $\{a_i^k\}_{k \geq 0}$ are bounded for all i , unless $\alpha_i^k - \langle a_i^k, y \rangle = 0$ for all $y \in Q$ and for all $k \geq 0$. We consider the index sets I_1 and I_2 separately and distinguish two possibilities for I_2 . Step 4 shows that the sequence $\{x^k\}_{k \geq 0}$ converges to

$$x^* = \lim_{t \rightarrow \infty} x^{k_t}, \tag{42}$$

where the sequence $\{x^{k_t}\}_{t \geq 0}$ is defined below, and proves that $x^* \in Q$. In Step 5 we show that for all $i, 1 \leq i \leq m$, the limit

$$\lim_{k \rightarrow \infty, k \in W} (\alpha_i^k - \langle a_i^k, x^* \rangle) = 0 \tag{43}$$

holds for a certain set of indices W (defined in (66)) and from this and other arguments presented there we obtain that Theorem 13 does indeed hold.

Step 1: In order to prove that $\{\varphi_k\}_{k \geq 0}$ is increasing, we show that $\varphi_{k+1} - \varphi_k \geq 0$ for all $k \geq 0$. By definition of φ_k , the fact that for $i \neq i(k)$ we have, by (28), $a_i^{k+1} = a_i^k$ and $\alpha_i^{k+1} = \alpha_i^k$, the definition of D_f and Lemma 11 we get

$$\begin{aligned} \varphi_{k+1} - \varphi_k &= f(x^{k+1}) - f(x^k) + \sum_{i \neq i(k)} \langle a_i^k, x^{k+1} - x^k \rangle \\ &\quad + \langle a_{i(k)}^{k+1}, x^{k+1} \rangle - \alpha_{i(k)}^{k+1} - (\langle a_{i(k)}^k, x^k \rangle - \alpha_{i(k)}^k) \\ &= D_f(x^{k+1}, x^k) + \langle \nabla f(x^k), x^{k+1} - x^k \rangle + \sum_{i=1}^m \langle a_i^k, x^{k+1} - x^k \rangle \\ &\quad - \langle a_{i(k)}^k, x^{k+1} - x^k \rangle + \langle a_{i(k)}^{k+1}, x^{k+1} \rangle \\ &\quad - \alpha_{i(k)}^{k+1} - (\langle a_{i(k)}^k, x^k \rangle - \alpha_{i(k)}^k) \\ &= D_f(x^{k+1}, x^k) + \langle a_{i(k)}^{k+1}, x^{k+1} \rangle - \alpha_{i(k)}^{k+1} \\ &\quad - (\langle a_{i(k)}^k, x^{k+1} \rangle - \alpha_{i(k)}^k). \end{aligned} \tag{44}$$

In order to complete Step 1 we prove the next two assertions. First we show that in all steps of Algorithm 8,

$$\langle a_{i(k)}^{k+1}, x^{k+1} \rangle - \alpha_{i(k)}^{k+1} = 0 \tag{45}$$

holds. Indeed in Step 3.3.1, (45) is true by (23) and (21). In Step 3.3.2, we obtain from (27) and (25):

$$\begin{aligned} \langle a_{i(k)}^{k+1}, x^{k+1} \rangle - \alpha_{i(k)}^{k+1} &= \lambda \langle a_{i(k)}^k, x^{k+1} \rangle + \mu \langle t^{i(k)}, x^{k+1} \rangle - \lambda \alpha_{i(k)}^k - \mu \theta_{i(k)} \\ &= \lambda (\langle a_{i(k)}^k, x^{k+1} \rangle - \alpha_{i(k)}^k) + \mu (\langle t^{i(k)}, x^{k+1} \rangle - \theta_{i(k)}) \\ &= 0. \end{aligned} \tag{46}$$

Secondly, in all steps of Algorithm 8 we have

$$\langle a_{i(k)}^k, x^{k+1} \rangle - \alpha_{i(k)}^k \leq 0. \tag{47}$$

This is true because in both Steps 3.3.1 and 3.3.2, $x^{k+1} \in L_{i(k)}^k$. By (44), (45) and (47) we have

$$\varphi_{k+1} - \varphi_k \geq D_f(x^{k+1}, x^k) \geq 0, \tag{48}$$

which shows that the sequence $\{\varphi_k\}_{k \geq 0}$ is increasing. We also get, using (44) and (45) in (47) that

$$\varphi_{k+1} - \varphi_k \geq \alpha_{i(k)}^k - \langle a_{i(k)}^k, x^{k+1} \rangle \geq 0. \tag{49}$$

Step 2: Let $y \in \text{dom } f$. Then, by the definition of the Bregman projection with respect to f , Lemma 11 and (40), we have

$$\begin{aligned}
 D_f(y, x^k) &= f(y) - f(x^k) - \langle \nabla f(x^k), y - x^k \rangle \\
 &= f(y) - f(x^k) + \left\langle \sum_{i=1}^m a_i^k, y - x^k \right\rangle \\
 &= f(y) - f(x^k) - \sum_{i=1}^m (\langle a_i^k, x^k \rangle - \alpha_i^k) + \sum_{i=1}^m (\langle a_i^k, y \rangle - \alpha_i^k) \\
 &= f(y) - \varphi_k + \sum_{i=1}^m (\langle a_i^k, y \rangle - \alpha_i^k). \tag{50}
 \end{aligned}$$

Since $y \in Q$, we have that, for all i , $\langle a_i^k, y \rangle - \alpha_i^k \leq 0$, by the definition of L_i^k and (16). Hence, from (50), the last inequality and Step 1,

$$D_f(y, x^k) \leq f(y) - \varphi_k \leq f(y) - \varphi_0 = f(y) - f(x^0) \tag{51}$$

and the sequence $\{x^k\}_{k \geq 0}$ is bounded by the definition of a Bregman function, (see [14, Definition 2.1.1(iv)]). By the left-hand side inequality of (51), we have

$$\varphi_k \leq f(y) \quad \text{for all } y \in Q \cap \text{dom } f. \tag{52}$$

Thus the sequence $\{\varphi_k\}_{k \geq 0}$ is bounded and $\lim_{k \rightarrow \infty} \varphi_k$ exists. This fact and (48) imply that (41) holds. Another inequality that follows from (50) is

$$\sum_{i=1}^m (\alpha_i^k - \langle a_i^k, y \rangle) \leq f(y) - \varphi_0 \quad \text{for all } y \in Q \cap \text{dom } f. \tag{53}$$

Since $\alpha_i^k - \langle a_i^k, y \rangle \geq 0$ for all i and for all $y \in Q \cap \text{dom } f$, we see that

$$\alpha_i^k - \langle a_i^k, y \rangle \leq f(y) - \varphi_0 \quad \text{for all } i \text{ and for all } y \in Q \cap \text{dom } f. \tag{54}$$

Observe that the derivation of (51) does not depend on δ and could have been reached at even if instead of $S_{Q_{i(k)}}$ in 3.3.2 of Algorithm 8 we would have taken any hyperplane that separates x^k from $Q_{i(k)}$. Therefore, (51) shows that $x^k \in G$, defined by (18), for all $k \geq 0$. Thus, δ is well defined.

Step 3: This step is divided into three cases.

Step 3(i): Assume that $i \in I_1$ and $\langle b^i, y \rangle - \beta_i = 0$ for all $y \in Q$. In this case we claim that

$$\langle a_i^k, y \rangle - \alpha_i^k = 0 \quad \text{for all } k \geq 0 \text{ and for all } y \in Q. \tag{55}$$

To show that (55) holds, recall that for $i \in I_1$, the functions q_i are affine functions, i.e.,

$$q_i(x) = \langle b^i, x \rangle - \beta_i, \quad \partial f(x) = \{b^i\}. \tag{56}$$

By Lemma 12, there exist nonnegative numbers γ_k such that

$$a_i^k = \gamma_k b^i, \quad \alpha_i^k = \gamma_k \beta_i. \tag{57}$$

If $\gamma_k > 0$, then

$$b^i = \frac{a_i^k}{\gamma_k}, \quad \beta_i = \frac{\alpha_i^k}{\gamma_k}. \tag{58}$$

Substituting b^i and β_i of (58) in $\langle b^i, x \rangle - \beta_i = 0$, we get that (55) holds, as claimed. If $\gamma_k = 0$ then $a_i^k = 0$ and $\alpha_i^k = 0$ and (55) holds too.

Step 3(ii): Assume that $i \in I_1$ and $\langle b^i, y \rangle - \beta_i \neq 0$ for some $y \in Q$. In this case the sequences $\{a_i^k\}_{k \geq 0}$ are bounded for all i . To show this, let $\tilde{y} \in Q \cap \text{dom } f$ satisfy $\beta_i - \langle b^i, \tilde{y} \rangle = \varepsilon > 0$. Using (54) with $y = \tilde{y}$ we have

$$\alpha_i^k - \langle a_i^k, \tilde{y} \rangle \leq f(\tilde{y}) - \varphi_0, \tag{59}$$

and by (57) we obtain

$$\alpha_i^k - \langle a_i^k, \tilde{y} \rangle = \gamma_k (\beta_i - \langle b^i, \tilde{y} \rangle) = \gamma_k \varepsilon. \tag{60}$$

Hence, by (59) and (60), $\gamma_k \varepsilon \leq f(\tilde{y}) - \varphi_0$, which means that the numbers γ_k are bounded by $(f(\tilde{y}) - \varphi_0)/\varepsilon$, so, by (57), the sequences $\{a_i^k\}_{k \geq 0}$ are indeed bounded.

Step 3(iii): Assume that $i \in I_2$. For all vectors t^i , generated in Step 3.3.2 of Algorithm 8, we have

$$\theta_i \geq \langle t^i, y \rangle \quad \text{for all } y \in Q_i. \tag{61}$$

Since \bar{y} , the existence of which is assumed in Assumption A5, is an interior point of Q_i , there exists a ball $B(\bar{y}, \epsilon_i)$ contained in Q_i . Therefore, $y = \bar{y} + \epsilon_i t^i / \|t^i\| \in B(\bar{y}, \epsilon_i) \subset Q_i$. Note that $t^i \neq 0$ by (24) and Definition 1.

Hence

$$\theta_i - \langle t^i, \bar{y} \rangle \geq \langle t^i, y \rangle - \langle t^i, \bar{y} \rangle = \langle t^i, y - \bar{y} \rangle = \epsilon_i \|t^i\|. \tag{62}$$

Using Lemma 12, we have

$$\alpha_i^k - \langle a_i^k, \bar{y} \rangle = \sum_{j=1}^r \gamma_j (\theta_j \langle t^j, y^j \rangle - \langle t^j, y^j, \bar{y} \rangle) \geq \epsilon_i \sum_{j=1}^r \gamma_j \|t^j(y^j)\|. \tag{63}$$

From (54) and the last inequality we obtain

$$f(\bar{y}) - \varphi_0 \geq \epsilon_i \sum_{j=1}^r \gamma_j \|t^j(y^j)\|, \tag{64}$$

which means that $\sum_{j=1}^r \gamma_j \|t^j(y^j)\|$ is bounded by $(f(\bar{y}) - \varphi_0)/\epsilon_i$.

From this fact and (34) we have

$$\|a_i^k\| = \left\| \sum_{j=1}^r \gamma_j t^i (y^j) \right\| \leq \sum_{j=1}^r \gamma_j \|t^i (y^j)\| \leq (f(\bar{y}) - \varphi_0) / \epsilon_i, \tag{65}$$

which proves that the sequences $\{a_i^k\}_{k \geq 0}$ are bounded when $i \in I_2$.

Step 4: From Step 2 we know that the sequence $\{x^k\}_{k \geq 0}$ is bounded, so it must have cluster points. Choose a convergent subsequence $\{x^{k_t}\}_{t \geq 0}$ of $\{x^k\}_{k \geq 0}$ such that $i(k_t) = 1$. Let $\{x^{k_t}\}$ converge to some point x^* . Since $D_f(x^{k_t+1}, x^{k_t}) \rightarrow 0$ as $k \rightarrow \infty$, the definition of Bregman functions (see [40]) implies that $\{x^{k_t+1}\}_{t \geq 0}$ converges to the point x^* . Repeating this, we get that all the subsequences $\{x^{k_t+1}\}_{t \geq 0}$, $\{x^{k_t+2}\}_{t \geq 0}, \dots, \{x^{k_t+T}\}_{t \geq 0}$ converge to x^* , where T is the almost cyclic control constant. Let

$$W = \bigcup_{t=1}^{\infty} \bigcup_{j=0}^T \{k_t + j\}, \tag{66}$$

i.e., W is the union of the indices belonging to all of the above sequences. It is clear that the sequence $\{x^k\}_{k \in W}$ also converges to x^* . Let us show next that $x^* \in Q$. If Step 3.3.1 appears infinitely many times for $\{x^{k_t}\}_{t \geq 0}$, that is, $q_1(x^{k_t}) \leq 0$ for some subsequence of $\{x^{k_t}\}_{t \geq 0}$, then $q_1(x^*) = \lim_{t \rightarrow \infty} q_1(x^{k_t}) \leq 0$. Hence we have $x^* \in Q_1$ by the definition of Q_1 . If Step 3.3.1 appears a finite number of times, then Step 3.3.2 appears infinitely many times. For Step 3.3.2 we know that $x^{k_t+1} \notin \text{int}(B(x^{k_t}, \delta q_1(x^{k_t})))$ because x^{k_t+1} is the Bregman projection of z^{k_t} onto the intersection of the two half-spaces $L_{i(k_t)}^{k_t}$ and $S_{Q_{i(k_t)}}$. Therefore,

$$\|x^{k_t+1} - x^{k_t}\| \geq \delta q_1(x^{k_t}), \tag{67}$$

and since $\lim_{t \rightarrow \infty} \|x^{k_t+1} - x^{k_t}\| = 0$ (because $\lim_{t \rightarrow \infty} x^{k_t+1} = \lim_{t \rightarrow \infty} x^{k_t}$) and $\delta > 0$, we have $\lim_{t \rightarrow \infty} q_1(x^{k_t}) \leq 0$. Hence $q_1(x^*) = \lim_{t \rightarrow \infty} q_1(x^{k_t}) \leq 0$, so $x^* \in Q_1$. Choosing a subsequence $\{x^{k_t+j_t}\}$ ($0 < j_t < T$) with $i(k_t + j_t) = 2$ which converges to the same point x^* , we see that $x^* \in Q_2$. Repeating this argument for $1 \leq i \leq m$ we obtain $x^* \in Q$.

Step 5: We now show that (43) holds. Take some $i, 1 \leq i \leq m$. If $\alpha_i^k - \langle a_i^k, y \rangle = 0$ for all $y \in Q$ and for all k , then (43) is true for this i . Otherwise, we know that $\{a_i^k\}_{k \geq 0}$ is bounded (by Step 3).

If $k \in W$, then W contains a set $W_k = \{p, p + 1, \dots, p + T\}$ containing k . We know that the set $\{i(p), i(p + 1), \dots, i(p + T - 1)\}$ contains i (by the choice of the almost cyclic control index). Let $r \in W_k, r \leq p + T - 1$, be the nearest integer to k such that $i(r) = i$. We distinguish between two cases according to the values of r and k .

Step 5(i): Assume that $r < k$. In this case $(r + 1) \in W_k$ (from the definition of W_k) and $a_i^k = a_i^{r+1}, \alpha_i^k = \alpha_i^{r+1}$ since there is no change in a_i^ℓ and α_i^ℓ for $r + 1 \leq \ell \leq k$ (by

(38)). Using the last two equations and (45), we obtain

$$\begin{aligned} \alpha_i^k - \langle a_i^k, x^* \rangle &= \alpha_i^{r+1} - \langle a_i^{r+1}, x^* \rangle \\ &= \alpha_{i(r)}^{r+1} - \langle a_i^{r+1}, x^{r+1} \rangle + \langle a_i^{r+1}, x^{r+1} - x^* \rangle \\ &= \langle a_{i(r)}^{r+1}, x^{r+1} - x^* \rangle. \end{aligned} \tag{68}$$

By definition of L_i^k and the last equation, we get

$$0 \leq \alpha_i^k - \langle a_i^k, x^* \rangle \leq \|a_i^{r+1}\| \|x^{r+1} - x^*\|. \tag{69}$$

Step 5(ii): Assume that $r \geq k$. In this case $a_i^k = a_i^r$ and $\alpha_i^k = \alpha_i^r$ because a_i^r and α_i^r do not change for $k \leq \ell \leq r$ (by 3.4, in the iterative step of Algorithm 8). Hence by the last two equalities and (49),

$$\begin{aligned} \alpha_i^k - \langle a_i^k, x^* \rangle &= \alpha_i^r - \langle a_i^r, x^* \rangle \\ &= \alpha_{i(r)}^r - \langle a_{i(r)}^r, x^{r+1} \rangle + \langle a_i^r, x^{r+1} - x^* \rangle \\ &\leq \varphi_{r+1} - \varphi_r + \langle a_i^r, x^{r+1} - x^* \rangle. \end{aligned} \tag{70}$$

Therefore

$$0 \leq \alpha_i^k - \langle a_i^k, x^* \rangle \leq \|a_i^k\| \|x^{r+1} - x^*\| + \varphi_{r+1} - \varphi_r. \tag{71}$$

Since r tends to infinity together with k , $r + 1 \in W$, the sequences $\{a_i^k\}_{k \geq 0}$ are bounded and $\{\varphi_k\}_{k \geq 0}$ converges, we see that (69) and (71) imply (43). Since (43) holds for all i , we have

$$\lim_{k \rightarrow \infty, k \in W} \sum_{i=1}^m (\alpha_i^k - \langle a_i^k, x^* \rangle) = 0. \tag{72}$$

Applying (50) with $y = x^*$, we get

$$D_f(x^*, x^k) = f(x^*) - \varphi_k + \sum_{i=1}^m (\langle a_i^k, x^* \rangle - \alpha_i^k). \tag{73}$$

By the definition of a Bregman function (see [14, Definition 2.1.1]), one has

$$\lim_{k \rightarrow \infty, k \in W} D_f(x^*, x^k) = 0. \tag{74}$$

Hence (72)–(74) imply that the subsequence $\{\varphi_k\}_{k \geq 0}$, $k \in W$, tends to $f(x^*)$, and since $\lim_{k \rightarrow \infty} \varphi_k$ exists,

$$\lim_{k \rightarrow \infty} \varphi_k = f(x^*). \tag{75}$$

Since, by (52), $\lim_{k \rightarrow \infty} \varphi_k \leq \min\{f(x) \mid x \in Q\}$, we obtain

$$f(x^*) = \min\{f(x) \mid x \in Q\}. \tag{76}$$

From the fact that f is strictly convex (because it is a Bregman function) and has a unique minimum in Q , it follows that the whole sequence $\{x^k\}_{k \geq 0}$ converges to x^* , and the proof is complete. \square

5. Particular cases

It is natural to ask, but quite complicated to answer, in what situations all assumptions made in the previous sections hold. We have no simple answer to this question at this time except for the particular cases discussed next. In these cases the choice of the half-spaces is constructively given. First, we show that the δ -SHS $S_{Q_{i(k)}}(x^k)$ can be defined via some subgradients at the current point x^k . The second special case deals with the construction of these δ -SHS's, via interior points in the convex sets (using the assumption that in each set we know an interior point). The idea of generating such a case is based on the (δ, η) -algorithm for convex inequalities with interior points (see, e.g., [14, Algorithm 5.5.3]).

5.1. Construction of $S_{Q_{i(k)}}(x^k)$ via subgradient vectors

We show here a specific choice of the δ -SHSs $S_{Q_{i(k)}}(x^k)$, made by constructing each of the δ -SHSs via subgradients. In this case we use underrelaxation parameters to define $S_{Q_{i(k)}}(x^k)$.

Let $\{\beta_k\}_{k \geq 0}$ be an infinite sequence of underrelaxation parameters such that $0 < \varepsilon \leq \beta_k \leq 1$ for all $k \geq 0$, with some arbitrarily small given ε . Let $v^{i(k)} = v^{i(k)}(x^k)$ be a subgradient of $q_{i(k)}$ at x^k .

Theorem 14. *Assume that $\partial q_i(G)$ is bounded for any bounded subset $G \subseteq \text{dom } q_i$, for all $i = 1, 2, \dots, m$. If in Algorithm 8 one uses, for all $k \geq 0$,*

$$0 \neq t^{i(k)}(x^k) = v^{i(k)} \in \partial q_{i(k)}(x^k) \tag{77}$$

and

$$\theta_{i(k)} = \langle v^{i(k)}, x^k \rangle - \beta_k q_{i(k)}(x^k) \tag{78}$$

to construct $S_{Q_{i(k)}}(x^k)$ by (24) whenever $q_{i(k)}(x^k) > 0$, then $S_{Q_{i(k)}}(x^k)$ is a δ -SHS.

Proof. Let $S_{Q_{i(k)}}(x^k)$ be the half-space defined by (77) and (78), i.e.,

$$S_{Q_{i(k)}}(x^k) = \{x \in R^n \mid \langle v^{i(k)}, x - x^k \rangle \leq -\beta_k q_{i(k)}(x^k)\}. \tag{79}$$

In order to conclude that the half-space $S_{Q_{i(k)}}(x^k)$ is a δ -SHS, we must show that

$$S_{Q_{i(k)}}(x^k) \cap \text{int } B(x^k, \delta q_{i(k)}(x^k)) = \emptyset \tag{80}$$

and, by Definition 1, that

$$S_{Q_{i(k)}}(x^k) \supseteq Q_{i(k)}. \tag{81}$$

First we show that (80) holds. In Step 3.3.2 of Algorithm 8, x^{k+1} is the Bregman projection of z^k onto the intersection $L_{i(k)}^k \cap S_{Q_{i(k)}}(x^k)$. This implies that $x^{k+1} \in S_{Q_{i(k)}}(x^k)$, that is,

$$\langle v^{i(k)}, x^{k+1} - x^k \rangle \leq -\beta_k q_{i(k)}(x^k). \tag{82}$$

It follows that

$$\beta_k q_{i(k)}(x^k) \leq |\langle v^{i(k)}, x^k - x^{k+1} \rangle| \leq \|v^{i(k)}\| \|x^k - x^{k+1}\|. \tag{83}$$

According to the comment in the last sentence of Step 2 of the proof, x^k belongs to the set G , defined by (18). Using the assumption on the boundedness of the subgradients, we have $\|v^{i(k)}\| \leq M$. Hence

$$\beta_k q_{i(k)}(x^k) \leq M \|x^k - x^{k+1}\|, \tag{84}$$

which implies that

$$\frac{1}{M} \beta_k q_{i(k)}(x^k) \leq \|x^k - x^{k+1}\|. \tag{85}$$

Taking any

$$\delta \leq \inf \left\{ \frac{1}{M} \beta_k \mid k \geq 0 \right\} = \frac{\varepsilon}{M}, \tag{86}$$

we get

$$x^{k+1} \notin \text{int } B(x^k, \delta q_{i(k)}(x^k)) \quad \text{for all } k \geq 0, \tag{87}$$

which implies that (80) is true. We now show that (81) also holds. Let $x \in Q_{i(k)}$, i.e., $q_{i(k)}(x) \leq 0$. By the *subgradient inequality* we have

$$q_{i(k)}(x) - q_{i(k)}(x^k) \geq \langle v^{i(k)}, x - x^k \rangle. \tag{88}$$

Thus

$$-q_{i(k)}(x^k) \geq \langle v^{i(k)}, x - x^k \rangle. \tag{89}$$

Since $q_{i(k)}(x^k) > 0$, both sides of (89) are negative. Hence

$$-\beta_k q_{i(k)}(x^k) \geq \langle v^{i(k)}, x - x^k \rangle, \tag{90}$$

i.e., $x \in S_{Q_{i(k)}}(x^k)$ by (79), which implies that (81) does indeed hold. This completes the proof. \square




Fig. 3. Geometric description of the construction of $S_{Q_{i(k)}}(x^k)$ via interior points in the sets.

5.2. Construction of $S_{Q_{i(k)}}(x^k)$ via interior points in the sets

In Case 3.3.2 of Algorithm 8 we can construct the half-spaces $S_{Q_{i(k)}}(x^k)$ for (24) by still another method.

Assumption C. There are m given interior points $y^i \in \text{int } Q_i, 1 \leq i \leq m$.

Method for the construction of the half-spaces $S_{Q_{i(k)}}(x^k)$ by (24):

If $q_{i(k)}(x^k) > 0$ (i.e., we are in Case 3.3.2 of Algorithm 8), choose some $0 \leq h \leq 1$, define

$$\bar{x}(h) = hy^{i(k)} + (1 - h)x^k, \tag{91}$$

and solve the nonlinear equation

$$q_{i(k)}(\bar{x}(h)) = 0. \tag{92}$$

Denote by h_k the smallest value of h for which $\bar{x}(h)$ solves (92) and set

$$\bar{x}^k = \bar{x}(h_k). \tag{93}$$

Then calculate a subgradient

$$t^{i(k)} \in \partial q_{i(k)}(\bar{x}^k) \tag{94}$$

(if $q_{i(k)}$ is differentiable at \bar{x}^k) then $t^{i(k)} = \nabla q_{i(k)}(\bar{x}^k)$) and

$$\theta_{i(k)} = \langle t^{i(k)}, \bar{x}^k \rangle, \tag{95}$$

and define $S_{Q_{i(k)}}(x^k)$ by

$$S_{Q_{i(k)}}(x^k) = \{x \in \mathbb{R}^n \mid \langle t^{i(k)}, x \rangle \leq \langle t^{i(k)}, \bar{x}^k \rangle\}. \tag{96}$$

See Fig. 3 for a geometric description of the construction of $S_{Q_{i(k)}}(x^k)$ via interior points in the sets.

Theorem 15. *Under Assumption C, whenever $q_{i(k)}(x^k) > 0$ in Algorithm 8 and we use the method described above to construct $S_{Q_{i(k)}}(x^k)$, then there exists a $\delta > 0$ such that $S_{Q_{i(k)}}(x^k)$ is a δ -SHS.*

Proof. We first show that (81) holds. If $x \in Q_{i(k)}$, then $q_{i(k)}(x) \leq 0$. Using the subgradient inequality

$$\langle t^{i(k)}, x - \bar{x}^k \rangle \leq q_{i(k)}(x) - q_{i(k)}(\bar{x}^k), \tag{97}$$

and the fact that $q_{i(k)}(x) \leq 0$ and $q_{i(k)}(\bar{x}^k) = 0$, we obtain

$$\langle t^{i(k)}, x - \bar{x}^k \rangle \leq 0. \tag{98}$$

In other words,

$$\langle t^{i(k)}, x \rangle \leq \langle t^{i(k)}, \bar{x}^k \rangle. \tag{99}$$

Thus $x \in S_{Q_{i(k)}}(x^k)$, by (96). We show now that (80) holds. Let

$$H_{i(k)} = \{x \in \mathbb{R}^n \mid \langle t^{i(k)}, x \rangle = \langle t^{i(k)}, \bar{x}^k \rangle\}. \tag{100}$$

Since $x^k \notin Q_{i(k)}$ (otherwise we do not use (96)), we have, by a simple geometric consideration,

$$\|P_{H_{i(k)}}(x^k) - x^k\| = \frac{\|x^k - \bar{x}^k\| \|y^{i(k)} - P_{H_{i(k)}}(y^{i(k)})\|}{\|y^{i(k)} - \bar{x}^k\|}, \tag{101}$$

where $P_{H_{i(k)}}(x^k)$ is the orthogonal projection of x^k onto $H_{i(k)}$ (see also [14, Fig. 5.8]). Since $\{x^k\}_{k \geq 0}$ is bounded (see the comment in the last sentence of Step 2), (101) implies that there is a positive M such that for all $k \geq 0$,

$$\|y^{i(k)} - \bar{x}^k\| \leq M. \tag{102}$$

By Bauschke and Borwein [4, Proposition 7.8 and Corollary 7.9], $q_{i(k)}$ is Lipschitz continuous. Hence there is a positive L such that

$$|q_{i(k)}(x^k) - q_{i(k)}(\bar{x}^k)| \leq L \|x^k - \bar{x}^k\|. \tag{103}$$

Since $q_{i(k)}(\bar{x}^k) = 0$ by (92) and (93), we obtain

$$q_{i(k)}(x^k) \leq L \|x^k - \bar{x}^k\|. \quad (104)$$

We also have

$$\|y^{i(k)} - P_{H_{i(k)}}(y^{i(k)})\| \geq d(y^{i(k)}, bdQ_{i(k)}) \geq d > 0, \quad (105)$$

where

$$d := \min\{d(y^i, bdQ_i) \mid 1 \leq i \leq m\}. \quad (106)$$

It follows from (101), (102), (104) and (105) that

$$\|P_{H_{i(k)}}(x^k) - x^k\| \geq \frac{q_{i(k)}(x^k)d}{ML}. \quad (107)$$

Let $x^{k+1} \in S_{Q_{i(k)}}(x^k)$. Then we also obtain

$$\|x^{k+1} - x^k\| \geq \|P_{H_{i(k)}}(x^k) - x^k\| \geq \frac{q_{i(k)}(x^k)d}{ML}. \quad (108)$$

Taking $\delta \leq \frac{d}{ML}$, we have that

$$x^{k+1} \notin \text{int } B(x^k, \delta q_{i(k)}(x^k)) \quad \text{for all } k \geq 0. \quad (109)$$

This completes the proof of Theorem 15. \square

Acknowledgments

We thank an anonymous referee for his insightful and constructive report which led to a much improved paper. Part of this work was done at the Center for Computational Mathematics and Scientific Computation (CCMSC) at the University of Haifa and was supported by research Grant 592/00 from the Israel Science Foundation founded by the Israel Academy of Sciences and Humanities. S. Reich's work was also partially supported by the Fund for the Promotion of Research at the Technion and by the Technion VPR Fund.

References

- [1] R. Aharoni, A. Berman, Y. Censor, An interior points algorithm for the convex feasibility problem, *Adv. Appl. Math.* 4 (1983) 479–489.
- [2] H.H. Bauschke, The approximation of fixed points of compositions of nonexpansive mappings in Hilbert space, *J. Math. Anal. Appl.* 202 (1996) 150–159.
- [3] H.H. Bauschke, J.M. Borwein, Dykstra's alternating projection algorithm for two sets, *J. Approx. Theory* 79 (1994) 418–443.

- [4] H.H. Bauschke, J.M. Borwein, On projection algorithms for solving convex feasibility problems, *SIAM Rev.* 38 (1996) 367–426.
- [5] H.H. Bauschke, J.M. Borwein, Legendre functions and the method of random Bregman projections, *J. Convex Anal.* 4 (1997) 27–67.
- [6] H.H. Bauschke, P.L. Combettes, A weak-to-strong convergence principle for Fejér-monotone methods in Hilbert spaces, *Math. Oper. Res.* 26 (2001) 248–264.
- [7] H.H. Bauschke, P.L. Combettes, Construction of best Bregman approximations in reflexive Banach spaces, *Proc. Amer. Math. Soc.* 131 (2003) 3757–3766.
- [8] H.H. Bauschke, A.S. Lewis, Dykstra’s algorithm with Bregman projections: a convergence proof, *Optimization* 48 (2000) 409–427.
- [9] J.P. Boyle, R.L. Dykstra, A method of finding projections onto the intersection of convex sets in Hilbert spaces, in: *Advances in Order Restricted Statistical Inference, Lecture Notes in Statistics*, Vol. 37, Springer, Berlin, Germany, 1986, pp. 28–47.
- [10] L.M. Bregman, The relaxation method of finding the common point of convex sets and its application to the solution of problems in convex programming, *USSR Comput. Math. Math. Phys.* 7 (1967) 200–217.
- [11] L.M. Bregman, Y. Censor, S. Reich, Dykstra’s algorithm as the nonlinear extension of Bregman’s optimization method, *J. Convex Analysis* 6 (1999) 319–333.
- [12] Y. Censor, A. Lent, An iterative row-action method for interval convex programming, *J. Optim. Theory Appl.* 34 (1981) 321–353.
- [13] Y. Censor, S. Reich, The Dykstra algorithm with Bregman projections, *Comm. Appl. Anal.* 2 (1998) 407–419.
- [14] Y. Censor, S.A. Zenios, *Parallel Optimization: Theory, Algorithms, and Applications*, Oxford University Press, New York, NY, USA, 1997.
- [15] P.L. Combettes, Signal recovery by best feasible approximation, *IEEE Trans. Image Process.* IP-2 (1993) 269–271.
- [16] P.L. Combettes, Strong convergence of block-iterative outer approximation methods for convex optimization, *SIAM J. Control Optim.* 38 (2000) 538–565.
- [17] G. Crombez, Finding projections onto the intersection of convex sets in Hilbert spaces, *Numer. Funct. Anal. Optim.* 16 (1995) 637–652.
- [18] D.A. D’Esopo, A convex programming procedure, *Naval Res. Logistics Quart.* 6 (1959) 33–42.
- [19] F. Deutsch, *Best Approximation in Inner Product Spaces*, Springer, New York, NY, USA, 2001.
- [20] F. Deutsch, H. Hundal, The rate of convergence of Dykstra’s cyclic projections algorithm: the polyhedral case, *Numer. Funct. Anal. Optim.* 15 (1994) 537–565.
- [21] R.L. Dykstra, An algorithm for restricted least squares regression, *J. Amer. Statist. Assoc.* 78 (1983) 837–842.
- [22] R.L. Dykstra, An iterative procedure for obtaining I -projections onto the intersection of convex sets, *Ann. Probab.* 13 (1985) 975–984.
- [23] M. Fukushima, An outer approximation algorithm for solving general convex programs, *Oper. Res.* 31 (1983) 101–113.
- [24] M. Fukushima, A relaxed projection method for variational inequalities, *Math. Programming* 35 (1986) 58–70.
- [25] N. Gaffke, R. Mathar, A cyclic projection algorithm via duality, *Metrika* 36 (1989) 29–54.
- [26] S.-P. Han, A successive projection method, *Math. Programming* 40 (1988) 1–14.
- [27] S.-P. Han, G. Lou, A parallel algorithm for a class of convex programs, *SIAM J. Control Optim.* 26 (1988) 345–355.
- [28] Y. Haugazeau, *Sur les Inéquations Variationnelles et la Minimisation de Fonctionnelles Convexes*, Ph.D. Thesis, Université de Paris, Paris, France, 1968.
- [29] C. Hildreth, A quadratic programming procedure, *Naval Res. Logistics Quart.* 4 (1957) 79–85 (Erratum, *ibid.*, p. 361).
- [30] H. Hundal, F. Deutsch, Two generalizations of Dykstra’s cyclic projections algorithm, *Math. Programming* 77 (1997) 335–355.

- [31] A.N. Iusem, A.R. De Pierro, A simultaneous iterative method for computing projections on polyhedra, *SIAM J. Control Optim.* 25 (1987) 231–243.
- [32] A.N. Iusem, A.R. De Pierro, On the convergence of Han’s method for convex programming with quadratic objective, *Math. Programming* 52 (1991) 265–284.
- [33] A.N. Iusem, B.F. Svaiter, A row-action method for convex programming, *Math. Programming* 64 (1994) 149–171.
- [34] A.N. Iusem, B.F. Svaiter, Primal-dual row-action method for convex programming, *J. Optim. Theory Appl.* 86 (1995) 73–112.
- [35] A. Lent, Y. Censor, Extensions of Hildreth’s row-action method for quadratic programming, *SIAM J. Control Optim.* 18 (1980) 444–454.
- [36] D.G. Luenberger, *Optimization by Vector Space Methods*, Wiley, New York, NY, USA, 1969.
- [37] G. Pierra, Eclatement de contraintes en parallèle pour la minimisation d’une forme quadratique, in: *Lecture Notes in Computer Science*, Vol. 41, Springer, New York, NY, USA, 1976, pp. 96–115.
- [38] G. Pierra, Decomposition through formalization in a product space, *Math. Programming* 28 (1984) 96–115.
- [39] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, NJ, USA, 1970.
- [40] M.V. Solodov, B.F. Svaiter, An inexact hybrid generalized proximal point algorithm and some new results on the theory of Bregman functions, *Math. Oper. Res.* 25 (2000) 214–230.
- [41] P. Tseng, Dual coordinate ascent methods for non-strictly convex minimization, *Math. Programming* 59 (1993) 231–247.