



ELSEVIER

Available at

www.ElsevierMathematics.com

POWERED BY SCIENCE @ DIRECT®

Journal of Approximation Theory 126 (2004) 198–217

JOURNAL OF
Approximation
Theory

http://www.elsevier.com/locate/jat

Ratio asymptotics and weak asymptotic measures for orthogonal polynomials on the real line

Barry Simon¹

*Division of Physics, Mathematics and Astronomy, California Institute of Technology,
Pasadena, CA 91125, USA*

Received 10 July 2003; accepted in revised form 1 December 2003

Communicated by Leonid Golinskii

Abstract

We study ratio asymptotics, that is, existence of the limit of $P_{n+1}(z)/P_n(z)$ ($P_n =$ monic orthogonal polynomial) and the existence of weak limits of $p_n^2 d\mu$ ($p_n = P_n/||P_n||$) as $n \rightarrow \infty$ for orthogonal polynomials on the real line. We show existence of ratio asymptotics at a single z_0 with $\text{Im}(z_0) \neq 0$ implies $d\mu$ is in a Nevai class (i.e., $a_n \rightarrow a$ and $b_n \rightarrow b$ where a_n, b_n are the off-diagonal and diagonal Jacobi parameters). For μ 's with bounded support, we prove $p_n^2 d\mu$ has a weak limit if and only if $\lim b_n, \lim a_{2n},$ and $\lim a_{2n+1}$ all exist. In both cases, we write down the limits explicitly.

© 2003 Elsevier Inc. All rights reserved.

MSC: 42C05; 47B36; 81Q10

Keywords: Polynomials; Asymptotics; Jacobi matrices

1. Introduction

In [8], Khrushchev asked two questions about orthogonal polynomials on the unit circle [5,14,15] and found the following remarkable theorems in terms of the monic orthogonal polynomials, Φ_n ; the orthonormal polynomials, $\varphi_n = \Phi_n/||\Phi_n||_{L^2}$; and the Verblunsky coefficients, $\alpha_n = -\overline{\Phi_{n+1}(0)}$.

E-mail address: bsimon@caltech.edu.

¹Supported in part by NSF Grant DMS-0140592.

Theorem A. $\Phi_{n+1}^*(z)/\Phi_n^*(z)$ has a limit uniformly in z over compact subsets of \mathbb{D} if and only if either

- (i) For $\ell = 1, 2, \dots, \lim_{n \rightarrow \infty} \alpha_{n+\ell} \alpha_n = 0$, or
- (ii) There is $a \in (0, 1]$ and $\lambda \in \partial\mathbb{D}$ so that $\lim_{n \rightarrow \infty} |\alpha_n| = a$, $\lim_{n \rightarrow \infty} \bar{\alpha}_{n+1} \alpha_n = a^2 \lambda$.

Theorem B. $|\varphi_n|^2 d\mu$ has a weak limit if and only if either

- (i) For $\ell = 1, 2, \dots, \lim_{n \rightarrow \infty} \alpha_{n+\ell} \alpha_n = 0$, or
- (ii) There exist $a, a' \in (0, 1]$, $\lambda \in \partial\mathbb{D}$, and integers $k \geq 1$ and $\ell \in \{0, 1, \dots, k - 1\}$ so that

$$\lim_{n \rightarrow \infty} |\alpha_{2nk+\ell+j}| = \begin{cases} a & \text{if } j = 0, \\ a' & \text{if } j = k, \\ 0 & \text{if } j = 1, \dots, k - 1, k + 1, \dots, 2k - 1, \end{cases}$$

$$\lim_{n \rightarrow \infty} \bar{\alpha}_{2nk+\ell} \alpha_{2nk+k+\ell} = a a' \lambda.$$

Khrushchev [8] also describes explicitly the limits in both cases. Our goal in this paper is to find the analogs of these theorems for orthogonal polynomials on the real line. The answers and proofs are much simpler—the methods of Khrushchev which depend heavily on Schur functions do not seem to extend, nor does mapping bounded intervals on \mathbb{R} to $\partial\mathbb{D}$ (as in [15, Section 11.5]) seem to allow direct transfer.

Before stating our results, let us set up notation. Given a measure $d\mu$ on \mathbb{R} with $\int x^{2n} d\mu < \infty$ for all n , we let $P_n(x)$ be the monic orthogonal and $p_n(x)$ the orthonormal polynomials. To define them, we suppose henceforth that $d\mu$ is nontrivial, that is, not supported on a finite set, and we will also assume throughout that $\mu(\mathbb{R}) = 1$. Thus P_n is determined by $P_n(x) = x^n +$ lower order and $\int x^j P_n(x) d\mu(x) = 0$ for $j = 0, 1, \dots, n - 1$. $p_n = P_n/\|P_n\|$ where $\|\cdot\|$ is the $L^2(\mathbb{R}, d\mu)$ norm.

It is well known [15] that the P_n 's obey a three-term recursion relation. There are $b_j \in \mathbb{R}$ and $a_j \in (0, \infty)$ so that

$$xP_n(x) = P_{n+1}(x) + b_{n+1}P_n(x) + a_n^2 P_{n-1}(x). \tag{1.1}$$

Our indexing of b_1, b_2, \dots and a_1, a_2, \dots is not common—often the labelling starts at b_0 and a_0 . We take this convention from [9] for reasons explained there. Eq. (1.1) implies inductively that

$$\|P_n\| = a_n \dots a_1 \tag{1.2}$$

and then that the p_n obey the recursion relation

$$xp_n(x) = a_{n+1}p_{n+1}(x) + b_{n+1}p_n(x) + a_n p_{n-1}(x). \tag{1.3}$$

In turn, (1.3) suggests we study the Jacobi matrix

$$J = \begin{pmatrix} b_1 & a_1 & 0 & 0 & \cdots \\ a_1 & b_2 & a_2 & 0 & \cdots \\ 0 & a_2 & b_3 & a_3 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \tag{1.4}$$

Since $\{p_n\}_{n=0}^\infty$ is an orthonormal set,

$$U : \sum_{j=0}^N v_j p_j \rightarrow \begin{pmatrix} v_1 \\ v_2 \\ \vdots \end{pmatrix}$$

is a unitary map of the closed span, S , of the p 's to $\ell^2(\mathbb{Z}_+)$ ($\mathbb{Z}_+ = \{1, 2, \dots\}$) and for $v \in S_0 = \text{span of } p$'s, we have $U^{-1}JUv = (\text{multiplication by } x) v$.

In case the moment problem is determinantal [1,13], J is self-adjoint, and $d\mu$ is just the spectral measure for J and vector $\delta = (1, 0, \dots)$.

We can now state our main results.

Theorem 1 (\equiv Theorems 2.1 and 2.2). *Suppose that at a single $z_0 \in \mathbb{C} \setminus \mathbb{R}$, we have*

$$\lim_{n \rightarrow \infty} \frac{P_{n+1}(z)}{P_n(z)} = f(z) \tag{1.5}$$

for $z = z_0$. Then, for some $a \in [0, \infty)$ and $b \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} a_n = a, \quad \lim_{n \rightarrow \infty} b_n = b. \tag{1.6}$$

Conversely, if (1.6) holds and $\text{spectrum}(J)$ is the spectrum of the operator J , then (1.5) holds for all $z \in \mathbb{C} \setminus \text{spectrum}(J)$ and

$$f(z) = \frac{(z - b) + \sqrt{(z - b)^2 - 4a^2}}{2}, \tag{1.7}$$

where the branch of the square root is taken with $\sqrt{\dots} = z + O(1/z)$ near $z = \infty$.

Theorem 2 (\equiv Theorems 3.1 and 3.2). *Let*

$$d\mu_n = p_n^2(x) d\mu(x). \tag{1.8}$$

Suppose that for $\ell = 1, 2$, and 4, $\lim_{n \rightarrow \infty} \int x^\ell d\mu_n$ exists. Then for $a, c \in [0, \infty)$ and $b \in \mathbb{R}$, we have

$$\lim_{n \rightarrow \infty} b_n = b, \quad \lim_{n \rightarrow \infty} a_{2n} = a, \quad \lim_{n \rightarrow \infty} a_{2n+1} = c. \tag{1.9}$$

Conversely, if (1.9) holds, the $d\mu_n$ have supports lying in a fixed bounded interval and there is a measure $d\rho_{b;a,c}$ so that for any continuous f on \mathbb{R} (including $f(x) = x^\ell$), we have

$$\int f(x) d\mu_n(x) \rightarrow \int f(x) d\rho(x). \tag{1.10}$$

$d\rho$ is a function of a, b, c only, and if $d\rho_{b;a,c} = d\rho_{b';a',c'}$, we have $b = b'$ and either $a = a'$, $c = c'$ or $a = c'$, $c = a'$.

Remark. We will show that the existence of the limit $\int x^\ell d\mu_n$ for $\ell = 2$ implies $d\mu$ has bounded support.

Theorem 1 is proven in Section 2 and Theorem 2 in Section 3. $d\rho$ is calculated in Section 5. Theorems 1 and 2 seem to be optimal in that the two pieces of real data (1.6) (i.e., a and b) correspond to one complex number $f(z_0)$ while the three moments in Theorem 2 correspond to the real numbers, a, b , and c .

There is previous work of Nevai [10] on the subjects of Theorems 1 and 2. He proved that (1.6) implies (1.5) with f given by (1.7), and conversely proved that if (1.5) holds for all $z \in \mathbb{C} \setminus \mathbb{R}$ and $f(z)$ given by (1.7), then (1.6) holds. He did not get a result depending on a single z_0 nor, more importantly, did he show that (1.7) are the only possible limits in (1.5).

Nevai [10] also proved that if $a_n \rightarrow a$ and $b_n \rightarrow b$, then $d\mu_n$ has a weak limit (he wrote down the explicit form of $d\rho_{b;a,a}$ for $a = c$, as we will in Section 5).

Barrios et al. [2] proved that if (1.5) holds at two points with f given by (1.7), then (1.6) holds. But, like Nevai, they did not address the fact that (1.7) are the only possible limits.

In [11], Nevai made a conjecture closely related to a special case of Theorem 2. Namely,

Nevai Conjecture 2.16 (Nevai [11]). *If (1.10) holds for all bounded uniformly continuous functions on \mathbb{R} with $d\rho(x) = \pi^{-1} \chi_{[-1,1]}(x)(1 - x^2)^{-1/2} dx$, then $a_n \rightarrow \frac{1}{2}$ and $b_n \rightarrow 0$.*

Corollary to Theorem 2. *If one supposes $\text{supp}(d\mu)$ is bounded, then Nevai’s conjecture holds.*

Proof. $x^\ell \upharpoonright \text{supp}(d\mu)$ is bounded, so $\int x^\ell d\mu_n$ converges for $\ell = 1, 2, 4$ to the same limit as for $a_n \equiv \frac{1}{2}$ and $b \equiv 0$. Uniqueness of the limit (and the fact that $a = c$) completes the proof. \square

Related to this is

Nevai Conjecture 2.17 (Nevai [11]). *If for some A , we have $\int_A^\infty d\mu_n \rightarrow 0$, then for every $\varepsilon > 0$, $[A + \varepsilon, \infty) \cap \text{supp}(d\mu)$ is finite.*

We mention

Weaker Nevai Conjecture 2.17. *If for some $A > 0$, $\mu_n(\{x \mid |x| > A\}) \rightarrow 0$, then $\text{supp}(d\mu)$ is bounded.*

Clearly, Nevai Conjecture 2.17 implies the weaker version. The point of this is that a positive solution of the Weaker Nevai Conjecture 2.17 plus the results of this paper would imply Nevai Conjecture 2.16.

It is a pleasure to thank Rowan Killip and Paul Nevai for cogent comments.

2. Ratio asymptotics

The two main theorems on limits of $P_{n+1}(x)/P_n(x)$ are as follows:

Theorem 2.1. *Suppose $a_n \rightarrow a \in [0, \infty)$ and $b_n \rightarrow b \in \mathbb{R}$. Then for all $z \in \mathbb{C} \setminus \text{spectrum}(J)$, we have that*

$$\lim_{n \rightarrow \infty} \frac{P_{n+1}(z)}{P_n(z)} = \frac{(z - b) + \sqrt{(z - b)^2 - 4a^2}}{2}. \tag{2.1}$$

Remark. 1. In (2.1), we take the branch of the square root with $\sqrt{\dots} \sim z$ for $|z|$ large, that is, as $z \rightarrow \infty$.

2. For $z \notin \mathbb{R}$, P_n is nonzero for all n . (2.1) for $z_0 \in \mathbb{R} \setminus \text{spectrum}(J)$ includes the fact that for z_0 fixed, $P_n(z_0) \neq 0$ for all large n .

3. One can also show that for $z \in \text{spectrum}(J) \setminus [b - 2a, b + 2a]$ so that z is an eigenvalue of J ,

$$\lim_{n \rightarrow \infty} \frac{P_{n+1}(z)}{P_n(z)} = \frac{(z - b) - \sqrt{(z - b)^2 - 4a^2}}{2}. \tag{2.2}$$

Theorem 2.2. *Suppose for one z_0 with $\text{Im } z_0 \neq 0$, $\lim_{n \rightarrow \infty} P_{n+1}(z)/P_n(z)$ exists (and is finite). Then there exists $a \in [0, \infty)$ with $b \in \mathbb{R}$ so that $a_n \rightarrow a$ and $b_n \rightarrow b$ so that (2.1) holds. In particular, the only functions that can occur as ratio asymptotics are the ones in (2.1).*

Theorem 2.1 is not new. In this generality, it is due to Nevai [10], who also proved a converse; namely, he showed that if (2.1) holds for all $z \in \mathbb{C} \setminus \mathbb{R}$, then $a_n \rightarrow a$ and $b_n \rightarrow b$. But we will sketch two proofs of Theorem 2.1 for the reader’s convenience. One uses transfer matrices and the other, operator theory.

As a preliminary, we need:

Proposition 2.3. *Let $\{x_{j,n}\}_{j=1}^n$ be the zeros of $P_n(x)$ with*

$$x_{1,n} < x_{2,n} < \dots < x_{n,n}. \tag{2.3}$$

Then

$$(i) \quad -\frac{P_{n-1}(z)}{P_n(z)} = \sum_{j=0}^n \frac{\alpha_{j,n}}{x_{j,n} - z}, \tag{2.4}$$

where $\alpha_{j,n} > 0$ and

$$\sum_{j=1}^n \alpha_{j,n} = 1. \tag{2.5}$$

(ii) If $\text{Im } z > 0$,

$$0 < -\text{Im} \left(\frac{P_{n-1}(z)}{P_n(z)} \right) \leq \frac{1}{\text{Im } z} \tag{2.6}$$

and

$$\left| \frac{P_{n-1}(z)}{P_n(z)} \right| \leq \frac{1}{\text{Im } z}. \tag{2.7}$$

(iii) If $\text{Im } z > 0$,

$$\text{Im} \left(\frac{P_{n+1}}{P_n} \right) \geq \text{Im } z. \tag{2.8}$$

Proof. (i) Since P_n is monic, $P_n(z) = \prod_{j=1}^n (z - x_{j,n})$. Since P_{n-1}/P_n has simple poles and goes to zero at infinity, (2.4) holds for some $\alpha_{j,n}$. Multiplying by $x_{i,n} - z$ and taking z to $x_{i,n}$, we find

$$\alpha_{j,n} = \frac{\prod_{\ell=1}^{n-1} (x_{j,n} - x_{\ell,n-1})}{\prod_{\ell=1; \ell \neq j}^n (z_{j,n} - x_{\ell,n})}. \tag{2.9}$$

For $j = n$, all factors in the right are positive. Since zeros of P_{n-1} and P_n interlace as we decrease j by one, both numerator and denominator each pick up a minus sign which cancel to prove $\alpha_{j,n} > 0$.

The left side of (2.4) is $-z^{-1} + O(z^{-2})$ as $z \rightarrow \infty$ since P_n is monic. The right side is $-z^{-1} \left(\sum_{j=0}^n \alpha_{j,n} \right) + O(z^{-2})$, so (2.5) holds.

(ii) This follows from (2.4) and (2.5) if one notes that for any $x \in \mathbb{R}$ and z with $\text{Im } z > 0$,

$$0 < \text{Im} \left(\frac{1}{x - z} \right) \leq \frac{1}{|x - z|} \leq \frac{1}{\text{Im } z}.$$

(iii) This follows immediately from

$$\frac{P_{n+1}(z)}{P_n(z)} = z - b_{n+1} - a_n^2 \frac{P_{n-1}(z)}{P_n(z)}. \tag{2.10}$$

Since $\text{Im } P_{n-1}/P_n < 0$, (2.10) implies (2.8). \square

Proof of Theorem 2.2. By replacing x by $(x - \text{Re } z_0)/\text{Im } z_0$ (i.e., translating and scaling the measure), we can suppose for notational simplicity that $z_0 = i$.

Let

$$\alpha = \lim_{n \rightarrow \infty} \frac{P_{n+1}(i)}{P_n(i)}. \tag{2.11}$$

By (2.8), $\text{Im } \alpha \geq 1$ so $-\text{Im}(\alpha^{-1}) = \text{Im } \alpha / |\alpha|^2 > 0$.

Taking imaginary parts of (2.10), we see that

$$a_n^2 = \frac{[\text{Im}(P_{n+1}/P_n) - 1]}{\text{Im}(-P_{n-1}/P_n)} \rightarrow \frac{(\text{Im } \alpha - 1)}{\text{Im}(-\alpha^{-1})} \equiv a^2 \geq 0 \tag{2.12}$$

proving a_n has a limit.

Taking real parts of (2.10) shows

$$b_{n+1} = -a_n^2 \text{Re}\left(\frac{P_{n-1}}{P_n}\right) - \text{Re}\left(\frac{P_{n+1}}{P_n}\right) \rightarrow -a^2 \text{Re}(\alpha^{-1}) - \text{Re}(\alpha) \equiv b. \quad \square \tag{2.13}$$

Our first proof of Theorem 2.1 is a simple consequence on the following theorem of Poincaré (see [4,12,14] for proofs):

Theorem of Poincaré. *Let $u_j \in \mathbb{C}$ solve the n th order difference equation*

$$u_{n+j} = a_{j,1}u_{n+j-1} + a_{j,2}u_{n+j-2} + \dots + a_{j,n}u_j, \tag{2.14}$$

$j = 1, 2, \dots$ Suppose

- (i) $a_{j,n} \neq 0$ for all j .
- (ii) $\lim_{j \rightarrow \infty} a_{j,\ell} = A_\ell$ exists for $\ell = 1, \dots, n$.

Let $\lambda_1, \dots, \lambda_n$ be the solutions of

$$A_n + A_{n-1}\lambda + \dots + A_1\lambda^{n-1} = \lambda^n. \tag{2.15}$$

Suppose $\{\lambda_j\}_{j=1}^n$ are distinct, and for $j \neq k$, $|\lambda_j| \neq |\lambda_k|$. Then, if u is not identically zero, we have for some k that

$$\lim_{j \rightarrow \infty} \frac{u_{j+1}}{u_j} = \lambda_k. \tag{2.16}$$

First Proof of Theorem 2.1. Eq. (1.1) is exactly of the form (2.14). Eq. (2.15) becomes

$$\lambda^2 = (z - b)\lambda - a^2 \tag{2.17}$$

whose solutions are

$$\lambda_{\pm} = \frac{(z - b) \pm \sqrt{(z - b)^2 - 4a^2}}{2}. \tag{2.18}$$

To prove (2.1), we must show that if $z \notin [b - 2a, b + 2a]$, then $|\lambda_+| \neq |\lambda_-|$ and then identify which root is taken by the ratio.

If α and β are complex numbers, $|\alpha + \beta| = |\alpha - \beta|$ if and only if α and β are orthogonal as vectors in $\mathbb{C} = \mathbb{R}^2$, if and only if $\beta = ic\alpha$ for $c \in \mathbb{R}$, if and only if

$\beta^2 = -c^2\alpha^2$. Taking $\alpha = (z - b)$ and $\beta = \sqrt{(z - b)^2 - 4a^2}$, we see $|\lambda_+| = |\lambda_-|$ if and only if

$$-c^2(z - b)^2 = (z - b)^2 - 4a^2$$

or

$$z - b = \pm \frac{2a}{\sqrt{1 + c^2}} \tag{2.19}$$

for $c \in \mathbb{R}$. Eq. (2.19) holds if and only if $z \in [b - 2a, b + 2a]$.

Since $|\lambda_+| \neq |\lambda_-|$, $\lambda_{\pm}(z)$ are analytic in $\mathbb{C} \setminus [b - 2a, b + 2a]$ (as is obvious from (2.18)). Since $\lambda_+\lambda_- = a^2$, $|\lambda_+||\lambda_-| = a^2$, and $\lambda_+ = z + O(1/z)$ as $|z| \rightarrow \infty$, $|\lambda_+| > a$ for $|z|$ large, and so since $|\lambda_+||\lambda_-| = a^2$ and $|\lambda_+| \neq |\lambda_-|$ implies $|\lambda_{\pm}| \neq a$, we see

$$|\lambda_+| > a \quad |\lambda_-| < a \quad \text{all } z \in \mathbb{C} \setminus [b - 2a, b + 2a]. \tag{2.20}$$

Thus for all $z \in \mathbb{C} \setminus [b - 2a, b + 2a]$, Poincaré’s theorem applies and P_{n+1}/P_n has a limit $\ell(z)$ where for each z , $\ell(z) = \lambda_+$ or $\ell(z) = \lambda_-$.

By (2.6), P_n/P_{n+1} is a normal family on $\mathbb{C} \setminus \mathbb{R}$, so $\ell(z)$ is analytic on $\mathbb{C} \setminus \mathbb{R}$. By (2.8), $|\ell(z)| > |\text{Im } z|$, so for $|z|$ large, $\ell(z) = \lambda_+(z)$ and thus, by analyticity, $\ell(z) = \lambda_+(z)$ for $z \in \mathbb{C} \setminus \mathbb{R}$. This establishes (2.1) there.

For $z \in \mathbb{R} \setminus [b - 2a, b + 2a]$,

$$\lim_{n \rightarrow \infty} \frac{p_{n+1}(z)}{p_n(z)} = \lim_{n \rightarrow \infty} \frac{1}{a} \frac{P_{n+1}(z)}{P_n(z)} = \frac{\lambda_{\pm}(z)}{a}.$$

If the limit is $\lambda_-(z)$, $\lim |p_{n+1}/p_n| < 1$, so $p_n \in \ell_2$ and $z \in \text{spectrum}(J)$. Conversely, if it is λ_+ , $\lim |p_{n+1}/p_n| > 1$, so $p_n \notin \ell_2$. Thus (2.1) holds for $z \in \mathbb{R} \setminus \text{spectrum}(J)$ and (2.2) holds for $z \in \text{spectrum}(J) \setminus [b - 2a, b + 2a]$. \square

Second Proof of Theorem 2.1. Let $J^{(n)}$ be the $n \times n$ matrix obtained from the first n rows and columns of J . As is well known,

$$\det(z - J^{(n)}) = P_n(z).$$

Thus, by Cramer’s rule,

$$\frac{P_{n-1}(z)}{P_n(z)} = (z - J^{(n)})_{nn}^{-1} = (z - \tilde{J}^{(n)})_{11}^{-1}, \tag{2.21}$$

where $\tilde{J}_{ij}^{(n)} = \tilde{J}_{n-i, n-j}^{(n)}$, that is,

$$\tilde{J}^{(n)} = \begin{pmatrix} b_n & a_{n-1} & 0 & \dots & \dots \\ a_{n-1} & b_{n-1} & a_{n-2} & \dots & \dots \\ 0 & a_{n-2} & b_{n-2} & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_1 & b_1 \end{pmatrix}. \tag{2.22}$$

If $\tilde{J}^{(\infty)}$ is the constant Jacobi matrix $a_n \equiv a$, $b_n \equiv b$, it is clear as operators on $\ell^2(\mathbb{Z}_+)$, $\tilde{J}^{(n)} \rightarrow \tilde{J}^{(\infty)}$ strongly. It follows that as operators on $\ell^2(\mathbb{Z})$, $(\tilde{J}^{(n)} - z)^{-1} \rightarrow (\tilde{J}^{(\infty)} - z)^{-1}$

strongly for $\text{Im } z \neq 0$. Thus

$$\frac{P_{n-1}(z)}{P_n(z)} = (z - \tilde{J}^{(n)})_{11}^{-1} \rightarrow (z - \tilde{J}^{(\infty)})_{11}^{-1}. \tag{2.23}$$

Let w solve

$$a(w + w^{-1}) + b = z \tag{2.24}$$

with $|w| < 1$. Let $u_n = w^n$. Thus $u_n \in \ell_2$ and

$$(z - \tilde{J}^{(\infty)})u = (aw^{-1})\delta_1,$$

so

$$(z - \tilde{J}^{(\infty)})_{11}^{-1} = aw^{-1}. \tag{2.25}$$

Solving (2.24), we see that $aw^{-1} = \text{RHS of (2.1)}$. This proves (2.1) on $\mathbb{C} \setminus \mathbb{R}$.

On $\mathbb{R} \setminus \text{spectrum}(J)$, one shows the eigenvalues of $\tilde{J}^{(n)}$ (equals zeros of $P_n(z)$) outside $[b - 2a, b + 2a]$ converge to the eigenvalues of J so for $z \in \mathbb{R} \setminus \text{spectrum}(J)$, $(\tilde{J}^{(n)} - z)^{-1} \rightarrow (\tilde{J}^{(\infty)} - z)^{-1}$ strongly also. \square

3. Weak asymptotic limits

Let $d\mu_n = p_n^2 d\mu$. In this section, the main theorems are

Theorem 3.1. *Suppose*

$$b_n \rightarrow b, \quad a_{2n} \rightarrow a, \quad a_{2n+1} \rightarrow c \tag{3.1}$$

for $b \in \mathbb{R}, a, c \in [0, \infty)$. Then as $n \rightarrow \infty$, $d\mu_n$ has a weak limit $d\rho_{b;a,c}(x)$, and for every ℓ ,

$$\int x^\ell d\mu_n \rightarrow \int x^\ell d\rho_{b;a,c}(x). \tag{3.2}$$

Remark. 1. The hypotheses imply that $d\mu$ is supported in $[\inf b_n - 2 \sup(a_n), \sup b_n + 2 \sup(a_n)]$ which is bounded, so weak convergence is equivalent to convergence of the moments.

2. We will see below that $d\rho_{b';a',c'} = d\rho_{b;a,c}$ implies $b' = b$ and either $a' = a, c' = c$, or $a' = c, c' = a$.

3. We will discuss the form of $d\rho_{b;a,c}$ in Section 5.

Theorem 3.2. *Suppose for $\ell = 1, 2$, and 4,*

$$\lim_{n \rightarrow \infty} \int x^\ell d\mu_n = A_\ell. \tag{3.3}$$

Then for some a, b, c , (3.1) holds. Moreover, A_1, A_2, A_4 determine $b, a + c, |a - c|$ (i.e., they determine b and the unordered pair (a, c)).

Remark. 1. We will see $A_2 < \infty$ implies $\sup(|b_n| + |a_n|) < \infty$.

2. The final assertion proves the second remark after Theorem 3.1.

Our proofs will depend on a graphical representation of $\int x^\ell d\mu_n$. Consider the lattice $\mathbb{Z}_+ = \{0, 1, \dots\}$. We will consider a random walk on \mathbb{Z}_+ where at each step, one either stays at the site one is at or one jumps by a single site. Paths have unnormalized weights, products over the steps: b_{k+1} if one stays at site k , a_{k+1} is one move from k to $k + 1$ or $k + 1$ to k . To be more precise, a path is a sequence $\rho_0, \rho_1, \dots, \rho_\ell \in \mathbb{Z}_+$ so that $|\rho_m - \rho_{m-1}| \leq 1$ and

$$W(\rho) = \prod_{j=0}^{\ell-1} w(\rho_j, \rho_{j+1}) \tag{3.4}$$

and

$$w(\rho_j, \rho_{j+1}) = \begin{cases} b_{k+1} & \text{if } \rho_{j+1} = \rho_j = k, \\ a_{k+1} & \text{if } \rho_{j+1} = \rho_j + 1 = k + 1, \\ a_k & \text{if } \rho_{j+1} = \rho_j - 1 = k - 1. \end{cases} \tag{3.5}$$

Here is the key tool:

Proposition 3.3.

$$\int x^\ell d\mu_n = \sum_{\rho \in Q_{n,\ell}} W(\rho), \tag{3.6}$$

where $Q_{n,\ell}$ is the set of all paths of length ℓ with $\rho_0 = \rho_\ell = n$.

Proof. Since

$$xp_n = a_{n+1}p_{n+1} + b_{n+1}p_n + a_n p_{n-1},$$

we see immediately that, by induction in j ,

$$x^j p_n = \sum c_{j,m,n} p_m, \tag{3.7}$$

where

$$c_{j,m,n} = \sum_{\rho \in Q_{n,m,j}} W(\rho)$$

and $Q_{n,m,j}$ is all paths of length j with $\rho_0 = n$ and $\rho_j = m$. (3.6) follows since $\int x^\ell d\mu_n = \langle p_n, x^\ell p_n \rangle = c_{\ell,n,n}$. \square

Proof of Theorem 3.1. Under hypothesis (3.1), J is bounded, so $d\mu$ has a bounded support, so weak convergence is equivalent to (3.2). By Proposition 3.3, $\int x^\ell d\mu_n$ is a finite sum over paths. This representation shows that if a_n, b_n , and \tilde{a}_n, \tilde{b}_n are two sets of Jacobi parameters and $\lim_{n \rightarrow \infty} |a_n - \tilde{a}_n| + |b_n - \tilde{b}_n| = 0$, then $|\int x^\ell d\mu_n - \int x^\ell d\tilde{\mu}_n| \rightarrow 0$. Thus we need only prove (3.2) for $b_n \equiv b, a_{2n} \equiv a, a_{2n+1} \equiv c$.

Fix ℓ . So long as $\ell < 2n$, there is a one-one correspondence between paths $\rho \in Q_{n,\ell}$ and $\rho \in Q_{n+1,\ell}$ by $U = TS$,

$$\begin{aligned} T(\rho)_j &= \rho_j + 1, \\ S(\rho)_j &= n - (\rho_j - n). \end{aligned}$$

S reflects the path in n , T translates by 1. $\ell < 2n$ is needed to assure paths do not get mapped into ones that have $\rho_j < 0$, which is forbidden (and that the inverse does not do this), showing U is a bijection of $Q_{n,\ell}$ and $Q_{n+1,\ell}$. The key point is that $W(U\rho) = W(\rho)$, for if $\rho_j = \rho_{j+1}$, the weight is always b and both S and T interchange links with weight a and those with weight c . It follows that if $b \equiv b, a_{2n} \equiv a, a_{2n+1} \equiv c$, then $\int x^\ell d\rho_n$ is independent of n once $\ell < 2n$, so the limit exists. Once the moments exist, they provide a measure since the nonnegative Hankel matrices converge to nonnegative Hankel matrices. \square

Proof of Theorem 3.2. $\int xp_n^2 d\mu = b_{n+1}$, so (3.4) for $\ell = 1$ implies $b_{n+1} \rightarrow A_1 \equiv b$. Let $d\tilde{\mu}(x) = d\mu(x + b)$. The Jacobi parameters of $\tilde{\mu}$ are given by

$$\tilde{a}_n = a_n \quad \tilde{b}_n = b_n - b. \tag{3.8}$$

Moreover,

$$\int x^\ell d\mu_n = \sum_{j=0}^{\ell} \binom{\ell}{j} (b)^{\ell-j} \int x^j d\tilde{\mu}_n \tag{3.9}$$

and

$$\int x^\ell d\tilde{\mu}_n = \sum_{j=0}^{\ell} \binom{\ell}{j} (-b)^{\ell-j} \int x^j d\mu_n. \tag{3.10}$$

Since $\tilde{b}_n \rightarrow 0$ and every odd ℓ random walk has a b_k factor in it, $\int x^\ell d\tilde{\mu}_n \rightarrow 0$ for all odd ℓ . Thus, (3.9) implies $\int x^3 d\mu_n$ exists and then (3.10) that $\int x^4 d\tilde{\mu}_n$ converges. Thus, without loss, we suppose $A_1 = 0$ and $b_n \rightarrow 0$.

If $b_n \rightarrow 0$, any path with $\rho_j = \rho_{j+1}$ contributes zero in the limit; so we can restrict to paths with $|\rho_{j+1} - \rho_j| = 1$. Thus, looking at the two such paths with $\rho_2 = \rho_0 = n$,

$$\lim_{n \rightarrow \infty} a_{n+1}^2 + a_n^2 = A_2. \tag{3.11}$$

In looking at paths with $\rho_0 = \rho_4 = n$, all those with $\rho_2 = n$ contribute $(\int x^2 d\mu_n)^2$, so

$$\lim_{n \rightarrow \infty} a_{n+2}^2 a_{n+1}^2 + a_n^2 a_{n-1}^2 = A_4 - A_2^2. \tag{3.12}$$

Thus, using $(x - y)^2 = (x + y)^2 - 4xy$,

$$\lim_{n \rightarrow \infty} (a_{n+2}^2 - a_{n+1}^2)^2 + (a_n^2 - a_{n-1}^2)^2 = 6A_2^2 - 4A_4. \tag{3.13}$$

Suppose a_n has a limit point, a , that is, $a_{n(j)} \rightarrow a$ as $j \rightarrow \infty$ for a subsequence. Define $c = \sqrt{A_2 - a^2}$. By (3.11), for any $\ell = 0, \pm 1, \pm 2, \dots$,

$$a_{n(j)+\ell} \rightarrow \begin{cases} a & \ell \text{ even,} \\ c & \ell \text{ odd.} \end{cases}$$

In particular, by (3.13),

$$|a^2 - c^2| = \sqrt{3A_2^2 - 2A_4}. \tag{3.14}$$

Since also

$$a^2 + c^2 = A_2 \tag{3.15}$$

there are at most two solutions of (3.14), (3.15):

$$a^2 = \frac{1}{2} \left[A_2 + \sqrt{3A_2^2 - 2A_4} \right], \tag{3.16}$$

$$c^2 = \frac{1}{2} \left[A_2 - \sqrt{3A_2^2 - 2A_4} \right] \tag{3.17}$$

and the one with a, c , reversed. Thus the right sides of (3.16) and (3.17) are the only limit points of a_n^2 .

The lemma below completes the proof. \square

Lemma 3.4. *Let x_n be a sequence so that for some $\alpha, \beta \in \mathbb{R}$,*

$$\lim_{n \rightarrow \infty} x_n + x_{n+1} = \alpha + \beta, \tag{3.18}$$

$$\lim_{n \rightarrow \infty} |x_n - x_{n+1}| = |\alpha - \beta|. \tag{3.19}$$

Then either

$$\lim_n x_{2n} = \alpha \quad \lim_n x_{2n+1} = \beta$$

or

$$\lim_n x_{2n} = \beta \quad \lim_n x_{2n+1} = \alpha.$$

Proof. By replacing x_n by $x_n - \frac{1}{2}(\alpha + \beta)$, we can suppose $\alpha = -\beta \geq 0$. If $\alpha = \beta = 0$, the result is trivial, so suppose $\alpha = -\beta > 0$. Pick N so that for $n > N$,

$$|x_n + x_{n+1}| < \alpha,$$

$$|x_n - x_{n+1}| > \alpha.$$

Thus, since $|x_n - x_{n+1}| > |x_n + x_{n+1}|$, x_n and x_{n+1} have opposite signs for all $n > N$, that is, for $n > N$, either $(-1)^n x_n > 0$ or $(-1)^{n+1} x_n > 0$. Since $\pm \alpha$ are the only allowed limit points if $(-1)^n x_n > 0$, $x_{2n} \rightarrow \alpha$, $x_{2n+1} \rightarrow \beta = -\alpha$, and if $(-1)^{n+1} x_n > 0$, $x_{2n} \rightarrow \beta$, $x_{2n+1} \rightarrow \alpha$. \square

4. Ratio asymptotics for p_{n+1}/p_n

In this section and the next, we discuss two further issues related to our results: what about $\lim p_{n+1}/p_n$ (p_n rather than P_n) and we calculate the measures $d\rho_{b;a,c}$ of (3.2) (already well known if $a = c$).

Let

$$R_n(z) = \frac{P_{n+1}(z)}{P_n(z)}, \quad r_n(z) = \frac{p_{n+1}(z)}{p_n(z)}. \tag{4.1}$$

Since $p_n = (a_1 \dots a_n)^{-1} P_n$,

$$r_n(z) = a_{n+1}^{-1} R_n(z), \tag{4.2}$$

so we immediately see with Theorem 2.1 that if $a_n \rightarrow a \neq 0$ and $b_n \rightarrow b$,

$$\lim_{n \rightarrow \infty} r_n(z) = \frac{(z - b) + \sqrt{(z - b)^2 - 4a^2}}{2a}. \tag{4.3}$$

We want to address a converse. One problem we find is that while $|R_n(z)| > \text{Im } |z|$, without an a priori upper bound on a_n , we do not have a bound for $|r_n(z)|$, so it is not obvious that existence of the limit implies a_n is bounded.

Example 4.1. Let $a_n = e^{n!}$, $b_n =$ arbitrary bounded sequence, especially one without a limit. Then, by (1.3) and

$$\begin{aligned} r_n(z) &= \frac{z - b_{n+1}}{a_{n+1}} - \frac{a_n}{a_{n+1}} [r_{n-1}(z)]^{-1} \\ &= \frac{z - b_{n+1}}{a_{n+1}} - \frac{a_n^2}{a_{n+1}} [R_{n-1}(z)]^{-1} \end{aligned} \tag{4.4}$$

by (4.2). By (2.7),

$$|r_n(z)| \leq \frac{|z - b_{n+1}|}{a_{n+1}} + \frac{a_n^2}{a_{n+1}} |\text{Im } z|^{-1} \rightarrow 0 \tag{4.5}$$

since a_n is chosen so $a_n^2/a_{n+1} \rightarrow 0$. Thus for this example, $r_n(z) \rightarrow 0$.

Because of this example, we will need to suppose that if $\lim r_n(z)$ exists, it has nonzero imaginary part. Here is a result that requires two points rather than one, with some extra conditions:

Theorem 4.2. Suppose $\sup_n a_n < \infty$. Suppose z_1, z_2 are in $\{z \mid \text{Im } z > 0\}$ and let

$$\lim_{n \rightarrow \infty} r_n(z_j) = \lambda_j \tag{4.6}$$

with

(a) $\text{Im } \lambda_j > 0$,

and either

$$(b1) \quad \frac{\operatorname{Im} \lambda_1}{\operatorname{Im} z_1} \neq \frac{\operatorname{Im} \lambda_2}{\operatorname{Im} z_2}$$

or

$$(b2) \quad \frac{\operatorname{Im}(\lambda_1^{-1})}{\operatorname{Im} z_1} \neq \frac{\operatorname{Im}(\lambda_2^{-1})}{\operatorname{Im} z_2}.$$

Then $a_n \rightarrow a \neq 0$ and $b_n \rightarrow b$.

Proof. By (1.3),

$$a_{n+1}r_n(z) = (z - b_{n+1}) - a_n[r_{n-1}(z)]^{-1}. \tag{4.7}$$

Since $\operatorname{Im}(-r_{n-1}(z))^{-1} > 0$, (4.7) implies

$$a_{n+1} \operatorname{Im} r_n(z_j) \geq \operatorname{Im} z_j,$$

which implies

$$\liminf a_n \geq \frac{\operatorname{Im} z_j}{\operatorname{Im} \lambda_j} > 0,$$

so the a 's are bounded above and below.

Let (a, c) be a limit point of (a_{n+1}, a_n) . By (4.7),

$$a \operatorname{Im} \lambda_j = \operatorname{Im} z_j + c(\operatorname{Im}(-\lambda_j)^{-1}). \tag{4.8}$$

If $\operatorname{Im} \lambda_1/\operatorname{Im} z_1 \neq \operatorname{Im} \lambda_2/\operatorname{Im} z_2$, (4.8) implies

$$a \left[\frac{\operatorname{Im} \lambda_1}{\operatorname{Im} z_1} - \frac{\operatorname{Im} \lambda_2}{\operatorname{Im} z_2} \right] = c \left[\frac{\operatorname{Im}(-\lambda_1)^{-1}}{\operatorname{Im} z_1} - \frac{\operatorname{Im}(-\lambda_2)^{-1}}{\operatorname{Im} z_2} \right],$$

so we can solve for a as a multiple of c , and then for c in (4.8) for $j = 1$. If $\operatorname{Im}(\lambda_1)^{-1}/\operatorname{Im} z_1 \neq \operatorname{Im}(\lambda_2)^{-1}/\operatorname{Im} z_2$, we solve for c as a multiple of a . Either way, we see (4.8) has a unique solution for (a, c) so $(a_{n+1}, a_n) \rightarrow (a, c)$. But then $(a_{n+2}, a_{n+1}) \rightarrow (a, c)$ so $a = c$ and $a_n \rightarrow a \neq 0$. This implies $\lim R_n(z_1) = \lim a_n c_n(z)$ exists. So, by Theorem 4.2, $b_n \rightarrow b$. \square

We have a second remark about the existence of $\lim_{n \rightarrow \infty} r_n(z)$. In the OPUC case, existence of $\varphi_{n+1}^*(z)/\varphi_n^*(z)$ for all $z \in \mathbb{D}$ implies the same of $\Phi_{n+1}^*(z)/\Phi_n^*(z)$ (and, by taking $\alpha_j = 0$ if $j \neq n^2$, $\alpha_{n^2} = \frac{1}{2}$, not conversely) for $\varphi_{n+1}^*(0)/\varphi_n^*(0) = \rho_n^{-1}$, so existence of the φ ratio limit at $z = 0$ implies $\rho_n \rightarrow \rho_\infty$ and then, since $\Phi_{n+1}^*(z)/\Phi_n^*(z) = \rho_n \varphi_{n+1}^*(0)/\varphi_n^*(0)$, we get the Φ_n^* ratio limits.

The same is true here, but alas, the analog of $z = 0$ for OPUC is $z = \infty$ here. The following captures the idea, without the need for hypotheses (b) of Theorem 4.2.

Theorem 4.3. *Suppose a_n and $|b_n|$ are bounded and $r_n(z)$ converges to a nonzero limit as $n \rightarrow \infty$ for all z in a small neighborhood of $z_0 \in \mathbb{C} \setminus \mathbb{R}$. Then $a_n \rightarrow a$ and $b_n \rightarrow b$ for some $a \neq 0, b \in \mathbb{R}$.*

Proof. By (4.4),

$$r_n(z) = \frac{z}{a_{n+1}} - \frac{b_{n+1}}{a_{n+1}} + O\left(\frac{1}{z^{-1}}\right). \tag{4.9}$$

Thus

$$r_n(z)^{-1} = \frac{a_{n+1}}{z} + \frac{b_{n+1}}{z^2} + O\left(\frac{1}{z^3}\right). \tag{4.10}$$

If $\text{supp}(d\mu) \subset [-c, c]$ (take $c = \sup_n |b_n| + 2 \sup |a_n|$), then for $\rho > c$, by the Cauchy formula for every $\ell \in \mathbb{Z}$ and the fact that r_n has its zeros in $[-c, c]$, $\frac{1}{2\pi i} \oint_{|z|=\rho} r_n(z)^{-1} z^\ell dz$ is ρ independent for $\rho > c$. Taking ρ to infinity, we get, by Eq. (4.9),

$$\frac{1}{a_{n+1}} = \frac{1}{2\pi i} \oint_{|z|=c+1} r_n(z)^{-1} dz, \tag{4.11}$$

$$-\frac{b_{n+1}}{a_{n+1}} = \frac{1}{2\pi i} \oint_{|z|=c+1} r_n(z)^{-1} z dz. \tag{4.12}$$

Thus uniform convergence of $r_n(z)$ to a limit on $|z| = c + 1$ implies convergence of a_n and b_n . Therefore, we are reduced to showing convergence of r_n^{-1} on a single compact subset of $\mathbb{C} \setminus \mathbb{R}$ implies convergence on all compact subsets of $\mathbb{C} \setminus [-c, c]$.

By (2.4) and (2.4) and $x_{i,n} \in [-c, c]$, we have

$$|R_n(z)^{-1}| \leq \sup_{x \in [-c, c]} |z - x|^{-1}$$

and thus, by (4.2),

$$|r_n(z)^{-1}| \leq \left[\sup_n |a_n| \right] \sup_{x \in [-c, c]} |z - x|^{-1}. \tag{4.13}$$

Thus convergence of $r_n(z)$ on a compact implies, by Vitali’s theorem and (4.13), uniform convergence of $r_n(z)^{-1}$ on compact sets in $\mathbb{C} \setminus [-c, c]$. \square

5. Calculation of $d\rho_{b;a,c}(x)$

In this section, we compute the weak limit $\rho_{b;a,c}$ of $p_n^2 d\mu$ for μ , the measure corresponding to the Jacobi matrix $a_{2n} = c, a_{2n+1} = a, b_n = b$. We will also find $d\mu_{b;a,c}$, the measure associated to the Jacobi matrix

$$J = \begin{pmatrix} b & a & & & \\ a & b & c & & \\ & c & b & a & \\ & & a & b & c \\ & & & \ddots & \ddots & \ddots \end{pmatrix}. \tag{5.1}$$

We begin by computing

$$f(z; b; a, c) = \int \frac{d\mu_{b;a,c}(x)}{z - x} \tag{5.2}$$

and

$$G(z; b; a, c) = \int \frac{d\rho_{b;a,c}(x)}{z - x}. \tag{5.3}$$

This calculation is not unrelated to calculations in Khrushchev [8] for the measure associated to a period 2 Verblunsky coefficient. As he does, we could ask for the Jacobi coefficients for the measure $d\rho$ and show they converge exponentially fast to those for $d\mu$ (with a need to interchange a and c depending on the sign of $a - c$).

We begin with a result about a finite Jacobi matrix

$$J^{[1,n]} = \begin{pmatrix} b_1 & a_1 & & & & & & & \\ a_1 & b_2 & a_2 & & & & & & \\ & & \ddots & \ddots & \ddots & & & & \\ & & & \ddots & \ddots & \ddots & & & \\ & & & & \ddots & \ddots & \ddots & & \\ & & & & & \ddots & \ddots & \ddots & \\ & & & & & & a_{n-1} & & \\ a_{n-1} & & & & & & & & b_n \end{pmatrix}. \tag{5.4}$$

We let $J^{[j,k]}$ for $1 \leq j \leq k \leq n$ denote the $(k - j + 1) \times (k - j + 1)$ matrix we get by keeping rows and columns between j and k (inclusive). We refer to the row number of $J^{[j,k]}$ as $j, j + 1, \dots$ so, for example, $(J^{[j,k]})_{jj} = b_j$.

Here is a key lemma that appears in [6] although closely related formulae have appeared elsewhere; in particular, the $k = 1, n$ results go back to Jacobi:

Proposition 5.1. *For $2 \leq k \leq n - 1$,*

$$\begin{aligned} & [(z - J^{[1,n]})^{-1}]_{kk} \\ & = 1 / \{z - b_k - a_{k-1}^2 [(z - J^{[1,k-1]})^{-1}]_{k-1,k-1} - a_k^2 [(z - J^{[k+1,n]})^{-1}]_{k+1,k+1}\}. \end{aligned} \tag{5.5}$$

For $k = 1$,

$$[(z - J^{[1,n]})^{-1}]_{11} = 1 / \{z - b_1 - a_1^2 [(z - J^{[2,n]})^{-1}]_{22}\}. \tag{5.6}$$

Proof. Let $2 \leq k \leq n - 1$. Here is a proof that is more direct than that in [6], although the essence is the same. If row and column k are removed, the resulting matrix is $J^{[1,k-1]} \oplus J^{[k+1,n]}$ so, by Cramer's rule,

$$[(z - J^{[1,n]})^{-1}]_{kk} = \frac{\det(z - J^{[1,k-1]}) \det(z - J^{[k+1,n]})}{\det(z - J^{[1,n]})}. \tag{5.7}$$

Expanding $\det(z - J^{[1,n]})$ in minors in row k ,

$$\det(z - J^{[1,n]}) = (z - b_k)d_1 - a_{k-1}^2 d_2 - a_k^2 d_3, \tag{5.8}$$

where

$$d_1 = \det(z - J^{[1,k-1]})\det(z - J^{[k+1,n]}), \tag{5.9}$$

$$d_2 = \det(z - J^{[1,k-2]})\det(z - J^{[k+1,n]}), \tag{5.10}$$

$$d_3 = \det(z - J^{[1,k-1]})\det(z - J^{[k+2,n]}) \tag{5.11}$$

(where $\det(z - J^{[1,0]})$ and $\det(z - J^{[n+1,n]})$, which occur if $k = 2$ or $k = n - 1$, are interpreted as 1).

Finally, note that, by Cramer’s rule again,

$$[(z - J^{[1,k-1]})^{-1}]_{k-1,k-1} = \frac{d_2}{d_1}, \tag{5.12}$$

$$[(z - J^{[k+1,n]})^{-1}]_{k+1,k+1} = \frac{d_3}{d_1}. \tag{5.13}$$

Eqs. (5.7)–(5.13) imply (5.5). Eq. (5.6) is proven in a similar way (but, e.g., the analog of (5.8) has only two terms). \square

Corollary 5.2. *For the functions f and G of (5.2)/(5.3), we have that for $z \in \mathbb{C} \setminus \mathbb{R}$,*

$$[G(z; b; a, c)]^{-1} = z - b - c^2 f(z; b; a, c) - a^2 f(z; b; c, a). \tag{5.14}$$

Remark. This formula makes it evident once again that G is symmetric in a and c .

Proof. Let J be given by (5.1). On $\ell^2(\mathbb{Z}^+)$, $J^{[1,n]} \oplus 0$ converges strongly to J . Thus

$$f(z; b; a, c) = [(z - J)^{-1}]_{11} = \lim_{n \rightarrow \infty} [(z - J^{[1,n]})^{-1}]_{11}. \tag{5.15}$$

Moreover,

$$\int \frac{p_k(x)^2 d\mu_{b;a,c}(x)}{z - x} = [(z - J)^{-1}]_{kk} = \lim_{n \rightarrow \infty} [(z - J^{[1,n]})^{-1}]_{kk}, \tag{5.16}$$

so

$$G(z; b; a, c) = \lim_{k \rightarrow \infty} \left[\lim_{n \rightarrow \infty} [(z - J^{[1,n]})^{-1}]_{kk} \right]. \tag{5.17}$$

Eq. (5.14) follows by using (5.5), (5.15), and (5.17) together with the structure of J (e.g., $J^{[1+2\ell,n+2\ell]} = J^{[1,n]}$ and $J^{[1+2\ell+1,n+2\ell+1]}$ is $\tilde{J}^{[1,n]}$ for \tilde{J} , the matrix with a and c reversed). \square

Remark. The limit of (5.5) as $n \rightarrow \infty$ is a precise analog of Khrushchev’s formula [7] that for the unit circle case, the Schur function of $|\varphi_n|^2 d\mu$ is $b_n f_n$. It would be

interesting to see if one could translate our proof here to a proof of Khrushchev’s formula using the CMV matrix [3,14] in place of the Jacobi matrix.

Corollary 5.3.

$$[f(z; b; a, c)]^{-1} = z - b - a^2 f(z; b; c, a). \tag{5.18}$$

Proof. This is identical to the last proof using (5.6) in place of (5.5). Of course, this is a special case of the well-known Stieltjes continued fraction expansion for the Stieltjes transform of the measure associated to a Jacobi matrix. \square

Henceforth, for simplicity, we take $b = 0$. Since $G(z; b; a, c) = G(z - b; 0; a, c)$, and similarly for f , it is easy to go from this case to the case of general b .

As a warmup, consider the case $a = c$. Then (5.18) becomes

$$1 = f(z - a^2 f), \tag{5.19}$$

which is solved by

$$f(z; 0; a, a) = \frac{z - \sqrt{z^2 - 4a^2}}{2a^2}, \tag{5.20}$$

where the branch of the square root is taken (with $\sqrt{\dots} = z + O(1/z)$ as $|z| \rightarrow \infty$ consistent with $f(z) \sim 1/z + O(z^{-2})$). By (5.14), we get

$$G(z; 0; a, a) = \frac{1}{\sqrt{z^2 - 4a^2}}. \tag{5.21}$$

$\lim_{\varepsilon \downarrow 0} \text{Im } G(x + i\varepsilon; 0; a, a)$ is only nonzero for $z \in [-2a, 2a]$ so

$$d\rho_{b;a,a}(x) = \frac{1}{\pi} \lim_{\varepsilon \downarrow 0} \text{Im } G(x + i\varepsilon; b; a, c) dx \tag{5.22}$$

implies the well-known

$$d\rho_{b;a,a} = \frac{1}{\pi \sqrt{4a^2 - (x - b)^2}} \chi_{[b-2a, b+2a]}(x) dx, \tag{5.23}$$

consistent with Nevai’s conjecture.

Here is the main result of this section:

Theorem 5.4. Define

$$I(z; a, c) = (c^2 - a^2 + z^2)^2 - 4z^2 c^2. \tag{5.24}$$

Then

$$(i) \quad f(z; 0; a, c) = \frac{c^2 - a^2 + z^2 - \sqrt{I(z; a, c)}}{2c^2 z}, \tag{5.25}$$

$$(ii) \quad G(z; 0; a, c) = \frac{z}{\sqrt{I(z; a, c)}}, \tag{5.26}$$

$$(iii) \quad d\rho_{b;a,c}(x) = [\chi_{J_2}(x) + \chi_{J_2}(x)]w(x) dx, \quad (5.27)$$

where

$$J_1 = [b + |c - a|, b + a + c] \quad J_2 = -J_1 \quad (5.28)$$

and for $x \in J_1 \cup J_2$,

$$w(x) = x\pi^{-1} \left[\sqrt{-I(x - b; a, c)} \right]^{-1}. \quad (5.29)$$

Proof. (i) Iterating (5.15) once, we get a quadratic equation for $f(z; 0; a, c)$ whose solution (the one with $f = 1/z + O(1/z^2)$ at infinity) is (5.25).

(ii) Follows from (5.14) and (5.25) if we note that $I(z; a, c) = z^4 - 2(a^2 + c^2)z^2 + (a^2 - c^2)^2$ is symmetric under interchange of a and c .

(iii) $I(z; a, c) = 0$ if and only if $c^2 - a^2 + z^2 = \pm 2zc$ if and only if $(z \pm c)^2 = a^2$ if and only if $z = \pm a \pm c$ (independent \pm) which say \sqrt{I} has a branch cut on $J_1 \cup J_2$ and G is purely imaginary there. (5.22) completes the proof. \square

We note there is another proof of the theorem using Floquet theory and the theory of periodic whole-line Jacobi matrices.

References

- [1] N.I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*, Hafner, New York, 1965 (Russian original, 1961).
- [2] D. Barrios, G. Lopes, E. Torrano, Distribution of zeros and the asymptotics of polynomials that satisfy three-term recurrent relations with complex coefficients, *Mat. Sb.* 184 (1993) 63–92 (translation in *Russian Acad. Sci. Sb. Math.* 80 (1995) 309–333).
- [3] M.J. Cantero, L. Moral, L. Velázquez, Five-diagonal matrices and zeros of orthogonal polynomials on the unit circle, *Linear Algebra Appl.* 362 (2003) 29–56.
- [4] A.O. Gel'fond, *Calculus of Finite Differences*, International Monographs on Advanced Mathematics and Physics, Hindustan Publishing, Delhi, 1971.
- [5] Ya.L. Geronimus, *Orthogonal Polynomials: Estimates, Asymptotic Formulas, and Series of Polynomials Orthogonal on the Unit Circle and on an Interval*, Consultants Bureau, New York, 1961.
- [6] F. Gesztesy, B. Simon, m -functions and inverse spectral analysis for finite and semi-infinite Jacobi matrices, *J. d'Analyse Math.* 73 (1997) 267–297.
- [7] S. Khrushchev, Schur's algorithm, orthogonal polynomials, and convergence of Wall's continued fractions in $L^2(\mathbb{T})$, *J. Approx. Theory* 108 (2001) 161–248.
- [8] S. Khrushchev, Classification theorems for general orthogonal polynomials on the unit circle, *J. Approx. Theory* 116 (2002) 268–342.
- [9] R. Killip, B. Simon, Sum rules for Jacobi matrices and their applications to spectral theory, *Ann. of Math.*, to appear.
- [10] P. Nevai, *Orthogonal polynomials*, *Memoirs of the American Mathematical Society*, Vol. 18, No. 213, American Mathematical Society, Providence, RI, 1979, 185pp.
- [11] P. Nevai, *Research problems in orthogonal polynomials*, *Approximation Theory VI*, Vol. II (College Station, TX, 1989), Academic Press, Boston, MA, 1989, pp. 449–489.

- [12] H. Poincaré, Sur les equations linéaires aux differentielles ordinaires et aux différences finies, *Amer. J. Math.* 7 (1885) 203–258.
- [13] B. Simon, The classical moment problem as a self-adjoint finite difference operator, *Adv. in Math.* 137 (1998) 82–203.
- [14] B. Simon, *Orthogonal Polynomials on the Unit Circle*, Colloquium Publications, American Mathematical Society, Providence, RI, expected 2004.
- [15] G. Szegő, *Orthogonal Polynomials*, American Mathematical Society Colloquium Publications, Vol. 23, American Mathematical Society, Providence, RI, 1939; 3rd Edition, 1967.