



ELSEVIER

Available online at www.sciencedirect.com



Journal of Approximation Theory 129 (2004) 102–117

JOURNAL OF
Approximation
Theory

<http://www.elsevier.com/locate/jat>

Erratum

Convergence of Hermite–Fejér type interpolation of higher order on an arbitrary system of nodes^{☆, ☆ ☆}

Ying Guang Shi*

Department of Mathematics, Hunan Normal University, Changsha, Hunan 410081, China

Received 17 January 2002; accepted in revised form 23 April 2003

Communicated by József Szabados

Abstract

A criterion of convergence for general Hermite–Fejér-type interpolation of higher order on an arbitrary system of nodes is given.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Convergence; Hermite–Fejér interpolation

1. Introduction

Let $\mathbb{N}_1 := \{1, 3, 5, \dots\}$ and $\mathbb{N}_2 := \{2, 4, 6, \dots\}$. Let $n \in \mathbb{N}$ ($n \geq 2$), $m_{kn} \in \mathbb{N}$ ($k = 1, 2, \dots, n$, $n = 2, 3, \dots$), and

$$x_{0n} := 1 \geq x_{1n} > x_{2n} > \dots > x_{m_n} \geq x_{n+1,n} := -1, \quad n = 2, 3, \dots \quad (1.1)$$

In what follows, we shall often omit the superfluous notations, i.e., m_{kn}, x_{kn}, \dots will be denoted by m_k, x_k, \dots , etc. Throughout this paper let $N := N_n := \sum_{k=1}^n m_{kn} - 1$

[☆]doi of original article 10.1016/S0021-9045(03)000844, 10.1016/j.jat.2004.05.001

^{☆☆}Project 19971089 supported by National Natural Science Foundation of China.

*Corresponding author. Institute of Computational Mathematics and Scientific/Engineering Computing, Chinese Academy of Sciences, No. A1, Nansijie, Zhongguancun, P. O. Box 2719, Beijing 100080, PR China.

E-mail address: syg@lsec.cc.ac.cn.

and $m := \max_{1 \leq k \leq n, n \geq 2} m_{kn} < +\infty$. Denote by \mathbf{P}_N the set of polynomials of degree at most N and by A_{jk} the fundamental polynomials for Hermite interpolation, i.e., $A_{jk} \in \mathbf{P}_N$ satisfy

$$A_{jk}^{(p)}(x_q) = \delta_{jp} \delta_{kq}, \quad p = 0, 1, \dots, m_q - 1, \quad j = 0, 1, \dots, m_k - 1, \quad q, k = 1, 2, \dots, n. \tag{1.2}$$

The Hermite–Fejér-type interpolation for $f \in C[-1, 1]$ is given by

$$H_{nm}(f, x) = \sum_{k=1}^n f(x_k) A_{0k}(x). \tag{1.3}$$

In [5] the author established a criterion of convergence for Hermite–Fejér-type interpolation of higher order on an arbitrary system of nodes as follows ($\|\cdot\|$ stands for the uniform norm on $[-1, 1]$ and $f_i(x) := x^i, i = 1, 2, \dots$):

Theorem A (Shi [5, Theorem 4.2]). *Let $m_{kn} \equiv m \in \mathbb{N}_2$. Then*

$$\lim_{n \rightarrow \infty} \|H_{nm}(f) - f\| = 0 \tag{1.4}$$

holds for all $f \in C[-1, 1]$ if and only if

$$\|H_{nm}\| := \left\| \sum_{k=1}^n |A_{0k}| \right\| = o(1) \tag{1.5}$$

is true and (1.4) holds for the two monomials $f = f_i, i = 1, 2$.

The main aim of this paper is to establish a criterion of convergence for general Hermite–Fejér-type interpolation of higher order on an arbitrary system of nodes, replacing the assumption that $m_{kn} \equiv m \in \mathbb{N}_2$ by the assumption that all $m_{kn} \in \mathbb{N}_2$. That is the following:

Theorem. *Let all $m_{kn} \in \mathbb{N}_2$. Then relation (1.4) holds for all $f \in C[-1, 1]$ if and only if relation (1.5) is true and (1.4) holds for the two monomials $f = f_i, i = 1, 2$.*

This extension is not trivial. To prove the theorem we have to prove a series of auxiliary lemmas, which are of independent interest and put in the next section. Then we give the proof of the theorem in the last section.

2. Auxiliary lemmas

First, we state some known results needed later.

Lemma A (Borwein and Erdélyi [1, p. 235]). *Let $P \in \mathbf{P}_n$. Then*

$$\begin{aligned} |P(y)| &\leq |T_n(y)| \|P\| \\ &= \frac{1}{2} \{ [y + (y^2 - 1)^{1/2}]^n + [y - (y^2 - 1)^{1/2}]^n \} \|P\|, \quad |y| > 1, \end{aligned} \tag{2.1}$$

where T_n stands for the n th Chebyshev polynomial of the first kind.

Lemma B (Shi [5, Lemma 4.1]). *Let $P_k \in \mathbf{P}_n$, $k = 1, 2, \dots, M$, and $1 \geq y_1 > y_2 > \dots > y_M \geq -1$. If*

$$\left\| \sum_{k=1}^M |(x - y_k)P_k(x)| \right\| = \mu_n$$

and

$$\sum_{k=1}^M |P_k(y_j)| \leq v_n, \quad j = 1, 2, \dots, M,$$

then

$$\left\| \sum_{k=1}^M |P_k| \right\| \leq 2(n^2 \mu_n + v_n).$$

In particular, if $M = 1$ and $P_1(y_1) = 0$, $|y_1| < 1$, then

$$\|P_1\| \leq \frac{4n\mu_n}{(1 - y_1^2)^{1/2}}.$$

Lemma C (Shi [5, Theorem 2.1]). *If for a fixed n , $m_k - j$ is odd and $j < i \leq m_k - 1$ then*

$$|A_{ik}(x)| \leq \frac{j!}{i!} d_k^{i-j} |A_{jk}(x)|, \quad x \in [-1, 1], \quad 1 \leq k \leq n, \tag{2.2}$$

where $d_k := \max\{|x_k - x_{k-1}|, |x_k - x_{k+1}|\}$, $k = 1, 2, \dots, n$.

Lemma 1. *Let all $m_{kn} \in \mathbb{N}_2$. If*

$$\|H_{nm}\| = \mu_n, \tag{2.3}$$

then

$$\left\| \sum_{k=1}^n |A_{1k}| \right\| \leq 8m^2 n^2 \mu_n. \tag{2.4}$$

Proof. By the same argument as that of [5, Theorem 2.3] we can get

$$\sum_{k=1}^n (x - x_k) A_{1k}(x) = \sum_{k=1}^n |(x - x_k) A_{1k}(x)| \leq \sum_{k=1}^n (x - x_k)^2 A_{0k}(x), \quad x \in \mathbb{R}. \tag{2.5}$$

Hence by (2.3) we obtain

$$\sum_{k=1}^n (x - x_k) A_{1k}(x) \leq 4\mu_n,$$

which according to Lemma B gives (2.4) (by $\deg A_{ik} < nm$). \square

Lemma 2. Let A_{jk}^* be the fundamental polynomials for Hermite interpolation based on the system of nodes

$$x_{kn}^* = ax_{kn}, \quad 0 < a < 1, \quad k = 1, 2, \dots, n, \quad n = 2, 3, \dots \quad (2.6)$$

Then with the notations $x = \cos \theta$ and $x^* = \cos \theta^*$

$$\theta_{kn}^* = \arccos(a \cos \theta_{kn}), \quad k = 1, 2, \dots, n, \quad (2.7)$$

$$A_{jk}^*(x) = a^j A_{jk}(x/a), \quad k = 1, 2, \dots, n, \quad j = 0, 1, \dots, m_k - 1, \quad (2.8)$$

$$\max_{|x| \leq 1} |A_{jk}^*(x)| = a^j \max_{|x| \leq 1/a} |A_{jk}(x)|, \quad k = 1, 2, \dots, n, \quad j = 0, 1, \dots, m_k - 1 \quad (2.9)$$

and

$$\|H_{nm}^*\| := \left\| \sum_{k=1}^n |A_{0k}^*| \right\| \leq [a^{-1} + (a^{-2} - 1)^{1/2}]^n \|H_{nm}\|. \quad (2.10)$$

Proof. Relations (2.7)–(2.9) may be obtained directly from the definition. Now let us prove (2.10). Assume that

$$\sum_{k=1}^n |A_{0k}(y)| = \max_{|x| \leq 1/a} \sum_{k=1}^n |A_{0k}(x)|, \quad |y| \leq 1/a.$$

This relation, together with (2.8), yields

$$\begin{aligned} \|H_{nm}^*\| &= \max_{|x| \leq 1} \sum_{k=1}^n |A_{0k}^*(x)| = \max_{|x| \leq 1} \sum_{k=1}^n |A_{0k}(x/a)| \\ &= \max_{|x| \leq 1/a} \sum_{k=1}^n |A_{0k}(x)| = \sum_{k=1}^n |A_{0k}(y)|. \end{aligned} \quad (2.11)$$

If $|y| \leq 1$ then by (2.11) we have that

$$\|H_{nm}^*\| = \sum_{k=1}^n |A_{0k}(y)| \leq \|H_{nm}\|,$$

if $|y| > 1$ then by (2.11) and (2.1) we have that

$$\begin{aligned} \|H_{nm}^*\| &= \sum_{k=1}^n |A_{0k}(y)| = \sum_{k=1}^n [\operatorname{sgn} A_{0k}(y)] A_{0k}(y) \\ &\leq \frac{1}{2} \{ [y + (y^2 - 1)^{1/2}]^n + [y - (y^2 - 1)^{1/2}]^n \} \left\| \sum_{k=1}^n [\operatorname{sgn} A_{0k}(y)] A_{0k} \right\| \\ &\leq [|y| + (y^2 - 1)^{1/2}]^n \|H_{nm}\| \\ &\leq [a^{-1} + (a^{-2} - 1)^{1/2}]^n \|H_{nm}\|. \quad \square \end{aligned}$$

In the sequel c, c', \dots will stand for positive constants depending only on m , unless otherwise indicated; their values may be different at different occurrences, even in subsequent formulas; in addition, $c_r, c_r' \geq 1$ will stand for positive constants depending only on r and increasing with respect to r , their values may also be different at different occurrences, even in subsequent formulas.

Lemma 3. Let $r \in \mathbb{N}, p > 0$, and

$$f(\theta) = (\sin p\theta)^r. \tag{2.12}$$

Then

$$|f^{(j)}(\theta)| \leq c_j (rp)^j |\sin p\theta|^{r-j}, \quad 0 \leq j \leq r. \tag{2.13}$$

Proof (By induction). Relation (2.13) with $j = 0$ is trivial. Suppose now, as an induction hypothesis, that relation (2.13) holds for all $j, j \leq q < r$. Differentiating (2.12) yields

$$f'(\theta) = rp(\sin p\theta)^{r-1} \cos p\theta$$

and then differentiating the above relation q times gives (using the induction hypothesis)

$$\begin{aligned} |f^{(q+1)}(\theta)| &= rp \left| \sum_{j=0}^q \binom{q}{j} [(\sin p\theta)^{r-1}]^{(j)} (\cos p\theta)^{(q-j)} \right| \\ &\leq rp \sum_{j=0}^q \binom{q}{j} c_j [(r-1)p]^j |\sin p\theta|^{r-1-j} p^{q-j} \\ &\leq c_q (rp)^{q+1} |\sin p\theta|^{r-1-q} \sum_{j=0}^q \binom{q}{j} \\ &= c_q (rp)^{q+1} |\sin p\theta|^{r-1-q} 2^q \\ &= c_{q+1} (rp)^{q+1} |\sin p\theta|^{r-(q+1)}, \end{aligned}$$

which shows that relation (2.13) is true for $j = q + 1$. By induction this proves (2.13). \square

Lemma 4. Let $r \in \mathbb{N}$, $p > 0$, and

$$g(\theta) := \left[\frac{\sin(p\theta/2)}{\sin(\theta/2)} \right]^r. \tag{2.14}$$

Then

$$|g^{(j)}(\theta)| \leq c_j (rp)^j |\sin(\theta/2)|^{-r-j}, \quad \theta \neq 0, \quad 0 \leq j \leq r. \tag{2.15}$$

Proof (By induction). Relation (2.14) implies

$$|g(\theta)| \leq |\sin(\theta/2)|^{-r},$$

which implies that relation (2.15) is true for $j = 0$. Suppose now, as an induction hypothesis, that relation (2.15) is true for all j , $j \leq q < r$. Rewrite (2.14) as

$$g(\theta) [\sin(\theta/2)]^r = [\sin(p\theta/2)]^r.$$

Differentiating this relation $(q + 1)$ times and then using (2.13) (replacing c_j by c_j' in it) and the induction hypothesis, we get

$$\begin{aligned} & |g^{(q+1)}(\theta) [\sin(\theta/2)]^r| \\ &= \left| \left[[\sin(p\theta/2)]^r \right]^{(q+1)} - \sum_{j=0}^q \binom{q+1}{j} g^{(j)}(\theta) \left[[\sin(\theta/2)]^r \right]^{(q+1-j)} \right| \\ &\leq c_{q+1}' (rp/2)^{q+1} |\sin(p\theta/2)|^{r-q-1} \\ &\quad + \sum_{j=0}^q \binom{q+1}{j} c_j (rp)^j |\sin(\theta/2)|^{-r-j} c_{q+1-j}' (r/2)^{q+1-j} |\sin(\theta/2)|^{r-(q+1-j)} \\ &\leq c_{q+1}' (rp)^{q+1} + c_q c_{q+1}' (rp)^{q+1} |\sin(\theta/2)|^{-q-1} \sum_{j=0}^q \binom{q+1}{j} \\ &\leq c_{q+1} (rp)^{q+1} |\sin(\theta/2)|^{-q-1} \sum_{j=0}^{q+1} \binom{q+1}{j} \\ &\leq c_{q+1} (rp)^{q+1} |\sin(\theta/2)|^{-q-1}. \end{aligned}$$

Then

$$|g^{(q+1)}(\theta)| \leq c_{q+1} (rp)^{q+1} |\sin(\theta/2)|^{-r-(q+1)},$$

which shows that relation (2.15) is true for $j = q + 1$. By induction this proves (2.15). \square

Lemma 5. Let $\psi \in \mathbf{P}_n$ and

$$\phi(\theta) = \psi(\cos \theta), \quad 0 \leq \theta \leq \pi. \tag{2.16}$$

Then

$$|\psi^{(j)}(\cos \theta)| \leq \frac{c_j}{(\sin \theta)^{j^2}} \sum_{v=1}^j |\phi^{(v)}(\theta)| (\sin \theta \neq 0), \quad j \geq 1. \tag{2.17}$$

Proof. First let us prove that

$$\phi^{(j)}(\theta) = \sum_{v=1}^j c_{vj}(\theta) \psi^{(v)}(\cos \theta), \quad j \geq 1, \tag{2.18}$$

where $c_{vj}(\theta)$ is a trigonometric polynomial of degree v and

$$c_{jj}(\theta) = (-\sin \theta)^j, \quad |c_{vj}(\theta)| \leq j!, \quad v = 1, 2, \dots, j. \tag{2.19}$$

We use induction. It is easy to see that relations (2.18) and (2.19) hold for $j = 1$. Suppose now, as an induction hypothesis, that relations (2.18) and (2.19) are true for all j , $1 \leq j \leq q$. By differentiation of (2.18) with $j = q$ we obtain

$$\begin{aligned} \phi^{(q+1)}(\theta) &= \sum_{v=1}^q [c_{vq}'(\theta) \psi^{(v)}(\cos \theta) - c_{vq}(\theta) \psi^{(v+1)}(\cos \theta) \sin \theta] \\ &= \sum_{v=1}^{q+1} [c_{vq}'(\theta) - c_{v-1,q}(\theta) \sin \theta] \psi^{(v)}(\cos \theta) \\ &= \sum_{v=1}^{q+1} c_{v,q+1}(\theta) \psi^{(v)}(\cos \theta), \end{aligned}$$

where $c_{0,q}(\theta) = c_{q+1,q}(\theta) = 0$ and

$$c_{v,q+1}(\theta) = c_{vq}'(\theta) - c_{v-1,q}(\theta) \sin \theta. \tag{2.20}$$

Hence by (2.19) $c_{q+1,q+1}(\theta) = -c_{qq}(\theta) \sin \theta = (-\sin \theta)^{q+1}$ and $\|c_{q+1,q+1}\| \leq 1$ (here $\|\cdot\|$ stands for the uniform norm on $[0, 2\pi]$); by Bernstein inequality it follows from (2.20) that $\|c'_{vq}\| \leq v \|c_{vq}\|$ and hence

$$\|c_{v,q+1}\| \leq v \|c_{vq}\| + \|c_{v-1,q}\| \leq (v+1)q! \leq (q+1)!, \quad 1 \leq v \leq q.$$

This shows that relations (2.18) and (2.19) are true for $j = q + 1$. By induction this proves (2.18) and (2.19).

For the proof of (2.17) we again use induction. It is easy to see that relation (2.17) is true for $j = 1$. Suppose now, as an induction hypothesis, that relation (2.17) is true for all j , $1 \leq j \leq q$. It follows from (2.18), (2.19), and the induction

hypothesis that

$$\begin{aligned}
 |\psi^{(q+1)}(\cos \theta)| &\leq \frac{1}{(\sin \theta)^{q+1}} \left[|\phi^{(q+1)}(\theta)| + (q+1)! \sum_{j=1}^q |\psi^{(j)}(\cos \theta)| \right] \\
 &\leq \frac{1}{(\sin \theta)^{q+1}} \left[|\phi^{(q+1)}(\theta)| + (q+1)! \sum_{j=1}^q \frac{c_j}{(\sin \theta)^{j^2}} \sum_{v=1}^j |\phi^{(v)}(\theta)| \right] \\
 &\leq \frac{1}{(\sin \theta)^{q+1}} \left[|\phi^{(q+1)}(\theta)| + \frac{(q+1)!c_q}{(\sin \theta)^{q^2}} \sum_{j=1}^q \sum_{v=1}^j |\phi^{(v)}(\theta)| \right] \\
 &\leq \frac{1}{(\sin \theta)^{q+1}} \left[|\phi^{(q+1)}(\theta)| + \frac{q(q+1)!c_q}{(\sin \theta)^{q^2}} \sum_{j=1}^q |\phi^{(j)}(\theta)| \right] \\
 &\leq \frac{c_{q+1}}{(\sin \theta)^{(q+1)^2}} \sum_{j=1}^{q+1} |\phi^{(j)}(\theta)|.
 \end{aligned}$$

This shows that relation (2.17) is true for $j = q + 1$ and by induction proves (2.17). \square

Lemma 6. Let $0 \leq \theta, \tau \leq \pi$, g be defined by (2.14), and

$$\phi(\theta) = p^{-r}[g(\theta + \tau) + g(\theta - \tau)]. \tag{2.21}$$

Let ψ be defined by (2.16). Then

$$|\psi^{(j)}(\cos \theta)| \leq \frac{c_j(rp)^j}{p^r(\sin \theta)^{j^2}} \left| \frac{3\pi}{\theta - \tau} \right|^{r+j}, \quad 0 < \theta < \pi, \theta \neq \tau, 0 \leq j \leq r. \tag{2.22}$$

Proof. Suppose without loss of generality that $0 < \theta \leq \pi/2$. By (2.21) and (2.15)

$$|\phi^{(j)}(\theta)| \leq \frac{c_j(rp)^j}{p^r} \left\{ \frac{1}{|\sin[(\theta + \tau)/2]|^{r+j}} + \frac{1}{|\sin[(\theta - \tau)/2]|^{r+j}} \right\}. \tag{2.23}$$

Since $0 < \theta \leq \pi/2$, we have $0 < (\theta + \tau)/2 \leq 3\pi/4$ and hence

$$\sin \frac{\theta + \tau}{2} \geq \begin{cases} \frac{2}{\pi} \left(\frac{\theta + \tau}{2} \right) = \frac{\theta + \tau}{\pi}, & \frac{\theta + \tau}{2} \leq \frac{\pi}{2}, \\ 2^{-1/2} \geq 2^{-1/2} \frac{2}{\pi} \left(\frac{\theta + \tau}{2} \right) \geq \frac{\theta + \tau}{2\pi}, & \frac{\pi}{2} < \frac{\theta + \tau}{2} \leq \frac{3\pi}{4}. \end{cases} \tag{2.24}$$

This implies

$$\sin \frac{\theta + \tau}{2} \geq \frac{\theta + \tau}{2\pi}, \quad 0 < \frac{\theta + \tau}{2} \leq \frac{3\pi}{4}. \tag{2.25}$$

Of course, we have

$$\left| \sin \frac{\theta - \tau}{2} \right| \geq \left| \frac{\theta - \tau}{\pi} \right|.$$

Substituting the above inequalities into (2.23), we obtain

$$\begin{aligned}
 |\phi^{(j)}(\theta)| &\leq \frac{c_j(rp)^j}{p^r} \left\{ \left| \frac{2\pi}{\theta + \tau} \right|^{r+j} + \left| \frac{\pi}{\theta - \tau} \right|^{r+j} \right\} \\
 &\leq \frac{c_j(rp)^j}{p^r} \left| \frac{3\pi}{\theta - \tau} \right|^{r+j}, \quad j \geq 0.
 \end{aligned}
 \tag{2.26}$$

inserting estimation (2.26) into (2.17), we have

$$\begin{aligned}
 |\psi^{(j)}(\cos \theta)| &\leq \frac{c_j}{(\sin \theta)^{j^2}} \sum_{v=1}^j |\phi^{(v)}(\theta)| \\
 &\leq \frac{c_j}{(\sin \theta)^{j^2}} \sum_{v=1}^j \frac{c_v(rp)^v}{p^r} \left| \frac{3\pi}{\theta - \tau} \right|^{r+v} \\
 &\leq \frac{c_j(rp)^j}{p^r(\sin \theta)^{j^2}} \left| \frac{3\pi}{\theta - \tau} \right|^{r+j}. \quad \square
 \end{aligned}$$

Lemma 7. *Let all $m_{kn} \in \mathbb{N}_1$ or all $m_{kn} \in \mathbb{N}_2$. If*

$$\|H_{nm}\| = \mu_n, \tag{2.27}$$

then

$$\theta_{k+1,n} - \theta_{k,n} \leq \frac{c \ln(n\mu_n)}{n}, \quad k = 0, 1, \dots, n, \quad n = 2, 3, \dots \tag{2.28}$$

Proof. The proof follows the line given by Erdős and Turán [3, p. 718; 2,4]. Let

$$n \geq \ln(n^{m^2+4}\mu_n) + 2, \tag{2.29}$$

$r \in \mathbb{N}_2$, and $p \in \mathbb{N}_1$ ($p \geq 3$) such that

$$\frac{1}{2}r(p - 1) \leq n - 1. \tag{2.30}$$

Suppose that

$$\begin{aligned}
 \max_{0 \leq k \leq n} (\theta_{k+1,n} - \theta_{k,n}) &= \theta_{i+1,n} - \theta_{i,n} := 2\delta_n, \quad \tau := \frac{1}{2}(\theta_{i+1,n} + \theta_{i,n}), \\
 \xi &:= \cos \tau.
 \end{aligned}
 \tag{2.31}$$

Let ϕ and ψ be defined by (2.21) and (2.16), respectively. Observing that $\psi(\cdot) \in \mathbf{P}_{n-1}$, we see

$$\phi(\theta) = \psi(\cos \theta) = \psi(x) = \sum_{k=1}^n \sum_{j=0}^{m_k-1} \psi^{(j)}(x_k) A_{jk}(x). \tag{2.32}$$

If all $m_{kn} \in \mathbb{N}_2$, then we have by (2.4) and (2.2)

$$\|A_{jk}\| \leq \frac{1}{j!} d_k^{j-1} \|A_{1k}\| \leq 8m^2n^2\mu_n, \quad j \geq 1, \quad 1 \leq k \leq n,$$

which, coupled with (2.27), gives

$$\|A_{jk}\| \leq 8m^2 n^2 \mu_n, \quad j \geq 0, \quad 1 \leq k \leq n. \tag{2.33}$$

If all $m_{kn} \in \mathbb{N}_1$, then we have by (2.2)

$$\|A_{jk}\| \leq \frac{1}{j!} d_k^j \|A_{0k}\| \leq 2\mu_n,$$

which also implies (2.33).

We separate two cases when $\min\{\theta_1, \pi - \theta_n\} \geq 1/n$ and when $\min\{\theta_1, \pi - \theta_n\} < 1/n$.

Case 1: $\min\{\theta_1, \pi - \theta_n\} \geq 1/n$. In [4, p. 537] it is proved that $\phi(\tau) \geq 1$. Using this inequality, and applying (2.31)–(2.33) and (2.22), we obtain

$$\begin{aligned} 1 \leq \phi(\tau) &= \sum_{k=1}^n \sum_{j=0}^{m_k-1} \psi^{(j)}(x_k) A_{jk}(\xi) \\ &\leq \sum_{k=1}^n \sum_{j=0}^{m_k-1} \frac{c_j (rp)^j}{p^r (\sin \theta_k)^{j^2}} \left| \frac{3\pi}{\theta_k - \tau} \right|^{r+j} 8m^2 n^2 \mu_n \\ &\leq \frac{c_m (rp)^m n^2 \mu_n}{p^r} \left(\frac{3\pi}{\delta_n} \right)^{r+m} \sum_{k=1}^n \sum_{j=0}^{m_k-1} \frac{1}{(\sin \theta_k)^{j^2}} \\ &\leq \frac{c_m (rp)^m n^{(m-1)^2+3} \mu_n}{p^r} \left(\frac{3\pi}{\delta_n} \right)^{r+m}. \end{aligned}$$

Since the inequality $p \geq 3$ implies that $p - 1 \geq 2p/3$, by (2.30) we get $rp \leq 3n$. Using this inequality and observing that $\delta_n \geq \pi/[2(n + 1)]$, we get

$$1 \leq c_m n^{m^2+4} \mu_n \left(\frac{3\pi}{p\delta_n} \right)^r$$

and hence

$$\delta_n \leq \frac{3\pi}{p} [c_m n^{m^2+4} \mu_n]^{1/r}. \tag{2.34}$$

Now choose

$$r = 2 \left[\frac{1}{2} \ln(n^{m^2+4} \mu_n) \right]$$

and

$$p = 1 + 2 \left\lceil \frac{n-1}{r} \right\rceil,$$

which obviously satisfy condition (2.30). From these definitions we can obtain estimations

$$r \leq \ln(n^{m^2+4} \mu_n) \tag{2.35}$$

and

$$r \geq 2 \left\{ \frac{1}{2} \ln(n^{m^2+4} \mu_n) - 1 \right\} \geq \frac{1}{2} \ln(n^{m^2+4} \mu_n), \tag{2.36}$$

because $m \geq 2$, $n \geq 2$, and $\mu_n \geq 1$; meanwhile, by (2.29) and (2.35)

$$p \geq 1 + 2 \left(\frac{n-1}{r} - 1 \right) = \frac{2(n-1) - r}{r} \geq \frac{n}{\ln(n^{m^2+4}\mu_n)}. \tag{2.37}$$

Inserting estimations (2.36) and (2.37) into (2.34) we see

$$\delta_n \leq \frac{3\pi \ln(n^{m^2+4}\mu_n)}{n} \{c_m n^{m^2+4} \mu_n\}^{2/\ln(n^{m^2+4}\mu_n)}. \tag{2.38}$$

It is clear that for $x > 0$ and $x \neq 1$

$$x^{1/\ln x} = e^{\ln(x^{1/\ln x})} = e.$$

Using this identity inequality (2.38) becomes

$$\delta_n \leq \frac{3\pi \ln(n^{m^2+4}\mu_n)e^2}{n} c_m^{2/\ln(n^{m^2+4}\mu_n)} \leq \frac{c \ln(n\mu_n)}{n}. \tag{2.39}$$

Case 2: $\min\{\theta_1, \pi - \theta_n\} < 1/n$. In this case let us consider the Hermite interpolation based on the system of nodes (2.6) with

$$a = \frac{n^2}{n^2 + 1}. \tag{2.40}$$

By (2.10)

$$\|H_{nm}^*\| \leq \left(\frac{n^2 + 1}{n^2} + \frac{2n + 1}{n^2} \right)^n \mu_n \leq \left(1 + \frac{3}{n} \right)^n \mu_n \leq c\mu_n. \tag{2.41}$$

By means of (2.6) we see that

$$a \geq a \cos \theta_1 = \cos \theta_1^* = 1 - 2 \sin^2 \frac{\theta_1^*}{2} \geq 1 - \frac{(\theta_1^*)^2}{2}$$

and hence according to (2.40)

$$\theta_1^* \geq [2(1 - a)]^{1/2} \geq \frac{1}{n}. \tag{2.42}$$

Similarly, we can conclude that

$$\pi - \theta_n^* \geq \frac{1}{n}. \tag{2.43}$$

Then we can apply the conclusion of Case 1 to the Hermite interpolation based on the system of nodes (2.6) with (2.40) to obtain (using (2.41))

$$\theta_{k+1,n}^* - \theta_{k,n}^* \leq \frac{c \ln(n\mu_n)}{n}, \quad k = 0, 1, \dots, n, \quad n = 2, 3, \dots$$

To prove (2.28) it remains to estimate the difference $\theta_k - \theta_k^*$. From (2.6) and (2.40) it follows that

$$\begin{aligned} \left| \sin \frac{\theta_k + \theta_k^*}{2} \sin \frac{\theta_k - \theta_k^*}{2} \right| &= \frac{1}{2} |\cos \theta_k - \cos \theta_k^*| = \frac{1}{2} |\cos \theta_k - a \cos \theta_k| \\ &= \frac{1}{2} (1 - a) |\cos \theta_k| \leq \frac{1}{2} (1 - a) = \frac{1}{2(n^2 + 1)}. \end{aligned} \tag{2.44}$$

Meanwhile, by virtue of (2.42) and (2.43) we have

$$\begin{aligned} \sin \frac{\theta_k + \theta_k^*}{2} &\geq \frac{2}{\pi} \min \left\{ \frac{\theta_k + \theta_k^*}{2}, \pi - \frac{\theta_k + \theta_k^*}{2} \right\} \\ &\geq \frac{2}{\pi} \min \left\{ \frac{\theta_k^*}{2}, \frac{\pi - \theta_k^*}{2} \right\} \geq \frac{1}{\pi n} \end{aligned}$$

and

$$\left| \sin \frac{\theta_k - \theta_k^*}{2} \right| \geq \frac{|\theta_k - \theta_k^*|}{\pi},$$

which, coupled with (2.44), yield

$$|\theta_k - \theta_k^*| \leq \frac{\pi^2 n}{2(n^2 + 1)} \leq \frac{\pi^2}{2n}.$$

At last, we conclude

$$\begin{aligned} \theta_{k+1} - \theta_k &\leq \theta_{k+1}^* - \theta_k^* + |\theta_{k+1}^* - \theta_{k+1}| + |\theta_k^* - \theta_k| \\ &\leq \frac{c \ln(n\mu_n)}{n} + \frac{\pi^2}{n} \leq \frac{c \ln(n\mu_n)}{n}. \end{aligned}$$

This completes the proof supposing (2.29). When condition (2.29) does not hold, the statement is obvious. \square

Lemma 8. *If relation (1.5) is true then*

$$\theta_{k+1,n} - \theta_{kn} \geq \frac{c}{n}, \quad k = 1, 2, \dots, n - 1, \quad n = 2, 3, \dots \quad (2.45)$$

Proof. The proof follows the line given by Erdős and Turán [3, p. 718; 4]. By Rolle theorem and Bernstein inequality

$$\frac{1}{|\theta_{k+1} - \theta_k|} = \left| \frac{A_{0k}(\cos \theta_k) - A_{0k}(\cos \theta_{k+1})}{\theta_k - \theta_{k+1}} \right| = \left| \frac{d[A_{0k}(\cos \theta)]}{d\theta} \right|_{\theta=\theta'} \leq c(mn - 1),$$

which is equivalent to (2.45). \square

3. Proof of theorem

It suffices to show the sufficiency according to Banach–Steinhaus theorem. To this end again applying Banach–Steinhaus theorem it is enough to show that (1.4) holds for every polynomial.

Assume that P is an arbitrary polynomial. Let N be so large that $P \in \mathbf{P}_N$. Then

$$P(x) = \sum_{k=1}^n \sum_{j=0}^{m_k-1} P^{(j)}(x_k) A_{jk}(x)$$

and hence

$$\begin{aligned}
 R_{nm}(P, x) &:= |H_{nm}(P, x) - P(x)| = \left| \sum_{k=1}^n \sum_{j=1}^{m_k-1} P^{(j)}(x_k) A_{jk}(x) \right| \\
 &= \left| \sum_{k=1}^n \sum_{j=1}^{m-1} P^{(j)}(x_k) A_{jk}(x) \right| \leq \sum_{j=1}^{m-1} \left| \sum_{k=1}^n P^{(j)}(x_k) A_{jk}(x) \right| := \sum_{j=1}^{m-1} S_j.
 \end{aligned}
 \tag{3.1}$$

(Here we agree that $A_{jk} \equiv 0$ for $j \geq m_k$.) To estimate S_j we separate three cases when $j = 1, j = 2,$ and $j \geq 3$ (if $m = 2$ then only the first case can occur).

First, by the same argument as that of [5, Theorem 4.1, pp. 73, 74] we can get the estimations

$$S_1 \leq 3 \|P\|^* r_{nm}(x)
 \tag{3.2}$$

and

$$S_2 \leq c \|P\|^* r_{nm}(x),
 \tag{3.3}$$

where

$$\|P\|^* := \max_{0 \leq j \leq m-1} \|P^{(j)}\|$$

and

$$r_{nm}(x) := R_{nm}(f_1, x) + R_{nm}(f_2, x).$$

Next, to estimate S_j ($j \geq 3$) we need some preliminaries. Relation (4.12) in [5] states that

$$\sum_{k=1}^n (x - x_k)^2 A_{0k}(x) \leq 2r_{nm}(x),$$

which, together with (2.5), gives

$$\sum_{k=1}^n (x - x_k) A_{1k}(x) \leq 2r_{nm}(x).
 \tag{3.4}$$

Using this estimate and applying Lemma B, we obtain estimates

$$\|A_{1k}\| \leq \frac{8mn \|r_{nm}\|}{\sin \theta_k} \quad (\sin \theta_k \neq 0), \quad k = 1, 2, \dots, n
 \tag{3.5}$$

and

$$\|A_{1k}\| \leq 4m^2 n^2 \|r_{nm}\|, \quad k = 1, 2, \dots, n.
 \tag{3.6}$$

Now for the estimate of S_j ($j \geq 3$) we use (2.2) and (3.4) to get that for a fixed x

$$\begin{aligned}
 S_j &\leq \|P^{(j)}\| \sum_{k=1}^n |A_{jk}(x)| \leq \|P^{(j)}\| \sum_{k=1}^n d_k^{j-1} |A_{1k}(x)| \leq c \|P^{(j)}\| \sum_{k=1}^n d_k^2 |A_{1k}(x)| \\
 &= c \|P^{(j)}\| \left[\sum_{|x_k-x| \geq d_k^2} d_k^2 |A_{1k}(x)| + \sum_{|x_k-x| < d_k^2} d_k^2 |A_{1k}(x)| \right] \\
 &\leq c \|P^{(j)}\| \left[\sum_{|x_k-x| \geq d_k^2} (x-x_k) A_{1k}(x) + \sum_{|x_k-x| < d_k^2} d_k^2 |A_{1k}(x)| \right] \\
 &\leq c \|P^{(j)}\| \left[\|r_{nm}\| + \sum_{|x_k-x| < d_k^2} d_k^2 |A_{1k}(x)| \right]. \tag{3.7}
 \end{aligned}$$

Here we have to estimate the term $d_k^2 |A_{1k}(x)|$. By (1.5) and (2.28),

$$\begin{aligned}
 |x_k - x_{k+1}| &= |\cos \theta_k - \cos \theta_{k+1}| \leq (\theta_{k+1} - \theta_k)(\sin \theta_k + \theta_{k+1} - \theta_k) \\
 &\leq c \frac{\ln n}{n} \left(\sin \theta_k + \frac{\ln n}{n} \right),
 \end{aligned}$$

similarly,

$$\begin{aligned}
 |x_k - x_{k-1}| &= |\cos \theta_k - \cos \theta_{k-1}| \leq (\theta_k - \theta_{k-1})(\sin \theta_k + \theta_k - \theta_{k-1}) \\
 &\leq c \frac{\ln n}{n} \left(\sin \theta_k + \frac{\ln n}{n} \right).
 \end{aligned}$$

Thus

$$d_k \leq c \frac{\ln n}{n} \left(\sin \theta_k + \frac{\ln n}{n} \right), \quad k = 1, 2, \dots, n. \tag{3.8}$$

We distinguish two cases.

Case 1: $\sin \theta_k \geq (\ln n)/n$. In this case by (3.5) and (3.8)

$$\begin{aligned}
 d_k^2 |A_{1k}(x)| &\leq c \left(\frac{\ln n}{n} \right)^2 \left(\sin \theta_k + \frac{\ln n}{n} \right)^2 \frac{8mn \|r_{nm}\|}{\sin \theta_k} \\
 &\leq c \left(\frac{\ln n}{n} \right)^2 (2 \sin \theta_k)^2 \frac{8mn \|r_{nm}\|}{\sin \theta_k} \\
 &\leq c \frac{(\ln n)^2 \|r_{nm}\|}{n}.
 \end{aligned}$$

Case 2: $\sin \theta_k < (\ln n)/n$. In this case by (3.6) and (3.8)

$$\begin{aligned} d_k^2 |A_{1k}(x)| &\leq c \left(\frac{\ln n}{n}\right)^2 \left(\sin \theta_k + \frac{\ln n}{n}\right)^2 4m^2 n^2 \|r_{nm}\| \\ &\leq c \left(\frac{\ln n}{n}\right)^2 \left(2 \frac{\ln n}{n}\right)^2 4m^2 n^2 \|r_{nm}\| \\ &\leq c \frac{(\ln n)^4 \|r_{nm}\|}{n^2} \\ &\leq c \frac{(\ln n)^2 \|r_{nm}\|}{n}. \end{aligned}$$

Thus in both cases

$$d_k^2 |A_{1k}(x)| \leq c \frac{(\ln n)^2 \|r_{nm}\|}{n}. \tag{3.9}$$

It remains to estimate $K_n := \text{card}\{k: |x_k - x| \leq d_k^2\}$. Assume without loss of generality that $\theta_k \leq \pi/2$, $1 \leq k \leq n - 1$. By (2.25) and (2.45)

$$\begin{aligned} |x_k - x_{k+1}| &= |\cos \theta_k - \cos \theta_{k+1}| = |2 \sin[(\theta_k + \theta_{k+1})/2] \sin[(\theta_k - \theta_{k+1})/2]| \\ &\geq c(\theta_k + \theta_{k+1})(\theta_k - \theta_{k+1}) \geq c(\theta_k - \theta_{k+1})^2 \geq c/n^2. \end{aligned} \tag{3.10}$$

Since (3.8) implies $d_k \leq c(\ln n)/n$, we can conclude that

$$K_n \leq c(\ln n)^2. \tag{3.11}$$

Then by (3.9) and (3.11)

$$\sum_{|x_k - x| \leq d_k^2} d_k^2 |A_{1k}(x)| \leq c \frac{(\ln n)^4 \|r_{nm}\|}{n} \leq c \|r_{nm}\|. \tag{3.12}$$

By (3.1)–(3.3), (3.7), and (3.12) we obtain

$$\|R_{nm}(P)\| \leq c \|P\|^* \|r_{nm}\|. \tag{3.13}$$

Therefore

$$\lim_{n \rightarrow \infty} \|R_{nm}(P)\| = 0.$$

This completes the proof. \square

Acknowledgments

Y.G.S. thanks the referee for carefully reading my manuscript and many helpful comments on improving the original manuscript.

References

- [1] P. Borwein, T. Erdélyi, *Polynomials and Polynomials Inequalities*, Springer, New York, 1995.
- [2] P. Erdős, On the uniform distribution of the roots of certain polynomials, *Ann. of Math.* 43 (1942) 59–64.
- [3] P. Erdős, P. Turán, On interpolation II, *Ann. of Math.* 39 (1938) 703–724.
- [4] P. Erdős, P. Turán, On interpolation III, *Ann. of Math.* 41 (1940) 510–553.
- [5] Y.G. Shi, On Hermite interpolation, *J. Approx. Theory* 105 (2000) 49–86.