



# Exact Markov inequalities for the Hermite and Laguerre weights<sup>☆</sup>

Lozko Milev\*, Nikola Naidenov

*Department of Mathematics, University of Sofia, Blvd. James Boucher 5, 1164 Sofia, Bulgaria*

Received 11 August 2004; accepted 26 September 2005

Communicated by Tamás Erdélyi  
Available online 7 December 2005

## Abstract

Denote by  $\pi_n$  the set of all real algebraic polynomials of degree at most  $n$  and let  $U_n := \{e^{-x^2} p(x) : p \in \pi_n\}$ ,  $V_n := \{e^{-x} p(x) : p \in \pi_n\}$ . We prove the following exact Markov inequalities:

$$\|u^{(k)}\|_{\mathbb{R}} \leq \|u_{*,n}^{(k)}\|_{\mathbb{R}} \|u\|_{\mathbb{R}}, \quad \forall u \in U_n, \quad \forall k \in \mathbb{N},$$

and

$$\|v^{(k)}\|_{\mathbb{R}_+} \leq \|v_{*,n}^{(k)}\|_{\mathbb{R}_+} \|v\|_{\mathbb{R}_+}, \quad \forall v \in V_n, \quad \forall k \in \mathbb{N},$$

where  $\|\cdot\|_{\mathbb{R}}$  ( $\|\cdot\|_{\mathbb{R}_+}$ ) is the supremum norm on  $\mathbb{R}$  ( $\mathbb{R}_+ := [0, \infty)$ ) and  $u_{*,n}$  ( $v_{*,n}$ ) is the Chebyshev polynomial from  $U_n$  ( $V_n$ ).

© 2005 Elsevier Inc. All rights reserved.

MSC: 41A17; 26D10

Keywords: Weighted polynomials; Markov inequality

## 1. Introduction

Denote by  $\pi_n$  the set of all real algebraic polynomials of degree not exceeding  $n$ , and by  $\|\cdot\|_I$  the supremum norm for a given interval  $I \subseteq \mathbb{R}$ ,  $\|f\|_I := \sup_{x \in I} |f(x)|$ .

<sup>☆</sup> Research was supported by the Bulgarian Ministry of Education and Science under Contract MM-1402/2004.

\* Corresponding author.

E-mail addresses: [milev@fmi.uni-sofia.bg](mailto:milev@fmi.uni-sofia.bg) (L. Milev), [nikola@fmi.uni-sofia.bg](mailto:nikola@fmi.uni-sofia.bg) (N. Naidenov).

In 1892 Markov [13] proved that if  $f \in \pi_n$  satisfies  $\|f\|_{[-1,1]} \leq 1$  then for every  $k = 1, \dots, n$

$$\|f^{(k)}\|_{[-1,1]} \leq T_n^{(k)}(1),$$

where the equality is attained only for the Chebyshev polynomial  $T_n(x) := \cos(n \arccos x)$  (up to a factor  $-1$ ).

It is well known that the Markov inequality and the Chebyshev polynomial play an important role in the theory of approximations with algebraic polynomials. There are a lot of results on Markov-type inequalities (see, e.g. [2,4,18,21,22], and the references therein). In connection with the research in the field of the weighted approximation by polynomials, Markov-type inequalities have been proved for various weights, norms and sets over which the norm is taken (cf. [16,23,25,24,9,20,7,17,14,12,10]). In the case of supremum norm on an infinite interval there are only two exact Markov-type inequalities (see [11,5]). They are of the form

$$\|(wp)'\| \leq C_n(w)\|wp\|, \quad \forall p \in \pi_n, \tag{1}$$

where  $w(x) = e^{-x^2}$  on  $\mathbb{R}$  or  $w(x) = e^{-x}$  on  $\mathbb{R}_+ := [0, \infty)$ . The equality in (1) is attained only for the corresponding weighted Chebyshev polynomial (up to a constant factor).

The aim of this paper is to extend the above inequalities to derivatives of arbitrary order. Note that such an extension was obtained in [15] for polynomials which have only real zeros.

Next we formulate our main results. Denote by  $U_n$  the space of all weighted polynomials of the form  $u(x) = e^{-x^2}p(x)$ , where  $p \in \pi_n$ . We shall use the notation  $u_{*,n}$  for the Chebyshev polynomial from  $U_n$ . Precisely,  $u_{*,n}$  is the unique polynomial from  $U_n$  which has norm equal to 1 and there exist  $n + 1$  points  $t_0 < \dots < t_n$  such that  $u_{*,n}(t_k) = (-1)^{n-k}$  for  $k = 0, \dots, n$ .

**Theorem 1.** *Let  $u \in U_n$ . Then for every natural number  $k$ , the inequality*

$$\|u^{(k)}\|_{\mathbb{R}} \leq \|u_{*,n}^{(k)}\|_{\mathbb{R}} \|u\|_{\mathbb{R}}$$

*holds. The equality is attained if and only if  $u(x) = cu_{*,n}(x)$ .*

Let  $V_n$  be the space of all weighted polynomials of the form  $v(x) = e^{-x}p(x)$ , where  $p \in \pi_n$ , and  $v_{*,n}$  be the Chebyshev polynomial from  $V_n$ .

**Theorem 2.** *Let  $v \in V_n$ . Then for every natural number  $k$ , the inequality*

$$\|v^{(k)}\|_{\mathbb{R}_+} \leq \|v_{*,n}^{(k)}\|_{\mathbb{R}_+} \|v\|_{\mathbb{R}_+}$$

*holds. The equality is attained if and only if  $v(x) = cv_{*,n}(x)$ .*

In the proofs of the above theorems we use some ideas of Bojanov [2], who gave a new proof of the inequality of Markov for algebraic polynomials.

## 2. Markov inequality for the weight $e^{-x^2}$ on $\mathbb{R}$

For the sake of simplicity in this section we shall write  $\|\cdot\|$  instead of  $\|\cdot\|_{\mathbb{R}}$ . To start with we note that every non-zero polynomial from  $U_n$  has at most  $n$  real zeros, counting the multiplicities and if  $u \in U_n$  then  $u' \in U_{n+1}$ . Next we list some of the results of [15], which will be needed in the sequel. Let  $\mathcal{U}_n := \{u \in U_n : u \text{ has } n \text{ simple real zeros}\}$ . It is easily seen that if  $u \in \mathcal{U}_n$  then

$u' \in \mathcal{U}_{n+1}$ . Moreover, if  $x_1 < \dots < x_n$  are the zeros of  $u$  and  $t_0 < \dots < t_n$  are the zeros of  $u'$ , then  $t_0 < x_1 < t_1 < \dots < t_{n-1} < x_n < t_n$ .

The following theorem from [15] gives the solution of a problem about interpolation at extremal points for polynomials from  $\mathcal{U}_n$  (cf. [6,19,8,1]).

**Theorem A.** *Given positive numbers  $h_0, \dots, h_n$ , there exists a unique  $u \in \mathcal{U}_n$  and a unique set of points  $t_0 < \dots < t_n$  such that*

$$\begin{aligned} u(t_k) &= (-1)^{n-k} h_k, \quad k = 0, \dots, n, \\ u'(t_k) &= 0, \quad k = 0, \dots, n. \end{aligned} \tag{2}$$

Since every  $u \in \mathcal{U}_n$  has exactly  $n + 1$  extremal points  $t_0 < \dots < t_n$ , Theorem A shows that the parameters  $h_i(u) := |u(t_i)|$ ,  $i = 0, \dots, n$ , determine  $u$  uniquely (up to multiplication by  $-1$ ). Given  $\mathbf{h} = (h_0, \dots, h_n)$  where  $h_j > 0$  for  $j = 0, \dots, n$ , we shall use the notation  $u(\mathbf{h}; \cdot)$  for the unique solution of (2). Clearly,  $u_{*,n} = u(\mathbf{1}; \cdot)$ , where  $\mathbf{1} = (1, 1, \dots, 1) \in \mathbb{R}^{n+1}$ .

In [3] Bojanov and Rahman proposed a method for derivation of estimates for functionals in the set of algebraic polynomials, having only real zeros. This method was applied in [15] to prove the following:

**Theorem B.** *Let  $u_1$  and  $u_2$  be polynomials from  $\mathcal{U}_n$ . Suppose that*

$$0 < h_i(u_1) \leq h_i(u_2) \quad \text{for } i = 0, \dots, n.$$

*Then for every natural number  $k$ , the inequalities*

$$0 < h_j(u_1^{(k)}) \leq h_j(u_2^{(k)}), \quad j = 0, \dots, n + k, \tag{3}$$

*hold. In particular,*

$$\|u_1^{(k)}\| \leq \|u_2^{(k)}\|. \tag{4}$$

*Moreover, the equality in (3) (for some  $j$ ) and (4) is attained if and only if  $h_i(u_1) = h_i(u_2)$  for all  $i = 0, \dots, n$ .*

Consequently, the absolute values of the local extrema of the  $k$ th derivative of a weighted polynomial  $u \in \mathcal{U}_n$  are strictly increasing functions of  $h_0(u), \dots, h_n(u)$ .

In the next lemma we study a Birkhoff-type interpolation problem for weighted polynomials.

**Lemma 1.** *Let  $k$  and  $m$  be natural numbers. Given points  $t_1 < \dots < t_m, \xi$  and arbitrary values  $\{a_j\}_1^{m+2}$ , there exists a unique polynomial  $g \in U_{m+1}$  for which*

$$g(t_j) = a_j, \quad j = 1, \dots, m, \quad g^{(k)}(\xi) = a_{m+1}, \quad g^{(k+1)}(\xi) = a_{m+2}. \tag{5}$$

**Proof.** Conditions (5) can be considered as a system of linear equations for the coefficients in the representation

$$g(x) = e^{-x^2} \sum_{i=0}^{m+1} b_i x^i.$$

In order to prove the existence and the uniqueness of the solution of (5), it is sufficient to prove that the corresponding homogeneous system

$$g(t_j) = 0, \quad j = 1, \dots, m, \quad g^{(k)}(\zeta) = 0, \quad g^{(k+1)}(\zeta) = 0 \tag{6}$$

has only the trivial solution. The proof goes by induction on  $k$ , for arbitrary  $m, t_1 < \dots < t_m$  and  $\zeta$ .

Let  $k = 1$ . If  $\zeta = t_j$  for some  $j \in \{1, \dots, m\}$  then  $g$  has  $m + 2$  zeros, counting the multiplicities, hence  $g \equiv 0$ . So, we may assume  $\zeta \notin \{t_1, \dots, t_m\}$ . By Rolle's theorem,  $g'(x)$  changes its sign at some points  $\zeta_i \in (t_i, t_{i+1})$  for  $i = 1, \dots, m - 1$ . But  $g(x) \rightarrow 0$  for  $x \rightarrow \pm\infty$ , hence  $g'(x)$  has also zeros  $\zeta_0 < t_0$  and  $\zeta_m > t_m$ .

If  $\zeta \notin \{\zeta_0, \dots, \zeta_m\}$ , then according to (6),  $\zeta$  is at least double zero of  $g'$ . Thus  $g' \in U_{m+2}$  has  $m + 3$  zeros counting the multiplicities. It follows that  $g$  is a constant, i.e.  $g \equiv 0$ , provided  $m \geq 1$ .

Otherwise, if  $\zeta = \zeta_j$  for some  $j \in \{0, \dots, m\}$  then  $g'$  must change its sign at  $\zeta$ . Taking in view (6), we conclude that  $g'$  has at least triple zero at  $\zeta$ , which also implies  $g \equiv 0$ .

Let  $k \geq 2$ . Assume the assertion holds for  $k - 1$ . Let  $g$  satisfy (6) for some  $t_1 < \dots < t_m$  and  $\zeta$ . Consider the polynomial  $g_1(x) := g'(x)$ . Clearly,  $g_1 \in U_{m+2}$ ,  $g_1$  vanishes at some points  $\zeta_0 < \dots < \zeta_m$  and  $g_1^{(k-1)}(\zeta) = g_1^{(k)}(\zeta) = 0$ . Then by the inductual hypothesis  $g_1 \equiv 0$ , hence  $g \equiv 0$ . The lemma is proved.  $\square$

**Lemma 2.** Let  $u \in U_n, \|u\| = 1$ . Let  $t_1 < \dots < t_m$  ( $m \leq n$ ) be the points for which  $|u(t_k)| = 1$ . If  $g \in U_n$  vanishes at  $t_1, \dots, t_m$  then

$$\|u + \varepsilon g\| = 1 + o(\varepsilon) \quad \text{as } \varepsilon \rightarrow 0.$$

**Proof.** We can choose  $\delta > 0$  so that

$$t_j \notin (t_i - \delta, t_i + \delta)$$

for  $i \neq j$  ( $i, j = 1, \dots, n$ ). Since  $u + \varepsilon g$  tends uniformly to  $u$  on  $\mathbb{R}$  as  $\varepsilon \rightarrow 0$  there exists an  $\varepsilon_0 > 0$  such that

$$|u(x) + \varepsilon g(x)| < 1 \quad \text{for } x \notin \bigcup_{i=1}^n [t_i - \delta, t_i + \delta],$$

provided  $0 < \varepsilon < \varepsilon_0$ . Hence

$$\|u + \varepsilon g\| = \max_{i=1, \dots, m} \|u + \varepsilon g\|_{[t_i - \delta, t_i + \delta]}.$$

Let  $i$  be a fixed number from  $\{1, \dots, m\}$ . Without loss of generality we may assume that  $u(t_i) = 1$ . We define  $x_i(\varepsilon) \in (t_i - \delta, t_i + \delta)$  as the solution of  $u(x) + \varepsilon g(x) = 1$ , farthest from  $t_i$ . (It is possible  $x_i(\varepsilon) = t_i$ .)

Let  $\Delta_i(\varepsilon) := \{x : |x - t_i| \leq |x_i(\varepsilon) - t_i|\}$ . Clearly

$$\|u + \varepsilon g\|_{[t_i - \delta, t_i + \delta]} = \|u + \varepsilon g\|_{\Delta_i(\varepsilon)}.$$

Let  $u'(t_i) = \dots = u^{(2l-1)}(t_i) = 0, u^{(2l)}(t_i) < 0$ . (Recall that  $t_i$  is a local maximum of  $u$ .) We can assume that  $u^{(2l)}(x) \leq c < 0$  for  $x \in [t_i - \delta, t_i + \delta]$ , provided  $\delta$  is sufficiently small. We have

$$u(t_i + x_i(\varepsilon) - t_i) + \varepsilon g(t_i + x_i(\varepsilon) - t_i) = 1$$

and by Taylor’s formula we get

$$1 + \frac{u^{(2l)}(\xi_i^1)}{(2l)!}(x_i(\varepsilon) - t_i)^{2l} + \varepsilon g'(\xi_i^2)(x_i(\varepsilon) - t_i) = 1,$$

where  $\xi_i^1, \xi_i^2 \in \Delta_i(\varepsilon)$ . Hence

$$(x_i(\varepsilon) - t_i)^{2l-1} = -\frac{(2l)!g'(\xi_i^2)\varepsilon}{u^{(2l)}(\xi_i^1)} = O(\varepsilon),$$

which implies  $x_i(\varepsilon) - t_i = O(\varepsilon^{\frac{1}{2l-1}})$ . For each  $x \in \Delta_i(\varepsilon)$  we have

$$u(x) + \varepsilon g(x) = 1 + \frac{u^{(2l)}(\eta_i^1)}{(2l)!}(x - t_i)^{2l} + \varepsilon g'(\eta_i^2)(x - t_i) = 1 + O(\varepsilon^{\frac{2l}{2l-1}}),$$

which finishes the proof of Lemma 2.  $\square$

In the next lemma we prove a property of the polynomials from  $\mathcal{U}_n$ , which is well known for algebraic polynomials.

**Lemma 3.** *Each zero  $\eta$  of the derivative of a weighted polynomial  $u(x) = ce^{-x^2}(x - x_1) \cdots (x - x_n)$  ( $c \neq 0$ ) is a strictly increasing function of  $x_k$  in the domain  $x_1 < \cdots < x_n$ .*

**Proof.** Denote for brevity  $\omega(x) = (x - x_1) \cdots (x - x_n)$ . Since

$$\frac{u'(x)}{u(x)} = -2x + \frac{\omega'(x)}{\omega(x)}$$

and  $u'(\eta) = 0$ , we get

$$-2\eta + \sum_{i=1}^n \frac{1}{\eta - x_i} = 0.$$

Differentiating the last identity with respect to  $x_k$  we obtain

$$\left(2 + \sum_{i=1}^n \frac{1}{(\eta - x_i)^2}\right) \frac{\partial \eta}{\partial x_k} = \frac{1}{(\eta - x_k)^2},$$

which implies  $\frac{\partial \eta}{\partial x_k} > 0$ . Lemma 3 is proved.  $\square$

An immediate consequence of Lemma 3 is the following:

**Corollary 3.** *Let  $u_1$  and  $u_2$  be two polynomials from  $\mathcal{U}_n$  having zeros  $x_1 < \cdots < x_n$  and  $y_1 < \cdots < y_n$ , respectively. Suppose that*

$$x_i \leq y_i, \quad i = 1, \dots, n,$$

*with at least one strict inequality. Then the zeros  $t_1 < \cdots < t_{n+1}$  of  $u'_1(x)$  and the zeros  $\tau_1 < \cdots < \tau_{n+1}$  of  $u'_2(x)$  satisfy*

$$t_i < \tau_i, \quad i = 1, \dots, n + 1.$$

Our next result is a weighted analogue of the famous Markov’s lemma concerning the zeros of the algebraic polynomials.

**Lemma 4.** *Assume that the zeros  $x_1 < \dots < x_n$  of  $u_1 \in \mathcal{U}_n$  and  $y_1 < \dots < y_{n-1}$  of  $u_2 \in \mathcal{U}_{n-1}$  satisfy the interlacing conditions*

$$x_1 \leq y_1 \leq x_2 \leq \dots \leq x_{n-1} \leq y_{n-1} \leq x_n.$$

*Then the zeros  $t_1 < \dots < t_{n+1}$  of  $u'_1$  and the zeros  $\tau_1 < \dots < \tau_n$  of  $u'_2$  interlace strictly, that is,*

$$t_1 < \tau_1 < t_2 < \dots < t_n < \tau_n < t_{n+1}.$$

**Proof.** We will prove only the inequalities

$$t_i < \tau_i \quad \text{for } i = 1, \dots, n. \tag{7}$$

(The remaining ones can be established in a similar way.) Set

$$y_k(\varepsilon) := \begin{cases} y_k & \text{for } k = 1, \dots, n - 1, \\ \frac{1}{\varepsilon} & \text{for } k = n. \end{cases}$$

The inequalities

$$x_1 \leq y_1(\varepsilon) \leq x_2 \leq \dots \leq y_{n-1}(\varepsilon) \leq x_n < y_n(\varepsilon) \tag{8}$$

hold true, provided  $\varepsilon$  is a sufficiently small positive number.

Let us define  $u_\varepsilon(x) := u_2(x)(1 - \varepsilon x)$ . Clearly,  $y_k(\varepsilon)$ ,  $k = 1, \dots, n$ , are the zeros of  $u_\varepsilon$  and let  $\tau_1(\varepsilon) < \dots < \tau_{n+1}(\varepsilon)$  be the zeros of  $u'_\varepsilon$ . Corollary 3 and (8) imply

$$t_i < \tau_i(\varepsilon) \quad \text{for } i = 1, \dots, n + 1. \tag{9}$$

Note that  $\tau_i(\varepsilon) \rightarrow \tau_i$ ,  $i = 1, \dots, n$ , because  $u_\varepsilon^{(k)}$  tends uniformly to  $u_2^{(k)}$  on  $\mathbb{R}$  as  $\varepsilon \rightarrow 0$ . According to Lemma 3, each of  $\tau_i(\varepsilon)$  increases strictly when  $\varepsilon$  decreases. Letting  $\varepsilon \downarrow 0$  in (9) we obtain (7). Lemma 4 is proved.  $\square$

In the next lemma we compare the norms of the derivatives of the weighted Chebyshev polynomials for different  $n$ .

**Lemma 5.** *For every natural number  $k$  the inequality*

$$\|u_{*,n-1}^{(k)}\| < \|u_{*,n}^{(k)}\| \tag{10}$$

*holds true.*

**Proof.** Let  $u_{*,n-1}(x) = e^{-x^2}(\alpha_{n-1}x^{n-1} + \dots)$ , where  $\alpha_{n-1} > 0$ . For every  $\varepsilon > 0$  we consider the polynomial  $u_\varepsilon(x) = u_{*,n-1}(x) - \varepsilon x^n e^{-x^2}$ . It is easily seen that for each  $j \geq 0$  we have

$$\|u_\varepsilon^{(j)} - u_{*,n-1}^{(j)}\| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \tag{11}$$

Let us fix a point  $b$  greater than all zeros of  $u_{*,n-1}$ . Clearly,  $u_{*,n-1}(b) > 0$ . Hence, for sufficiently small  $\varepsilon$ ,  $u_\varepsilon$  has  $n - 1$  simple zeros in  $(-\infty, b)$  (close to the zeros of  $u_{*,n-1}$ ) and  $u_\varepsilon(b) > 0$ . But

the leading coefficient of  $u_\varepsilon(x)$  is negative, hence  $u_\varepsilon$  must have another real zero  $x(\varepsilon) > b$ . Since  $b$  can be arbitrarily large, it follows that  $x(\varepsilon) \rightarrow \infty$  as  $\varepsilon \rightarrow 0$ .

Let us denote the points of the local extrema of the oscillating polynomial  $u_\varepsilon$  by  $t_0(\varepsilon) < \dots < t_n(\varepsilon)$  and those of  $u_{*,n-1}$  by  $t_0 < \dots < t_{n-1}$ . We have  $t_n(\varepsilon) \rightarrow \infty$  while (from (11))  $t_i(\varepsilon) \rightarrow t_i$  as  $\varepsilon \rightarrow 0$  for  $i = 0, \dots, n - 1$ . Also  $u_\varepsilon(t_n(\varepsilon)) \rightarrow -0$  and  $u_\varepsilon(t_i(\varepsilon)) \rightarrow (-1)^{n-1-i}$  for  $i = 0, \dots, n - 1$ .

According to Theorem A,  $u_\varepsilon(x) = -u(\mathbf{h}_0(\varepsilon); x)$ , where  $\mathbf{h}_0(\varepsilon) := (h_0(u_\varepsilon), \dots, h_n(u_\varepsilon))$ . If  $\mathbf{h}_1(\varepsilon) := (h_0(u_\varepsilon), \dots, h_{n-1}(u_\varepsilon), 1/2)$  then by Theorem B  $\|u_\varepsilon^{(k)}\| < \|u^{(k)}(\mathbf{h}_1(\varepsilon); \cdot)\|$ , provided  $\varepsilon$  is sufficiently small. Letting  $\varepsilon \rightarrow 0$  we obtain

$$\|u_{*,n-1}^{(k)}\| \leq \|u^{(k)}(\mathbf{h}_1; \cdot)\|, \tag{12}$$

where  $\mathbf{h}_1 = (1, \dots, 1, 1/2) \in \mathbb{R}^{n+1}$ . Using again the strict monotonicity we get

$$\|u^{(k)}(\mathbf{h}_1; \cdot)\| < \|u_{*,n}^{(k)}\|. \tag{13}$$

Inequality (10) is a direct consequence from (12) and (13). Lemma 5 is proved.  $\square$

**Proof of Theorem 1.** An equivalent setting is to prove that  $u_{*,n}$  is the unique solution of the extremal problem

$$\|u^{(k)}\| \rightarrow \sup \text{ over all } u \in U_n, \|u\| \leq 1. \tag{14}$$

Let  $u$  be a fixed extremal polynomial to problem (14). Note that  $\|u\| = 1$ . We claim that  $|u(x)|$  attains its maximal value at least at  $n$  points. Indeed, assume that  $t_1 < \dots < t_m$  ( $m \leq n - 1$ ) are all points such that  $|u(t_k)| = 1$ . Let  $M_k := \|u^{(k)}\| = |u^{(k)}(\xi)|$ . According to Lemma 1 there exists  $g \in U_{m+1} \subseteq U_n$  satisfying the conditions

$$g(t_j) = 0, \quad j = 1, \dots, m, \quad g^{(k)}(\xi) = \text{sign } u^{(k)}(\xi). \tag{15}$$

(For  $g^{(k+1)}(\xi)$  we can take any value.)

Consider the polynomial  $u_\varepsilon(x) := (u(x) + \varepsilon g(x))/\|u + \varepsilon g\|$ . Clearly,  $u_\varepsilon \in U_n$  and  $\|u_\varepsilon\| = 1$ . It follows from Lemma 2 and (15) that

$$|u_\varepsilon^{(k)}(\xi)| = \frac{|u^{(k)}(\xi) + \varepsilon g^{(k)}(\xi)|}{1 + o(\varepsilon)} = \frac{M_k + \varepsilon}{1 + o(\varepsilon)} > M_k,$$

provided  $\varepsilon$  is a sufficiently small positive number. The last inequality contradicts with the extremality of  $u$ . The claim is proved.

Note that the equation

$$|u(t)| = 1 \tag{16}$$

cannot have more than  $n + 1$  solutions. Otherwise,  $u'(x)$  would have  $n + 2$  zeros, so  $u'(x) \equiv 0$ , a contradiction.

Furthermore, if there exist exactly  $n + 1$  points at which (16) holds, then it is easily seen that  $u \equiv \pm u_{*,n}$ , so Theorem 1 will be proved.

It remains to exclude the case when (16) has exactly  $n$  solutions. Assume the contrary and let  $t_1 < \dots < t_n$  be all the points at which  $|u(x)|$  attains its maximal value.

Our next goal is to show that they are alternation points for  $u$ , i.e.  $u(t_k) = \sigma(-1)^k$  for  $k = 1, \dots, n$ , where  $\sigma \in \{-1, 1\}$ . Assume the contrary. Then there exists an index  $i$  for which

$u(t_i)u(t_{i+1}) > 0$ , hence  $u'$  has a zero  $\gamma \in (t_i, t_{i+1})$ . Consequently,  $\{t_k\}_1^n$  and  $\gamma$  are all the zeros of  $u' \in U_{n+1}$ . If  $\omega(x) := e^{-x^2}(x - t_1) \cdots (x - t_n)$ , then the zeros of  $u'$  and  $\omega$  interlace, hence by Lemma 4, the zeros of  $u^{(k+1)}$  and  $\omega^{(k)}$  interlace strictly. As  $u^{(k+1)}(\xi) = 0$ , we conclude that  $\omega^{(k)}(\xi) \neq 0$ . Then, for sufficiently small  $\varepsilon > 0$ , one of the polynomials  $(u \pm \varepsilon\omega)/\|u \pm \varepsilon\omega\|$  will have larger norm of the  $k$ th derivative than  $u$ , which is a contradiction.

So, the extremal polynomial  $u$  has  $n$  alternation points, hence at least  $n - 1$  simple zeros. If  $u \in U_{n-1}$  then  $u$  has to coincide with  $\pm u_{*,n-1}$ , but this is impossible in view of Lemma 5. It follows that  $u$  is a weighted polynomial of exact degree  $n$ , hence  $u$  must have  $n$  simple real zeros. Taking into account Theorem B, we conclude that  $u = \pm u_{*,n}$ , which is a contradiction. Theorem 1 is proved.  $\square$

### 3. Markov inequality for the weight $e^{-x}$ on $\mathbb{R}_+$

In this section we abbreviate the notation  $\|\cdot\|_{\mathbb{R}_+}$  to  $\|\cdot\|$ . The approach is similar to that in Section 2, but the analysis is somewhat simpler, due to the translation invariance property of  $V_n$ , that is,  $v(x + c) \in V_n$  for every  $v \in V_n$  and  $c \in \mathbb{R}$ .

**Lemma 6.** *Let  $k$  and  $m$  be natural numbers. Given points  $t_1 < \cdots < t_m$  in  $[0, \infty)$  and values  $\{a_j\}_0^m$ , there exists a unique polynomial  $g \in V_m$  for which*

$$g(t_j) = a_j, \quad j = 1, \dots, m, \quad g^{(k)}(0) = a_0.$$

**Proof.** As in Lemma 1, we will show that the homogeneous system of equations

$$v(t_j) = 0, \quad j = 1, \dots, m, \quad v^{(k)}(0) = 0 \tag{17}$$

admits only the trivial solution  $v \equiv 0$  in  $V_m$ .

Let  $v$  be a solution of (17). By Rolle’s theorem,  $v'(x)$  has at least one zero  $\xi_i \in (t_i, t_{i+1})$  for  $i = 1, \dots, m$ , where  $t_{m+1} := \infty$ . Repeating this argument, we conclude that  $v^{(k)}$  vanishes at some points  $\xi_1^{(k)} < \cdots < \xi_m^{(k)}$  in  $(0, \infty)$ . Because of (17),  $v^{(k)} \in V_m$  has  $m + 1$  zeros in  $[0, \infty)$ , which implies  $v^{(k)} \equiv 0$ .

Now, let  $v(x) = e^{-x}p(x)$ , where  $p(x)$  is an algebraic polynomial of degree  $\leq m$ . It is easily seen that  $v^{(k)}(x) = e^{-x}q(x)$ , where  $q(x) = \sum_{s=0}^k (-1)^{k-s} \binom{k}{s} p^{(s)}(x)$ . But  $q \equiv 0$ , hence the degree of  $p$  is less than  $m$ . Taking in view (17), we conclude that  $p \equiv 0$ . The lemma is proved.  $\square$

**Lemma 7.** *Let  $v \in V_n, \|v\| = 1$ . Let  $m \leq n$  and  $t_1 < \cdots < t_m$  be the points for which  $|v(t_k)| = 1$ . If  $g \in V_n$  vanishes at  $t_1, \dots, t_m$  then*

$$\|v + \varepsilon g\| = 1 + o(\varepsilon) \quad \text{if } \varepsilon \rightarrow 0.$$

**Proof.** As in Lemma 2, it is sufficient to consider  $v + \varepsilon g$  on small neighbourhoods of the points  $t_i, i = 1, \dots, m$ . If  $t_i > 0$  then the estimation of the norm of  $v + \varepsilon g$  around  $t_i$  is completely analogous to that in Lemma 2. It remains to estimate  $v + \varepsilon g$  around  $t_1$  if  $t_1 = 0$ . Let  $\delta < t_2$  be a sufficiently small, fixed positive number. Our goal is to prove that  $\|v + \varepsilon g\|_{[0, \delta]} = 1 + o(\varepsilon)$  as  $\varepsilon \rightarrow 0$ . Without loss of generality we may assume  $v(0) = 1$  and, as a consequence,  $v'(0) \leq 0$ . If  $v'(0) < 0$  then it is easy to see that  $\|v + \varepsilon g\|_{[0, \delta]} = 1$ .



Suppose now  $v'(0) = 0$ . Set  $x(\varepsilon) := \sup\{x \in [0, \delta] : v(x) + \varepsilon g(x) = 1\}$ . It follows that  $\|v + \varepsilon g\|_{[0, \delta]} = \|v + \varepsilon g\|_{[0, x(\varepsilon)]}$ . Furthermore, arguing as in Lemma 2, we get  $x(\varepsilon) = O(\varepsilon^{\frac{1}{s-1}})$ , provided  $v'(0) = \dots = v^{(s-1)}(0) = 0$ ,  $v^{(s)}(0) \neq 0$  for some  $s \geq 2$ . Consequently, if  $x \in [0, x(\varepsilon)]$  then  $v(x) + \varepsilon g(x) = 1 + O(\varepsilon^{\frac{s}{s-1}})$ , which finishes the proof of Lemma 7.  $\square$

**Proof of Theorem 2.** As in Theorem 1, it is sufficient to prove that  $v_{*,n}$  is the unique solution of the extremal problem

$$\|v^{(k)}\| \rightarrow \sup \quad \text{over all } v \in V_n, \quad \|v\| \leq 1. \quad (18)$$

Let  $v$  be a fixed extremal polynomial to problem (18). Clearly,  $\|v\| = 1$  and the equation

$$|v(t)| = 1 \quad (19)$$

cannot have more than  $n + 1$  solutions on  $[0, \infty)$ . We claim that  $|v(x)|$  attains its maximal value at exactly  $n + 1$  points. On the contrary, we assume that Eq. (19) has exactly  $m \leq n$  solutions  $t_1 < \dots < t_m$  in  $[0, \infty)$ . There exists a point  $\xi \in [0, \infty)$  such that  $M_k := \|v^{(k)}\| = |v^{(k)}(\xi)|$ . Without loss of generality we suppose that  $\xi = 0$ . (Otherwise, we can consider  $v_1(x) := v(x + \xi) \in V_n$ . We have  $\|v_1\| \leq \|v\| = 1$  and  $|v_1^{(k)}(0)| = M_k$ , hence  $v_1$  is also extremal in (18), which implies  $\|v_1\| = 1$ . In addition, the equation  $|v_1(x)| = 1$  also has less than  $n + 1$  solutions in  $[0, \infty)$ .) Lemma 6 ensures the existence of a  $g \in V_m \subseteq V_n$  such that

$$g(t_j) = 0, \quad j = 1, \dots, m, \quad g^{(k)}(0) = \text{sign } v^{(k)}(0). \quad (20)$$

If  $v_\varepsilon(x) := (v(x) + \varepsilon g(x))/\|v + \varepsilon g\|$  then  $v_\varepsilon \in V_n$  and  $\|v_\varepsilon\| = 1$ . Using Lemma 7 and (20) (as in the proof of Theorem 1) we conclude that  $|v_\varepsilon^{(k)}(0)| > M_k$ , provided  $\varepsilon > 0$  is sufficiently small. This is a contradiction, which proves the claim.

Let us denote the points at which  $|v(x)|$  attains its maximal value by  $t_0 < \dots < t_n$ . Next we will prove that they are alternation points for  $v$ , which implies  $v = \pm v_{*,n}$ . Assume the contrary, i.e. there exists  $i \in \{0, \dots, n-1\}$  such that  $v(t_i)v(t_{i+1}) > 0$ . Then  $v'$  has a zero in  $(t_i, t_{i+1})$ . Since  $v'(t_k) = 0$  for  $k = 1, \dots, n$  and  $v' \in V_n$ , we conclude that  $v' \equiv 0$ , which is a contradiction. Theorem 2 is proved.  $\square$

**Remark.** In fact  $\|v_{*,n}^{(k)}\| = |v_{*,n}^{(k)}(0)|$ . Otherwise, a proper translation of  $v_{*,n}$  will produce a different extremal polynomial in (18).

## Acknowledgment

The authors thank Professor Borislav Bojanov for his valuable remarks regarding this paper.

## References

- [1] B. Bojanov, A generalization of Chebyshev polynomials, *J. Approx. Theory* 26 (1979) 293–300.
- [2] B. Bojanov, Markov-type inequalities for polynomials and splines, in: C.K. Chui, L.L. Schumaker, J. Stöckler (Eds.), *Approximation Theory X: Abstract and Classical Analysis*, Vanderbilt University Press, Nashville, TN, 2002, pp. 31–90.
- [3] B.D. Bojanov, Q.I. Rahman, On certain extremal problems for polynomials, *J. Math. Anal. Appl.* 189 (1995) 781–800.
- [4] P. Borwein, T. Erdélyi, *Polynomials and polynomial inequalities*, Graduate Texts in Mathematics, vol. 161, Springer, New York, Berlin, Heidelberg, 1995.

- [5] H. Carley, X. Li, R.N. Mohapatra, A sharp inequality of Markov type for polynomials associated with Laguerre weight, *J. Approx. Theory* 113 (2001) 221–228.
- [6] C. Davis, Problem 4714, *Amer. Math. Monthly* 63 (1956) 729;  
C. Davis, Solution, *Amer. Math. Monthly* 64 (1957) 679–680.
- [7] P. Dörfler, New inequalities of Markov type, *SIAM J. Math. Anal.* 18 (1987) 490–494.
- [8] C.H. Fitzgerald, L.L. Schumaker, A differential equation approach to interpolation at extremal points, *J. Analyse Math.* 22 (1969) 117–134.
- [9] G. Freud, On Markov–Bernstein type inequalities and their applications, *J. Approx. Theory* 19 (1977) 22–37.
- [10] A.L. Levin, D.S. Lubinsky,  $L_p$  Markov–Bernstein inequalities for Freud weights, *J. Approx. Theory* 77 (1994) 229–248.
- [11] X. Li, R.N. Mohapatra, R.S. Rodriguez, On Markov’s inequality on  $R$  for the Hermite weight, *J. Approx. Theory* 75 (1993) 267–273.
- [12] D.S. Lubinsky, E.B. Saff, Markov–Bernstein and Nikolskii inequalities, and Christoffel functions for exponential weights on  $[-1, 1]$ , *SIAM J. Math. Anal.* 24 (1993) 528–556.
- [13] V.A. Markov, On the Functions of Least Deviation from Zero in a Given Interval, St. Petersburg, 1892 (in Russian);  
W. Markoff, Über Polynome die in einem gegebenen Intervalle möglichst wenig von Null abweichen, *Math. Ann.* 77 (1916) 213–258 (German translation with condensation).
- [14] H.N. Mhaskar, General Markov–Bernstein and Nikolskii type inequalities, *Approx. Theory Appl.* 6 (1990) 107–117.
- [15] L. Milev, Weighted polynomial inequalities on infinite intervals, *East J. Approx.* 5 (1999) 449–465.
- [16] W. Milne, On the maximum absolute value of the derivative of  $e^{-x^2} P_n(x)$ , *Trans. Amer. Math. Soc.* 33 (1931) 143–146.
- [17] G.M. Milovanović, Various extremal problems of Markov’s type for algebraic polynomials, *Facta Univ. Ser. Math. Inform.* 2 (1987) 7–28.
- [18] G.M. Milovanović, D.S. Mitrinović, Th.M. Rassias, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific, Singapore, 1994.
- [19] J. Mycielski, S. Paszkowski, A generalization of Chebyshev polynomials, *Bull. Acad. Polonaise Sci. Série Math. Astr. et Phys.* 8 (1960) 433–438.
- [20] P. Nevai, V. Totik, Weighted polynomial inequalities, *Constr. Approx.* 2 (1986) 113–127.
- [21] Q.I. Rahman, G. Schmeisser, *Analytic Theory of Polynomials*, Clarendon Press, Oxford, 2002.
- [22] T.J. Rivlin, *Chebyshev Polynomials*, second ed., Wiley, New York, 1990.
- [23] E. Schmidt, Über die nebst ihren Ableitungen orthogonalen Polynomensysteme und das zugehörige Extremum, *Math. Ann.* 119 (1944) 165–204.
- [24] G. Szegő, On some problems of approximations, *Magyar Tud. Akad. Mat. Kutató Int. Közl.* 9 (1964) 3–9.
- [25] P. Turan, Remark on a theorem of Erhard Schmidt, *Mathematica* 2 (1960) 373–378.