

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713454007>

Non-taxane compounds from the bark of *Taxus yunnanensis*

Sheng-Hong Li^a; Hong-Jie Zhang^b; Ping Yao^b; Xue-Mei Niu^a; Wei Xiang^a; Han-Dong Sun^a

^a Laboratory of Phytochemistry, Kunming Institute of Botany, The Chinese Academy of Sciences, Kunming, Yunnan, China ^b Program for Collaborative Research in the Pharmaceutical sciences, College of Pharmacy, University of Illinois, Chicago, IL, USA

Online publication date: 09 September 2010

To cite this Article Li, Sheng-Hong , Zhang, Hong-Jie , Yao, Ping , Niu, Xue-Mei , Xiang, Wei and Sun, Han-Dong(2002) 'Non-taxane compounds from the bark of *Taxus yunnanensis*', *Journal of Asian Natural Products Research*, 4: 2, 147 — 154

To link to this Article: DOI: 10.1080/10286020290027443

URL: <http://dx.doi.org/10.1080/10286020290027443>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

NON-TAXANE COMPOUNDS FROM THE BARK OF *TAXUS YUNNANENSIS*

SHENG-HONG LI^a, HONG-JIE ZHANG^{b,*}, PING YAO^b, XUE-MEI NIU^a,
WEI XIANG^a and HAN-DONG SUN^{a,†}

^aLaboratory of Phytochemistry, Kunming Institute of Botany, the Chinese Academy of Sciences,
Kunming 650204, Yunnan, China;

^bProgram for Collaborative Research in the Pharmaceutical sciences, College of Pharmacy,
University of Illinois at Chicago, 833 Wood Street, Chicago, IL 60612, USA

(Received 18 September 2001; Revised 29 September 2001; In final form 23 October 2001)

From the bark of *Taxus yunnanensis*, 15 non-taxane compounds were isolated. Through spectroscopic methods such as 1D and 2D NMR and MS experiments, one of them was determined as a new abietane-type diterpenoid named taxayunnin (**1**). The other 14 known compounds were identified as taxamairin C (**2**), taxamairin A (**3**), 3 β -hydroxy-sandaracopimaric acid (**4**), (+)-3-hydroxy-isodrimenin (**5**), rubrosterone (**6**), ponasterone A (**7**), ecdysterone (**8**), 20-hydroxy-echysone-20,22-monoacetonide (**9**), 7-oxositosterol (**10**), stigmast-4-en-6 β -ol-3-one (**11**), 5 α ,6 β -dihydroxy-daucosterol (**12**), β -sitosterol (**13**), daucosterol (**14**), 1-*O*- β -D-glucopyranosyl-(2*S*, 3*R*, 4*E*, 8*Z*)-2-*N*-(2'-hydroxypalmitoyl)-octadeca-sphinga-4,8-dienine (**15**), respectively. Compounds **4–6**, **9–12** and **15** were isolated from *Taxus* plants for the first time.

Keywords: Taxaceae; *Taxus yunnanensis*; Bark; Non-taxane compounds

INTRODUCTION

Besides the major and well-known taxane diterpenoids [1], *Taxus* plants also contain abundant non-taxane compounds such as non-taxane diterpenoids, steroids, lignans, flavonoids, sugar derivatives and so on [2,3]. Our previous work on the chemical and biological constituents of the root and bark of *Taxus yunnanensis*, an evergreen tree endemic to Yunnan of China, has led to the isolation of many new and known taxane diterpenoids including paclitaxel [4–14]. In this paper, we report the isolation and structure characterization of a new abietane diterpenoid, taxayunnin (**1**), and fourteen known non-taxane compounds (**2–15**) from the bark of this plant.

RESULTS AND DISCUSSION

Taxayunnin (**1**) was obtained as pale yellow needles with mp 215°C. It showed a molecular ion peak at m/z 346 [M^+] in its HREI mass spectrum, which in combination with ¹H and ¹³C

*Corresponding author. Tel.: +1-312-996-7868. Fax: +1-312-413-5894. E-mail: zhanghj@uic.edu

†Tel.: +86-871-5223251. Fax: +86-871-5216343. E-mail: hdsun@mail.kib.ac.cn

NMR (including DEPT) spectra suggested a molecular formula $C_{21}H_{30}O_4$. Its 1H NMR spectrum (see Table I) displayed three methyls linked to quaternary carbon atoms, an isopropyl group, and a deshielded aromatic proton at δ 7.62. These evidences indicated **1** was an abietane diterpenoid. Comparison of 1H and ^{13}C NMR spectra (Tables I and II) of **1** with those of **2** revealed that **1** was similar to **2** except for A-ring moiety. The difference could be rationalized to an oxygen bridge, which occurred between C-3 and C-20 in **2**. Further comparison of the 1H NMR data of **1** with those of 3β -hydroxy-demethyl cryptojaponol [15], an abietane diterpenoid isolated from *Salvia pubescens*, suggested that **1** was a mono-methyl ether of 3β -hydroxy-demethyl cryptojaponol. The methoxy group attached to C-12 was supported by the simultaneous HMBC correlations of C-12 with both of 12-OCH₃ and H-15 (Fig. 1). The β -orientation of the hydroxy group at C-3 was established by the NOE correlations of H-3 with H-5 and CH₃-18. Due to the γ -gauche deshielding effect of 3β -OH, the chemical shift of C-19 moved upfield obviously (about 3 ppm comparing with that of **2**). Thus the structure of compound **1** was determined as 3,11-dihydroxy-12-methoxy-8,11,13-abietatrien-7-one, and was given the trivial name taxayunnin.

Along with the new diterpenoid, fourteen known non-taxane compounds were also obtained. On the basis of the comparison of their spectral data with those reported in the





TABLE I ^1H NMR data of compounds **1** and **2**

Proton	1 *	2 †
H-1 α	1.58 (1H, m)	1.45 (1H, ddd, 3.0, 12.4, 18.0)
H-1 β	3.30 (1H, dt, 3.6, 13.9)	3.43 (1H, ddd, 5.4, 12.6, 18.0)
H-2 α	1.81 (2H, m)	1.91 (1H, ddd, 3.7, 12.7, 16.5)
H-2 β	–	2.24 (1H, ddd, 5.4, 12.3, 17.9)
H-3	3.35 (1H, dd, 5.2, 6.9)	–
H-5	1.83 (1H, dd, 3.1, 13.9)	2.11 (1H, dd, 2.9, 14.9)
H-6 α	2.60 (1H, dd, 14.2, 16.9)	2.53 (1H, dd, 4.0, 14.8)
H-6 β	2.68 (1H, dd, 3.1, 16.9)	2.64 (1H, t, 14.9)
H-14	7.62 (1H, s)	7.50 (1H, s)
H-15	3.20 (1H, spt, 6.9)	3.18 (1H, spt, 6.8)
CH ₃ -16	1.24 (3H, d, 6.9)	1.20 (3H, d, 6.8)
CH ₃ -17	1.26 (3H, d, 6.9)	1.22 (3H, d, 6.8)
CH ₃ -18	1.07 (3H, s)	1.05 (3H, s)
CH ₃ -19	0.95 (3H, s)	1.10 (3H, s)
CH ₃ -20	1.39 (3H, s)	4.75 (1H, dd, 1.4, 8.9, H-20a)
/H ₂ -20		4.15 (1H, dd, 2.4, 9.0, H-20b)
11-OH	6.12 (1H, s)	6.29 (1H, s)
12-OCH ₃	3.81 (3H, s)	3.79 (3H, s)

* 500 MHz, CDCl₃, *J* in Hz, δ in ppm.† 400 MHz, CDCl₃, *J* in Hz, δ in ppm.TABLE II ^{13}C NMR data of compounds **1–4**

Carbon	1 *	2 †	3 †	4 †
C-1	34.7 t	29.9 t	188.1 s	36.9 t
C-2	27.9 t	35.9 t	120.9 s	27.0 t
C-3	78.0 d	98.1 s	146.1 s	75.6 d
C-4	39.2 s	40.8 s	133.7 d	53.4 s
C-5	49.7 d	47.2 d	130.2 s	49.7 d
C-6	35.3 t	35.9 t	147.8 d	24.5 t
C-7	198.5 s	198.3 s	123.8 d	35.3 t
C-8	128.6 s	130.1 s	200.8 s	136.0 s
C-9	137.3s	127.2 s	50.5 s	50.4 d
C-10	39.9 s	37.0 s	151.4 s	37.7 s
C-11	146.5 s	147.6 s	131.4 d	18.8 t
C-12	149.3 s	149.1 s	26.8 q	34.5 t
C-13	139.4 s	140.5 s	27.5 q	37.5 s
C-14	117.5 d	117.2 d	146.6 s	129.6 d
C-15	26.8 d	26.7 d	148.3 s	148.7 d
C-16	23.4 q	23.4 q	136.6 s	110.3 t
C-17	23.5 q	23.4 q	119.4 d	26.1 q
C-18	28.0 q	26.7 q	27.5 d	182.5 s
C-19	15.3 q	18.2 q	23.4 q	11.0 q
C-20	18.0 q	65.3 t	23.4 q	12.9 q
OCH ₃	61.8 q	61.9 q	62.0 q	

* 125 MHz, CDCl₃, δ in ppm.† 100 MHz, CDCl₃, δ in ppm.FIGURE 1 Selected HMBC correlations of compound **1** (from H to C).

literatures, these compounds were identified as an abietane-type diterpenoid, taxamairin C (**2**) [16]; a deformed abietane-type diterpenoid, taxamairin A (**3**) [16]; an isopimarane-type diterpenoid, 3 β -hydroxysandaracopimaric acid (**4**) [17]; a drimane-type sesquiterpenoid, (+)-3-hydroxy-isodrimenin (**5**) [18]; nine steroids, namely rubrosterone (**6**) [19], ponasterone A (**7**) [20], ecdysterone (**8**) [20], 20-hydroxy-echysone-20,22-monoacetone (**9**) [21], 7-oxositosterol (**10**) [22], stigmast-4-en-6 β -ol-3-one (**11**) [22], 5 α ,6 β -dihydroxy-daucosterol (**12**) [23], β -sitosterol (**13**) [22], daucosterol (**14**) [24]; and a cerebroside, 1-*O*- β -D-glucopyranosyl-(2*S*, 3*R*, 4*E*, 8*Z*)-2-*N*-(2'-hydroxypalmitoyl)-octadeca-sphinga-4,8-dienine (**15**) [25] respectively. The structures of compounds **4–7**, **9**, **12** and **15** were further confirmed by 2D NMR experiments. Full assignments of compound **7** were also achieved. Among these known compounds, **4–6**, **9–12** and **15** were isolated from *Taxus* plants for the first time.

EXPERIMENTAL SECTION

General Experimental Procedures

Melting points were obtained on a XRC-1 micro melting point apparatus and are uncorrected. 1D and 2D NMR experiments were performed either on a Bruker AM-400 or DRX-500 spectrometer. Unless otherwise indicated, chemical shifts (δ) were expressed in ppm with reference to the solvent signals. FABMS and HRFABMS were taken on a VG Auto Spec-3000 or on a Finnigan MAT 90 instrument. IR spectra were recorded on a Bio-Rad FTS-135 spectrometer with KBr pellets. UV spectral data were obtained on a UV 210A spectrometer. Optical rotations were carried out on a HORIBA SEPA-300 High Sensitive Polarimeter or Perkin-Elmer model 241 Polarimeter. Column chromatography was performed either on Si gel (200–300 mesh, Qingdao Marine Chemical Incorporation, China), Si gel H (10–40 μ , Qingdao Marine Chemical Incorporation, China), Lichroprep Rp₁₈ gel (40–63 μ m, Merck, Darmstadt, Germany), or on MCI gel (70–150 μ , Mitsubishi Chemical Corporation, Tokyo, Japan). Fractions were monitored by TLC and spots were visualized by heating Si gel plates sprayed with 10% H₂SO₄ in EtOH.

Plant Material

The barks of *T. yunnanensis* Cheng et L. K. Fu (Taxaceae) were collected in Lijiang Prefecture of Yunnan Province of People's Republic of China in January, 1997. The plant material was identified by Prof Xi-Wen Li at Kunming Institute of Botany. A voucher specimen (No. YAF-97-18) has been deposited at the Yunnan Academy of Forestry, Kunming, Yunnan, People's Republic of China.

Extraction and Isolation

Dried bark (50 kg) was milled and extracted by maceration in EtOH for one week, the extract was concentrated *in vacuo* to a syrup, diluted with H₂O and partitioned with CHCl₃. The CHCl₃ layer was evaporated *in vacuo* to afford 500 g of a residue, which was absorbed on 800 g of Si gel and chromatographed on a pre-packed (2 kg) Si gel column. Gradient elution was accomplished with CHCl₃–Me₂CO (1:0, 9:1, 8:2, 7:3, 0:1). 102 mg of compound **7** was crystallized from the Me₂CO fraction. The CHCl₃ fraction was rechromatographed on Si gel eluting with petroleum ether–Me₂CO and petroleum ether–*i*-PrOH to afford compounds **1** (15 mg), **3** (400 mg), **5** (10 mg), **10** (20 mg) and **13** (24 mg), respectively. Part of the 9:1 CHCl₃–Me₂CO fraction was subjected to repeated chromatography on Si gel sequentially eluting with CHCl₃–Me₂CO, petroleum ether–EtOAc, petroleum ether–Me₂CO and cyclohexane–EtOAc to yield compounds **2** (24 mg), **4** (12 mg) and **11** (5 mg), respectively.

Part of the Me₂CO fraction was further chromatographed on Si gel using CHCl₃–MeOH and CHCl₃–*i*-PrOH as eluents, and on RP₁₈ and MCI gel using MeOH–H₂O and acetonitrile–H₂O as eluents to provide compounds **6** (20 mg), **8** (8 mg), **9** (22 mg), **12** (4 mg), **14** (15 mg) and **15** (30 mg), respectively.

Taxayunnin (1)

C₂₁H₃₀O₄, pale yellow needles, mp 215°C; [α]_D^{16.4} + 7.33° (*c* 0.75, CH₃OH); UV (MeOH) λ_{max} (log ε): 312 (3.7), 296.5 (3.5), 269 (4.3) nm; IR ν_{max} (KBr): 3443, 2972, 2944, 2876, 1734, 1700, 1675, 1671, 1636, 1598, 1560, 1507, 1450, 1419, 1363, 1321, 1212, 1172, 1140, 1085, 1049, 1014, 996, 948, 760, 659, 627 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) data see Table I, ¹³C NMR (125 MHz, CDCl₃) data see Table II; EIMS *m/z* (%): 346 [M]⁺ (100), 331 (20), 313 (51), 304 (5), 287 (20), 271 (15), 259 (18), 245 (64), 233 (20), 219 (18), 203 (17), 193 (16), 173 (10), 157 (7), 145 (8), 128 (9), 115 (10), 91 (11), 77 (7), 69 (9), 57 (14); HREIMS *m/z* 346.2134 [M]⁺, Calcd 346.2144.

Taxamairin C (2)

C₂₁H₂₈O₅, colorless needles; IR ν_{max} (KBr): 3490, 2967, 2931, 2873, 1679, 1608, 1561, 1474, 1423, 1398, 1366, 1325, 1253, 1226, 1176, 1102, 1055, 1039, 995, 918, 887, 748, 551 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) data see Table I; ¹³C NMR (100 MHz, CDCl₃) data see Table II; Positive FABMS *m/z* (%): 361 [M + H]⁺ (100), 343 [M – H₂O + H]⁺ (17), 304 (6), 245 (2), 85 (3), 60 (11).

Taxamairin A (3)

C₂₁H₂₂O₄, yellow needles; ¹H NMR (400 MHz, CDCl₃) δ 7.75 (1H, s, H-4), 7.28 (1H, d, *J* = 9.9 Hz, H-6), 6.08 (1H, d, *J* = 9.8 Hz, H-7), 6.92 (1H, s, H-11), 1.43 (3H, s, CH₃-12), 1.43 (3H, s, CH₃-13), 6.64 (1H, br s, 14-OH), 7.91 (1H, s, H-17), 3.32 (1H, heptet, *J* = 6.8 Hz, H-18), 1.29 (3H, d, *J* = 7.0 Hz, CH₃-19), 1.29 (3H, d, *J* = 7.0 Hz, CH₃-20), 3.87 (3H, s, 15-OCH₃); ¹³C NMR (100 MHz, CDCl₃) data see Table II; EIMS *m/z* (%): 338 [M]⁺ (87), 323 [M – CH₃]⁻ (6), 267 [M – (CH₃)₂CH–CO]⁻ (55), 252 (10), 237 (13), 165 (16), 115 (7).

3β-Hydroxysandaracopimaric Acids (4)

C₂₀H₃₀O₃, colorless needles; ¹H NMR (400 MHz, CDCl₃) δ 1.81 (1H, dd, *J* = 2.3, 12.1 Hz, H-1a), 1.27 (1H, overlap, H-1b), 1.70 (1H, m, H-2a), 1.58 (1H, m, H-2b), 4.10 (1H, dd, *J* = 4.0, 11.7 Hz, H-3), 1.78 (1H, dd, *J* = 2.3, 12.2 Hz, H-5), 1.54 (1H, m, H-6a), 1.27 (1H, overlap, H-6b), 2.23 (1H, dd, *J* = 3.1, 14.5 Hz, H-7a), 2.09 (1H, m, H-7b), 1.74 (1H, br d, *J* = 8.5 Hz, H-9), 1.63 (1H, m, H-11a), 1.52 (1H, m, H-11b), 1.46 (1H, m, H-12a), 1.37 (1H, m, H-12b), 5.23 (1H, br s, H-14), 5.76 (1H, dd, *J* = 10.6, 17.4 Hz, H-15), 4.90 (1H, dd, *J* = 1.3, 17.4 Hz, H-16a), 4.88 (1H, dd, *J* = 1.3, 10.6, H-16b), 1.04 (3H, s, H-17), 1.16 (3H, s, H-19), 0.83 (3H, s, H-20), 1.17 (1H, br s, 3'-OH); ¹³C NMR (100 MHz, CDCl₃) data see Table II; Negative FABMS *m/z* (%): 317 [M – H]⁻ (100), 301 (2), 97 (1); FABMS *m/z* (%): 318 [M]⁺ (5), 301 (54), 283 (15), 249 (9), 219 (16), 191 (49), 177 (11), 157 (100), 139 (12), 122 (6), 106 (3), 92 (1), 79 (3).

(+)-3-Hydroxyisodrimenin (5)

C₁₅H₂₂O₃, white needles; ¹³C NMR (100 MHz, CDCl₃) δ 32.7 (t, C-1), 27.4 (t, C-2), 78.7 (d, C-3), 38.9 (s, C-4), 51.7 (d, C-5), 18.1 (t, C-6), 25.5 (t, C-7), 158.9 (s, C-8), 135.4 (s, C-9), 34.7 (s, C-10), 70.6 (t, C-11), 172.5 (s, C-12), 15.4 (q, C-13), 28.3 (q, C-14), 20.1 (q, C-15); EIMS *m/z* (%): 250 [M]⁺ (61), 232 (18), 217 (56), 207 (100), 194 (51), 189 (45), 175 (10), 163 (31), 151 (57), 138 (21), 125 (16), 119 (26), 105 (41), 91 (63), 77 (46), 69 (66), 55 (37).

Rubrosterone (6)

C₁₉H₂₆O₅, amorphous solid; ¹³C NMR (100 MHz, CD₃OD) δ 37.4 (t, C-1), 68.5 (d, C-2), 68.7 (d, C-3), 32.9 (t, C-4), 52.0 (d, C-5), 205.9 (s, C-6), 122.5 (d, C-7), 164.6 (s, C-8), 35.9 (d, C-9), 39.4 (s, C-10), 20.8 (t, C-11), 25.0 (t, C-12), 54.1 (s, C-13), 80.5 (s, C-14), 29.1 (t, C-15), 34.0 (t, C-16), 220.2 (s, C-17), 17.6 (q, CH₃-18), 24.6 (q, CH₃-19); Negative FABMS *m/z* (%): 333 [M - H]⁻ (100), 316 (55), 172 (1), 155 (2), 125 (4), 111 (6), 97 (17), 80 (16).

Ponasterone A (7)

C₂₇H₄₄O₆, white needles; IR ν_{\max} (KBr): 3387, 2958, 2873, 1645, 1381, 1051, 874 cm⁻¹; ¹H NMR (400 MHz, Pyridine-*d*₅) δ 2.13–2.26 (1H, m, H-1a), 1.87–1.97 (1H, m, H-1b), 4.18 (1H, br dd, *J* = 3.7, 11.5 Hz, H-2), 4.23 (1H, br s, H-3), 2.02–2.10 (1H, m, H-4a), 1.69–1.86 (1H, m, H-4b), 3.02 (1H, dd, *J* = 3.6, 13.0 Hz, H-5), 6.27 (1H, d, *J* = 2.0 Hz, H-7), 3.60 (1H, br t, *J* = 8.4 Hz, H-9), 1.69–1.86 (1H, m, H-11a), 1.87–1.97 (1H, m, H-11b), 2.02–2.10 (1H, m, H-12a), 2.62 (1H, ddd, *J* = 4.0, 8.5, 12.5 Hz, H-12b), 1.87–1.97 (1H, m, H-15a), 2.13–2.26 (1H, m, H-15b), 2.02–2.10 (1H, m, H-16a), 2.46 (1H, ddd, *J* = 3.9, 9.9, 13.7 Hz, H-16b), 2.93 (1H, t, *J* = 9.2 Hz, H-17), 1.23 (3H, s, CH₃-18), 1.06 (3H, s, CH₃-19), 1.58 (3H, s, CH₃-21), 3.81 (1H, d, *J* = 10.1 Hz, H-22), 1.54 (1H, d, *J* = 9.9 Hz, H-23a), 1.69–1.86 (1H, m, H-23b), 1.37–1.46 (1H, m, H-24a), 1.69–1.86 (1H, m, H-24b), 1.46–1.53 (1H, m, H-25), 0.81 (3H, d, *J* = 4.5 Hz, CH₃-26), 0.80 (1H, d, *J* = 4.5 Hz, CH₃-27); ¹³C NMR (100 MHz, Pyridine-*d*₅) δ 38.0 (t, C-1), 68.1 (d, C-2), 68.1 (d, C-3), 32.5 (t, C-4), 51.4 (d, C-5), 203.4 (s, C-6), 121.8 (d, C-7), 166.0 (s, C-8), 34.5 (d, C-9), 38.7 (s, C-10), 21.2 (t, C-11), 32.1 (t, C-12), 48.2 (s, C-13), 84.3 (s, C-14), 31.8 (t, C-15), 21.5 (t, C-16), 50.1 (d, C-17), 17.9 (q, CH₃-18), 24.5 (q, CH₃-19), 76.8 (s, C-20), 21.6 (q, CH₃-21), 76.8 (d, C-22), 30.3 (t, C-23), 37.2 (t, C-24), 28.2 (d, C-25), 22.4 (q, CH₃-26), 23.3 (q, CH₃-27); Positive FABMS *m/z* (%): 465 [M + H]⁺ (29), 385 (77), 343 (39), 283 (19), 85 (59).

Ecdysterone (β-edysone) (8)

C₂₇H₄₄O₇, colorless block crystals; Positive FABMS *m/z* (%): 481 [M + H]⁺ (100), 463 (35), 445 (40), 427 (180), 347 (16), 303 (14), 250 (11), 143 (23), 69 (340).

20-Hydroxyechysone 20,22-monoacetone (9)

C₃₀H₄₈O₇, white powders; Negative FABMS *m/z* (%): 519 [M - H]⁻ (100), 503 (48), 443 (7), 318 (2), 249 (3), 125 (5).

7-Oxositosterol (10)

C₂₉H₄₈O₂, white needles; EIMS *m/z* (%): 428 [M]⁺ (100), 414 (14), 395 (9), 287 (14), 247 (10), 205 (10), 192 (21), 161 (18), 135 (16), 95 (18), 81 (22), 69 (29).

Stigmast-4-en-6 β -ol-3-one (11)

C₂₉H₄₈O₂, white needles; Positive FABMS *m/z* (%): 429 [M + H]⁺ (19), 373 (100), 355 (180), 263 (28), 235 (38), 206 (56), 171 (69), 157 (81), 133 (66), 115 (44), 101 (36), 79 (42), 65 (14).

5 α ,6 β -Dihydroxydaucosterol (12)

C₃₅H₆₂O₈, pale yellow needles; ¹H NMR (400 MHz, pyridine-*d*₅) δ 5.02 (1H, m, H-3), 2.83 (1H, t, *J* = 11.7 Hz, H-4a), 2.50 (1H, dd, *J* = 4.6, 12.5 Hz, H-4b), 4.14 (1H, br s, H-6), 0.71 (3H, s, CH₃-18), 1.53 (3H, s, CH₃-19), 0.97 (3H, d, *J* = 6.4 Hz, CH₃-21), 0.85 (3H, d, *J* = 6.8 Hz, CH₃-26), 0.86 (3H, d, *J* = 6.8 Hz, CH₃-27), 0.88 (3H, t, *J* = 7.4 Hz, CH₃-29), 4.95 (1H, d, *J* = 7.6 Hz, H-1'), 4.05 (1H, t, *J* = 8.2 Hz, H-2'), 4.19 (1H, t, *J* = 8.9 Hz, H-3'), 4.28 (1H, t, *J* = 9.0 Hz, H-4'), 3.75 (1H, m, H-5'), 4.49 (1H, dd, *J* = 2.2, 11.9 Hz, H-6'a), 4.38 (1H, dd, *J* = 5.0, 11.8, H-6'b); ¹³C NMR (100 MHz, pyridine-*d*₅) δ 33.1 (t, C-1), 29.6 (t, C-2), 75.4 (d, C-3), 38.7 (t, C-4), 75.9 (s, C-5), 76.4 (d, C-6), 35.7 (t, C-7), 31.3 (d, C-8), 45.9 (d, C-9), 39.2 (s, C-10), 21.8 (t, C-11), 40.7 (t, C-12), 44.2 (s, C-13), 56.7 (d, C-14), 24.7 (t, C-15), 28.7 (t, C-16), 56.7 (d, C-17), 12.4 (q, CH₃-18), 17.0 (q, CH₃-19), 36.6 (d, C-20), 19.1 (q, CH₃-21), 34.4 (t, C-22), 27.0 (t, C-23), 46.2 (d, C-24), 29.6 (d, C-25), 19.3 (q, CH₃-26), 20.0 (q, CH₃-27), 23.8 (t, C-28), 12.2 (q, CH₃-29), 102.4 (d, C-1'), 75.4 (d, C-2'), 78.6 (d, C-3'), 71.8 (d, C-4'), 78.3 (d, C-5'), 62.9 (t, C-6'); Negative FABMS *m/z* (%): 609 [M - H]⁻ (87), 571 (32), 519 (31), 505 (56), 479 (100), 463 (23), 421 (30), 325 (75), 311 (52), 294 (38), 245 (12), 159 (22), 119 (37), 99 (11), 80 (28).

1-O- β -D-Glucopyranosyl-(2S, 3R, 4E, 8Z)-2-N-(2'-hydroxypalmitoyl)-octadecaspinga-4,8,-dienine (15)

C₄₀H₇₅NO₉, white powders; ¹H NMR (400 MHz, CD₃OD) δ 5.72 (1H, dt, *J* = 15.4, 5.8 Hz, H-5), 5.48 (1H, dd, *J* = 7.4, 15.5 Hz, H-4), 5.37 (2H, m, H-8 and H-9), 4.26 (1H, d, *J* = 7.8 Hz, H-1''), 0.90 (6H, t, *J* = 7.0 Hz, CH₃-18 and CH₃-16'); ¹H NMR (400 MHz, Pyridine-*d*₅) δ 4.70 (1H, dd, *J* = 5.8, 10.3 Hz, H-1a), 4.21 (1H, overlap, H-1b), 4.76 (1H, m, H-2), 4.57 (1H, br t, *J* = 4.3 Hz, H-3), 5.95 (2H, m, H-4 and H-5), 5.48 (2H, m, H-8 and H-9), 0.86 (6H, t, *J* = 6.8 Hz, CH₃-18 and CH₃-16'), 8.36 (1H, d, *J* = 8.6 Hz, 2-NH). 4.76 (1H, overlap, H-2'), 4.90 (1H, d, *J* = 7.7 Hz, H-1''), 4.03 (1H, br t, *J* = 6.4 Hz, H-2''), 4.21 (2H, m, H-3'' and H-4''), 3.09 (1H, m, H-5''), 4.50 (1H, d, *J* = 10.1 Hz, H-6''a), 4.34 (1H, dd, *J* = 5.2, 11.7 Hz, H-6''b); ¹³C NMR (100 MHz, Pyridine-*d*₅, values may be interchanged) δ 70.1 (t, C-1), 54.7 (d, C-2), 72.6 (d, C-3), 132.1 (2C, d, C-4 and C-5), 32.9 (t, C-6), 27.6 and 27.4 (each t, C-7 and C-10), 130.7 and 129.5 (each d, C-8 and C-9), 14.3 (2C, q, CH₃-18 and CH₃-16'), 175.7 (s, C-1'), 72.4 (d, C-2'), 105.6 (d, C-1''), 75.1 (d, C-2''), 78.5 (2C, d, C-3'' and C-5''), 71.7 (d, C-4''), 62.8 (t, C-6''); Positive FABMS *m/z* (%): 715 (53), 697 (67), 534 (78), 262 (92); HRFABMS *m/z* 714.5497, calcd. 714.550.

Acknowledgements

This project was supported by grants from the National Science Foundation of China (3950081), the Young Academic and Technical Leader Raising Foundation of Yunnan Province (awarded to H.-J. Z.), and the Special Supported Biosciences and Biotechniques Foundation of Academia Sinica (STZ-11).

References

- [1] Baloglu, E. and Kingston, D.G.I. (1999), *J. Nat. Prod.* **62**, 1448–1472.
- [2] Appendino, G. (1995), *Nat. Prod. Rep.* **12**, 349–360.
- [3] Parmar, V.S., Jha, A., Bisht, K.S., Taneja, P., Singh, S.K., Kumar, A., Poonam, J.R. and Olsen, C.E. (1999), *Phytochemistry* **50**, 1267–1304.
- [4] Zhang, H.J., Takeda, Y., Minami, Y., Yoshida, K., Unemi, N., Mu, Q., Xiang, W. and Sun, H.D. (1993), *Acta Bot. Yunnanica* **15**, 424–426.
- [5] Zhang, H.J., Takeda, Y., Minami, Y., Yoshida, K., Matsumoto, T., Xiang, W., Mu, Q. and Sun, H.D. (1994), *Chem. Lett.*, 957–960.
- [6] Zhang, H.J., Takeda, Y., Matsumoto, T., Minami, Y., Yoshida, K., Xiang, W., Mu, Q. and Sun, H.D. (1994), *Heterocycles* **38**, 975–980.
- [7] Zhang, H.J., Takeda, Y. and Sun, H.D. (1995), *Phytochemistry* **39**, 1147–1151.
- [8] Zhang, H.J., Sun, H.D. and Takeda, Y. (1995), *J. Nat. Prod.* **58**, 1153–1159.
- [9] Zhang, H.J., Mu, Q., Xiang, W., Yao, P., Sun, H.D. and Takeda, Y. (1997), *Phytochemistry* **44**, 911–915.
- [10] Xiang, W., Zhang, H.J., Sun, H.D., Yao, P. and Li, L. (1999), *Chin. Trad. Herb. Drugs* **30**, 46–48.
- [11] Li, S.H., Zhang, H.J., Yao, P. and Sun, H.D. (1998), *Chin. Chem. Lett.* **9**, 1017–1020.
- [12] Li, S.H., Zhang, H.J., Yao, P., Sun, H.D. and Fong, H.H.S. (2000), *J. Nat. Prod.* **63**, 1488–1491.
- [13] Li, S.H., Zhang, H.J., Yao, P., Sun, H.D. and Fong, H.H.S. (2001), *Phytochemistry* **58**, 369–374.
- [14] Li, S.H., Zhang, H.J., Niu, X.M., Yao, P., Sun, H.D. and Fong, H.H.S. (2002), *Planta Med.*, in press.
- [15] Galicia, M.A., Esquivel, B., Sánchez, A.-A., Cárdenas, J., Ramamoorth, T.P. and Rodríguez-Hahn, L. (1988), *Phytochemistry* **27**, 217–219.
- [16] Liang, J.Y., Min, Z.D., Tanaka, T., Mizuno, M. and Inuma, M. (1988), *Huaxue Xuebao* **46**, 21–25.
- [17] Kozo, D. and Takami, K. (1972), *Phytochemistry* **11**, 841–842.
- [18] Maurs, M., Azerad, R., Cortes, M., Aranda, G., Delahaye, M.B. and Ricard, L. (1999), *Phytochemistry* **52**, 291–296.
- [19] Shiobara, Y., Inoue, S.S., Kato, K., Nishiguchi, Y., Oishi, Y., Nishimoto, N., Oliveira, F.D., Akisue, G., Akisue, M.K. and Hashimoto, G. (1993), *Phytochemistry* **32**, 1527–1530.
- [20] Imai, S., Fujioka, S., Nakanishi, K., Koreeda, M. and Kurokawa, T. (1967), *Steroids* **10**, 557–565.
- [21] Budesinsky, J.P.M., Vokac, K., Laudova, V. and Harmatha, J. (1994), *Phytochemistry* **37**, 707–711.
- [22] Greca, M.D., Monaco, P. and Previtiera, L. (1990), *J. Nat. Prod.* **53**, 1430–1435.
- [23] Kitajima, J., Ida, Y., Noda, N., Komori, T. and Kawasaki, T. (1982), *Yakugaku Zasshi* **102**, 1016–1022.
- [24] Yang, H., Xue, J.L. and Sun, H.D. (1997), *Acta Bot. Yunnanica* **19**, 92–96.
- [25] Sun, Y.K., Young-Hee, C., Hoon, H., Jinwoong, K., Young, C.K. and Heum, S.L. (1997), *J. Nat. Prod.* **60**, 274–276.