



Measurement of three-bond, $^{13}\text{C}'$ - $^{13}\text{C}^\beta$ J couplings in human ubiquitin by a triple resonance, E. COSY-type NMR technique

Jin-Shan Hu & Ad Bax

Laboratory of Chemical Physics, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892-0520, U.S.A.

Received 21 July 1997; Accepted 23 October 1997

Key words: carbon–carbon J coupling, E. COSY, Karplus equation, ϕ angle, 3D NMR, ubiquitin

Abstract

A [CO]HN(CA)CB-E.COSY pulse scheme is described for measurement of three-bond couplings, $^3J_{\text{C}'\text{C}^\beta}$, between carbonyl and aliphatic C^β carbons in ubiquitin, uniformly enriched with ^{13}C and ^{15}N . A Karplus relation, $^3J_{\text{C}'\text{C}^\beta} = 1.28 \cos^2(\phi - 120^\circ) - 1.02 \cos(\phi - 120^\circ) + 0.30$ Hz, is obtained by correlating the $^3J_{\text{C}'\text{C}^\beta}$ values measured for human ubiquitin with backbone ϕ angles from its crystal structure. As predicted, the new Karplus parametrization yields $^3J_{\text{C}'\text{C}^\beta}$ values slightly larger than previously obtained by quantitative J correlation [Hu, J.-S. and Bax, A. (1997) *J. Am. Chem. Soc.*, **119**, 6360–6368], but considerably smaller than what has been reported on the basis of other E.COSY-type measurements carried out on flavodoxin.

The protein backbone torsion angle ϕ is characterized by six three-bond J couplings, $^3J_{\text{HNH}\alpha}$, $^3J_{\text{HNH}\beta}$, $^3J_{\text{HNC}'}$, $^3J_{\text{C}'\text{H}\alpha}$, $^3J_{\text{C}'\text{C}^\beta}$ and $^3J_{\text{C}'\text{C}'}$. All these couplings have been measured in human ubiquitin by quantitative J correlation methods (Hu and Bax, 1996, 1997; Wang and Bax, 1996) whereas E. COSY-type techniques have also been used to measure three of them: $^3J_{\text{HNH}\alpha}$, $^3J_{\text{HNC}'}$, and $^3J_{\text{H}\alpha\text{C}'}$ (Wang and Bax, 1996). For these latter J couplings, excellent agreement is observed between the results obtained with the two different measuring techniques. Karplus-type equations (Karplus, 1959) have been parametrized or reparametrized which correlate these couplings with the intervening dihedral ϕ angle (Hu and Bax, 1996, 1997; Wang and Bax, 1996). For any given residue, there is a strong correlation between the differences between the six measured J couplings and values predicted by their Karplus curves on the basis of the X-ray structure. Indeed, any one of these J couplings can be predicted more accurately on the basis of the remaining five J couplings than on the basis of the

ϕ angle obtained from the crystal structure (Hu and Bax, 1997). This suggests that the methods used for measuring these 3J couplings are highly accurate and self-consistent.

Recently, Löhner et al. (1997) described an elegant new E. COSY-type experiment for measurement of three-bond $^3J_{\text{C}'\text{C}^\beta}$ couplings in the oxidized flavodoxin from *Desulfovibrio vulgaris*. Surprisingly, the magnitude of the $^3J_{\text{C}'\text{C}^\beta}$ couplings and the corresponding Karplus coefficients measured for flavodoxin are considerably ($\sim 40\%$) larger than those from our quantitative J measurements. We predicted (Hu and Bax, 1997) that the values obtained from the HN(CO)C quantitative J correlation experiment would be smaller by only 5–10% relative to the true J values, as no correction for the $^{13}\text{C}^\beta$ T_1 value had been made. In order to eliminate the possibility of a systematic error in our previous measurement of $^3J_{\text{C}'\text{C}^\beta}$, we use a technically straightforward E. COSY-type experiment to remeasure, $^3J_{\text{C}'\text{C}^\beta}$ couplings in human ubiquitin. Although the precision obtainable with this new measurement is lower compared to that of the original HN(CO)C experiment, the new data confirm that the HN(CO)C experiment underestimates $^3J_{\text{C}'\text{C}^\beta}$ by only $\sim 7\%$.

Supplementary material available from the authors: one table with $^3J_{\text{C}'\text{C}^\beta}$ coupling constants in ubiquitin, measured with the [CO]HN(CA)CB-E.COSY experiment.




Figure 2. Regions of (F_1, F_2) strips from the 600 MHz 3D [CO]HN(CA)CB-E.COSY spectrum of 3.5 mg $^{13}\text{C}/^{15}\text{N}$ -enriched ubiquitin (pH 4.7; 30 °C) in a 220 μl Shigemi microcell (1.8 mM) at 600 MHz, taken orthogonal to the F_3 ($^1\text{H}^{\text{N}}$) axis at the $^1\text{H}^{\text{N}}$ and ^{15}N (F_1) frequencies of Arg 54 -Asp 58 . The horizontal displacement for each doublet corresponds to $^1J_{\text{C}^{\text{N}}}$, the small vertical displacement to $^3J_{\text{C}^{\text{C}}\beta}$ (intraresidue), or $^2J_{\text{C}^{\text{C}}\beta}$ for cases where $^{13}\text{C}_{i-1}^{\beta}$ is observed (strips of Thr 55 -Asp 58). The full $^{13}\text{C}^{\beta}$ spectral window is 10 ppm, and extensive aliasing occurs in this dimension.

mension. The applicability of the experiment is limited to relatively small proteins; for larger proteins rapid transverse ^{13}C relaxation during the relatively long $^{13}\text{C}^{\alpha}$ de- and rephasing periods and the long $^{13}\text{C}^{\beta}$ constant-time evolution period will result in very low sensitivity.

The experiment is applied to 1.5 mM $^{13}\text{C}/^{15}\text{N}$ -enriched human ubiquitin, pH 4.7, at 30°, using a Bruker DMX-600 spectrometer, equipped with a triple-resonance probehead containing a self-shielded three-axis gradient coil. The 3D [CO]HN(CA)CB-E. COSY spectrum of human ubiquitin was recorded at 600 MHz as a $88^* \times 43^* \times 512^*$ (n^* refers to n complex points) data matrix with acquisition times of 90.0 (t_1 , ^{15}N), 28.5 (t_2 , ^{13}C), and 61.0 ms (t_3 , $^1\text{H}^{\text{N}}$). The spectral windows were 16 ppm (^{15}N), 10 ppm ($^{13}\text{C}^{\beta}$) and 13.9 ($^1\text{H}^{\text{N}}$) ppm, and extensive aliasing in both the ^{15}N and $^{13}\text{C}^{\beta}$ dimensions was used to maximize the digital resolution. The total measuring time was 48 h. The experiment was carried out twice, and the averages of the two measurements are used for fitting the Karplus curve. The pairwise root-mean-square difference between the two sets of $^3J_{\text{C}^{\text{C}}\beta}$ values was 0.5 Hz. The excitation profile of each of the $180^\circ_{\phi_4}$, $90^\circ_{\phi_5/\phi_7}$, $180^\circ_{\phi_6}$ and $180^\circ_{\phi_8}$ pulses in the $^{13}\text{C}^{\alpha/\beta}$ pulses was

checked to ensure that these pulses would not cause spurious partial inversion of the $^{13}\text{C}'$ pulse. These tests confirmed the lack of any detectable inversion of the $^{13}\text{C}'$ spin, and only minimal conversion of longitudinal to transverse $^{13}\text{C}'$ magnetization (corresponding to a $\leq 4^\circ$ flip angle for each of the $^{13}\text{C}^{\alpha/\beta}$ pulses).

Acquired data were apodized with a 60° -shifted squared sine-bell in the H^{N} dimension, truncated at 10%, with a cosine-bell in the ^{15}N dimension, truncated at 10%, and with a 55° -shifted sine-bell in the $^{13}\text{C}^{\beta}$ dimension, truncated at 5%. Data were zero-filled to yield a digital resolution of 1.9 (F_1), 2.9 (F_2), and 8.2 Hz (F_3). Data were processed using the package NMRPipe (Delaglio et al., 1995). Resonance assignments for human ubiquitin follow those reported previously (Di Stefano and Wand, 1987; Weber et al., 1987; Schneider et al., 1992; Wang et al., 1995).

Figure 2 shows F_1/F_2 ($^{15}\text{N}/^{13}\text{C}$) strips for residues Arg 54 -Asp 58 taken orthogonal to the F_3 (H^{N}) axis of the 3D spectrum. Each amide shows an intense correlation to its intraresidue C^{β} and a weaker one to the C^{β} of the preceding residue, resulting from magnetization transfer via $^2J_{\text{N}^{\text{C}}\alpha}$. The splitting in the horizontal dimension (^{15}N) corresponds to the $^1J_{\text{C}^{\text{N}}}$ coupling whereas the relative displacement of the intraresidue




Figure 3. Relation between ${}^3J_{C'C\beta}$ values measured with the [CO]HN(CA)CB-E.COSY experiment and protein backbone ϕ angles in human ubiquitin. Lines correspond to the Karplus curve ${}^3J_{C'C\beta} = A \cos^2(\phi - 120^\circ) + B \cos(\phi - 120^\circ) + C$ Hz. The thick solid line ($A = 1.28$; $B = -1.02$; $C = 0.30$) represents the best fit between [CO]HN(CA)CB-E.COSY-derived ${}^3J_{C'C\beta}$ values and ϕ angles derived from the crystal structure. The values of the Karplus curve for $\phi > 0^\circ$ are ill determined because of a lack of data points. The thick dashed line is the curve obtained with HN(CO)C-derived values (Hu and Bax, 1997) ($A = 1.61$; $B = -0.66$; $C = 0.26$). The thin dashed line is the curve reported by Löhr et al. (1997) ($A = 2.54$; $B = -0.55$; $C = 0.37$).

C^β doublet components in the ${}^{13}C^\beta$ dimension (F_2) equals ${}^3J_{C'C\beta}$. The displacement of the C^β doublet components of the preceding residues corresponds to ${}^2J_{C'C\beta}$. Correlations to C^β of Asp (e.g., Asp⁵⁸ in Figure 2), Asn, and aromatic residues exhibit an additional splitting in the ${}^{13}C^\beta$ dimension, resulting from ${}^1J_{C\beta C\gamma}$. Note that the difference in the ${}^{15}N$ dimension of these pairs of doublets corresponds to ${}^3J_{NC\gamma}$.

For ubiquitin, ${}^{13}C'T_1$ values at 151 MHz fall in the 1–1.4 s range (N. Tjandra and A. Bax, unpublished results). Considering that this T_1 is much longer than the ${}^{13}C^\beta$ constant-time evolution period and the ${}^{13}C^\alpha$ rephasing period, 2δ , this finite ${}^{13}C'T_1$ value has a negligible effect on the relative displacement of the multiplet components in the ${}^{13}C^\beta$ dimension. No correction for the finite ${}^{13}C'T_1$ is therefore necessary (Wang and Bax, 1996). Peak positions were determined using two different approaches: the commonly used polynomial interpolation method and the average position of ellipsoids best fitted to the calculated contours between 45 and 75% of the peak

maximum (Wang and Bax, 1996), using the program PIPP (Garrett et al., 1991). The two methods yield small (pairwise rmsd 0.21 Hz) random differences and the average of the two ${}^3J_{C'C\beta}$ values measured in this manner is used for all further analysis. Results are available as Supplementary Material.

${}^3J_{C'C\beta}$ couplings measured in ubiquitin with the [CO]HN(CA)CB-E.COSY experiment fall in the -0.2 to 3.1 Hz range. Figure 3 shows the relation between these couplings and the corresponding crystal structure ϕ angles (Vijay-Kumar et al., 1987). A best fit of the measured values to a Karplus equation, ${}^3J = A \cos^2 \theta + B \cos \theta + C$, is calculated using singular value decomposition (SVD), yielding

$${}^3J_{C'C\beta} = 1.28 \cos^2(\phi - 120^\circ) - 1.02 \cos(\phi - 120^\circ) + 0.30 \text{ Hz.} \quad (1)$$

For $\phi > 0^\circ$, the new Karplus curve (thick solid line in Figure 3) is ill-parametrized because of a lack of data. For $-180^\circ < \phi < 0^\circ$, the new parametrization is very similar to the curve obtained previously with the HN(CO)C experiment (thick dashes in Fig-

ure 3), but predicts ${}^3J_{C'C\beta}$ values slightly larger than our earlier curve. This agrees with our previous estimate (Hu and Bax, 1997) that the HN(CO)C-derived ${}^3J_{C'C\beta}$ values are smaller by 5–10% relative to the true values, because no correction for the finite ${}^{13}\text{C}^\beta$ T_1 values was made. The parametrization by Löhrl et al. (thin dashes in Figure 3) deviates substantially from both our earlier and our present parametrization, and predicts substantially larger *trans* ${}^3J_{C'C\beta}$ values ($\phi \approx -60^\circ$).

The rmsd between ${}^3J_{C'C\beta}$ values measured by [CO]HN(CA)CB-E. COSY and those predicted by Equation 1 is 0.4 Hz when using ϕ angles from the crystal structure. This rmsd is considerably poorer than that calculated with ${}^3J_{C'C\beta}$ values measured with the HN(CO)C experiment (0.24 Hz). This larger rmsd reflects the higher random uncertainty in the ${}^3J_{C'C\beta}$ values measured by [CO]HN(CA)CB-E. COSY, owing to the inherently lower sensitivity of the experiment, and the difficulty to measure peak positions in the ${}^{13}\text{C}^\beta$ dimension with an accuracy higher than 0.4 Hz. Note that the acquisition of the time domain data in this dimension is limited to $1/{}^1J_{CC} \approx 28$ ms.

The [CO]HN(CA)CB-E. COSY data reported here indicate the absence of large systematic errors in our previous measurement of ${}^3J_{C'C\beta}$ values in ubiquitin, and confirm that the previously reported Karplus parameters are quite accurate. Our original parametrization also agrees very well with values previously measured for $C'-N-C-C$ cis-form model compounds (Kao and Barfield, 1985). Although ${}^3J_{C'C\beta}$ values measured with the original HN(CO)C slightly underestimate the true values (by $\sim 7\%$ for ubiquitin, and somewhat less for slower tumbling proteins with longer ${}^{13}\text{C}^\beta$ T_1 values), this is accounted for in the original parametrization of this Karplus curve. The precision obtainable for ${}^3J_{C'C\beta}$ is considerably better for the original HN(CO)C experiment than for the [CO]HN(CA)CB-E. COSY experiment and the HN(CO)C experiment is therefore recommended for routine measurement of this coupling. Although the original and present parametrizations are virtually indistinguishable for negative ϕ angles, for positive ϕ angles the new parametrization is essentially undetermined and deviates very significantly. However, it is interesting to note that both in our original HN(CO)C derived Karplus parametrization and in Figure 3 the average of the *trans* ${}^3J_{C'C\beta}$ coupling ($\phi \approx -60^\circ$) falls slightly above the best-fit Karplus curve. This raises the intriguing question whether the assumption underlying the empirical Karplus curve, namely that ${}^3J_{C'C\beta}$ is dominated

by contributions from valence s orbitals to the Fermi contact term, is strictly valid for this type of coupling (Edison et al., 1993; Fukui, 1994). If not, Karplus parameters obtained for a given protein will depend on the fraction of residues with a *trans* ${}^3J_{C'C\beta}$ coupling, and therefore could vary from one protein to another.

Acknowledgements

We thank F. Delaglio and D. Garrett for software support. This work was supported by the AIDS Targeted Anti-Viral Program of the Office of the Director of the National Institutes of Health. J.-S.H. is supported by a postdoctoral fellowship from the Cancer Research Institute, New York, NY, U.S.A.

References

- Delaglio, F., Grzesiek, S., Vuister, G.W., Zhu, G., Pfeifer, J. and Bax, A. (1995) *J. Biomol. NMR*, **6**, 277–293.
- Di Stefano, D.L. and Wand, A.J. (1987) *Biochemistry*, **26**, 7272–7281.
- Edison, A.S., Markley, J.L. and Weinhold, F. (1993). *J. Phys. Chem.*, **97**, 11657–11665.
- Emsley, L. and Bodenhausen, G. (1992) *J. Magn. Reson.*, **97**, 135–148.
- Fukui, H. (1994) *Nucl. Magn. Reson. Spec. Periodic Rep.*, **24**, 133–161.
- Garret, D.S., Powers, R., Gronenborn, A.M. and Clore, G.M. (1991) *J. Magn. Reson.*, **95**, 214–240.
- Hu, J.-S. and Bax, A. (1996) *J. Am. Chem. Soc.*, **118**, 8170–8171.
- Hu, J.-S. and Bax, A. (1997) *J. Am. Chem. Soc.*, **119**, 6360–6368.
- Kao, L.-F. and Barfield, M. (1985) *J. Am. Chem. Soc.*, **107**, 2323–2330.
- Karplus, M. (1959) *J. Chem. Phys.*, **30**, 11–15.
- Kay, L.E., Keifer, P. and Saarinen, T. (1992) *J. Am. Chem. Soc.*, **114**, 1063–1064.
- Löhrl, F., Blümel, M., Schmidt, J.M. and Rüterjans, H. (1997) *J. Biomol. NMR*, **10**, 107–118.
- Palmer, A.G., Cavanagh, J., Wright, P.E. and Rance, M. (1991) *J. Magn. Reson.*, **93**, 151–170.
- Schneider, D.M., Dellwo, M.J. and Wand, A.J. (1992) *Biochemistry*, **31**, 3645–3652.
- Silver, M.S., Joseph, R.I. and Hoult, D.I. (1984) *J. Magn. Reson.*, **59**, 347–351.
- Vijay-Kumar, S., Bugg, C.E. and Cook, W.J. (1987) *J. Mol. Biol.*, **194**, 531–544.
- Wang, A.C. and Bax, A. (1996) *J. Am. Chem. Soc.*, **118**, 2483–2494.
- Wang, A.C., Grzesiek, S., Tschudin, R. and Bax, A. (1995) *J. Biomol. NMR*, **5**, 376–382.
- Weber, P.L., Brown, S.C. and Mueller, L. (1987) *Biochemistry*, **26**, 7282–7290.
- Wittekind, M. and Mueller, L. (1993) *J. Magn. Reson.*, **101**, 201–205.