

Contents of Volume 12, 1998

Vol. 12 No. 1 1998

Research papers

- Recommendations for the presentation of NMR structures of proteins and nucleic acids. IUPAC-IUBMB-IUPAB Inter-Union Task Group on the Standardization of Data Bases of Protein and Nucleic Acid Structures Determined by NMR Spectroscopy
J.L. Markley, A. Bax, Y. Arata, C.W. Hilbers, R. Kaptein, B.D. Sykes, P.E. Wright and K. Wüthrich 1–23
- NMR with ^{13}C , ^{15}N -doubly-labeled DNA: The *Antennapedia* homeodomain complex with a 14-mer DNA duplex
C. Fernández, T. Szyperski, A. Ono, H. Iwai, Shin-ichi Tate, M. Kainosho and K. Wüthrich 25–37
- Experiments and strategies for the assignment of full $^{13}\text{C}/^{15}\text{N}$ -labelled polypeptides by solid state NMR
S.K. Straus, T. Bremi and R.R. Ernst 39–50
- Chemical shift as a probe of molecular interfaces: NMR studies of DNA binding by the three amino-terminal zinc finger domains from transcription factor IIIA
M.P. Foster, D.S. Wuttke, K.R. Clemens, W. Jahnke, I. Radhakrishnan, L. Tennant, M. Reymond, J. Chung and P.E. Wright 51–71
- Internal and overall motions of the translation factor eIF4E: Cap binding and insertion in a CHAPS detergent micelle
A.M. McGuire, H. Matsuo and G. Wagner 73–88
- Complete assignment of ^1H , ^{13}C and ^{15}N chemical shifts for bovine β -lactoglobulin: Secondary structure and topology of the native state is retained in a partially unfolded form
S. Uhrínová, D. Uhrín, H. Denton, M. Smith, L. Sawyer and P.N. Barlow 89–107
- A new general method for the biosynthesis of stable isotope-enriched peptides using a decahistidine-tagged ubiquitin fusion system: An application to the production of mastoparan-X uniformly enriched with ^{15}N and $^{15}\text{N}/^{13}\text{C}$
T. Kohno, H. Kusunoki, K. Sato and K. Wakamatsu 109–121
- Conformational analysis of *Chlamydia*-specific disaccharide α -Kdo-(2→8)- α -Kdp-(2→O)-allyl in aqueous solution and bound to a monoclonal antibody: Observation of intermolecular transfer NOEs
T. Sokolowski, T. Haselhorst, K. Scheffer, R. Weisemann, P. Kosma, H. Brade, L. Brade and T. Peters 123–133
- Comparison of ^{15}N - and ^{13}C -determined parameters of mobility in melittin
L. Zhu, F.G. Prendergast and M.D. Kemple 135–144
- NMR detection of slow conformational dynamics in an endonuclease toxin
S.B.-M. Whittaker, R. Boetzel, C. MacDonald, Lu-Yun Lian, A.J. Pommer, A. Reilly, R. James, C. Kleanthous and G.R. Moore 145–159
- Secondary H/D isotope effect on hydrogen-bonded hydroxyl groups as a tool for recognizing distance constraints in conformational analysis of oligosaccharides
J. Dabrowski, H. Grosskurth, C. Baust and N.E. Nifant'ev 161–172
- Short communications*
- Sequence-specific ^1H assignment and secondary structure of the bacteriocin AS-48 cyclic peptide
G.M. Langdon, M. Bruix, A. Gálvez, E. Valdivia, M. Maqueda and M. Rico 173–175
- An empirical relationship between rotational correlation time and solvent accessible surface area
V. Krishnan and M. Cosman 177–182

| | |
|--|---------|
| Local helix content and RNA-binding activity of the N-terminal leucine-repeat region of hepatitis delta antigen Jya-Wei Cheng, I.-Jin Lin, Yuan-Chou Lou, Ming-Tao Pai and Huey-Nan Wu | 183–188 |
| <i>Assignment notes</i> | |
| Sequential assignment of the triple labelled 30.1 kDa cell-adhesion domain of intimin from enteropathogenic <i>E. coli</i> G. Kelly, S. Prasanna, S. Daniell, G. Frankel, G. Dougan, I. Connerton and S. Matthews | 189–191 |
| Assignment of ^1H , ^{13}C , and ^{15}N signals of turkey ovomucoid third domain at pH 2.0 S. Choe, Z. Dzakula, E.S. Kuloglu and J.L. Markley | 193–195 |
| ^1H , ^{15}N and ^{13}C resonance assignments and secondary structure of apo liver fatty acid-binding protein H. Wang, Yan He, Kuo Tung Hsu, J.F. Magliocca, J. Storch and R.E. Stark | 197–199 |
| ^1H , ^{13}C and ^{15}N NMR backbone assignments of 25.5 kDa metallo- β -lactamase from <i>Bacteroides fragilis</i> S.D.B. Scrofani, P.E. Wright and H.J. Dyson | 201–202 |
| <i>Instructions for authors</i> | 203–207 |

Vol. 12 No. 2 1998

Research papers

| | |
|---|---------|
| Hydrogen bonding geometry of a protein-bound carbohydrate from water exchange-mediated cross-relaxation E.W. Sayers, J.L. Weaver and J.H. Prestegard | 209–222 |
| A new experiment for the measurement of $^n\text{J}(\text{C},\text{P})$ coupling constants including $^3\text{J}(\text{C}_i', \text{P}_i)$ and $^3\text{J}(\text{C}_i', \text{P}_{i+1})$ in oligonucleotides C. Richter, B. Reif, K. Wörner, S. Quant, J.P. Marino, J.W. Engels, C. Griesinger and H. Schwalbe | 223–230 |
| Rotational-echo double-resonance in complex biopolymers: a study of <i>Nephila clavipes</i> dragline silk C.A. Michal and L.W. Jelinski | 231–241 |
| Nuclear magnetic resonance characterization of a paramagnetic DNA-drug complex with high spin cobalt; assignment of the ^1H and ^{31}P NMR spectra, and determination of electronic, spectroscopic and molecular properties M. Gochin | 243–257 |
| Some NMR experiments and a structure determination employing a $\{^{15}\text{N}, ^2\text{H}\}$ enriched protein T.K. Mal, S.J. Matthews, H. Kovacs, I.D. Campbell and J. Boyd | 259–276 |
| Practical model fitting approaches to the direct extraction of NMR parameters simultaneously from all dimensions of multidimensional NMR spectra R.A. Chylla, B.F. Volkman and J.L. Markley | 277–297 |
| An asymmetric deuterium labeling strategy to identify interprotomer and intraprotomer NOEs in oligomeric proteins A. Jasanoff | 299–306 |
| Identification of slow motions in the reduced recombinant high-potential iron sulfur protein I (HiPIP I) from <i>Ectothiorhodospira halophila</i> via ^{15}N rotating-frame NMR relaxation measurements L. Banci, I.C. Felli and D. Koulougliotis | 307–318 |
| <i>Short communications</i> | |
| Rapid measurement of scalar three-bond $^1\text{H}^N$ - $^1\text{H}^\alpha$ spin coupling constants in ^{15}N -labelled proteins H. Pongstingl and G. Otting | 319–324 |
| An HNCO-based pulse scheme for the measurement of $^{13}\text{C}^\alpha$ - $^1\text{H}^\alpha$ one-bond dipolar couplings in ^{15}N , ^{13}C labeled proteins D. Yang, J.R. Tolman, N.K. Goto and L.E. Kay | 325–332 |

| | |
|--|---------|
| Full matrix analysis of cross-relaxation fails in fractionally deuterated molecules Z. Zolnai, N. Juranić and S. Macura | 333–337 |
| Editing of multidimensional NMR spectra of partially deuterated proteins. Measurement of amide deuterium isotope effects on the chemical shifts of protein backbone nuclei A. Meissner, J. Briand and O.W. Sørensen | 339–343 |
| Single transition-to-single transition polarization transfer (ST2-PT) in [¹⁵ N, ¹ H]-TROSY K.V. Pervushin, G. Wider and K. Wüthrich | 345–348 |
| <i>Assignment notes</i> | |
| ¹ H, ¹³ C, ¹⁵ N resonance assignment of the 20 kDa double stranded RNA binding domain of PKR S. Nanduri, B. Carpick, Y. Yang, B.R.G. Williams and J. Qin | 349–351 |
| ¹ H, ¹³ C, and ¹⁵ N NMR resonance assignments of vaccinia glutaredoxin-1 in the fully reduced form J.J. Kelley III and J.H. Bushweller | 353–355 |
| ¹ H, ¹³ C and ¹⁵ N assignment of the Isl-1 homeodomain J.H. Ippel, G. Larsson, G. Behravan, M. Lundqvist, P.-O. Lycksell, J. Schleucher, J. Zdunek and S.S. Wijmenga | 357–359 |

Vol. 12 No. 3 1998

Research papers

| | |
|--|---------|
| Characterization of magnetically oriented phospholipid micelles for measurement of dipolar couplings in macromolecules M. Ottiger and A. Bax | 361–372 |
| Sensitivity optimized HCN and HCNCH experiments for ¹³ C/ ¹⁵ N labeled oligonucleotides R. Fiala, F. Jiang and V. Sklenář | 373–383 |
| Dynamics of the carbohydrate chains attached to the Fc portion of immunoglobulin G as studied by NMR spectroscopy assisted by selective ¹³ C labeling of the glycans Y. Yamaguchi, K. Kato, M. Shindo, S. Aoki, K. Furusho, K. Koga, N. Takahashi, Y. Arata and I. Shimada | 385–394 |
| CAMRA: Chemical shift based computer aided protein NMR assignments W. Gronwald, L. Willard, T. Jellard, R.F. Boyko, K. Rajarathnam, D.S. Wishart, F.D. Sönnichsen and B.D. Sykes | 395–405 |
| The solution structure of a gallium-substituted putidaredoxin mutant: GaPdx C85S T.C. Pochapsky, M. Kuti and S. Kazanis | 407–415 |
| Effects of sample preparation conditions on biomolecular solid-state NMR lineshapes D.L. Jakeman, D.J. Mitchell, W.A. Shuttleworth and J.N.S. Evans | 417–421 |
| On the NMR structure determination of a 44n RNA pseudoknot: Assignment strategies and derivation of torsion angle restraints M.H. Kolk, S.S. Wijmenga, H.A. Heus and C.W. Hilbers | 423–433 |
| <i>Short communications</i> | |
| Spin-state selection filters for the measurement of heteronuclear one-bond coupling constants P. Andersson, J. Weigelt and G. Otting | 435–441 |
| A liquid crystalline medium for measuring residual dipolar couplings over a wide range of temperatures H. Wang, M. Eberstadt, E.T. Olejniczak, R.P. Meadows and S.W. Fesik | 443–446 |
| Improved dilute bicelle solutions for high-resolution NMR of biological macromolecules J.A. Losonczi and J.H. Prestegard | 447–451 |

Assignment notes

- NMR backbone assignments of the cyanobacterial transcriptional factor, SmtB, that senses the zinc concentration in the cell
E.H. Morita, T. Kosada, T. Yamazaki, Y. Kyogoku and H. Hayashi 453–454
- Main-chain signal assignment for the PDZ2 domain from human protein tyrosine phosphatase hPTP1E and its complex with a C-terminal peptide from the Fas receptor
I. Ekiel, D. Banville, S.H. Shen, J.J. Slon-Usakiewicz, A. Koshy and K. Gehring 455–456
- Sequence specific ^1H , ^{13}C and ^{15}N resonance assignment of rat CD2 domain 1
H.A. Chen, M. Pfuhl, B. Davis and P.C. Driscoll 457–458
- Complete heteronuclear NMR resonance assignments and secondary structure of core binding factor β (1-141)
X. Huang, A. Speck and J.H. Bushweller 459–460
- Assignment of ^1H and ^{15}N resonances of Murine Tec SH3 domain
S.E. Pursglove, T.D. Mulhern, M.G. Hinds, R.S. Norton and G.W. Booker 461–462
- Resonance assignment and secondary structure of the cold shock domain of the human YB-1 protein
C.P.A.M. Kloks, A. Hoffmann, J.G. Omichinski, G.W. Vuister, C.W. Hilbers and S. Grzesiek 463–464
- Sequence-specific ^1H , ^{13}C and ^{15}N assignment of the EH1 domain of mouse Eps15
B. Whitehead, M. Tessari, H.H. Versteeg, S. van Delft, P.M.P. van Bergen en Henegouwen and G.W. Vuister 465–466
- ^1H , ^{15}N and ^{13}C chemical shift assignment of the guanine nucleotide exchange domain of human Elongation Factor-one beta
J.M.J. Pérez, J. Kriek, J. Dijk, W. Möller, G. Siegal, K. Hård, A.P. Kalverda and G.W. Canters 467–468
- Erratum 469

Vol. 12 No. 4 1998*Research papers*

- Propagation of experimental uncertainties using the Lipari-Szabo model-free analysis of protein dynamics
D. Jin, M. Andrec, G.T. Montelione and R.M. Levy 471–492
- Microdrop screening: A rapid method to optimize solvent conditions for NMR spectroscopy of proteins
C.A. Lepre and J.M. Moore 493–499
- On using time-averaging restraints in molecular dynamics simulation
W.R.P. Scott, A.E. Mark and W.F. van Gunsteren 501–508
- Local mobility of ^{15}N labeled biomolecules characterized through cross-correlation rates: Applications to paramagnetic proteins
I.C. Felli, H. Desvaux and G. Bodenhausen 509–521
- The NMR solution structure and characterization of pH dependent chemical shifts of the β -elicitin, cryptogein
P.R. Gooley, M.A. Keniry, R.A. Dimitrov, D.E. Marsh, D.W. Keizer, K.R. Gayler and B.R. Grant 523–534
- High pressure NMR study of a small protein gurmarin
K. Inoue, H. Yamada, T. Imoto and K. Akasaka 535–541

Short communications

- Conformational analysis of protein and nucleic acid fragments with the new grid search algorithm FOUND
P. Güntert, M. Billeter, O. Ohlenschläger, L.R. Brown and K. Wüthrich 543–548

| | |
|--|---------|
| Determination of the equilibrium micelle-inserting position of the fusion peptide of gp41 of human immunodeficiency virus type 1 at amino acid resolution by exchange broadening of amide proton resonances D.-K. Chang and S.-F. Cheng | 549–552 |
| PSEUDYANA for NMR structure calculation of paramagnetic metalloproteins using torsion angle molecular dynamics L. Banci, I. Bertini, M.A. Cremonini, G. Gori-Savellini, C. Luchinat, K. Wüthrich and P. Güntert | 553–557 |
| 3D HCCH-COSY-TOCSY experiment for the assignment of ribose and amino acid side chains in ¹³ C labeled RNA and protein W. Hu, L.T. Kakalis, L. Jiang, F. Jiang, X. Ye and A. Majumdar | 559–564 |
| <i>Assignment notes</i> | |
| ¹ H and ¹⁵ N resonance assignment of the calcium-bound form of the <i>Nereis diversicolor</i> sarcoplasmic Ca ²⁺ -binding protein C.T. Craescu, B. Prêcheur, A. van Riel, H. Sakamoto, J.A. Cox and Y. Engelborghs | 565–566 |
| Assignment of ¹ H, ¹³ C and ¹⁵ N signals of the inhibitor protein Im9 bound to the DNase domain of colicin E9 R. Boetzel, M. Czisch, C.J. MacDonald, R. Kaptein, A.M. Hemmings, R. James, C. Kleanthous and G.R. Moore | 567–568 |
| ¹ H and ¹⁵ N resonance assignment of neural cell adhesion molecule module-2 P.H. Jensen, N.K. Thomsen, V. Soroka, V. Berezin, E. Bock and F.M. Poulsen | 569–570 |
| Assignments of ¹ H and ¹³ C resonances in the complex of palmitate and a non-specific lipid transfer protein (ns-LTP) isolated from barley seeds M.H. Lerche and F.M. Poulsen | 571–572 |
| <i>Author index</i> | 573–575 |
| <i>Keyword index</i> | 576–578 |
| <i>Volume contents</i> | 579–583 |
| <i>Instructions for authors</i> | 584–588 |