

## Letters to the Editor

### NMR backbone assignment of the mitogen-activated protein (MAP) kinase p38

DOI 10.1007/s10858-005-2449-x

MAP kinase p38 is an important anti-inflammatory drug target (Lee et al., 1994). The construct p38(-2-349) of mouse p38 is very stable and suitable for applications in drug discovery. We have partially assigned p38 using principles developed for the catalytic domain of protein kinase A (Langer et al., 2004). 3D heteronuclear NMR experiments with  $^2\text{H}$ ,  $^{13}\text{C}$ ,  $^{15}\text{N}$  p38, 2D NMR experiments with selectively  $^{15}\text{N}$ -Tyr, Phe, Met, Ile, Val, Leu labeled p38 and distance information from a paramagnetic adenosine derivate were used. 254 (75.4%) of the expected 337 amide signals were found. Backbone and  $\text{C}_\beta$  shifts for 216 of these (85.0, 64.0% based on the sequence) were assigned and are mapped onto the sequence in Figure 1 (supplementary material), BMRB deposit with Accession number 6468.

References: Langer, et al. (2004) *Chembiochem.*, **5**, 1508–1516; Lee, et al. (1994) *Nature.*, **372**, 739–746.

Martin Vogtherr, Krishna Saxena, Susanne Grimme, Marco Betz, Ulrich Schieborr, Barbara Pescatore, Thomas Langer & Harald Schwalbe  
*Institut für Organische Chemie und Chemische Biologie, Zentrum für Biomolekulare Magnetische Resonanz, Goethe-Universität Frankfurt, D-60439, Frankfurt*

**Supplementary material** to this paper is available in electronic format at <http://dx.doi.org/10.1007/s10858-005-2449-x>.

### Resonance assignments for the DNA binding domain of ERCC-1/XPF heterodimer

DOI 10.1007/s10858-005-5275-2

Nucleotide excision repair (NER) is a sophisticated DNA repair mechanism that eliminates a wide variety of DNA lesions. The endonuclease ERCC-1/XPF (Excision Repair Cross Complementation group 1/ Xeroderma Pigmentosum group F) complex is the structure-specific endonuclease that participates in the repair of DNA damage by making the 5' incision. In the ERCC-1/XPF heterodimer, C-terminal domains of both subunits contain two tandemly repeated helix-hairpin-helix DNA binding motifs. The heterodimeric DNA binding domain in the ERCC-1/XPF complex plays a crucial role in positioning it at the appropriate location containing irregular DNA structures. We initiated a NMR structure determination of the heterodimeric DNA binding domain of the ERCC-1/XPF to explore its binding mode to unusual DNA structures. Here, we report nearly complete  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{15}\text{N}$  resonance assignment of the DNA binding domain in human ERCC-1/XPF heterodimer; ERCC-1 and XPF subunits consist of 79 residues (residues 219–297) and 96 residues (residues 810–905), respectively. BMRB deposit with accession number 6502.

Mika Masuyama<sup>a,b</sup>, Sonoko Ishino<sup>a</sup>, Tatsuya Nishino<sup>a</sup>, Hiroshi Moriuchi<sup>a</sup>, Hiroshi Ueno<sup>b</sup>, Kousuke Morikawa<sup>a</sup> & Shin-ichi Tate<sup>a,\*</sup>

<sup>a</sup>*Biomolecular Engineering Research Institute (BERI), Osaka, 565-0874, Japan;* <sup>b</sup>*Nara Women's University, Nara, 630-8506, Japan*

\*To whom correspondence should be addressed. E-mail: [tate@beri.or.jp](mailto:tate@beri.or.jp)

**Supplementary material** to this paper is available in electronic format at <http://dx.doi.org/10.1007/s10858-005-5275-2>.