

Letters to the Editor

¹H, ¹³C, and ¹⁵N sequence-specific resonance assignment and secondary structure of *Plasmodium falciparum* thioredoxin*

DOI 10.1007/s10858-005-0458-4

The malaria parasite *Plasmodium falciparum* spends part of its life cycle in human erythrocytes, where it is challenged with enhanced oxidative stress, and where it needs efficient anti-oxidants to maintain its reducing milieu and to protect itself against damage (Rahlfs et al., 2002). Thioredoxin (*Pf*Trx) is one of the major redox-regulatory molecules which, because of its dithiol/disulfide exchange activity, determines the oxidation state of protein cysteines, being therefore a potential target in research towards anti-malarial drugs (Kanzok et al., 2000). We initiated the structure determination of the recombinant 104-amino acid polypeptide, using multidimensional NMR spectroscopy. Sequential backbone resonance assignments were carried out on the basis of HNCA, CBCA(CO)NH, CBCANH, HNCO, ¹H–¹⁵N-HSQC and ¹H–¹⁵N-NOESY-HSQC experiments. Side-chain assignments were done using the HCCH-TOCSY experiment with the sample in ²H₂O. The chemical shifts for 99% of the backbone ¹H^N, ¹³C^α, ¹⁵N^H (without considering the proline residues), for 89% of the ¹³C^γ, 99% of the ¹H^γ, and 88% of all side-chain atoms (except the ¹³C shifts of the aromatic rings) of the *Pf*Trx were assigned. Data were deposited in the BioMagResBank under accession number 6442.

References: Kanzok et al. (2000) *J. Biol. Chem.*, **275**, 40181–40186; Rahlfs et al. (2002) *Cell. Mol. Life Sci.*, **59**, 1024–1041.

Claudia Elisabeth Munte^a, Katja Becker^b, Rolf Heiner Schirmer^c & Hans Robert Kalbitzer^{d,*}

^aInstituto de Física de São Carlos, Universidade de São Paulo, 13560-970, São Carlos, SP, Brazil;

^bInterdisziplinäres Forschungszentrum, Universität Giessen, 35392, Giessen, Germany; ^cBiochemie-Zentrum der Universität, 69120, Heidelberg, Germany; ^dInstitut für Biophysik und Physikalische Biochemie, Universität Regensburg, 93040, Regensburg, Germany

*To whom correspondence should be addressed. E-mail: hans-robot.kalbitzer@biologie.uni-regensburg.de
Supplementary material to this paper is available in electronic format at <http://dx.dio.org/10.1007/10858-005-0458-4>.

NMR assignment of human ubiquitin conjugating enzyme Ubc7

10.1007/s10858-005-1257-7

Human Ubc7, which is involved in the targeting of ER-associated proteins for degradation by the 26S proteasome, is unique among class I enzymes in that it can catalyze the linkage of ubiquitin to ubiquitin in the absence of other substrates and ubiquitin protein ligases (Kikkert et al., 2004). The presence of this activity indicates that Ubc7 contains a specific binding site for ubiquitin and has catalytic features that are distinct from other E2 enzymes. In order to gain an understanding of this activity at molecular detail, we have initiated NMR structural studies of Ubc7. 2D and 3D heteronuclear NMR experiments were carried out with uniformly ¹³C, ¹⁵N-labeled Ubc7, either fully protonated or ~70% perdeuterated. The ¹H, ¹³C, and ¹⁵N chemical shift assignments of Ubc7 are nearly complete, with exception of residues M1, A2, P20, P68 and aromatic carbon resonances. BMRB deposit with accession number 6711.

Reference: Kikkert et al. (2004) *J. Biol. Chem.*, **279**, 3525–3534.

Kathryn B. Briggman^a, Ananya Majumdar^a, Catherine S. Coleman^b, Vincent Chau^b & Joel R. Tolman^{a,*}

^aDepartment of Chemistry, Johns Hopkins University, Baltimore, 21218 MD, U.S.A.; ^bDepartment of Cellular and Molecular Physiology, Milton S. Hershey Medical Center, Pennsylvania State University College of Medicine, Hershey, PA 17033, U.S.A.

*To whom correspondence should be addressed. E-mail: tolman@jhu.edu

Supplementary material to this paper is available in electronic format at <http://dx.doi.org/10.1007/s10858-005-1257-7>