

## Letters to the Editor

### <sup>1</sup>H, <sup>13</sup>C and <sup>15</sup>N backbone and side chain resonance assignments of *Haloferax volcanii* DHFR1

DOI 10.1007/s10858-005-0459-3

*Haloferax volcanii* is an archaeal species which exploits the niche of hyper-saline environments. To understand how enzymes can adapt to function in very different salt concentrations, the salt-dependence of the properties of dihydrofolate reductases (DHFRs) from *H. volcanii* (*hv*DHFR1) and *E. coli* (*ec*DHFR) are being studied (Wright et al., 2002). To test the relationship between stability, activity and flexibility, we propose to investigate the flexibility of *ec*DHFR and *hv*DHFR1 through NMR relaxation studies at varying salt concentrations. Nearly complete chemical shift assignments of 50% <sup>2</sup>H and uniformly <sup>15</sup>N and <sup>13</sup>C labeled *hv*DHFR1 in 3.5 M NaCl have been accomplished through 2D and 3D heteronuclear NMR experiments. The high viscosity of the solvent makes the protein's correlation time more like a 25 kDa protein. Therefore we used ~50% <sup>2</sup>H labeling, which resulted in interpretable data. Chemical shift assignments were accomplished for 97% of the backbone amides, the residues A10, E11, R13, D18, and R69 were unassigned. In total, 98% of <sup>13</sup>C<sup>α</sup>, 96% of <sup>13</sup>C<sup>β</sup>, 92% of <sup>13</sup>CO, 97% of <sup>1</sup>H<sup>α</sup> and 97% of <sup>1</sup>H<sup>β</sup> were assigned. BMRB accession Nr. 6645.

Reference: Wright et al. (2002) *J. Mol. Biol.*, **323**, 327–344.

Bulent Binbuga &amp; John K. Young\*

*Department of Chemistry, Mississippi State University, Box 9573 MS, 39762, USA*

\*To whom correspondence should be addressed. E-mail: jky1@ra.msstate.edu

**Supplementary material** is available in electronic format at <http://dx.dio.org/10.1007/s10858-005-0459-3>.

### NMR Assignment of HI1506, a novel two-domain protein from *Haemophilus influenzae*

DOI 10.1007/s10858-005-3869-3

HI1506 is a 128-residue hypothetical protein of unknown function from *H. influenzae*. It was originally annotated as a shorter 85-residue protein but a sequence correction conducted in our laboratory revealed that the full-length protein has an additional 43 residues on the C-terminus, corresponding with a region initially ascribed to HI1507. As part of a larger effort to understand the functions of hypothetical proteins from Gram-negative bacteria, and *H. influenzae* in particular, we report here the NMR assignments for the corrected full-length HI1506 protein. Sequence-specific <sup>1</sup>H, <sup>13</sup>C, and <sup>15</sup>N assignments were completed for more than 95% of the backbone atoms. Most side chain atoms have also been assigned. Our results indicate that HI1506 has two domains connected by a short linker. The assignments provide the basis for structure determination and, potentially, further insight into function. The chemical shifts have been deposited in the BMRB (accession number BMRB-6780).

Nese Sari, Deok Cheon Yeh, Victoria Doseeva, Karen Surabian, Osnat Herzberg, &amp; John Orban\*

*Center for Advanced Research in Biotechnology, University of Maryland Biotechnology Institute, 9600 Gurdelsky Drive, Rockville, MD, 20850, USA*

\*To whom correspondence should be addressed. E-mail: orban@umbi.umd.edu

**Supplementary material** is available in electronic format at <http://dx.dio.org/10.1007/s10858-005-3869-3>.