

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

Efficient Conversion of a 1,6-Anhydro Chitobiose Derivative into the Corresponding Tetradecyl β -Glycoside Derivative by Means of Participation of a Neighboring Tetradecanamide Group

Atsushi Umino^a; Hiroshi Hinou^a; Hiroyoshi Kuzuhara^a; Koji Matsuoka^a; Daiyo Terunuma^b; Shunya Takahashi^b

^a Department of Functional Materials Science, Faculty of Engineering, Saitama University, Urawa-shi, Japan ^b The Institute of Physical and Chemical Research, Wako-shi, Saitama, Japan

To cite this Article Umino, Atsushi , Hinou, Hiroshi , Kuzuhara, Hiroyoshi , Matsuoka, Koji , Terunuma, Daiyo and Takahashi, Shunya(1998) 'Efficient Conversion of a 1,6-Anhydro Chitobiose Derivative into the Corresponding Tetradecyl β -Glycoside Derivative by Means of Participation of a Neighboring Tetradecanamide Group', *Journal of Carbohydrate Chemistry*, 17: 2, 231 – 239

To link to this Article: DOI: 10.1080/07328309808002324

URL: <http://dx.doi.org/10.1080/07328309808002324>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

**EFFICIENT CONVERSION OF A 1,6-ANHYDRO CHITOBIOSE
DERIVATIVE INTO THE CORRESPONDING TETRADECYL β -
GLYCOSIDE DERIVATIVE BY MEANS OF PARTICIPATION OF A
NEIGHBORING TETRADECANAMIDE GROUP**

Atsushi Umino,^a Hiroshi Hinou,^a Koji Matsuoka,^a Daiyo Terunuma,^a Shunya Takahashi,^b
and Hiroyoshi Kuzuhara^{a*}

^a Department of Functional Materials Science, Faculty of Engineering, Saitama University,
255 Shimo-okubo, Urawa-shi 338, Japan, ^b The Institute of Physical and Chemical
Research, Wako-shi, Saitama 351-01, Japan

Received July 11, 1997 - Final Form November 7, 1997


ABSTRACT

In the course of our studies on the synthesis of amphiphilic chitoheptaose derivatives carrying binary long hydrocarbon chains at the reducing sugar, tetradecyl 4-*O*-(2-amino-2-deoxy- β -D-glucopyranosyl)-2-deoxy-2-tetradecanamido- β -D-glucopyranoside was prepared as a model. For general applicability, the 1,6-anhydro-2-deoxy-2-tetradecanamido- β -D-glucopyranosyl moiety was employed as a precursor of the reducing end. Acetolysis of the 1,6-anhydro ring using triethylsilyl triflate gave an oxazoline intermediate as a major product, accompanied by α -glycosyl acetate as a by-product. Immediate treatment of the mixture with 1-tetradecanol and protic acid followed by a separation work-up efficiently led to tetradecyl β -glycoside derivative.

INTRODUCTION


Chitin, one of the abundant polysaccharides, and its deacetylated derivative, chitosan, have attracted much attention from the view point of their diverse biological possibilities.¹ Small fragments of chitin such as *N*-acetylchitohexaose and larger

oligosaccharides of this type as well as their deacetylated compounds are also interesting because of their biological activities in both mammals and plants. For example, chitohexaose and its *N*-peracetylated derivative show a growth-inhibitory effect against solid tumors implanted in mice,² while *N*-acetylchitooligosaccharides of more than dp 6 induce phytoalexin formation in rice cells.³ Unexpectedly, glycolipid type derivatives of chitooligosaccharide of dp 2-4 possessing a long fatty *N*-acyl group at the reducing D-glucosamine moiety were recently found to have immunostimulating and antitumor activities.⁴ Because of this biological relevance, we planned to introduce binary hydrophobic tails into the reducing end of the originally bioactive chitoheptaose, mimicking phospholipids of cell membranes. One hydrophobic group was to be introduced as an *N*-acyl group, while the other as an aglycon of a glycoside. Prior to synthesis of the heptasaccharide derivative, a general way to introduce different types of hydrophobic groups into the terminal monosaccharide moiety had to be established. This paper describes the synthesis of tetradecyl 4-*O*-(2-amino-2-deoxy- β -D-glucopyranosyl)-2-deoxy-2-tetradecanamido- β -D-glucopyranoside (**1**) as a disaccharidic model compound, which includes an efficient conversion of 1,6-anhydro-2-deoxy-2-tetradecanamido- β -D-glucopyranose moiety, a precursor of the reducing end, into the corresponding tetradecyl β -glycopyranoside moiety.




RESULTS AND DISCUSSION

1,6-Anhydro-2-azido-3-*O*-benzyl-2-deoxy- β -D-glucopyranoside (**2**)⁵ was glycosylated at the 4-position with phenyl 3,4,6-tri-*O*-acetyl-1,2-dideoxy-2-phthalimido-1-thio- β -D-glucopyranoside (**3**)⁶ activated by *N*-iodosuccinimide and a catalytic amount of trifluoromethanesulfonic acid, giving 1,6-anhydrodisaccharide derivative (**4**) in good yield. The azido group of **4** was selectively reduced with hydrogen sulfide in pyridine-triethylamine mixture and the resulting amino group was immediately treated with tetradecanoyl chloride in pyridine for *N*-acylation to give 4-*O*-(3,4,6-tri-*O*-acetyl-2-deoxy-



2-phthalimido- β -D-glucopyranosyl)-1,6-anhydro-3-*O*-benzyl-2-deoxy-2-tetradecanamido- β -D-glucopyranose (**5**) in 96% yield as crystals.


The initial attempt to convert the 1,6-anhydro derivative **5** into the corresponding tetradecanyl β -glycoside derivative **8** was conducted in three steps: i.e., 1) acetolysis of the 1,6-anhydro ring into 1,6-diacetate, 2) tentative formation of an oxazoline intermediate, and 3) coupling with 1-tetradecanol for β -glycoside formation as shown on the next page. Acetolysis was carried out as follows. When 20 equivalents of trifluoroacetic acid was employed as catalyst, the reaction proceeded very sluggishly, giving α -glycosyl acetate **6** ($J_{1,2} = 3.5$ Hz in ^1H NMR spectrum) in only 43% yield after 3 days. On the other hand, 5 equivalents of boron trifluoride diethyl etherate remarkably improved the acetolysis reaction, which was complete within 10 minutes giving **6** in 61% yield. Compound **6** was then treated with trimethylsilyl triflate at 50 °C for 1 hour for conversion into the oxazoline intermediate **7**,⁷ which, without isolation, was soon condensed with 1-tetradecanol in the presence of a catalytic amount of (+)-10-camphorsulfonic acid, giving tetradecanyl 6-*O*-acetyl-4-*O*-(3,4,6-tri-*O*-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl)-3-*O*-benzyl-2-deoxy-2-tetradecanamido- β -D-glucopyranoside (**8**) in 74% yield from **6**. Thus, the overall yield of **8** from **5** via acetolysis with $\text{BF}_3\text{-Et}_2\text{O}$ was 45%. At this stage, we noticed a paper reported by B. Fraser-Reid et al.⁸ about a mild acetolysis procedure of 1,6-anhydro sugars using triethylsilyl triflate as catalyst. Applying this catalyst to acetolysis of **5** at 0 °C \rightarrow room temperature showed that **5** underwent cleavage of its anhydro ring within 30 min, giving the oxazoline derivative **7** as the major product and the glycosyl acetate **6** as the minor product. After neutralization with triethylamine, the mixture was subjected to



treatment with 1-tetradecanol in the presence of (+)-10-camphorsulfonic acid and subsequent work-up to give **8** in 60% overall yield from **5**. The intermediate **6** was also obtained in 22% yield. Therefore, a one-pot synthesis employing triethylsilyl triflate proved to be a more efficient way for conversion of **5** into **8**.

Compound **8** then underwent step-wise deprotections. A successive treatment of **8** with sodium methoxide in methanol at room temperature and then with ethylenediamine in 1-butanol at 90 °C completed de-*O*-acetylation and de-*N*-phthaloylation, leaving the tetradecanamido group unaffected. The product, tetradecyl 4-*O*-(2-amino-2-deoxy-β-D-glucopyranosyl)-3-*O*-benzyl-2-deoxy-2-tetradecanamido-β-D-glucopyranoside (**9**), had extremely poor solubility in most solvents, making the ¹H NMR spectrum measurement difficult. Debenzoylation of **9** was successful by catalytic hydrogenation with palladium hydroxide in a large volume of acetic acid. The solubility of the resulting **1** with a free amino group at the 2'-position was also very poor in most solvents but its ¹H NMR spectrum could be recorded using a solution of CDCl₃-CD₃COOD-D₂O (3:3:1 v/v) as the NMR solvent. For purification of **1**, it was converted into its hydrochloride salt.

Hydrochloric acid was added in small portions to a suspension of **1** in chloroform-methanol, giving **hydrochloride salt of 1** in 51% yield, which was recrystallizable from a large quantity of methanol. This hydrochloride salt was essentially insoluble in protic, aprotic, and even dipolar aprotic solvents although slightly soluble in a chloroform-methanol mixture.



EXPERIMENTAL

General methods. Melting points were determined with a Laboratory Devices MEL-TEMP II apparatus and are uncorrected. Optical rotations were measured with a JASCO DIP-1000 polarimeter, using a 10-cm micro cell. IR spectra were recorded with a JASCO FT/IR-300E spectrophotometer. ^1H NMR spectra were recorded at 200 MHz or 400 MHz with Bruker AC-200 or Bruker AM-400 spectrometers for solutions in CDCl_3 unless otherwise specified. Column chromatography was performed on Silica Gel 60 (70-230 mesh, E. Merck) with the solvent systems specified. Solvent extracts were dried over anhyd Na_2SO_4 , and solutions were concentrated under diminished pressure below 50°C . Analytical samples were dried at $50\text{--}60^\circ\text{C}$ over P_2O_5 for 4-5 h *in vacuo*.

4-*O*-(3,4,6-Tri-*O*-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl)-1,6-anhydro-2-azido-3-*O*-benzyl-2-deoxy- β -D-glucopyranose (4). To a solution of compounds **2** (3.31 g, 11.94 mmol), **3** (5.26 g, 9.96 mmol), and *N*-iodosuccinimide (5.60 g, 24.91 mmol) in dichloromethane (80 mL) was added dropwise a dichloromethane solution (13.56 mL) saturated with trifluoromethanesulfonic acid at -8°C under an argon atmosphere in the presence of molecular sieves 4A (13 g). The reaction mixture was kept at room temperature for another 30 min, poured into saturated aqueous NaHCO_3 , and extracted with chloroform. The extract was successively washed with aqueous sodium thiosulfate, sodium hydrogencarbonate, and brine, dried, and concentrated. The resulting residue was chromatographed with toluene-ethyl acetate (5:1 v/v) as eluent to give **4** (6.7 g, 96%) as an amorphous powder: $[\alpha]_{\text{D}}^{22} +34.2^\circ$ (*c* 1.09, CHCl_3); IR (KBr) 2102 (N_3), 1749 (OAc), and 1716 cm^{-1} (NPhth); ^1H NMR (400 MHz)

δ 3.08 (bs, 1H, H-2), 3.69 (dd, 1H, $J =$, H-6a), 3.73 (bs, 1H, H-3), 3.80-3.84 (m, 2H, H-4, H-5'), 4.04 (bd, 1H, H-6b), 4.17 (dd, 1H, $J_{6'a,6'b}=12.3$ Hz, $J_{5',6'a}=2.3$ Hz, H-6'a), 4.26 (dd, 1H, $J_{5',6'b}=4.6$ Hz, H-6'b), 4.40 (dd, 1H, $J_{1',2'}=8.5$ Hz, $J_{2',3'}=10.7$ Hz, H-2'), 4.45 (bd, 1H, H-5), 4.61 (m, 2H, PhCH₂), 5.20 (t, 1H, $J_{3',4'}=J_{4',5'}=10.0$ Hz, H-4'), 5.26 (s, 1H, H-1), 5.70 (d, 1H, H-1'), 5.71 (dd, 1H, H-3').

Anal. Calcd for C₃₃H₃₄N₄O₁₃ (694.65): C, 57.06; H, 4.93; N, 8.07. Found: C, 56.86; H, 4.90; N, 7.86.

4-O-(3,4,6-Tri-O-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl)-1,6-anhydro-3-O-benzyl-2-deoxy-2-tetradecanamido- β -D-glucopyranose (5). Hydrogen sulfide gas was slowly bubbled for 30 min into a solution of **4** (2.00 g, 2.87 mmol) in pyridine-triethylamine (7:3 v/v, 10 mL) and the resulting mixture was kept at room temperature overnight in a capped flask. After concentration of the mixture, the residual product was chromatographed on silica gel with toluene-ethyl acetate (1:5 v/v) as eluent to give the intermediate 2-amino derivative (1.76 g, 92%), which showed no IR absorption around 2100 cm⁻¹ due to the azido group. The resulting 2-amino compound (1.66 g, 2.48 mmol) was dissolved in dry pyridine (10 mL) and the solution cooled in an ice bath. Tetradecanoyl chloride (0.95 mL, 3.49 mmol) was added to the solution and the mixture was stirred at ambient temperature for 30 min. After addition of methanol (1 mL), the mixture was diluted with chloroform, washed with water, dried, and concentrated. The residue was chromatographed with toluene-ethyl acetate (2:1 v/v) to give **5** (2.09 g, 96%): mp 64-66 °C; $[\alpha]_D^{27}$ -58.6° (*c* 3.36, CHCl₃); IR (KBr) 3421 (NH), 1753 (OAc), 1718 (NPhth), 1674 (NHCO), and 1506 cm⁻¹ (NHCO); ¹H NMR (200 MHz) δ 0.88 (m, 3H, CH₃), 1.27 (s, 22H, CH₂), 1.89, 2.04, 2.09 (s, each 3H, OAc), 2.27 (m, 2H, NHCOCH₂), 3.53-3.59 (m, 2H, H-3, H-6'a), 3.74-3.82 (m, 2H, H-4, H-5'), 4.05 (dd, 1H, $J_{5,6a}=2.1$ Hz, $J_{6a,6b}=12.4$ Hz, H-6a), 4.14-4.20 (m, 3H, H-2, H-5, H-6b), 4.33 (dd, 1H, $J_{5',6'b}=4.3$ Hz, $J_{6'a,6'b}=12.4$ Hz, H-6'b), 4.34 (dd, 1H, $J_{1',2'}=8.6$ Hz, $J_{2',3'}=10.6$ Hz, H-2'), 4.59 (m, 2H, PhCH₂), 4.99 (s, 1H, H-1), 5.15 (dd, 1H, $J_{3',4'}=9.1$ Hz, $J_{4',5'}=9.9$ Hz, H-4'), 5.37 (d, 1H, H-1'), 5.75 (bd, 1H, NH), 5.91 (dd, 1H, H-3').

Anal. Calcd for C₄₇H₆₂N₂O₁₄ (879.01): C, 64.22; H, 7.11; N, 3.19. Found: C, 64.43; H, 7.18; N, 3.13.

1,6-Di-O-acetyl-4-O-(3,4,6-tri-O-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl)-3-O-benzyl-2-deoxy-2-tetradecanamido- α -D-glucopyranose (6). *Method A* Trifluoroacetic acid (0.82 mL, 10.6 mmol) was added at 0 °C under an argon atmosphere to a solution of **5** (500 mg, 0.56 mmol) in acetic anhydride (20 mL). The mixture was kept at room temperature for 3 days and concentrated. The resulting

residue was chromatographed with toluene-ethyl acetate (4:1 v/v) to give **6** (240 mg, 43%); mp 169-170 °C; $[\alpha]_{\text{D}}^{25} +49^{\circ}$ (c 0.36, CHCl_3); IR (KBr) 1750 (OAc), 1714 (NPhth), 1651 (NHCO), and 1532 cm^{-1} (NHCO); $^1\text{H NMR}$ (200 MHz) δ 0.88 (t, 3H, CH_3), 1.25 (s, 22H, CH_2), 1.83-2.12 (m, 17H, NHCOCH_2 , OAc x 5), 3.60-3.72 (m, 4H), 3.91-4.04 (m, 2H), 4.14-4.30 (m, 3H), 4.33 (dd, 1H, $J_{1',2'}=8.4\text{ Hz}$, $J_{2',3'}=10.5\text{ Hz}$, H-2'), 4.76 (m, 2H, PhCH_2), 4.92 (bd, 1H, NH), 5.15 (dd, 1H, $J_{3',4'}=9.5\text{ Hz}$, $J_{4',5'}=9.8\text{ Hz}$, H-4'), 5.54 (d, 1H, H-1'), 5.76 (dd, 1H, H-3'), 6.05 (d, 1H, $J_{1,2}=3.5\text{ Hz}$, H-1).

Method B Boron trifluoride diethyl etherate (0.14 mL, 1.13 mmol) was added at 0 °C under an argon atmosphere to a solution of **5** (200 mg, 0.23 mmol) in acetic anhydride (2 mL) and the mixture was kept at 0 °C for 10 min. After addition of triethylamine (1 mL), the mixture was diluted with toluene and concentrated to dryness. The resulting residue was chromatographed as described in method A, giving **6** (136 mg, 61%).

Anal. Calcd for $\text{C}_{51}\text{H}_{68}\text{N}_2\text{O}_{17}$ (981.10): C, 62.44; H, 6.99; N, 2.86. Found: C, 62.25; H, 7.00; N, 2.84.

Tetradecanyl 6-O-Acetyl-4-O-(3,4,6-tri-O-acetyl-2-deoxy-2-phthalimido- β -D-glucopyranosyl)-3-O-benzyl-2-deoxy-2-tetradecanamido- β -D-glucopyranoside (8). **Method A (from 6)** Trimethylsilyl triflate (0.07 mL, 0.36 mmol) was added at 0 °C under an argon atmosphere to a solution of **6** (340 mg, 0.35 mmol) in 1,2-dichloroethane (5 mL) and the mixture was stirred at 50 °C for 1 h. After addition of triethylamine (0.5 mL), the mixture was placed on a short column of silica gel and eluted with toluene-ethyl acetate (3:1 v/v). The eluate was concentrated to give the crude oxazoline intermediate (**7**), $^1\text{H NMR}$ (400 MHz) δ 5.86 (d, 1H, $J_{1,2}=7.5\text{ Hz}$, H-1), which was immediately used for the next glycosidation reaction without further purification. A solution of **7**, 1-tetradecanol (1.11 g, 5.19 mmol), and (+)-10-camphorsulfonic acid (8 mg, 0.034 mmol) in 1,2-dichloroethane was heated at 90 °C for 30 min with stirring under an argon atmosphere and diluted with chloroform. The mixture was successively washed with aqueous sodium hydrogencarbonate and water, dried, and concentrated to dryness. The resulting residue was chromatographed with toluene-ethyl acetate (3:1 v/v), giving **8** (292 mg, 74% from **6**) which was recrystallized from methanol: mp 149-150 °C; $[\alpha]_{\text{D}}^{26} +1.2^{\circ}$ (c 1.15, CHCl_3); IR (KBr) 1747 (OAc), 1714 (NPhth), 1647 (NHCO), 1536 cm^{-1} (NHCO); $^1\text{H NMR}$ (400 MHz) δ 0.88 (m, 6H, CH_3), 1.25 (s, 46H, CH_2), 1.84, 1.95, 2.00, 2.01 (s, each 3H, OAc), 2.03-2.05 (m, 2H, NHCOCH_2), 3.24 (m, 1H, one of OCH_2), 3.42-3.48 (m, 2H, H-2, H-5), 3.63-3.71 (m, 2H, H-5', one of OCH_2), 3.81-3.87 (m, 2H, H-4, H-6a), 3.97 (dd, 1H, $J_{5',6'a}=ca\ 1\text{ Hz}$, $J_{6'a,6'b}=12.2\text{ Hz}$, H-6'a), 4.07 (t, 1H, $J_{2,3}=J_{3,4}=7.5\text{ Hz}$, H-3), 4.22 (dd, 1H, $J_{5',6'a}=4.4\text{ Hz}$, H-6'b), 4.25-4.31 (m, 2H, H-2', H-6b), 4.68 (d, 1H, $J_{1,2}=6.4\text{ Hz}$, H-1), 4.75 (m, 2H,

PhCH₂), 5.13 (dd, 1H, J_{3',4'}=9.4 Hz, J_{4',5'}=9.8 Hz, H-4'), 5.50 (d, 1H, J_{1',2'}=8.4 Hz, H-1'), 5.61 (bd, 1H, NH), 5.78 (dd, 1H, J_{2',3'}=10.6 Hz, H-3')

Anal. Calcd for C₆₃H₉₄N₂O₁₆ (1135.44): C, 66.64; H, 8.34; N, 2.47. Found: C, 66.45; H, 8.34; N, 2.46.

Method B (from 5) Triethylsilyl triflate (0.18 mL, 0.79 mmol) was added at 0 °C under argon atmosphere to a solution of **5** (583 mg, 0.66 mmol) in acetic anhydride (5 mL) and the mixture was kept at room temperature for 30 min. After addition of triethylamine (1 mL), the mixture was co-evaporated with toluene to give **7** as a major product and **6** as a minor one. The mixture was dissolved in 1,2-dichloroethane (10 mL) and treated with 1-tetradecanol (1.0 g, 4.66 mmol) and (+)-10-camphorsulfonic acid (10 mg, 0.043 mmol) as described in Method A. A similar subsequent work-up and chromatographic purification gave **8** (447 mg, 59% from **5**) and **6** (169 mg, 22% from **5**).

Tetradecanyl 4-O-(2-Amino-2-deoxy-β-D-glucopyranosyl)-3-O-benzyl-2-deoxy-2-tetradecanamido-β-D-glucopyranoside (9). To a suspension of **8** (531 mg, 0.467 mmol) in methanol (15 mL) was added methanolic sodium methoxide (28%, 0.03 mL) and the mixture was stirred at room temperature for 4.5 h under an argon atmosphere. The resulting clear solution was neutralized with Amberlite IR-120B (H⁺, 2 mL), filtered, and concentrated. The residue was suspended in a mixture of 1-butanol (15 mL) and ethylenediamine (1 mL) and the suspension was heated at 90 °C with stirring under an argon atmosphere. The mixture was concentrated and then co-evaporated with toluene to give **9**, which was recrystallized from a large volume of 2-propanol, yielding **9** (390 mg, 99%): mp 200-201 °C; [α]_D²⁸ -2.60° {c 0.17, CHCl₃-CH₃OH (1:1 v/v)}; IR (KBr): around 3283 (NH₂, OH), 1648 (NHCO), 1556 cm⁻¹ (NHCO); ¹H NMR spectrum could not be recorded due to the poor solubility of **9**.

Anal. Calcd for C₄₇H₈₄N₂O₁₀•0.5H₂O (846.20): C, 66.71; H, 10.12; N, 3.31. Found: C, 66.83; H, 10.11; N, 3.56.

Tetradecyl 4-O-(2-Amino-2-deoxy-β-D-glucopyranosyl)-2-deoxy-2-tetradecanamido-β-D-glucopyranoside (1) and its hydrochloride. A solution of **9** (326 mg, 0.38 mmol) in acetic acid (30 mL) was shaken in H₂ for 8 h with palladium hydroxide on carbon (Aldrich). After dilution with chloroform-methanol, the mixture was filtered and concentrated to give **1**: mp 200 °C (dec); IR (KBr) around 3269 (NH₂, OH), 1654 (NHCO), 1550 (NHCO); ¹H NMR (400 MHz, CDCl₃-CD₃COOD-D₂O) δ 0.88 (t, 6H, CH₃), 1.26 (s, 46H, CH₂), 2.02-2.07 (m, 2H, NHCOCH₂), 3.21-3.26 (m, 1H, H-2'), 3.40-3.92 (m, 13H), 4.68 (d, 1H, J_{1,2}=8.3 Hz, H-1), 4.96 (d, 1H, J_{1',2'}=8.4 Hz, H-1'). To a suspension of **1** in a large volume (ca. 50 mL) of chloroform-methanol (1:1) was added 0.5 M aqueous HCl in small portions until the mixture became a clear solution.

After evaporation of the solvents, the resulting residue was recrystallized from a large volume of methanol, giving **hydrochloride of 1** (153 mg, 51%): mp 176 °C (dec).

Anal. Calcd for $C_{40}H_{79}N_2O_{10}Cl$ (783.53): C, 61.32; H, 10.16; N, 3.58. Found: C, 61.21; H, 10.35; N, 3.51.

ACKNOWLEDGMENT

The authors express their thanks to Mr. Tetsuo Koyama of this department for technical help.

REFERENCES AND NOTES

1. R. A. A. Muzzarelli, C. Jeuniaux, and G. W. Gooday, *Chitin in Nature and Technology*; Plenum Publishing Corp., New York 1986.
2. K. Suzuki, T. Mikami, Y. Okawa, A. Tokoro, S. Suzuki, and M. Suzuki, *Carbohydr. Res.*, **151**, 403 (1986).
3. A. Yamada, N. Shibuya, O. Kodama, and T. Akatsuka, *Biosci., Biotechnol., Biochem.*, **57**, 405 (1993).
4. V. I. Gorbach, I. N. Krasikova, P. A. Luk'yanov, Y. N. Loenko, T. F. Solov'eva, Y. S. Ovodov, V. V. Deev, and A. A. Pimenov, *Carbohydr. Res.*, **260**, 73 (1994).
5. N. Sakairi, S. Takahashi, F. Wang, Y. Ueno, and H. Kuzuhara, *Bull. Chem. Soc. Jpn.*, **67**, 1756 (1994).
6. O. Kanie, Y. Ito, and T. Ogawa, *J. Am. Chem. Soc.*, **116**, 12073 (1994) and references cited therein.
7. The structure of this labile oxazoline derivative was confirmed by the NMR data described in EXPERIMENTAL and the behavior on TLC (Rf 0.70, toluene-ethyl acetate, 1:1 v/v).
8. M. Zottola, B. V. Rao, and B. Fraser-Reid, *J. Chem. Soc., Chem. Commun.*, 969 (1991).