

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

### DE-O-BENZYLATION OF STERICALLY HINDERED BENZYL ETHERS[1]

John G. Riley<sup>a</sup>; T. Bruce Grindley<sup>a</sup>

<sup>a</sup> Department of Chemistry, Dalhousie University, hali, Canada

Online publication date: 31 March 2001

**To cite this Article** Riley, John G. and Grindley, T. Bruce(2001) 'DE-O-BENZYLATION OF STERICALLY HINDERED BENZYL ETHERS[1]', *Journal of Carbohydrate Chemistry*, 20: 2, 159 – 169

**To link to this Article:** DOI: 10.1081/CAR-100103955

**URL:** <http://dx.doi.org/10.1081/CAR-100103955>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## DE-*O*-BENZYLATION OF STERICALLY HINDERED BENZYL ETHERS<sup>1</sup>

John G. Riley and T. Bruce Grindley\*

Department of Chemistry, Dalhousie University, Halifax, NS,  
Canada B3H 4J3

### ABSTRACT

Sterically hindered benzyl ethers that cannot be removed by hydrogenolysis with a variety of catalysts are removed readily by the reaction with *N*-bromosuccinimide and light in the presence of aqueous calcium carbonate, a reaction developed by Binkley and Hehemann. It was found that these conditions are compatible with the presence of phthalimides and glycosyl sulfides and fluorides, in addition to the groups previously shown to be inert.


### INTRODUCTION

Benzyl ethers are probably the most commonly used protecting groups for alcohols. The normal technique for deprotection is hydrogenolysis, either catalytic or by transfer of hydrogen.<sup>2–6</sup> Palladium on charcoal is the most commonly used catalyst but some benzyl groups resist hydrogenolysis with this catalyst and other catalysts, such as Pearlman's catalyst (Pd(OH)<sub>2</sub>), are often tried. Many other methods have been utilized, such as dissolving metal reductions,<sup>7–9</sup> various oxidative methods which convert benzyl ethers to benzoyl esters,<sup>10–15</sup> or cleavage with Lewis acids. However, few of these alternative methods are compatible with esters and/or thioglycosides or other sensitive functional groups.

The method discovered by Binkley and Hehemann,<sup>16</sup> which uses light-catalysed  $\alpha$ -bromination of benzyl ethers in the presence of aqueous base, has only been

---

\*Corresponding author.



*Scheme 1.*

used preparatively three times in the ten years since the initial publication.<sup>17-19</sup> We now report that this method is very effective for the removal of hindered benzyl ethers and also note that it is compatible with many sensitive functional groups, such as thioglycosides and glycosyl fluorides.

## RESULTS AND DISCUSSION


A key part of a synthetic sequence required the removal of a benzyl group on an oxygen atom flanked by two pivaloyloxy groups, that is, the benzyl group in compound **2a** in Scheme 1. Despite repeated attempts, this benzyl group could not be removed by hydrogenolysis at 4 atm using a variety of catalysts, including Pd/C, Pd(OH)<sub>2</sub> and Pd(OAc)<sub>2</sub>. Because most alternatives were not compatible with ester protecting groups, we tried photochemical  $\alpha$ -bromination,<sup>16,20</sup> and this method proved to be very effective.

Scheme 1 shows the de-*O*-benzylation of some substituted 3-*O*-benzyl-D-galactopyranoside derivatives. Removal of the *O*-benzyl groups proceeds in short times under mild conditions in excellent yields as summarized in the Table. The conditions of Binkley and Hehemann were used,<sup>16</sup> that is, irradiation with a 375 W white light lamp of a vigorously magnetically stirred two-phase system consisting

*Table 1.* Reaction Times and Yields in de-*O*-benzylation Reactions

Substrate	Reaction Time (min)	Yield (%)
<b>2a</b>	15	95
<b>3a</b>	15	87
<b>4a</b>	15	95
<b>5a</b>	30	76
<b>6a</b>	45	83
7a	15	72





Scheme 2.

of the compound and *N*-bromosuccinimide in carbon tetrachloride and aqueous calcium carbonate. It was necessary to purge the reaction mixtures thoroughly with nitrogen before irradiation to avoid the conversion of the benzyl ethers to benzoate esters, a process which has been noted as being a difficulty with this reaction.<sup>18</sup> Recrystallization of the *N*-bromosuccinimide resulted in improved yields. The inexpensive 375 W lamps, usually called heat lamps, were originally used by Binkley and Hehemann<sup>16,17</sup> and are available in speciality lighting stores. The reactions were monitored by TLC and stopped when the substrate had been consumed. Reaction times for the benzyl ether to disappear completely ranged from 15 to 45 min.

Scheme 2 shows the application of the reaction to a di-*O*-benzyl derivative.

As can be seen from the Table, this procedure is remarkably effective for the de-*O*-benzylation of sterically hindered benzyl ethers. Moreover, a number of protecting groups and reactive glycosidic groups that had not been studied by previous workers were shown to be unaffected; these include glycosyl thiophenyl groups and fluorides, and phthalimide protecting groups.

## EXPERIMENTAL

**General Methods.** <sup>1</sup>H and <sup>13</sup>C NMR spectra were run at 300 K on Bruker AC-250 or AMX-400 NMR instruments operating at 250.13 and 62.9 MHz, and 400.16 and 100 MHz, respectively. The samples were made up in concentrations between 10 and 15 mM in chloroform-*d* unless otherwise specified. Chemical shifts are given in parts per million (ppm)( $\pm 0.01$  ppm) relative to TMS (tetramethylsilane) in the case of <sup>1</sup>H NMR spectra, and to the central line of chloroform-*d* ( $\delta$  77.23) for the <sup>13</sup>C NMR spectra. Spectral assignments were made by first-order analysis and COSY and HETCOR experiments. Mass spectra were measured on a CEC 21-110B mass spectrometer using electron ionization (70.0 eV). The probe and source temperature was 143°C. Pyridine was dried by refluxing over calcium hydride before distillation. Dichloromethane was first dried with calcium chloride, then refluxed over calcium hydride for one hour, fractionally distilled and stored over molecular sieves (4 Å). *N,N*-Dimethylformamide was stored over activated molecular sieves (4 Å) for 72 h before distillation under reduced pressure from freshly activated molecular sieves (4 Å). Toluene was stored over calcium hydride for 3 h then distilled onto activated molecular sieves (4 Å). Benzyl chloride was dried with MgSO<sub>4</sub>, refluxed over calcium hydride and then fractionally distilled under reduced pressure, collecting the middle fraction and then storing over calcium hydride. *N*-Bromosuccinimide was recrystallized from water then dried for 12 h at room temperature under vacuum (1 torr). Thin-layer chromatography



(TLC) was performed on 0.25 mm thick Whatman G/UV silica gel on aluminum plates. Components were visualized by spraying with a 2% ceric sulfate solution in 1 M H<sub>2</sub>SO<sub>4</sub> followed by heating on a hot plate until discoloration occurred. Flash chromatography was performed on silica gel TLC standard grade (230–400 mesh) with mixed solvents of ethyl acetate: hexanes at volume ratios indicated. Irradiation was performed with a Phillips 375 W reflector lamp.

**Phenyl 3-*O*-Benzyl-1-thio-β-D-galactopyranoside (1a).** Phenyl 1-thio-β-D-galactopyranoside<sup>21</sup> (20.13 g, 0.0740 mol) and dibutyltin oxide (19.57 g, 0.0786 mol, 1 equiv) were refluxed for 12 h in toluene (500 mL) with azeotropic removal of water. The solvent was removed and the residue was dried at 0.1 torr for 1 h. Cesium fluoride (16.60 g, 0.109 mol, 1.45 equiv) was added followed by a solution of distilled benzyl bromide (15.82 g, 0.0925 mol, 1.0 equiv) in *N,N*-dimethylformamide (300 mL) and the reaction mixture was stirred until TLC (eluant, 4:1 ethyl acetate: acetone) indicated that the starting material had been completely consumed (3.5 h). Distilled water (2 mL) was added, the solvent evaporated and the residue was dissolved in ethanol and filtered through celite. The filtrate was reconcentrated to a dark brown syrup that was purified by flash chromatography using 1:1 ethyl acetate: hexanes as eluant. Crystallization (ethyl acetate) produced fine white needles (15.9 g, 60 %): mp 156–160°C, lit<sup>22</sup> 161–163°C; [α]<sub>D</sub> –8.4° (*c* 2.0, methanol), lit<sup>22</sup> [α]<sub>D</sub> –8.5°; <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ 3.36 (dd, 1H, J<sub>3,4</sub> = 3.36 Hz, J<sub>2,3</sub> = 9.16 Hz, H-3), 3.64 (td, 1H, J<sub>4,5</sub> = 0.92 Hz, J<sub>5,6</sub> = J<sub>5,6'</sub> = 5.95 Hz, H-5), 3.76 (s, 1H, OH-4), 3.77 (d, 2H, J<sub>5,6</sub> = 5.80 Hz, H-6,6'), 3.84 (t, 1H, J<sub>1,2</sub> = J<sub>2,3</sub> = 9.5 Hz, H-2), 4.22 (td, 1H, J<sub>4,5</sub> = 1.0 Hz, J<sub>3,4</sub> = 3.35 Hz, H-4), 4.42, 4.40 (2s, 2H, OH-2 and OH-3), 4.67 (d, 1H, J<sub>1,2</sub> = 9.77 Hz, H-1), 4.69, 4.79 (2d, 2H, J = –12.2 Hz, OCH<sub>2</sub>Ph), 7.22–7.60 (m, 10H, CH<sub>2</sub>Ph, SPh); <sup>13</sup>C NMR (acetone-*d*<sub>6</sub>): δ 61.8 (C-6), 66.4 (C-4), 69.3 (C-2), 71.6 (CH<sub>2</sub>Ph), 79.6 (C-5), 83.4 (C-3), 88.9 (C-1), 127.2–131.6 (Ph, SPh).

**Phenyl 3-*O*-Benzyl-2,4,6-tri-*O*-pivaloyl-1-thio-β-D-galactopyranoside (2a).** Compound **1a** (7.19 g, 0.0199 mol) was dissolved in dry pyridine (50 mL) and pivaloyl chloride (8.81 g, 9.0 mL, 0.0731 mol, 3.7 equiv) was added dropwise. The reaction was stirred at a bath temperature of 75°C for 12 h. The reaction mixture was poured into ice water (150 mL) that was then extracted with dichloromethane (150 mL). The organic layer was washed with 1 M hydrochloric acid (3 × 150 mL), saturated sodium hydrogen carbonate solutions (3 × 150 mL), and water (2 × 150 mL), then dried, and concentrated. The residue was applied to a flash chromatography column and eluted with ethyl acetate:hexanes 1:4. The title compound was obtained as colorless crystals (10.356 g, 0.0168 mol, 85 %), recrystallized from methanol-water: mp 101–2°C; [α]<sub>D</sub> +21.3° (*c* 2.0, dichloromethane); <sup>1</sup>H NMR: δ 1.14, 1.21, 1.22 (3s, 3 × 9H, 3 C(CH<sub>3</sub>)<sub>3</sub>), 3.64 (dd, 1H, J<sub>3,4</sub> = 3.3 Hz, J<sub>2,3</sub> = 9.3 Hz, H-3), 3.93 (br t, 1H, J<sub>5,6</sub> = J<sub>5,6'</sub> = 6.7 Hz, H-5), 4.16 (m, 2H, H-6,6'), 4.36, 4.67 (2d, 2H, J = –11.29 Hz, OCH<sub>2</sub>Ph), 4.68 (d, 1H, J<sub>1,2</sub> = 10.07 Hz, H-1), 5.16 (t, 1H, J<sub>1,2</sub> = J<sub>2,3</sub> = 9.8 Hz, H-2), 5.54 (br d, 1H, J<sub>3,4</sub> = 3.21 Hz, H-4), 7.2–7.6 (m, 10H, CH<sub>2</sub>Ph, SPh); <sup>13</sup>C NMR: δ 26.8, 26.9, 27.0 (3



C(CH<sub>3</sub>)<sub>3</sub>, 38.3, 38.5, 38.8 (3 C(CH<sub>3</sub>)<sub>3</sub>), 62.0 (C-6), 65.5 (C-4), 67.9 (C-2), 71.2 (CH<sub>2</sub>Ph), 74.6 (C-5), 78.2 (C-3), 86.1 (C-1), 127.4, 127.8, 127.9, 132.2, 132.8, 137.0 (Ph, SPh), 176.4, 177.2, 177.8 (3 OCO).

Anal. Calcd for C<sub>34</sub>H<sub>46</sub>O<sub>8</sub>S: C, 66.42; H, 7.54; S, 5.22. Found: C, 66.57; H, 7.52; S, 5.68.

### 3-O-Benzyl-2,4,6-tri-O-pivaloyl- $\alpha$ -D-galactopyranosyl Fluoride (3a).

Compound **2a** (0.5215 g, 0.848 mmol) was dissolved in dry dichloromethane (20 mL) and cooled with a calcium chloride ice bath that maintained the temperature at 0°C. Diethylaminosulfur trifluoride (0.155 mL, 1.27 mmol) was added dropwise to the stirred solution followed by *N*-bromosuccinimide (NBS) (0.196 g, 1.10 mmol). After 40 min, the reaction mixture was poured into a saturated solution of sodium hydrogen carbonate (50 mL). The aqueous mixture was extracted with ether (4  $\times$  25 mL) and the combined extracts were dried (MgSO<sub>4</sub>) and concentrated. The residue purified by flash chromatography on silica gel (eluant, ethyl acetate: hexanes 1:5) to give the title compound (R<sub>f</sub>, 0.94) as pure colorless crystals (0.3675 g, 88%) that were recrystallized from ether-hexanes: mp 103–104°C; [ $\alpha$ ]<sub>D</sub> +73.0° (*c* 0.9, dichloromethane); <sup>1</sup>H NMR:  $\delta$  1.21 (s, 3  $\times$  9H, 3 C(CH<sub>3</sub>)<sub>3</sub>), 3.99 (dd, 1H, J<sub>3,4</sub> = 3.20 Hz, J<sub>2,3</sub> = 10.20 Hz, H-3), 4.13 (d, 2H, H-6,6'), 4.34 (br t, 1H, J<sub>5,6</sub> = J<sub>5,6'</sub> = 6.71 Hz, H-5), 4.46, 4.71 (2d, 2H, J = 11.14 Hz, OCH<sub>2</sub>Ph), 5.16 (ddd, 1H, J<sub>1,2</sub> = 2.59 Hz, J<sub>2,3</sub> = 10.22 Hz, J<sub>2,F</sub> = 24.72 Hz, H-2), 5.64 (br d, 1H, J<sub>3,4</sub> = 2.51 Hz, H-4), 5.64, 5.86 (2d, 1H, J<sub>1,2</sub> = 2.59, J<sub>1,F</sub> = 53.86 Hz, H-1), 7.2–7.6 (m, 5H, CH<sub>2</sub>Ph); <sup>13</sup>C NMR:  $\delta$  27.21, 27.24, 27.34 (3 C(CH<sub>3</sub>)<sub>3</sub>), 39.0, 39.3 (3 C(CH<sub>3</sub>)<sub>3</sub>), 61.6 (C-6), 66.0 (C-4), 69.4 (d, <sup>2</sup>J<sub>C,F</sub> = 24.0 Hz, C-2), 69.4 (C-5), 71.9 (CH<sub>2</sub>Ph), 73.0 (C-3), 104.7 (d, <sup>1</sup>J<sub>C,F</sub> = 226.7 Hz, C-1), 127.2, 127.5, 127.8, 128.3, 129.1, 137.3 (Ph), 177.3, 177.9, 177.9 (3 OCO); <sup>19</sup>F NMR:  $\delta$  -150.52 (dd, J<sub>H1,F</sub> = 53.4 Hz, J<sub>H2,F</sub> = 25.1 Hz, F); EIMS *m/z*: 524 (1.0%, M<sup>+</sup>), 489 (0.5%, M<sup>+</sup> - HF - CH<sub>3</sub>), 439 (57%, M<sup>+</sup> - *t*-BuC=O).

Anal. Calcd for C<sub>28</sub>H<sub>41</sub>O<sub>8</sub>F: C, 64.10; H, 7.88. Found: C, 64.22; H, 7.71.

**Ethyl 6-(3-O-Benzyl-2,4,6-tri-O-pivaloyl- $\beta$ -D-galactopyranosyloxy)hexanoate (4a).** Compound **2a** (0.81 g, 1.317 mmol) was dissolved in dry dichloromethane (40 mL) under nitrogen in a glove bag and ethyl 6-hydroxyhexanoate (0.428 mL, 2.635 mmol) was added, followed by *N*-iodosuccinimide (0.341 g, 1.515 mmol), silver triflate (0.677 g, 2.635 mmol), and powdered 4Å molecular sieves. The reaction vessel was then stoppered with a rubber septum and covered with aluminum foil. After 3 h, the mixture was filtered over celite and the filtrate was washed with saturated sodium hydrogen carbonate solutions (3  $\times$  30 mL) and water (3  $\times$  30 mL). The organic layer was dried and concentrated to a residue that was applied to a flash chromatography column. Elution with ethyl acetate: hexanes 1:5 gave the title compound (R<sub>f</sub>, 0.43) as a colorless syrup (0.739 g, 84.4%): [ $\alpha$ ]<sub>D</sub> +18.0° (*c* 3.7, chloroform); <sup>1</sup>H NMR:  $\delta$  1.17, 1.20, 1.21 (3s, 3  $\times$  9H, 3 C(CH<sub>3</sub>)<sub>3</sub>), 1.24 (t, 3H, J = 7.0 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.29–1.41 (complex m, 2H, OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.5–1.7 (complex m, 4H, OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.28 (t, 2H, J = 7.3 Hz, OCOCH<sub>2</sub>), 3.42 (complex m, 1H, one OCH<sub>2</sub>CH<sub>2</sub>), 3.58 (dd, 1H, J<sub>3,4</sub> = 3.4 Hz, J<sub>2,3</sub> = 9.9 Hz,



H-3), 3.78–3.89 (complex m, 2H, one  $\text{OCH}_2\text{CH}_2$ , H-5), 4.05–4.22 (complex m, 4H,  $\text{OCH}_2\text{CH}_3$ , H-6,6'), 4.40 (d, 1H,  $J_{1,2} = 8.1$  Hz, H-1), 4.37, 4.68 (2d, 2H,  $J = -11.3$  Hz,  $\text{OCH}_2\text{Ph}$ ), 5.13 (dd, 1H,  $J_{1,2} = 8.2$  Hz,  $J_{2,3} = 10.1$  Hz, H-2), 5.50 (br d, 1H,  $J_{3,4} = 2.8$  Hz, H-4), 7.2–7.4 (m, 5H, Ph);  $^{13}\text{C}$  NMR:  $\delta$  14.4 ( $\text{OCH}_2\text{CH}_3$ ), 24.8 ( $\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$ ), 25.7 ( $\text{OCH}_2\text{CH}_2\text{CH}_2$ ), 27.1–27.7 (9  $\text{C}(\text{CH}_3)_3$ ), 29.4 ( $\text{OCH}_2\text{CH}_2$ ), 34.3 ( $\text{CH}_2\text{CO}_2\text{Et}$ ), 38.9, 38.9, 39.2 (3  $\text{COC}(\text{CH}_3)_3$ ), 60.4, ( $\text{OCH}_2\text{CH}_3$ ), 61.9 (C-6), 65.6 (C-4), 69.6 ( $\text{OCH}_2\text{CH}_2$ ), 70.3 (C-2), 71.1 (C-5), 71.6 ( $\text{CH}_2\text{Ph}$ ), 77.5 (C-3), 101.6 (C-1), 127.7, 127.8, 128.3, 128.3, 137.5 (Ph), 173.8 ( $\text{COCH}_2$ ), 176.9, 177.7, 178.1 ( $\text{COC}(\text{CH}_3)_3$ ); EIMS  $m/z$ : 664 (not observed,  $\text{M}^+$ ), 505 (2.4%,  $\text{M}^+$  -  $\text{O}(\text{CH}_2)_5\text{CO}_2\text{Et}$ ); HRMS for  $\text{C}_{28}\text{H}_{41}\text{O}_8$ : Calcd 505.2801. Found: 505.2806.

**Phenyl 3-*O*-Benzyl-2,4,6-tri-*O*-acetyl-1-thio- $\beta$ -D-galactopyranoside (5a).**

Acetic anhydride (470 mL) was added slowly to a stirred solution of compound **1a** (15.67 g, 0.04353 mol) in pyridine (470 mL) at  $0^\circ\text{C}$  and the solution was left for 12 h, then poured into an ice-cold saturated solution of sodium hydrogen carbonate (900 mL). The resulting mixture was extracted with dichloromethane ( $3 \times 500$  mL) and the combined extracts were washed with 0.8 M hydrochloric acid ( $3 \times 250$  mL), dried and concentrated to a dark yellow solid (25.69 g). Recrystallization (ethyl acetate: hexanes 1:3) yielded the title compound as colourless needles (19.05 g, 90.2%):  $[\alpha]_{\text{D}}^{25} 63.4^\circ$  ( $c$  3.07, chloroform); mp  $100\text{--}101^\circ\text{C}$ ;  $^1\text{H}$  NMR  $\delta$ : 2.06, 2.08, 2.16 (3s,  $3 \times 3\text{H}$ , 3  $\text{COCH}_3$ ), 3.57 (dd, 1H,  $J_{3,4} = 3.36$  Hz,  $J_{2,3} = 9.46$  Hz, H-3), 3.83 (td, 1H,  $J_{4,5} = 1.7$  Hz,  $J_{5,6} = J_{5,6'} = 6.26$  Hz, H-5), 4.17 (d, 2H,  $J_{5,6} = 6.41$  Hz, H-6,6'), 4.62 (d, 1H,  $J_{1,2} = 10.07$  Hz, H-1), 4.40, 4.68 (2d, 2H,  $J = -12.26$  Hz,  $\text{OCH}_2\text{Ph}$ ), 5.16 (t, 1H,  $J_{1,2} = J_{2,3} = 9.8$  Hz, H-2), 5.54 (dd, 1H,  $J_{4,5} = 1.07$  Hz,  $J_{3,4} = 3.36$  Hz, H-4), 7.23–7.55 (m, 10H,  $\text{CH}_2\text{Ph}$ , SPh);  $^{13}\text{C}$  NMR  $\delta$ : 20.8, 20.8, 21.0 (3  $\text{COCH}_3$ ), 62.3 (C-6), 66.0 (C-4), 68.8 (C-2), 71.3 ( $\text{CH}_2\text{Ph}$ ), 74.7 (C-5), 77.6 (C-3), 86.6 (C-1), 127.9, 127.8, 128.4, 128.8, 132.3, 133.0, 137.3 (Ph, SPh), 169.4, 170.4, 170.5 (CO); EIMS  $m/z$ : 488 (not observed,  $\text{M}^+$ ), 379 (60.6%,  $\text{M}^+$  - SPh); HRMS for  $\text{C}_{19}\text{H}_{23}\text{O}_8$ : Calcd 379.1393. Found: 379.1377.

Anal. Calcd for  $\text{C}_{25}\text{H}_{28}\text{O}_8\text{S}$ : C, 61.46; H, 5.78. Found: C, 61.50; H, 5.62.

**Ethyl 6-(2,4,6-Tri-*O*-acetyl-3-*O*-benzyl- $\beta$ -D-galactopyranosyloxy)hexanoate (6a).** Compound **5a** (21.28 g, 0.0441 mol) was dissolved in distilled, dry dichloromethane (200 mL) containing activated 3 $\text{\AA}$  powdered molecular sieves in a 2 necked round bottomed flask wrapped in aluminum foil. Ethyl 6-hydroxyhexanoate (14.78 g, 15.0 mL, 0.09222 mol, 2.0 equiv) was added, followed by *N*-iodosuccinimide (11.46 g, 0.0511 mol, 1.15 equiv), and then a solution of silver triflate (12.76 g, 0.04966 mol, 1.10 equiv) in 100 mL of dry toluene. The reaction mixture was stirred at room temperature under a stream of  $\text{N}_2$  (g) until starting material was consumed (1.5 h). The reaction mixture was filtered through celite, and the filtrate was washed with saturated sodium hydrogen carbonate solutions ( $2 \times 250$  mL), followed by water ( $2 \times 250$  mL). The combined aqueous layers were washed with dichloromethane ( $2 \times 250$  mL). The organic fractions were combined, dried and concentrated to a dark yellowish residue. Flash chromatography



on silica gel using 1:2 ethyl acetate: hexanes as eluant gave the title compound as a syrup (4.58 g, 19.5%):  $[\alpha]_D +36.6^\circ$  (*c* 4.18, chloroform);  $^1\text{H NMR}$  1.24 (t, 3H,  $J = 7.0$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 1.29–1.41 (complex m, 2H,  $\text{OCH}_2\text{CH}_2\text{CH}_2$ ), 1.5–1.7 (complex m, 4H,  $\text{OCH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.02, 2.06, 2.13 (3s,  $3 \times 3\text{H}$ , 3  $\text{COCH}_3$ ), 2.27 (t, 2H,  $J = 7.3$  Hz,  $\text{OCOCH}_2$ ), 3.45 (complex m, 1H, one  $\text{OCH}_2\text{CH}_2$ ), 3.54 (dd, 1H,  $J_{3,4} = 3.5$  Hz,  $J_{2,3} = 9.9$  Hz, H-3), 3.79 (t, 1H,  $J_{5,6} = J_{5,6'} = 6.76$  Hz, H-5), 3.82–3.94 (m, 1H, one  $\text{OCH}_2\text{CH}_2$ ), 4.11 (q, 2H,  $J = 7.0$  Hz,  $\text{OCH}_2\text{CH}_3$ ), 4.17 (AB part of ABX, 2H, H-6,6'), 4.35 (d, 1H,  $J_{1,2} = 8.2$  Hz, H-1), 4.40, 4.69 (2d, 2H,  $J = -12.2$  Hz,  $\text{OCH}_2\text{Ph}$ ), 5.11 (dd, 1H,  $J_{1,2} = 8.1$  Hz,  $J_{2,3} = 10.0$  Hz, H-2), 5.51 (br d, 1H,  $J_{3,4} = 3.4$  Hz, H-4), 7.2–7.4 (m, 5H, Ph);  $^{13}\text{C NMR}$   $\delta$ : 14.3 ( $\text{OCH}_2\text{CH}_3$ ), 20.7, 20.8, 20.8 (3  $\text{COCH}_3$ ), 24.6 ( $\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$ ), 25.4 ( $\text{OCH}_2\text{CH}_2\text{CH}_2$ ), 29.1 ( $\text{OCH}_2\text{CH}_2$ ), 34.2 ( $\text{CH}_2\text{CO}_2\text{Et}$ ), 60.3 ( $\text{OCH}_2\text{CH}_3$ ), 62.0 (C-6), 65.9 (C-4), 69.6 ( $\text{OCH}_2\text{CH}_2$ ), 70.5 (C-2), 70.9 (C-5), 71.3 ( $\text{OCH}_2\text{Ph}$ ), 76.6 (C-3), 101.4 (C-1), 127.8, 127.9, 128.4, 137.6 (Ph), 169.4, 170.5, 170.6 ( $\text{COCH}_3$ ), 173.6 ( $\text{COCH}_2$ ); EIMS  $m/z$ : 538 (not observed,  $\text{M}^+$ ), 379 (11.6%,  $\text{M}^+ - \text{O}(\text{CH}_2)_5\text{CO}_2\text{Et}$ ); HRMS for  $\text{C}_{19}\text{H}_{23}\text{O}_8$ : Calcd 379.1393. Found: 379.1369.

**Phenyl 6-*O*-*tert*-Butyldimethylsilyl-2-deoxy-2-phthalimido-1-thio- $\beta$ -D-glucopyranoside (7b).** *Tert*-butyldimethylsilyl chloride (0.903 g, 5.99 mmol) in dry pyridine (15 mL) was added dropwise (3 mL / 20 min) via a syringe pipet to a stirred solution of phenyl 2-deoxy-2-phthalimido-1-thio- $\beta$ -D-glucopyranoside<sup>23</sup> (2.004 g, 4.99 mmol) in dry pyridine (30 mL). The reaction mixture was stirred 12 h, then poured into water (200 mL). The mixture was extracted with diethyl ether ( $5 \times 40$  mL). The combined extracts were dried ( $\text{MgSO}_4$ ), concentrated, and the residue purified by flash chromatography on silica gel (eluant ethyl acetate: hexanes 2:1) to give the title compound ( $R_f$ , 0.66) as a colorless syrup (1.402 g, 70.0%), crystallized from ethanol-water to give very fine colorless needles: mp 68–70°C;  $[\alpha]_D +26.1^\circ$  (*c* 2.1, dichloromethane);  $^1\text{H NMR}$   $\delta$  0.11 (d,  $2 \times 3\text{H}$ ,  $\text{Si}(\text{CH}_3)_2$ ), 0.91 (d, 9H,  $\text{SiC}(\text{CH}_3)_3$ ), 3.55–3.65 (br m, 2H, H-4, H-5), 3.88, 3.97 (AB part of ABX pattern, 2H,  $J_{6,6'} = -10.5$  Hz,  $J_{5,6} = 5.1$  Hz,  $J_{5,6'} = 4.5$  Hz, H-6,6'), 4.18 (t, 1H,  $J_{1,2} = J_{2,3} = 10.38$  Hz, H-2), 4.35 (dd, 1H,  $J_{2,3} = 10.24$  Hz,  $J_{3,4} = 7.94$  Hz, H-3), 5.60 (d, 1H,  $J_{1,2} = 10.23$  Hz, H-1), 7.2–7.6 (4m, 9H, SPh, Phth):  $^{13}\text{C NMR}$ :  $\delta$  -5.4 ( $\text{SiC}(\text{CH}_3)_3$ ), 18.2 ( $\text{SiC}(\text{CH}_3)_3$ ), 25.8 ( $\text{SiC}(\text{CH}_3)_3$ ), 55.2 (C2), 64.7 (C6), 72.7, 74.3, 78.0 (C3, C4, C5), 83.6 (C1), 127.8, 128.9, 131.6, 132.3, 134.2 (SPh, Phth), 210.6 (2 CO, Phth). EIMS  $m/z$ : 515 ( $\text{M}^+$ , not observed), 406 (10%, M - SPh), 388 (72%, 406 -  $\text{H}_2\text{O}$ ).

Anal. Calcd for  $\text{C}_{26}\text{H}_{33}\text{NO}_6\text{SSi.H}_2\text{O}$ : C, 58.51; H, 6.61; N, 2.62. Found: C, 58.69; H, 6.54; N, 2.55.

**Phenyl 6-*O*-*tert*-Butyldimethylsilyl-2-phthalimido-3,4-di-*O*-benzyl-2-deoxy-1-thio- $\beta$ -D-glucopyranoside (7a).** Compound 7b (0.100 g, 0.194 mmol) was dissolved in dry benzyl chloride (10 mL, 71.8 mmol). Sodium hydride (60% in mineral oil, 0.017 g, 0.423 mmol) was then added and the reaction was allowed to proceed under nitrogen at 70°C for 12 h. The reaction mixture was then poured into water (20 mL) and extracted with dichloromethane ( $3 \times 30$  mL). The com-





bined organic extracts were dried ( $\text{MgSO}_4$ ), concentrated, and the residue purified by flash chromatography on silica gel (eluant ethyl acetate: hexanes 1:2) to give the title compound ( $R_f$ , 0.65) as a colorless syrup (0.068 g, 50%):  $[\alpha]_D +22.3^\circ$  ( $c$  0.9, dichloromethane);  $^1\text{H NMR}$ :  $\delta$  0.11 (d,  $2 \times 3\text{H}$ ,  $\text{Si}(\text{CH}_3)_2$ ), 0.91 (d, 9H,  $\text{SiC}(\text{CH}_3)_3$ ), 3.52 (broad dt, 1H,  $J_{4,5} = 10.38$ ,  $J_{5,6} = J_{5,6'} = 2.59$  Hz, H-5), 3.78 (broad t, 1H,  $J_{3,4} = J_{4,5} = 8.54$  Hz, H-4), 3.92 (broad s, 2H, H-6,6'), 5.16 (t, 1H,  $J_{1,2} = J_{2,3} = 10.22$  Hz, H-2), 4.35 (t, 1H,  $J_{3,4} = J_{2,3} = 8.85$  Hz, H-3), 4.43, 4.78 (2d, 2H,  $J = 12.05$  Hz,  $\text{OCH}_2\text{Ph}$ ), 4.77, 4.86 (2d, 2H,  $J = 10.99$  Hz,  $\text{OCH}_2\text{Ph}$ ), 5.51 (d, 1H,  $J_{1,2} = 10.23$  Hz, H-1), 7.2–7.6 (4m, 19H,  $2\text{CH}_2\text{Ph}$ ,  $\text{SPh}$ ,  $\text{Npth}$ ):  $^{13}\text{C NMR}$ :  $\delta$  -4.9 ( $\text{SiC}(\text{CH}_3)_3$ ), 26.0 ( $\text{SiC}(\text{CH}_3)_3$ ), 29.7 ( $\text{SiC}(\text{CH}_3)_3$ ), 54.9 (C2), 61.5 (C3), 62.1 (C6), 72.1, 75.0 (2  $\text{OCH}_2\text{Ph}$ ), 79.0 (C4), 80.2 (C5), 83.1 (C1), 127.4, 127.8, 128.0, 128.4, 128.5, 128.8, 132.7, (2  $\text{CH}_2\text{Ph}$ ,  $\text{SPh}$ ,  $\text{Npth}$ ), 138.2, 138.4 (2 CO,  $\text{Npth}$ ).

**General Method for De-*O*-benzylation: Phenyl 2,4,6-Tri-*O*-pivaloyl-1-thio- $\beta$ -D-galactopyranoside (2b).** Compound **2a** (0.100 g, 0.16 mmol) was dissolved in carbon tetrachloride (10 mL) and water (5 mL). NBS (2.5 equiv, 0.049 g, 0.276 mmol) was then added, followed by calcium carbonate (4.4 equiv, 0.072 g, 0.716 mmol). The solution was then purged with nitrogen for one hour. Using a 375 W incandescent light, the reaction mixture was irradiated for 15 min while being monitored by TLC. The mixture was then poured into water (25 mL) and extracted with dichloromethane ( $3 \times 30$  mL). The combined organic extracts were dried ( $\text{MgSO}_4$ ), concentrated, and the residue was purified by flash chromatography on silica gel using ethyl acetate: hexanes 1:5 as eluant, to give the title compound (0.081 g, 95%) as a colorless syrup ( $R_f$ , 0.23). The product was crystallized from hexane to give fine colorless needles: mp 105–106°C;  $[\alpha]_D -5.0^\circ$  ( $c$  1.2, chloroform);  $^1\text{H NMR}$ :  $\delta$  1.27, 1.20 (2s,  $3 \times 9\text{H}$ ,  $3 \text{C}(\text{CH}_3)_3$ ), 3.91 (m, 2H, H-3, H5), 4.12 (m, 2H, H-6,6'), 4.70 (d, 1H,  $J_{1,2} = 10.07$  Hz, H-1), 4.92 (t, 1H,  $J_{1,2} = J_{2,3} = 9.76$  Hz, H-2), 5.32 (br dd, 1H,  $J_{3,4} = 3.35$  Hz,  $J_{4,5} = 0.9$ , H-4), 7.2–7.6 (m, 5H,  $\text{SPh}$ );  $^{13}\text{C NMR}$ :  $\delta$  27.1 (3  $\text{C}(\text{CH}_3)_3$ ), 38.7, 38.9, 39.2 (3  $\text{C}(\text{CH}_3)_3$ ), 62.1 (C-6), 69.5 (C-4), 70.2 (C-2), 73.0, 74.9 (C-3, C-5), 8.6 (C-1), 128.3, 128.9, 131.9, 133.2, ( $\text{SPh}$ ), 178.0, 178.1, 178.6 (3 OCO); EIMS  $m/z$ : 524 (0.1%,  $\text{M}^+$ ), 509 (0.1%, M-Me), 415 (100%, M-SPh); HRMS for  $\text{C}_{27}\text{H}_{40}\text{O}_8\text{S}$ : Calcd 524.2444. Found: 524.2437.

**2,4,6-Tri-*O*-pivaloyl- $\beta$ -D-galactopyranosyl Fluoride (3b).** Compound **3a** (0.229 g, 0.56 mmol) was reacted as outlined for compound **2a** to give the title compound (0.187 g, 87%) as a colorless syrup:  $R_f$ , 0.50 (ethyl acetate: hexanes 1:5);  $[\alpha]_D +52.5^\circ$  ( $c$  3.9, dichloromethane);  $^1\text{H NMR}$ :  $\delta$  1.20, 1.28, 1.29 (3s,  $3 \times 9\text{H}$ ,  $3 \text{C}(\text{CH}_3)_3$ ), 4.11 (d, 2H, H-6,6'), 4.28 (dd, 1H,  $J_{3,4} = 3.36$  Hz,  $J_{2,3} = 10.37$  Hz, H-3), 4.37 (br t, 1H,  $J_{5,6} = J_{5,6'} = 6.72$  Hz, H-5), 4.98 (ddd, 1H,  $J_{1,2} = 2.60$  Hz,  $J_{2,3} = 10.38$  Hz,  $J_{2,F} = 24.42$  Hz, H-2), 5.44 (br d, 1H,  $J_{3,4} = 3.36$  Hz, H-4), 5.63, 5.84 (2d, 1H,  $J_{1,2} = 2.89$ ,  $J_{1,F} = 53.86$  Hz, H-1);  $^{13}\text{C NMR}$ :  $\delta$  27.1, 27.2, 27.4 (3  $\text{C}(\text{CH}_3)_3$ ), 38.9, 39.2, 39.5 (3  $\text{C}(\text{CH}_3)_3$ ), 61.4 (C-6), 67.0 (C-3), 69.5 (2C, C-4, C-5), 70.5 (d,  $^2J_{C,F} = 23.8$  Hz, C-2), 104.8 (d,  $^1J_{C,F} = 227.7$  Hz, C-1), 178.1, 178.4, 178.9 (3 OCO); EIMS  $m/z$ : 434 (not observed,  $\text{M}^+$ ), 414 (7.6%,  $\text{M}^+ - \text{HF}$ ), 399



(1.5%,  $M^+$  - HF -  $CH_3$ ), 333 (2.8 %,  $M^+$  -  $OCOtBu$ ); HRMS for  $C_{21}H_{34}O_8$ : Calcd 414.2253. Found: 414.2245.

**Ethyl 6-(2,4,6-Tri-*O*-pivaloyl- $\beta$ -D-galactopyranosyloxy)hexanoate (4b).**

Compound **4a** (0.068 g, 0.10 mmol) was reacted as outlined for compound **2a** to give the title compound (0.057 g, 95%) as a colorless syrup:  $[\alpha]_D +3.2^\circ$  (*c* 6.1, dichloromethane);  $^1H$  NMR:  $\beta$  1.20, 1.22, 1.27 (3s,  $3 \times 9H$ , 3  $C(CH_3)_3$ ), 1.25 (t, 3H,  $J = 6.8$  Hz,  $OCH_2CH_3$ ), 1.29–1.41 (complex m, 2H,  $OCH_2CH_2CH_2$ ), 1.5–1.8 (complex m, 4H,  $OCH_2CH_2CH_2CH_2$ ), 2.30 (t, 2H,  $J = 7.3$  Hz,  $OCOCH_2$ ), 3.41–3.50 (complex m, 1H, one  $OCH_2CH_2$ ), 3.80–3.92 (m, 3H, one  $OCH_2CH_2$ , H-3, H-5), 4.10 (q, 2H,  $J = 7.0$  Hz,  $OCH_2CH_3$ ), 4.12 (AB part of ABX, 2H, H-6,6'), 4.46 (d, 1H,  $J_{1,2} = 8.1$  Hz, H-1), 4.90 (dd, 1H,  $J_{1,2} = 7.9$  Hz,  $J_{2,3} = 10.1$  Hz, H-2), 5.30 (br d, 1H,  $J_{3,4} = 3.5$  Hz, H-4);  $^{13}C$  NMR:  $\delta$  14.3 ( $OCH_2CH_3$ ), 21.1 ( $CH_2CH_2COOEt$ ), 24.7 ( $OCH_2CH_2CH_2$ ), 25.6 ( $OCH_2CH_2$ ), 27.2–27.3 (9  $C(CH_3)_3$ ) 29.4 ( $CH_2COOEt$ ), 39.8, 39.0, 39.4 (3  $C(CH_3)_3$ ) 60.3 ( $OCH_2CH_3$ ), 61.9 (C-6), 69.4, 69.7 (C-4,  $OCH_2CH_2$ ), 71.1, 72.1 (C-3, C-5), 72.9 (C-2), 101.1 (C-1), 173.7 ( $COCH_2$ ), 178.0, 178.2, 179.2 ( $COC(CH_3)_3$ ); EIMS  $m/z$ : 574 (not observed,  $M^+$ ), 415 (5.3%,  $M^+$  -  $O(CH_2)_5CO_2Et$ ); HRMS for  $C_{21}H_{35}O_8$ : Calcd 415.2332. Found: 415.2333.

**Phenyl 2,4,6-Tri-*O*-acetyl-1-thio- $\beta$ -D-galactopyranoside (5b).**

Compound **5a** (0.197 g, 0.404 mmol) was reacted as outlined for compound **2a** to give the title compound (0.122 g, 76%) as a colorless syrup, crystallized from hexanes: ethyl acetate to give fine colorless needles: mp 120–121°C;  $R_f$  0.21 (ethyl acetate: hexanes 2:3);  $[\alpha]_D +14.6^\circ$  (*c* 1.0, chloroform);  $^1H$  NMR:  $\delta$  2.05, 2.15, 2.73 (3s,  $3 \times 3H$ , 3  $COCH_3$ ), 3.89 (m, 2H, H-3, H-5), 4.13, 4.16 (AB part of ABX pattern, 2H,  $J_{5,6} = 7.4$  Hz,  $J_{5,6'} = 5.5$  Hz,  $J_{6,6'} = -11.5$  Hz, H-6,6'), 4.68 (d, 1H,  $J_{1,2} = 9.92$  Hz, H-1), 5.04 (t, 1H,  $J_{1,2} = J_{2,3} = 9.77$  Hz, H-2), 5.36 (dd, 1H,  $J_{3,4} = 3.53$  Hz,  $J_{4,5} = 0.94$  Hz, H-4), 7.20–8.10 (3m, 6H, SPh);  $^{13}C$  NMR:  $\delta$  20.8, 21.2, 29.7 (3  $COCH_3$ ), 62.4 (C-6), 70.1 (C-4), 70.8 (C-2), 72.1, 74.9 (C-5, C-3), 86.3 (d, C-1), 128.1, 129.0, 129.0, 130.2, 132.4, 132.4 (SPh), 171.2, 171.2, 178.5 (3  $OCO$ ); EIMS  $m/z$ : 398 (0.5%,  $M^+$ ), 289 (100,  $M^+$  - SPh); HRMS for  $C_{18}H_{22}O_8S$ : Calcd 398.1035. Found: 398.1054.

**Ethyl 6-(2,4,6-Tri-*O*-acetyl- $\beta$ -D-galactopyranosyloxy)hexanoate (6b).**

Compound **6a** (2.00 g, 3.43 mmol) was reacted as outlined for compound **2a** to give the title compound (1.26g, 83%) as a colorless syrup:  $[\alpha]_D +7.7^\circ$  (*c* 1.3, dichloromethane);  $^1H$  NMR:  $\delta$  1.25 (t, 3H,  $J = 6.8$  Hz,  $OCH_2CH_3$ ), 1.29–1.41 (complex m, 2H,  $OCH_2CH_2CH_2$ ), 1.5–1.8 (complex m, 4H,  $OCH_2CH_2CH_2CH_2$ ), 2.02, 2.04, 2.13 (3s,  $3 \times 3H$ , 3  $COCH_3$ ), 2.29 (t, 2H,  $J = 7.3$  Hz,  $OCOCH_2$ ), 3.48 (X of ABXY, 1H,  $OCH_2CH_2$ ), 3.79–3.93 (complex m, 3H,  $OCH_2CH_2$ , H-3, H-5), 4.12 (q, 2H,  $J = 7.2$  Hz,  $OCH_2CH_3$ ), 4.15 (AB part of ABX, 2H, H-6,6'), 4.41 (d, 1H,  $J_{1,2} = 7.9$  Hz, H-1), 4.97 (dd, 1H,  $J_{1,2} = 8.4$  Hz,  $J_{2,3} = 9.8$  Hz, H-2), 5.33 (br d, 1H,  $J_{3,4} = J_{4,5} = 3.2$ , H-4);  $^{13}C$  NMR:  $\delta$  14.4 ( $OCH_2CH_3$ ), 20.8, 20.9, 21.1 (3  $COCH_3$ ), 24.7 ( $CH_2CH_2CO_2Et$ ), 25.5 ( $OCH_2CH_2CH_2$ ), 29.2 ( $OCH_2CH_2$ ), 34.23



(CH<sub>2</sub>CO<sub>2</sub>Et), 60.4 (OCH<sub>2</sub>CH<sub>3</sub>), 62.1 (C-6), 69.9 (C-4, OCH<sub>2</sub>CH<sub>2</sub>), 71.0, 71.4 (C-3, C-5), 72.8 (C-2), 101.1 (C-1), 170.7, 171.1, 171.1 (COCH<sub>3</sub>), 173.6 (COCH<sub>2</sub>); EIMS *m/z*: 448 (not observed, M<sup>+</sup>), 403 (0.9%, M<sup>+</sup> - OEt), 289 (14.8%, M<sup>+</sup> - O(CH<sub>2</sub>)<sub>5</sub>CO<sub>2</sub>Et); HRMS for C<sub>18</sub>H<sub>27</sub>O<sub>10</sub> and C<sub>12</sub>H<sub>17</sub>O<sub>8</sub>: Calcd 403.1604 and 289.0923. Found: 403.1633 and 289.0987, respectively.

**Phenyl 6-*O*-*tert*-Butyldimethylsilyl-2-deoxy-2-*N*-phthalimido-1-thio-β-D-glucopyranoside (7b).** Compound **7a** (0.095 g, 0.137 mmol) was reacted as outlined for compound **2a** to give the title compound (0.51 g, 72.4%) as a colorless syrup, with R<sub>f</sub> and spectral characteristics identical to **7b**, prepared above.

### ACKNOWLEDGMENTS

We thank Drs. L. R. Ramaley and J. S. Grossert for accurate mass determinations and D. Johnston and M. Safatli for preparation of some starting materials. We are grateful for financial support from NSERC.

### REFERENCES

1. Presented at the 20<sup>th</sup> International Carbohydrate Symposium, Hamburg, Germany, August 27–September 1, 2000.
2. Wuts, P.G.M.; Greene, T.W. *Protective Groups in Organic Synthesis*, 3rd; Wiley: New York, 1999.
3. Kocienski, P.J. *Protecting Groups*; Thieme: Stuttgart, 1994.
4. Grindley, T.B. Protecting Groups; In *Modern Methods in Carbohydrate Synthesis*; Khan, S.H., O'Neill, R.A., Eds. Harwood Academic Publishers: Amsterdam, 1995.
5. Ziegler, T. Protecting Group Strategies for Carbohydrates; In *Carbohydrate Chemistry*; Boons, G.-J., Ed. Blackie: London, 1998; 21–45.
6. Jarowicki, K.; Kocienski, P. Protecting Groups. *J. Chem. Soc., Perkin Trans. 1* **2000**, 2495–2527.
7. McCloskey, C.M. Benzyl Ethers of Carbohydrates. *Adv. Carbohydr. Chem.* **1957**, *12*, 137–156.
8. Holick, S.A.; Anderson, L. On the Debenzylation of *O*-Benzyl-protected Aralkyl Thioglucosides with Sodium-liquid Ammonia. *Carbohydr. Res.* **1974**, *34*, 208–213.
9. Hwu, J.R.; Chua, V.; Schroeder, J.E.; Barrans, R.E., Jr.; Khoudary, K.P.; Wang, N.; Wetzal, J.M. Calcium in Liquid Ammonia for the Reduction of Benzyl Ethers. Mechanistic Clues Derived from Chemoselectivity Studies. *J. Org. Chem.* **1986**, *51*, 4733–4734.
10. Angyal, S.J.; James, K. Oxidative Demethylation with Chromium Trioxide in Acetic Acid. *Carbohydr. Res.* **1970**, *12*, 147–149.
11. Schmidt, W.; Steckhan, E. Oxidative Removal of the Benzyl Ether Protecting Group by Homogeneous Electron Transfer. *Angew. Chem. Int. Ed.* **1979**, *18*, 801–802.
12. Schuda, P.F.; Cichowicz, M.B.; Heimann, M.R. A Facile Method for the Oxidative Removal of Benzyl Ethers: the Oxidation of Benzyl Ethers to Benzoates by Ruthenium Tetraoxide. *Tetrahedron Lett.* **1983**, *24*, 3829–3830.



13. Csuk, R.; Dorr, P. Convenient Oxidative Debenzylation Using Dimethyldioxirane. *Tetrahedron* **1994**, *50* (33), 9983–9988.
14. Angibeault, P.; Defaye, J.; Gadelle, A.; Utile, J.-P. Mild Deprotection of Benzyl Ether Protective Groups with Ozone. *Synthesis* **1999**, 1123–1125.
15. Adinolfi, M.; Barone, G.; Guariniello, L.; Iadonisi, A. Facile Cleavage of Carbohydrate Benzyl Ethers and Benzylidene Acetals Using the NaBrO<sub>3</sub>/Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> Reagent Under Two-phase Conditions. *Tetrahedron Lett.* **1999**, *40* (48), 8439–8441.
16. Binkley, R.W.; Hehemann, D.G. A Light-Initiated Process for Rapid Debenzylation of Carbohydrates. *J. Org. Chem.* **1990**, *55*, 378–380.
17. Binkley, R.W. Aglycon-Disaccharide Coupling in Mithramycin Analog Synthesis. *J. Carbohydr. Chem.* **1990**, *9*, 507–511.
18. Giuliano, R.M.; Villani, F.J. Stereoselectivity of Addition of Organometallic Reagents to Pentodialdo-1,4-furanoses - Synthesis of L-Axenose and D- Evermicose from a Common Intermediate. *J. Org. Chem.* **1995**, *60* (1), 202–211.
19. Maryanoff, B.E.; Costanzo, M.J.; Nortey, S.O.; Greco, M.N.; Shank, R.P.; Schupsky, J.J.; Ortegon, M.P.; Vaught, J.L. Structure-activity Studies on Anticonvulsant Sugar Sulfamates Related to Topiramate. Enhanced Potency with Cyclic Sulfate Derivatives. *J. Med. Chem.* **1998**, *41* (8), 1315–1343.
20. Hashimoto, H.; Kawa, M.; Saito, Y.; Date, T.; Horito, S.; Yoshimura, J. A Convenient One-pot Preparation of Glycosyl Bromides as well as Glycosyl Acetates from Benzyl Glycosides using *N*-Bromosuccinimide. *Tetrahedron Lett.* **1987**, *28*, 3505–3508.
21. Janaki, N.; Patil, J.R.; Bose, J.L. Synthesis of Some Aryl  $\beta$ -Thioglycosides. *Ind. J. Chem.* **1969**, *7*, 227–228.
22. Malet, C.; Hindsgaul, O. Generation of Molecular Diversity on *N*-Acetyllactosamine via *O*-Cyanomethyl Ethers. *Carbohydr. Res.* **1997**, *303*, 51–65.
23. Ogawa, T.; Nakabayashi, S.; Sasajima, K. Synthesis of Phenyl 6-*O*-Acyl-3-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- $\beta$ -D-glucopyranoside. *Carbohydr. Res.* **1981**, *95*, 308–312.

Accepted December 6, 2000



## Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

**[Order now!](#)**

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081CAR100103955>