

This article was downloaded by:

On: 23 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

### Synthesis of Tetra-*O*-acetyl-1-thio- $\alpha$ -d-glucopyranose by Reaction of Tetra-*O*-acetyl- $\alpha$ -d-glucopyranosyl Bromide with *N,N*-Dimethylthioformamide

Takayoshi Fujihira<sup>a</sup>; Gen Arakawa<sup>b</sup>; Haruo Kamijo<sup>b</sup>; Toshio Takido<sup>b</sup>; Manabu Seno<sup>b</sup>

<sup>a</sup> Technical Department and Research Laboratory, Japan Sugar Refiners' Association, Chiyoda-ku, Japan <sup>b</sup> Department of Materials Applied Chemistry, College of Science and Technology, Nihon University, Chiyoda-ku, Japan

Online publication date: 04 August 2003

**To cite this Article** Fujihira, Takayoshi , Arakawa, Gen , Kamijo, Haruo , Takido, Toshio and Seno, Manabu(2003) 'Synthesis of Tetra-*O*-acetyl-1-thio- $\alpha$ -d-glucopyranose by Reaction of Tetra-*O*-acetyl- $\alpha$ -d-glucopyranosyl Bromide with *N,N*-Dimethylthioformamide', Journal of Carbohydrate Chemistry, 22: 2, 73 – 78

**To link to this Article:** DOI: 10.1081/CAR-120020478

**URL:** <http://dx.doi.org/10.1081/CAR-120020478>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Synthesis of Tetra-*O*-acetyl-1-thio- $\alpha$ -D-glucopyranose by Reaction of Tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl Bromide with *N,N*-Dimethylthioformamide

Takayoshi Fujihira,<sup>1,\*</sup> Gen Arakawa,<sup>2</sup> Haruo Kamijo,<sup>2</sup>  
Toshio Takido,<sup>2</sup> and Manabu Seno<sup>2</sup>

<sup>1</sup>Technical Department and Research Laboratory, Japan Sugar  
Refiners' Association, Tokyo, Japan

<sup>2</sup>Department of Materials Applied Chemistry, College of Science  
and Technology, Nihon University, Tokyo, Japan

### ABSTRACT

A reaction system was found to prepare tetra-*O*-acetyl-1-thio-D-glycopyranose in both  $\alpha$  and  $\beta$ -forms. Methanolysis of the adduct prepared from the reaction of tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with *N,N*-dimethylthioformamide afforded the corresponding tetra-*O*-acetyl-1-thio-D-glucopyranose with an anomer ratio  $\alpha/\beta$  of 52:48 in 98% yield. The anomer mixture was easily separated by column chromatography to obtain the product of  $\alpha$ -form. This synthetic method is very convenient to proceed by one-pot reaction under ordinary conditions.

\*Correspondence: Takayoshi Fujihira, Technical Department and Research Laboratory, Japan Sugar Refiners' Association, 5-7 Sanbancho, Chiyoda-ku, Tokyo 102-0075, Japan; E-mail: fujihira@dream.com.



## INTRODUCTION

Tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose is usually prepared by the reaction of tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with thiourea followed by hydrolysis in alkaline solution of the isothiuronium bromide. This method gives tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose selectively.<sup>[1]</sup> In a preceding paper,<sup>[2]</sup> we reported that the reaction of tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with thioacetamide instead of thiourea gives an adduct identified as 2-(tetra-*O*-acetyl-D-glucopyranosylthio)ethaniminium bromide, methanolysis of which affords tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose. This method also gives tetra-*O*-acetyl-1-thio- $\beta$ -D-galactopyranose, tetra-*O*-acetyl-1-thio- $\alpha$ -D-mannopyranose, hepta-*O*-acetyl-1-thio- $\beta$ -lactose, hepta-*O*-acetyl-1-thio- $\beta$ -maltose or hepta-*O*-acetyl-1-thio- $\beta$ -cellobiose starting with tetra-*O*-acetyl-D-glucopyranosyl bromide or chloride.


The reaction of acylated sugar affords generally per-*O*-acetyl-1,2-*trans*-glycoside owing to the participation of a neighboring acyl group.<sup>[3,4]</sup> Utilizing this characteristic a variety of per-*O*-acetyl-1,2-*trans*-1-thioglycoses have been successfully prepared, but it is generally difficult to prepare per-*O*-acetyl-1,2-*cis*-1-thiosugars by this method in a good yield.<sup>[4]</sup> Since the work of P. Brigl<sup>[5]</sup> that reported the preparation of tetra-*O*-acetyl-1-thio- $\alpha$ -D-glucopyranose from 3,4,6-tri-*O*-acetyl-1,2-anhydro- $\alpha$ -D-glucopyranose, some attempts for synthesis of 1,2-*cis*-1-thioglycoses have been explored using, for example, alkyl or benzyl xanthates<sup>[6]</sup> and thioacetic acid.<sup>[7-10]</sup> It was also reported that tetra-*O*-acetyl-1-thio- $\alpha$ -D-glucopyranose is prepared through the reaction of the phenylmercury(II)thio derivative of tetra-*O*-acetyl-1-*S*-acetyl-1-thio- $\alpha$ -D-glucopyranose with hydrogen sulfide.<sup>[11]</sup>

In the course of our investigations, we found that a modification of our method,<sup>[2]</sup> in which *N,N*-dimethylthioformamide is used, gives an anomeric mixture of tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranoses, from which the  $\alpha$ -anomer can be readily separated.

## RESULTS AND DISCUSSION

Initially, the reaction of tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with *N,N*-dimethylthioformamide instead of thioacetamide under anhydrous conditions was not successful. As described previously,<sup>[2]</sup> the starting materials (tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide and *N,N*-dimethylthioformamide in this case) are soluble in a warm nonpolar organic solvent (e.g., benzene), but the thioiminium salt is barely soluble. K. Hattori<sup>[12]</sup> and Y. Kobayashi<sup>[13]</sup> reported that thioiminium salts can be prepared from the reaction of alkyl or aryl halides with *N,N*-dimethylthioformamide, so we reexamined our reaction under modified conditions. As a result, it was found that the reaction takes place if a slight amount of water is present, at approximately 100°C. The reaction system was estimated to contain ca. ~0.2% of water. The product was mainly *N,N*-dimethyl(tetra-*O*-acetyl-D-glucopyranosylthio)methaniminium bromide as indicated by FAB-MS data; the fragment positive ion peaks of  $m/z$  420 ( $[M - Br^-]^+$ ) and 331 (pyranosyl ring cation) (Figure 1) and the negative ion peaks of  $m/z$  578, 580, 582 ( $[M + Br^-]^-$ ) and 363 (glucosylthiolate anion) and 79, 81 ( $Br^-$ ) were clearly observed (Figure 2). The methanolysis of the thioiminium salt gave an anomeric mixture of tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose with the anomer






**Figure 1.** FAB spectrum (positive ion mode) of thioiminium salt from tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with *N,N*-dimethylthioformamide.

ratio of  $\alpha$ : $\beta$  = 52:48 in 98% yield. This anomer mixture was easily separated by column chromatography.

Little is known about the effect of a slight amount of water in these reactions. It was reported that the anomeric mixture of pyridinium bromide derivatives obtained from the reaction of tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide in pyridine is 1:1.<sup>[14,15]</sup> Although the effect of moisture was not mentioned, the anomer ratio of the product



**Figure 2.** FAB spectrum (negative ion mode) of thioiminium salt from tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide with *N,N*-dimethylthioformamide.



varied with the concentration of tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide in pyridine.<sup>[15]</sup> This reaction is probably a first example of the reaction which is not affected by the participation of neighboring groups in the absence of a catalyst.

The present method was unsuccessful when tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl chloride instead of bromide was used. The present method gave a similar result by using *N,N*-dimethylthioacetamide instead of *N,N*-dimethylthioformamide and, therefore, this reaction is considered to require a tertiary thioamide. The use of primary or secondary thioamides gives a different result from that of tertiary thioamide. Primary or secondary thioamides react with tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide under completely anhydrous conditions to give exclusively tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose. On the other hand, the reaction system containing a slight amount of water affords an anomeric mixture of the product, but the content of  $\alpha$ -form product is rather low (20 ~ 30%).

## EXPERIMENTAL

<sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded with a JEOL GX-400 instrument using tetramethylsilane as a standard. FAB MS spectra were recorded by direct inlet probe with a JEOL MS 700 instrument using glycerin as a matrix and Xe as a reaction gas. Chemical ionization mass spectra were recorded by direct inlet probe with a JEOL JMS-AMII 150 instrument using trimethylmethane as a reaction gas. Optical rotation was determined with a Horiba SEPA-200 polarimeter at 20°C. HPLC analysis (RI monitor) was accomplished by using a Senshu Pak silica-3301-N column (8 × 300 mm, Senshu Kagaku Co. Ltd.), with hexane and ethanol (9:1) as an eluent at a flow rate of 1.5 mL/min.

**General method.** Freshly prepared tetra-*O*-acetyl- $\alpha$ -D-glucopyranosyl bromide (2.5 mmol, 1.03 g) and *N,N*-dimethylthioformamide (2.6 mmol, 0.23 g) were stirred mechanically under an argon stream at approximately 100°C for 5 min. Dry methanol (20 mL) was added after cooling to 20°C, and the mixture was stirred for about 10 min (until the solid dissolved). Solvent was evaporated under reduced pressure, and the precipitated product was separated and purified by silica gel chromatography (benzene–chloroform, 9:1). The products are tetra-*O*-acetyl-1-thio- $\alpha$ -D-glucopyranose (0.46 g, 51%) and tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose (0.42 g, 47%) as a colorless syrup.

**Tetra-*O*-acetyl-1-thio- $\alpha$ -D-glucopyranose.** *R*<sub>f</sub> 0.59 (benzene:chloroform 9:1); mp 93–94°C (Lit. 92–93°C<sup>[7]</sup>);  $[\alpha]_{\text{D}}^{20} + 171^{\circ}$  (Lit. + 168<sup>o[7]</sup>) (*c* 0.8, chloroform); <sup>1</sup>H NMR (chloroform-*d*):  $\delta$  1.91 (d, 1H, *J*<sub>1,SH</sub> 5.77 Hz, SH), 4.11 (dd, 1H, *J*<sub>5,6a</sub> 2.20, *J*<sub>6a,6b</sub> 12.50 Hz, H-6a), 4.30 (dd, 1H, *J*<sub>5,6b</sub> 4.12, *J*<sub>6a,6b</sub> 12.50 Hz, H-6b), 4.44 (ddd, 1H, *J*<sub>5,6a</sub> 2.20, *J*<sub>5,6b</sub> 4.12, *J*<sub>4,5</sub> 10.16 Hz, H-5), 5.03 (dd, 1H, *J*<sub>1,2</sub> 5.77, *J*<sub>2,3</sub> 9.89 Hz, H-2), 5.07 (dd, 1H, *J*<sub>3,4</sub> 9.89, *J*<sub>4,5</sub> 10.16 Hz, H-4), 5.39 (dd, 1H, *J*<sub>3,4</sub> 9.89, *J*<sub>2,3</sub> 9.89 Hz, H-3), 5.94 (dd, 1H, *J*<sub>1, SH</sub> 5.77, *J*<sub>1,2</sub> 5.77 Hz, H-1); <sup>13</sup>C NMR (chloroform-*d*):  $\delta$  61.6 (C-6), 68.9 (C-4), 69.9 (C-3), 70.3 (C-2), 77.1 (C-5), 77.2 (C-1), CIMS: *m/z* 365 (M + 1).

**Tetra-*O*-acetyl-1-thio- $\beta$ -D-glucopyranose.** *R*<sub>f</sub> 0.62 (benzene:chloroform 9:1); mp 113–114°C (Lit. 115°C<sup>[11]</sup>);  $[\alpha]_{\text{D}}^{20} + 11^{\circ}$  (*c* 1.0, chloroform); <sup>1</sup>H NMR (chloroform-*d*):



$\delta$  2.32 (d, 1H,  $J_{1,SH}$  9.61 Hz, SH), 3.73 (ddd, 1H,  $J_{5,6a}$  2.20,  $J_{5,6b}$  4.94,  $J_{4,5}$  9.75 Hz, H-5), 4.13 (dd, 1H,  $J_{5,6a}$  2.20,  $J_{6a,6b}$  12.36 Hz, H-6a), 4.25 (dd, 1H,  $J_{5,6b}$  4.94,  $J_{6a,6b}$  12.36 Hz, H-6b), 4.55 (dd, 1H,  $J_{1,SH}$  9.61,  $J_{1,2}$  9.61 Hz, H-1), 4.98 (dd, 1H,  $J_{1,2}$  9.61,  $J_{2,3}$  9.34 Hz, H-2), 5.11 (dd, 1H,  $J_{3,4}$  9.61,  $J_{4,5}$  9.75 Hz, H-4), 5.20 (dd, 1H,  $J_{3,4}$  9.61,  $J_{2,3}$  9.34 Hz, H-3);  $^{13}\text{C}$  NMR (chloroform-*d*):  $\delta$  62.0 (C-6), 68.1 (C-4), 73.5 (C-2), 73.6 (C-3), 76.3 (C-5), 78.7 (C-1), CIMS:  $m/z$  365 (M + 1).

### ACKNOWLEDGMENT

We thank Ms. Fujita for measurements of NMR spectra.

### REFERENCES

1. Horton, D. 1-Thio-D-glucose. In *Methods in Carbohydrate Chemistry*; Whistler, R.L., Ed.; Academic Press, Inc.: New York, 1963; Vol. 2, 433–437.
2. Fujihira, T.; Chida, M.; Kamijo, H.; Takido, T.; Seno, M. Novel synthesis of 1-thioglycopyranoses via thioiminium salts. *J. Carbohydr. Chem.* **2002**, *21*, 287–292.
3. Igarashi, K. The Koenigs–Knorr reaction. In *Advances in Carbohydrate Chemistry and Biochemistry*; Tipson, R.S., Horton, D., Eds.; Academic Press, Inc.: New York, 1977; Vol. 34, 243–283.
4. Schmidt, R.R. Synthesis of glycosides. In *Comprehensive Heterocyclic Chemistry: The Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds*; Katritzky, A.R., Rees, C.W., Eds.; Pergamon Press, Inc.: London, 1984; Vol. 6, 33–64.
5. Horton, D.; Huston, D.H. Developments in the chemistry of thio sugars. In *Advances in Carbohydrate Chemistry*; Wolform, M.L., Tipson, R.S., Eds.; Academic Press, Inc.: New York, 1963; Vol. 18, 123–199, and references cited therein.
6. Sakata, M.; Haga, M.; Tejima, S.; Akagi, M. Thiosugars. VI. Reaction products of potassium alkyl- and benzylxanthates with acetylated glucosyl halides. *Chem. Pharm. Bull.* **1964**, *12*, 652–656.
7. Blanc-Muesser, M.; Defaye, J.; Driguez, H. Synthesis stereoselectives de 1-thioglycosides. *Carbohydr. Res.* **1978**, *67*, 305–328.
8. Blanc-Muesser, M.; Defaye, J.; Driguez, H. Stereoselective thioglycoside syntheses. Part 4. A new approach to 1,4-linked 1-thio-disaccharides and a synthesis of thiomaltose. *J. Chem. Soc., Perkin Trans. 1*, **1982**, 15–18.
9. Defaye, J.; Driguez, H.; Ohleyer, E.; Orgeret, C.; Viet, C. Stereoselective syntheses of 1,2-*trans*-related 1-thioglycoses. *Carbohydr. Res.* **1984**, *130*, 317–321.
10. Gadelle, A.; Defaye, J. A simple preparation of 2,3,4,6-tetra-*O*-acetyl-1-*S*-acetyl-1-thio- $\alpha$ -D-glucopyranose. *Carbohydr. Res.* **1990**, *200*, 497–498.
11. Defaye, J.; Driguez, H.; Poncet, S.; Chambert, R.; Glatron, M.F.P. Synthesis of 1-thiosucrose and anomers, and the behavior of levansucrase and invertase with this substrate analog. *Carbohydr. Res.* **1984**, *130*, 299–315.
12. Hattori, K.; Takido, T.; Itabashi, K. The synthesis of thiols by use of *N,N*-dimethylthioformamide. *Nippon Kagaku Kaishi* **1979**, 101–105.



13. Kobayashi, Y.; Kawamoto, N.; Sato, Y. Fast atom bombardment mass spectral analysis of thioimidate-ester salts. *Bunseki Kagaku* **1992**, *41*, 441–446.
14. Fischer, E.; Raske, K. Verbindung von Acetobromglucose und Pyridin. *Bericht* **1910**, *43*, 1750–1753.
15. Lemieux, R.U.; Morgan, A.R. The mechanism for the formation of 1,2-*cis*-pyridine nucleosides from 1,2-*cis*-acetohalogenosugars. A novel rearrangement. *J. Am. Chem. Soc.* **1963**, *85*, 1889–1890.
16. Durette, P.L.; Shen, T.Y. Synthesis of 1-thio- $\beta$ -cellobiosides. *Carbohydr. Res.* **1978**, *67*, 484–490.

Received July 17, 2002

Accepted December 23, 2002

