

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

Synthesis of Nitrogen Heterocycles from Methyl α - and β -d-Glucopyranosides

Maria Teresa Capanema Pedrosa^a; Rosemeire Brondi Alves^a; Maria Auxiliadora Fontes Prado^b; José Dias de Souza Filho^a; Ricardo José Alves^b; Norma Beatriz D'Accorso^c

^a Departamento de Química, Instituto de Ciências Exatas, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil ^b Departamento de Produtos Farmacêuticos, Faculdade de Farmácia, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil ^c Departamento de Química Orgânica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

Online publication date: 30 September 2003

To cite this Article Pedrosa, Maria Teresa Capanema , Alves, Rosemeire Brondi , Prado, Maria Auxiliadora Fontes , Filho, José Dias de Souza , Alves, Ricardo José and D'Accorso, Norma Beatriz(2003) 'Synthesis of Nitrogen Heterocycles from Methyl α - and β -d-Glucopyranosides', *Journal of Carbohydrate Chemistry*, 22: 6, 433 – 442

To link to this Article: DOI: 10.1081/CAR-120025329

URL: <http://dx.doi.org/10.1081/CAR-120025329>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Synthesis of Nitrogen Heterocycles from Methyl α - and β -D-Glucopyranosides

Maria Teresa Capanema Pedrosa,¹ Rosemeire Brondi Alves,^{1,*}
Maria Auxiliadora Fontes Prado,² José Dias de Souza Filho,¹
Ricardo José Alves,² and Norma Beatriz D'Accorso³

¹Departamento de Química, Instituto de Ciências Exatas and

²Departamento de Produtos Farmacêuticos, Faculdade de Farmácia,
Universidade Federal de Minas Gerais, Belo Horizonte, Brazil

³Departamento de Química Orgânica, Facultad de Ciencias Exactas y Naturales,
Universidad de Buenos Aires, Buenos Aires, Argentina

ABSTRACT

Eight nitrogen heterocycles, mono and disubstituted tetrazoles and oxadiazoles, were synthesized from methyl D-glucopyranoside anomers. The monosubstituted tetrazoles resulted from the reaction of 6-cyanoglucopyranoside derivatives with sodium azide. By alkylation of the monosubstituted tetrazoles, the 1,5 and 2,5 disubstituted tetrazoles were obtained. The monosubstituted tetrazoles were reacted with acetic anhydride to give the oxadiazoles.

Key Words: Tetrazole; Oxadiazole; Heterocycle.

*Correspondence: Rosemeire Brondi Alves, Departamento de Química, Instituto de Ciências Exatas, Universidade Federal de Minas Gerais, Avenida Antônio Carlos 6627, Belo Horizonte 31.270-901, Brazil; E-mail: brondi@dedalus.lcc.ufmg.br.



INTRODUCTION

Tetrazoles are an increasingly popular structural unit, often used as metabolically stable surrogates for carboxylic acid groups, and as convenient lipophilic spacers in pharmaceuticals.^[1] This unit also has roles in coordination chemistry as a ligand, and in various material science applications, including in specialty explosives.^[2] There have been considerable achievements in the practical application of tetrazoles, especially in medicine and biochemistry.^[3–5] Recent publications include the use of tetrazoles as inhibitors of monoamine oxidase,^[6] as antiviral and antibacterial agents,^[7,8] and antagonists of cerebellum-specific GABA_A receptors^[9] and angiotensin II.^[10] The oxadiazoles also display biological activities such as fungicidal,^[11] antibacterial^[12] and glycosidase inhibitors.^[13] We are interested in the synthesis of heterocycles using carbohydrates as starting materials, a strategy that offers the possibility of synthesis of compounds possessing chiral centers with the necessary enantiomeric purity required for active biological compounds. In this context, we synthesized some heterocycles using carbohydrates as starting materials.^[14–17]


RESULTS AND DISCUSSION

The initial work centered on the preparation of methyl 2,3,4-tri-*O*-benzyl-6-cyano-6-deoxy-*D*-glucopyranoside anomers **1 α** and **1 β** in five conventional synthesis steps starting from methyl $\alpha(\beta)$ -*D*-glucopyranosides (Scheme 1). The starting materials were protected as 4,6-di-*O*-benzylidene derivatives^[18] and then the C-2 and C-3 hydroxyl groups were *O*-benzylated.^[19] Removal of the benzylidene group under LiAlH₄ reduction conditions^[20] and replacement of the hydroxy group at C-6 by iodine^[21] afforded the methyl 2,3,4-tri-*O*-benzyl-6-deoxy-6-iodo- $\alpha(\beta)$ -*D*-glucopyranosides (**4 α** and **4 β**). Treatment of **4 α** and **4 β** with potassium cyanide in DMF^[22] gave nitriles **1 α** and **1 β** , respectively. After the preparation of these compounds, we went on to obtain the target tetrazoles.

Several methods of tetrazole synthesis are known,^[2,4,5,23] but the most convenient method of synthesis of substituted tetrazoles involves the reaction of nitriles with salts of hydrazoic acid.^[2,5] Thus, products **1 α** and **1 β** were submitted to reaction with sodium azide and ammonium chloride in DMF^[16,17] to give 5-(methyl α/β -2,3,4-tri-*O*-benzyl-6-deoxy-*D*-glucopyranos-6-yl)tetrazole **2 α** and **2 β** in 81% and 73% yields, respectively.

Distinct oxadiazoles can be obtained by many methods.^[12,24] In this work, we synthesized 1,3,4-oxadiazoles **3 α** and **3 β** in 84% and 60% yields, respectively, by the reaction of tetrazoles **2 α** and **2 β** with acetic anhydride and pyridine^[12,16,17] at 110°C. To construct disubstituted tetrazoles, the alkylation reactions of tetrazole anions are often used.^[25] However, due to the existence of the tetrazole tautomeric forms, alkylation gives mixtures of *N*(1)- and *N*(2)-alkylation isomers.^[25] The ratio of these regioisomers is affected by the electronic and steric effect of the substituent.^[25] Due to the two tautomeric forms of tetrazole, tetrazole **2 α** when reacted with iodine derivative **4 α** in the presence of anhydrous potassium carbonate in acetone,^[6,17] gave a mixture of the two alkylated tetrazoles: 2,5-bis(methyl 2,3,4-tri-*O*-benzyl-6-deoxy- α -*D*-glucopyranos-6-yl)-2*H*-tetrazole (**5 α**) (31%) and 1,5-bis(methyl 2,3,4-tri-*O*-benzyl-6-deoxy- α -*D*-





Scheme 1. Reagents, conditions and yields: i, NaN_3 , DMF, NH_4Cl , 95°C , 192 h (α) and 144 h (β), 81% (α) and 73% (β); ii, acetic anhydride, pyridine, 110°C , 96 h (α) and 144 h (β), 84% (α) and 60% (β); iii, K_2CO_3 , acetone, compounds **2 α** and **4 α** , 70°C , 120 h, **5 α** (31%), **6 α** (34%); compounds **2 β** e **4 β** , 70°C , 64 h, **5 β** (50%) and **6 β** (25%).

glucopyranos-6-yl)-1*H*-tetrazole (**6 α**) (34%) in an approximate 1:1 ratio. Tetrazole **2 β** by reaction with **4 β** under the same conditions described for **2 α** gave the regioisomers **5 β** (50%) and **6 β** (25%) in a 2:1 ratio. In a preliminary systematic conformational research (Hyperchem Pro 6.0 – Molecular Mechanics), it was observed that when the anomeric methoxy group is in the β position, there is a steric repulsion with the nucleophilic glycoside moiety, and the alkylation on the tetrazolic ring occurs preferentially at position 2. In contrast, when the anomeric methoxy group is α , the cited repulsive interaction is absent, and alkylation occurs at positions 1 and 2 in the same ratio (Scheme 1).

The structures of all the compounds obtained were confirmed by NMR spectroscopy. However, it is interesting to remark on the unequivocal characterization of the isomeric tetrazoles **5** and **6**. The literature reports that tetrazole isomers with methyl or alkyl substituents at positions 1 and 2 are readily distinguished by the ^1H and ^{13}C chemical shifts of the *N*-alkyl group. Alkyl groups bonded to *N*-1 are more shielded by *ca.* 0.15–0.35 ppm in the ^1H spectra and by *ca.* 2–6 ppm in the ^{13}C spectra relative to their corresponding *N*-2 isomers. However, HMBC experiments were carried out for confirmation and revealed that the cross correlation of $\text{H}6''$ with tetrazolic carbon occurs solely in derivatives **6** because of the 3J coupling constant.



EXPERIMENTAL

General procedures. Melting points were determined on a Mettler FP80HT apparatus and are uncorrected. Optical rotation was determined with a Perkin-Elmer 341 Polarimeter at 25°C. ^{13}C and ^1H NMR spectra were recorded on a Bruker Avance DRX-400 spectrometer. Chemical shifts are reported in δ units downfield from Me_4Si . Elemental analyses were carried out using a Perkin-Elmer 2400 CHN apparatus. Column chromatography was performed with silica gel 60, 70-230 mesh (Merck). The term “standard work-up” means that the organic layer was washed with satd NaCl solution, dried over anhydrous Na_2SO_4 , filtered, and the solvent was removed under reduced pressure. Dry solvents were prepared as follows. DMF was dried over potassium hydroxide and stirred for 24 hours at room temperature. The solution was filtered and distilled under reduced pressure. Dry pyridine was prepared after agitation with KOH for 17 h at room temperature, and then distilled. Potassium permanganate was added to refluxing acetone until its color remained purple. The solution was refluxed for 6 h more and distilled. Dry acetone was collected under desiccated K_2CO_3 .

Methyl 2,3,4-tri-*O*-benzyl-6-cyano-6-deoxy- β -D-glucopyranoside **1 β .**^[22] To a solution of methyl 2,3,4-tri-*O*-benzyl-6-deoxy-6-iodo- β -D-glucopyranoside (**4 β**) (1.00 g, 1.74 mmol) in dry DMF (25 mL), KCN (0.38 g, 5.85 mmol) was added and the reaction mixture stirred at rt for 65 h. DMF was removed under reduced pressure, giving a residue. Water was added, the aqueous solution extracted with CH_2Cl_2 (4×30 mL), and the combined organic extracts submitted to standard work-up. The crude product was purified by column chromatography (10% EtOAc in hexane, gradually increasing the percentage of EtOAc) to give product **1 β** (0.61 g, 74%) as a white solid: $[\alpha]_{\text{D}} + 32$ (*c* 1.3, CHCl_3); mp 69–73°C; ^1H NMR (CDCl_3 , 400 MHz): δ 7.37–7.23 (m, 15H, Ar), 4.96 (d, 1H, J_{gem} 10.9 Hz, $1 \times \text{PhCH}_2$), 4.92 (d, 1H, J_{gem} 11.0 Hz, $1 \times \text{PhCH}_2$), 4.91 (d, 1H, J_{gem} 11.2 Hz, $1 \times \text{PhCH}_2$), 4.78 (d, 1H, J_{gem} 10.9 Hz, $1 \times \text{PhCH}_2$), 4.70 (d, 1H, J_{gem} 11.0 Hz, $1 \times \text{PhCH}_2$), 4.60 (d, 1H, J_{gem} 11.2 Hz, $1 \times \text{PhCH}_2$), 4.35 (d, 1H, $J_{1,2}$ 7.8 Hz, H1), 3.65 (t, 1H, $J_{3,4}$ 9.1 Hz, H3), 3.58 (s, 3H, MeO), 3.49 (ddd, 1H, $J_{5,4}$ 9.1, $J_{5,6'}$ 7.7, $J_{5,6''}$ 3.2 Hz, H5), 3.45 (dd, 1H, H2), 3.36 (t, 1H, H4), 2.70 (dd, 1H, J_{gem} 16.8 Hz, H6'), 2.43 (dd, 1H, H6''); ^{13}C NMR (CDCl_3 , 100 MHz): δ 138.3, 138.2, 137.5 ($3 \times \text{C ipso}$), 128.6, 128.4, 128.3, 128.2, 128.1, 127.9, 127.7 ($15 \times \text{Ar}$), 116.8 (CN), 104.6 (C1), 84.2 (C3), 82.2 (C2), 79.8 (C4), 75.6, 75.2, 74.8 ($3 \times \text{PhCH}_2$), 70.5 (C5), 57.2 (MeO), 21.0 (C6).

Anal. Calcd for $\text{C}_{29}\text{H}_{31}\text{NO}_5$: C, 73.55; H, 6.60; N, 2.96. Found: C, 73.14; H, 6.59; N, 2.86.

Treating the epimer **4 α** (3.15 g, 6.79 mmol) with KCN as described above for **4 β** , product **1 α** (2.36 g, 91%) was obtained as a whitish oil: $[\alpha]_{\text{D}} + 46.8$ (*c* 1.1, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz): δ 7.36–7.24 (m, 15H, Ar), 5.01 (d, 1H, J_{gem} 10.9 Hz, $1 \times \text{PhCH}_2$), 4.93 (d, 1H, J_{gem} 11.2 Hz, $1 \times \text{PhCH}_2$), 4.80 (d, 1H, J_{gem} 10.9 Hz, $1 \times \text{PhCH}_2$), 4.79 (d, 1H, J_{gem} 12.1 Hz, $1 \times \text{PhCH}_2$), 4.65 (d, 1H, J_{gem} 12.1 Hz, $1 \times \text{PhCH}_2$), 4.60 (d, 1H, J_{gem} 11.2 Hz, $1 \times \text{PhCH}_2$), 4.57 (d, 1H, $J_{1,2}$ 3.6 Hz, H1), 3.98 (t, 1H, $J_{3,4}$ 9.6 Hz, H3), 3.79 (ddd, 1H, $J_{5,4}$ 9.6, $J_{5,6'}$ 7.0, $J_{5,6''}$ 3.4 Hz, H5), 3.55 (dd, 1H, H2), 3.39 (s, 3H, MeO), 3.33 (t, 1H, H4), 2.62 (dd, 1H, J_{gem} 16.8 Hz, H6'), 2.42 (dd, 1H, H6''); ^{13}C NMR (CDCl_3 , 50 MHz): δ 138.3, 137.7, 137.5 ($3 \times \text{C ipso}$), 128.5, 128.4, 128.3, 128.1, 128.0, 127.9, 127.8, 127.7 ($15 \times \text{Ar}$), 116.9 (CN),



98.0 (C1), 81.5 (C3), 79.8 (C2 or C4), 79.7 (C2 or C4), 75.6, 75.1, 73.4 (3 × PhCH₂), 66.2 (C5), 55.4 (MeO), 20.7 (C6).

Anal. Calcd for C₂₉H₃₁NO₅: C, 73.55; H, 6.60; N, 2.96. Found: C, 72.95; H, 6.59; N, 2.99.

5-(Methyl 2,3,4-tri-*O*-benzyl-6-deoxy-β-D-glucopyranos-6-yl)tetrazole 2β.^[16,17]

To a solution of methyl 2,3,4-tri-*O*-benzyl-6-cyano-6-deoxy-β-D-glucopyranoside (**1β**) (0.17 g, 0.36 mmol) in dry DMF (10 mL), NaN₃ (0.28 g, 4.31 mmol) and NH₄Cl (0.23 g, 4.30 mmol) were added. The solution was stirred at 95°C for 144 h. DMF was removed under reduced pressure, giving a residue. Aqueous H₂SO₄ (3 mol.L⁻¹) was added until pH 1 was reached. Water was added, the aqueous solution extracted with CH₂Cl₂ (4 × 25 mL), and the combined organic extracts submitted to standard work-up. The crude product was purified by column chromatography (30% EtOAc in hexane, gradually increasing the percentage of EtOAc) to give product **2β** (0.14 g, 73%) as a white solid: [α]_D + 2 (c 1.2, Me₂SO); mp 217–219°C; ¹H NMR (Me₂SO-*d*₆, 400 MHz): δ 16.03 (br s, 1H, NH), 7.89–7.26 (m, 15H, Ar), 4.83 (d, 1H, *J*_{gem} 11.1 Hz, 1 × PhCH₂), 4.82 (d, 1H, *J*_{gem} 11.1 Hz, 1 × PhCH₂), 4.78 (d, 1H, *J*_{gem} 11.5 Hz, 1 × PhCH₂), 4.73 (d, 2H, *J*_{gem} 11.1 Hz, 2 × PhCH₂), 4.62 (d, 1H, *J*_{gem} 11.5 Hz, 1 × PhCH₂), 4.32 (d, 1H, *J*_{1,2'} 7.8 Hz, H1'), 3.72 (td, 1H, *J*_{5',6'} 9.1, *J*_{5',6'} 3.0 Hz, H5'), 3.65 (t, 1H, *J*_{3',4'} 9.1 Hz, H3'), 3.42 (t, 1H, H4'), 3.36 (dd, 1H, *J*_{gem} 15.1 Hz, H6'), 3.29–3.25 (m, 1H, H2'), 3.25 (s, 3H, MeO), 3.09 (dd, 1H, H6''); ¹³C NMR (Me₂SO-*d*₆, 100 MHz): δ 153.0 (C5), 138.5, 138.4, 138.1 (3 × C *ipso*), 128.2, 128.1, 128.0, 127.9, 127.8, 127.6, 127.5, 127.3 (15 × Ar), 103.3 (C1'), 83.5 (C3'), 81.6 (C2'), 80.6 (C4'), 74.4, 74.0, 73.5 (3 × PhCH₂), 71.7 (C5'), 56.0 (MeO), 25.8 (C-6').

Anal. Calcd for C₂₉H₃₂N₄O₅: C, 67.43; H, 6.24; N, 10.85. Found: C, 67.13; H, 6.36; N, 11.25.

Treatment of product **1α** (2.21 g, 4.66 mmol) as described above for **1β** gave product **2α** (1.94 g, 81%) as a white solid: [α]_D + 13.6 (c 1.2, CHCl₃); mp 218–223°C; ¹H NMR (CDCl₃, 400 MHz): δ 13.23 (br s, 1H, NH), 7.34–7.24 (m, 15H, Ar), 4.97 (d, 1H, *J*_{gem} 10.9 Hz, 1 × PhCH₂), 4.89 (d, 1H, *J*_{gem} 11.0 Hz, 1 × PhCH₂), 4.80 (d, 1H, *J*_{gem} 10.9 Hz, 1 × PhCH₂), 4.76 (d, 1H, *J*_{gem} 12.0 Hz, 1 × PhCH₂), 4.65 (d, 1H, *J*_{gem} 11.0 Hz, 1 × PhCH₂), 4.62 (d, 1H, *J*_{gem} 12.0 Hz, 1 × PhCH₂), 4.60 (d, 1H, *J*_{1,2'} 3.6 Hz, H1'), 4.00–3.93 (m, 1H, H5'), 3.99 (t, 1H, *J*_{3',4'} 9.7 Hz, H3'), 3.56 (dd, 1H, H2'), 3.48 (dd, 1H, *J*_{gem} 15.3, *J*_{6',5'} 3.6 Hz, H6'), 3.22 (s, 3H, MeO), 3.19 (t, 1H, *J*_{4',5'} 9.7 Hz, H4'), 3.14 (dd, 1H, *J*_{6',5'} 9.7 Hz, H6''); ¹³C NMR (CDCl₃, 50 MHz): δ 153.3 (C5), 138.3, 137.8, 137.6 (3 × C *ipso*), 128.6, 128.5, 128.4, 128.3, 128.1, 128.0, 127.9, 127.8, 127.7 (15 × Ar), 98.0 (C1'), 81.5 (C3'), 80.3 (C4'), 79.9 (C2'), 75.7, 75.1, 73.3 (3 × PhCH₂), 67.9 (C5'), 55.4 (MeO), 25.9 (C6').

Anal. Calcd for C₂₉H₃₂N₄O₅: C, 67.43; H, 6.24; N, 10.85. Found: C, 67.19; H, 5.99; N, 10.89.

2-Methyl-5-(methyl 2,3,4-tri-*O*-benzyl-6-deoxy-β-D-glucopyranos-6-yl)-1,3,4-oxadiazole 3β.^[12,16,17]

To a stirred solution of 5-(methyl 2,3,4-tri-*O*-benzyl-6-deoxy-β-D-glucopyranos-6-yl)tetrazole (**2β**) (0.15 g, 0.29 mmol) in dry pyridine (3 mL), acetic anhydride (9 mL, 99.10 mmol) was added. The solution was stirred at 110°C for 144 h, then cooled to rt and cold aq HCl (19 mL, 3 mol.L⁻¹) was added. Water was added, the aqueous solution extracted with CH₂Cl₂ (4 × 25 mL), and the combined organic



extracts submitted to standard work-up. The crude product was purified by column chromatography (40% EtOAc in hexane, gradually increasing the percentage of EtOAc) to give product **3β** (0.09 g, 60%) as a white solid: $[\alpha]_D + 13.2$ (*c* 0.6, CHCl₃); mp 101–105°C; ¹H NMR (CDCl₃, 400 MHz): δ 7.34–7.25 (m, 15H, Ar), 4.95 (d, 1H, J_{gem} 10.9 Hz, 1 × PhCH₂), 4.93 (d, 1H, J_{gem} 11.5 Hz, 1 × PhCH₂), 4.90 (d, 1H, J_{gem} 11.7 Hz, 1 × PhCH₂), 4.78 (d, 1H, J_{gem} 10.9 Hz, 1 × PhCH₂), 4.69 (d, 1H, J_{gem} 11.7 Hz, 1 × PhCH₂), 4.66 (d, 1H, J_{gem} 11.5 Hz, 1 × PhCH₂), 4.30 (d, 1H, $J_{1',2'}$ 7.8 Hz, H1'), 3.75 (td, 1H, $J_{5',6''}$ 8.8 Hz, $J_{5',6'}$ 3.8, H5'), 3.68 (t, 1H, $J_{3',4'}$ 8.8 Hz, H3'), 3.45 (s, 3H, MeO), 3.45–3.39 (m, 1H, H2'), 3.41 (t, 1H, H4'), 3.26 (dd, 1H, J_{gem} 15.4 Hz, H6'), 2.93 (dd, 1H, H6''), 2.43 (s, 3H, Me); ¹³C NMR (CDCl₃, 100 MHz): δ 164.2 (C5), 163.7 (C2), 138.5, 138.4, 137.9 (3 × C *ipso*), 128.5, 128.4, 128.3, 128.1, 128.0, 127.9, 127.8, 127.7 (15 × Ar), 104.5 (C1'), 84.5 (C3'), 82.4 (C2'), 80.7 (C4'), 75.7, 74.9, 74.7 (3 × PhCH₂), 72.0 (C5'), 56.9 (MeO), 28.3 (C6'), 10.8 (Me).

Anal. Calcd for C₃₁H₃₄N₂O₆: C, 70.17; H, 6.46; N, 5.28. Found: C, 69.97; H, 6.29; N, 5.32.

The treatment of tetrazole **2α** (0.20 g, 0.39 mmol) as described above for **2β** gave product **3α** (0.17 g, 84%) as a whitish oil: $[\alpha]_D + 42.7$ (*c* 1.2, CHCl₃); ¹H NMR (CDCl₃, 400 MHz): δ 7.36–7.25 (m, 15H, Ar), 5.01 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.95 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.81 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.78 (d, 1H, J_{gem} 12.2 Hz, 1 × PhCH₂), 4.66 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.63 (d, 1H, J_{gem} 12.2 Hz, 1 × PhCH₂), 4.50 (d, 1H, $J_{1',2'}$ 3.6 Hz, H1'), 4.12–4.02 (m, 1H, H5'), 4.01 (t, 1H, $J_{3',4'}$ 9.6 Hz, H3'), 3.50 (dd, 1H, H2'), 3.34 (t, 1H, $J_{4',5'}$ 9.6 Hz, H4'), 3.29 (s, 3H, MeO), 3.19 (dd, 1H, J_{gem} 15.4, $J_{6',5'}$ 3.7 Hz, H6'), 2.89 (dd, 1H, $J_{6'',5'}$ 8.4 Hz, H6''), 2.42 (s, 3H, Me); ¹³C NMR (CDCl₃, 50 MHz): δ 164.2 (C5), 163.6 (C2), 138.4, 137.9, 137.8 (3 × C *ipso*), 128.3, 128.0, 127.9, 127.8, 127.7, 127.6 (15 × Ar), 97.8 (C1'), 81.7 (C3'), 80.6 (C4'), 79.8 (C2'), 75.6, 74.8, 73.2 (3 × PhCH₂), 67.5 (C5'), 55.2 (MeO), 27.9 (C6'), 10.7 (Me).

Anal. Calcd for C₃₁H₃₄N₂O₆: C, 70.17; H, 6.46; N, 5.28. Found: C, 69.98; H, 6.44; N, 5.33.

2,5-Bis(methyl 2,3,4-tri-*O*-benzyl-6-deoxy-β-D-glucopyranos-6-yl)-2H-tetrazole 5β and 1,5-Bis(methyl 2,3,4-tri-*O*-benzyl-6-deoxy-β-D-glucopyranos-6-yl)-1H-tetrazole 6β.^[6,17] To a solution of 5-(methyl 2,3,4-tri-*O*-benzyl-6-deoxy-β-D-glucopyranos-6-yl)tetrazole (**2β**) (0.35 g, 0.68 mmol) and methyl 2,3,4-tri-*O*-benzyl-6-deoxy-6-iodo-β-D-glucopyranoside (**4β**) (0.39 g, 0.68 mmol) in dry acetone (120 mL), anhydrous K₂CO₃ (0.94 g, 6.81 mmol) was added. The solution was stirred at 70°C for 64 h in a pressure reactor. After cooling to rt, the solvent was removed, water was added, the aqueous solution was extracted with EtOAc (4 × 20 mL) and the combined organic extracts were submitted to standard work-up. The crude product was purified by column chromatography (10% EtOAc in hexane, gradually increasing the percentage of EtOAc) to give product **5β** (0.33 g, 50%) and **6β** (0.16 g, 25%) as white solids.

Compound **5β**: $[\alpha]_D + 18$ (*c* 1.0, CHCl₃); mp 122–123°C; ¹H NMR (CDCl₃, 400 MHz): δ 7.32–7.27 (m, 30H, Ar), 4.95 (d, 1H, J_{gem} 10.9 Hz, 1 × PhCH₂), 4.94 (d, 1H, J_{gem} 10.9 Hz, 1 × PhCH₂), 4.93 (d, 1H, J_{gem} 10.9 Hz, 1 × PhCH₂), 4.92 (d, 1H, J_{gem} 11.2 Hz, 1 × PhCH₂), 4.88 (d, 1H, J_{gem} 11.1 Hz, 1 × PhCH₂), 4.86 (d, 1H, J_{gem} 11.1 Hz, 1 × PhCH₂), 4.78 (d, 2H, J_{gem} 10.9 Hz, 2 × PhCH₂), 4.73 (dd, 1H, $J_{6b'',6a''}$ 13.9, $J_{6b'',5''}$ 3.1 Hz, H-6b''), 4.72 (d, 1H, J_{gem} 11.2 Hz, 1 × PhCH₂), 4.71 (d, 1H, J_{gem} 10.9

Hz, 1 × PhCH₂), 4.67 (d, 1H, J_{gem} 11.1 Hz, 1 × PhCH₂), 4.65 (d, 1H, J_{gem} 11.1 Hz, 1 × PhCH₂), 4.56 (dd, 1H, $J_{6a'',5''}$ 7.6 Hz, H-6a''), 4.27 (d, 1H, $J_{1,2}$ 7.8 Hz, H1' or H1''), 4.21 (d, 1H, $J_{1,2}$ 7.7 Hz, H1' or H1''), 3.86–3.80 (m, 2H, H5', H-5''), 3.68 (t, 1H, $J_{3,4}$ 9.1 Hz, H3' or H3''), 3.67 (t, 1H, $J_{3,4}$ 9.1, H3' or H3''), 3.48 (t, 1H, $J_{4',5'}$ 9.1 Hz, H4'), 3.47 (t, 1H, $J_{4',5'}$ 9.1 Hz, H4''), 3.42–3.34 (m, 2H, H2', H2''), 3.38 (s, 3H, MeO' or MeO''), 3.35 (dd, 1H, $J_{6b',6a'}$ 15.3, $J_{6b',5'}$ 3.6 Hz, H6b'), 3.32 (s, 3H, MeO' or MeO''), 3.08 (dd, 1H, $J_{6a',5'}$ 8.1 Hz, H6a'); ¹³C NMR (CDCl₃, 100 MHz): δ 163.4 (C-5), 138.6, 138.4, 138.3, 138.2, 137.8 (6 × C *ipso*), 128.6, 128.5, 128.4, 128.3, 128.2, 128.1, 128.0, 127.9, 127.8, 127.7, 127.6, 127.5 (30 × Ar), 104.5, 104.4 (C1', C1''), 84.7, 84.5 (C3', C3''), 82.5, 82.2 (C2', C2''), 81.0, 78.4 (C4', C4''), 75.7, 75.6, 75.0, 74.9, 74.7 (6 × PhCH₂), 72.6 (C5'), 72.4 (C5''), 56.8, 56.7 (MeO', MeO''), 53.5 (C6''), 28.2 (C6').

Anal. Calcd for C₅₇H₆₂N₄O₁₀: C, 71.08; H, 6.49; N, 5.82. Found: C, 69.97; H, 6.29; N, 5.32.

Compound **6β**: [α]_D + 14 (*c* 0.9, CHCl₃); mp 110–114°C; ¹H NMR (CDCl₃, 400 MHz): δ 7.36–7.25 (m, 30H, Ar), 4.97 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.96 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.95 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.94 (d, 1H, J_{gem} 10.5 Hz, 1 × PhCH₂), 4.87 (d, 1H, J_{gem} 11.0 Hz, 1 × PhCH₂), 4.86 (d, 1H, J_{gem} 11.0 Hz, 1 × PhCH₂), 4.81 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.79 (d, 1H, J_{gem} 10.5 Hz, 1 × PhCH₂), 4.78 (d, 2H, J_{gem} 10.8 Hz, 2 × PhCH₂), 4.71 (dd, 1H, $J_{6b'',6a''}$ 14.5, $J_{6b'',5''}$ 2.8 Hz, H6b''), 4.67 (d, 1H, J_{gem} 11.0 Hz, 1 × PhCH₂), 4.66 (d, 1H, J_{gem} 11.0 Hz, 1 × PhCH₂), 4.38 (dd, 1H, $J_{6a'',5''}$ 7.5 Hz, H6a''), 4.22 (d, 1H, $J_{1',2'}$ 7.9 Hz, H1'), 4.18 (d, 1H, $J_{1'',2''}$ 7.9 Hz, H1''), 3.74 (ddd, 1H, $J_{5',4'}$ 9.7, $J_{5',6a'}$ 8.2, $J_{5',6b'}$ 3.2 Hz, H5'), 3.72–3.66 (m, 1H, H5''), 3.69 (t, 1H, $J_{3,4}$ 9.0 Hz, H3' or H3''), 3.68 (t, 1H, $J_{3,4}$ 9.1 Hz, H3' or H3''), 3.47 (dd, 1H, $J_{6b',6a'}$ 15.1 Hz, H6b'), 3.45–3.34 (m, 4H, H2', H2'', H4', H4''), 3.33 (s, 3H, MeO' or MeO''), 3.29 (s, 3H, MeO' or MeO''), 3.07 (dd, 1H, H6a'); ¹³C NMR (CDCl₃, 100 MHz): δ 153.8 (C5), 138.4, 138.3, 138.2, 138.1, 138.0, 137.8 (6 × C *ipso*), 128.6, 128.5, 128.4, 128.3, 128.1, 128.0, 127.9, 127.8, 127.7, 127.6, 127.5 (30 × Ar), 104.6, 104.5 (C1', C1''), 84.4, 84.3 (C3', C3''), 82.2, 82.0, 80.2, 78.1 (C2', C2'', C4', C4''), 75.6, 74.9, 74.8, 74.7 (6 × PhCH₂), 73.4 (C5''), 73.1 (C5'), 57.0 (MeO'), 57.0 (MeO''), 47.7 (C6''), 25.7 (C6').

Anal. Calcd for C₅₇H₆₂N₄O₁₀: C, 71.08; H, 6.49; N, 5.82. Found: C, 70.60; H, 6.31; N, 5.76.

By treatment of tetrazole **2α** (1.00 g, 1.94 mmol) with **4α** (1.10 g, 1.92 mmol) as described above for **2β**, product **5α** (0.57 g, 31%) and **6α** (0.64g, 34%) were obtained as whitish oils.

Compound **5α**: [α]_D + 47.1 (*c* 1.0, CHCl₃); ¹H NMR (CDCl₃, 400 MHz): δ 7.33–7.28 (m, 30H, Ar), 5.00 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.98 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.96 (d, 1H, J_{gem} 11.1 Hz, 1 × PhCH₂), 4.95 (d, 1H, J_{gem} 11.7 Hz, 1 × PhCH₂), 4.81 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.80 (d, 1H, J_{gem} 10.8 Hz, 1 × PhCH₂), 4.75 (d, 1H, J_{gem} 12.1 Hz, 1 × PhCH₂), 4.74 (d, 1H, J_{gem} 12.0 Hz, 1 × PhCH₂), 4.72 (d, 1H, J_{gem} 11.1 Hz, 1 × PhCH₂), 4.71 (d, 1H, J_{gem} 11.7 Hz, 1 × PhCH₂), 4.70 (dd, 1H, $J_{6b'',6a''}$ 14.7, $J_{6b'',5''}$ 2.8 Hz, H6b''), 4.61 (d, 1H, J_{gem} 12.1 Hz, 1 × PhCH₂), 4.59 (d, 1H, J_{gem} 12.0 Hz, 1 × PhCH₂), 4.57 (dd, 1H, $J_{6a'',5''}$ 8.3 Hz, H6a''), 4.44 (d, 1H, $J_{1',2'}$ 3.5 Hz, H1'), 4.43 (d, 1H, $J_{1'',2''}$ 3.5 Hz, H1''), 4.17–4.07 (m, 2H, H5' and H5''), 4.01 (t, 1H, $J_{3'',4''}$ 9.4 Hz, H3''), 4.00 (t, 1H, $J_{3',4'}$ 9.4 Hz, H3'), 3.45 (dd, 1H, H2''), 3.41 (dd, 1H, H2'), 3.35 (t, 1H, $J_{4',5'}$ 9.4 Hz, H4'), 3.37–3.31 (m, 1H, H4''), 3.30 (dd, 1H, $J_{6b',6a'}$ 14.7, $J_{6b',5'}$ 3.4 Hz, H6b'), 3.19 (s, 3H, MeO' or MeO''), 3.11



(s, 3H, MeO' or MeO''), 3.00 (dd, 1H, $J_{6a',5'}$ 8.5 Hz, H6a'); ^{13}C NMR (CDCl_3 , 100 MHz): δ 163.5 (C5), 138.6, 138.3, 138.2, 138.0, 137.8 ($6 \times \text{C ipso}$), 128.6, 128.5, 128.4, 128.3, 128.2, 128.0, 127.9, 127.8, 127.7, 127.6, 127.5, 127.4 ($30 \times \text{Ar}$), 97.8 (C1' or C1''), 97.7 (C1' or C1''), 81.9 (C3''), 81.8 (C3'), 81.0 (C4'), 80.0 (C2'), 79.7 (C2''), 78.3 (C4''), 75.8, 75.7, 74.9, 74.8, 73.4, 73.3 ($6 \times \text{PhCH}_2$), 68.4 (C5'), 68.2 (C5''), 55.2 (MeO' or MeO''), 55.1 (MeO' or MeO''), 53.4 (C6''), 27.9 (C6').

Anal. Calcd for $\text{C}_{57}\text{H}_{62}\text{N}_4\text{O}_{10}$: C, 71.08; H, 6.49; N, 5.82. Found: C, 70.89; H, 6.32; N, 5.65.

Compound **6a**: $[\alpha]_{\text{D}} + 51.3$ (c 1.0, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz): δ 7.34–7.26 (m, 30H, Ar), 4.99 (d, 1H, J_{gem} 10.9 Hz, $1 \times \text{PhCH}_2$), 4.98 (d, 1H, J_{gem} 11.0 Hz, $1 \times \text{PhCH}_2$), 4.96 (d, 1H, J_{gem} 12.3 Hz, $1 \times \text{PhCH}_2$), 4.95 (d, 1H, J_{gem} 12.5 Hz, $1 \times \text{PhCH}_2$), 4.81 (d, 1H, J_{gem} 10.9 Hz, $1 \times \text{PhCH}_2$), 4.80 (d, 1H, J_{gem} 11.0 Hz, $1 \times \text{PhCH}_2$), 4.76 (d, 1H, J_{gem} 11.7 Hz, $1 \times \text{PhCH}_2$), 4.74 (d, 1H, J_{gem} 11.7 Hz, $1 \times \text{PhCH}_2$), 4.73 (d, 1H, J_{gem} 12.3 Hz, $1 \times \text{PhCH}_2$), 4.72 (d, 1H, J_{gem} 12.5 Hz, $1 \times \text{PhCH}_2$), 4.63 (d, 1H, J_{gem} 11.7 Hz, $1 \times \text{PhCH}_2$), 4.60 (d, 1H, J_{gem} 11.7 Hz, $1 \times \text{PhCH}_2$), 4.48 (dd, 1H, $J_{6b'',6a''}$ 14.4, $J_{6b'',5''}$ 2.5 Hz, H6b''), 4.39 (d, 1H, $J_{1',2'}$ 3.6 Hz, H1'), 4.38 (d, 1H, $J_{1'',2''}$ 3.6 Hz, H1''), 4.15 (dd, 1H, $J_{6a'',5''}$ 8.0 Hz, H6a''), 4.00–3.86 (m, 2H, H5' and H5''), 3.96 (t, 2H, $J_{3',4'}$ 9.4 Hz, H3' and H3''), 3.43 (dd, 1H, H2'), 3.40 (dd, 1H, H2''), 3.32–3.23 (m, 3H, H4', H4'' and H6b'), 3.09 (s, 3H, MeO' or MeO''), 2.98 (s, 3H, MeO' or MeO''), 2.81 (dd, 1H, $J_{6a',6b'}$ 15.1, $J_{6a',5'}$ 8.8 Hz, H6a'); ^{13}C NMR (CDCl_3 , 100 MHz): δ 153.7 (C5), 138.5, 138.3, 138.2, 138.0, 137.9, 137.8 ($6 \times \text{C ipso}$), 128.6, 128.5, 128.4, 128.3, 128.0, 127.9, 127.8, 127.7, 127.6 ($30 \times \text{Ar}$), 97.8 (C1' or C1''), 97.7 (C1' or C1''), 81.8 (C3' and C3''), 80.3 (C4'), 79.8 (C2' or C2''), 79.7 (C2' or C2''), 78.1 (C4''), 75.7, 75.6, 74.7, 74.6, 73.3, 73.2 ($6 \times \text{PhCH}_2$), 69.1 (C5''), 68.7 (C5'), 55.1 (MeO' or MeO''), 55.0 (MeO' or MeO''), 47.5 (C6''), 25.3 (C6').

Anal. Calcd for $\text{C}_{57}\text{H}_{62}\text{N}_4\text{O}_{10}$: C, 71.08; H, 6.49; N, 5.82. Found: C, 70.91; H, 6.35; N, 5.72.

ACKNOWLEDGMENTS

The authors thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the fellowships (Maria T. C. Pedrosa, Rosemeire B. Alves and Maria A. F. Prado).

REFERENCES

- Demko, Z.P.; Sharpless, K.B. An intramolecular [2 + 3] cycloaddition route to fused 5-heterosubstituted tetrazoles. *Org. Lett.* **2001**, 3 (25), 4091–4094.
- Demko, Z.P.; Sharpless, K.B. Preparation of 5-substituted *1H*-tetrazoles from nitriles in water. *J. Org. Chem.* **2001**, 66, 7945–7950.
- Singh, H.; Chawla, A.S.; Kapoor, V.K.; Paul, D.; Malhotra, R.K. Medicinal chemistry of tetrazoles. In *Progress in Medicinal Chemistry*; Ellis, G.P., West, G.B., Eds.; North-Holland Biomedical Press: Holland, 1980; Vol. 17, 151–176.
- Butler, R.N. Tetrazoles. In *Comprehensive Heterocyclic Chemistry: The Structure*,

- Reactions, Synthesis and Uses of Heterocyclic Compounds*; Katritzky, A.R., Rees, C.W., Eds.; Imprenta Pergamon Press: Oxford, 1984; Vol. 4, 791–838.
- Koldobskii, G.I.; Ostrovskii, V.A. Tetrazoles. *Russ. Chem. Rev.* **1994**, *63* (10), 797–814.
 - Harfenist, M.; Heuser, D.J.; Joyner, C.T.; Batchelor, J.F.; White, H.L. Selective inhibitors of monoamine oxidase. 3. Structure-activity relationship of tricyclics bearing imidazoline, oxadiazole, or tetrazole groups. *J. Med. Chem.* **1996**, *39* (9), 1857–1863.
 - Ogilvie, W.W.; Yoakim, C.; Dô, F.; Haché, B.; Lagacé, L.; Naud, J.; O'Meara, J.A.; Déziel, R. Synthesis and antiviral activity of monobactams inhibiting the human cytomegalovirus protease. *Bioorg. Med. Chem.* **1999**, *7*, 1521–1531.
 - Genin, M.J.; Allwine, D.A.; Anderson, D.J.; Barbachyn, M.R.; Emmert, D.E.; Garmon, S.A.; Graber, D.R.; Grega, K.C.; Hester, J.B.; Hutchinson, D.K.; Morris, J.; Reischer, R.J.; Ford, C.W.; Zurenko, G.E.; Hamel, J.C.; Schaadt, R.D.; Stapert, D.; Yagi, B.H. Substituent effects on the antibacterial activity of nitrogen-carbon-linked (azolyphenyl)oxazolidinones with expanded activity against the fastidious gram-negative organisms *Haemophilus influenzae* and *Moraxella catarrhalis*. *J. Med. Chem.* **2000**, *43* (5), 953–970.
 - Lüddens, H.; Lang, H.J.; Korpi, E.R. Structure-activity relationship of furosemide-derived compounds as antagonists of cerebellum-specific GABA_A receptors. *Eur. J. Pharm.* **1998**, *344*, 269–277.
 - Middlemiss, D.; Watson, S.P. A medicinal chemistry case study: an account of an angiotensin II antagonist drug discovery programme. *Tetrahedron* **1994**, *50* (46), 13049–13080.
 - Cannizzaro, C.E.; Thiel, I.M.E.; D'Accorso, N.B. Synthesis and debenzoylation products of two perbenzoylated 2-substituted 5-D-galactosyl-1,3,4-oxadiazoles. *J. Heterocyclic Chem.* **1998**, *35* (2), 481–484.
 - Fascio, M.L.; D'Accorso, N.B. Synthesis of some oxadiazole derivatives of D-mannose. *J. Heterocyclic Chem.* **1995**, *32*, 815–818.
 - Martins Alho, M.A.; D'Accorso, N.B. Synthesis and characterization of some heterocyclic derivatives from 1,2:3,4-di-O-isopropylidene- α -D-galacto-1,6-hexadialdo-1,5-pyranose oxime. *J. Heterocyclic Chem.* **1999**, *36*, 177–182.
 - Binatti, I.; Prado, M.A.F.; Alves, R.J.; Souza Filho, J.D. Synthesis of benzomacrolactam by 11-endo selective aryl radical cyclization of 2-iodobenzamide derived from D-galactose. *J. Br. Chem. Soc.* **2002**, *13* (5), 570–575.
 - Prado, M.A.F.; Alves, R.J.; Souza Filho, J.D.; Alves, R.B.; Pedrosa, M.T.C.; Prado, R.F.; Faraco, A.A.G. Synthesis of benzolactams by 11-endo selective aryl radical cyclisation of 2-iodobenzamides. *J. Chem. Soc., Perkin Trans. 1* **2000**, *12*, 1853–1857.
 - Faraco, A.A.G.; Prado, M.A.F.; D'Accorso, N.B.; Alves, R.J.; Souza Filho, J.D.; Prado, R.F. Synthesis and characterization of some nitrogen heterocycles from D-galactose derivatives. *J. Heterocyclic Chem.* **1999**, *36*, 1129–1133.
 - Klier, A.H.; Alves, R.J.; Prado, M.A.F.; Souza Filho, J.D.; D'Accorso, N.B. Synthesis of new five membered nitrogen containing heterocycles bearing D-galactose side chains. *Synth. Commun.* **2000**, *30* (23), 4361–4374.
 - Hall, D.M. A practical synthesis of methyl 4,6-O-benzylidene- α - and - β -D-glucopyranoside. *Carbohydr. Res.* **1980**, *86*, 158–160.



19. Pietraszkiewicz, M.; Jurczak, J. Synthesis of chiral diaza-crown ethers incorporating carbohydrate units. *Tetrahedron* **1984**, *40* (15), 2967–2970.
20. Lipták, A.; Jodál, I.; Nánási, P. Stereoselective ring-cleavage of 3-*O*-benzyl- and 2,3-di-*O*-benzyl-4,6-*O*-benzylidenehexopyranoside derivatives with the LiAlH₄-AlCl₃ reagent. *Carbohydr. Res.* **1975**, *44*, 1–11.
21. Kohn, A.; Schmidt, R.R. Untersuchungen zur synthese von C₁-verzweigten cycliten aus D-glucose. *Liebigs Ann. Chem.* **1987**, *12*, 1045–1054.
22. Sugihara, J.M.; Teerlink, W.J.; MacLeod, R.; Dorrence, S.M.; Springer, C.H. Direct syntheses of some cyano and nitro derivatives of carbohydrates by nucleophilic displacement. *J. Chem. Soc.* **1963**, *28*, 2079–2082.
23. Duncia, J.V.; Pierce, M.E.; Santella, J.B. Three synthetic routes to a sterically hindered tetrazole. A new one-step mild conversion of an amide into a tetrazole. *J. Org. Chem.* **1991**, *56*, 2395–2400.
24. Clapp, L.B. 1,2,3- and 1,2,4-oxadiazoles. In *Comprehensive Heterocyclic Chemistry: The Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds*; Katritzky, A.R., Rees, C.W., Eds.; Imprenta Pergamon Press: Oxford, 1984; Vol. 6, 365–391.
25. Gyöngy, Y.S.; Shim, J.; Yamamoto, Y. Regiospecific synthesis of 2-allylated-5-substituted tetrazoles via palladium-catalyzed reaction of nitriles, trimethylsilyl azide, and allyl acetates. *Tetrahedron Lett.* **2000**, *41*, 4193–4196.

Received April 10, 2003

Accepted July 14, 2003

