

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

Montmorillonite K-10 as a Reusable Catalyst for Fischer Type of Glycosylation under Microwave Irradiation

Dipak K. Roy^a; Manobjyoti Bordoloi^a

^a Natural Products Chemistry Division, North East Institute of Science and Technology, Jorhat, Assam, India

To cite this Article Roy, Dipak K. and Bordoloi, Manobjyoti(2008) 'Montmorillonite K-10 as a Reusable Catalyst for Fischer Type of Glycosylation under Microwave Irradiation', *Journal of Carbohydrate Chemistry*, 27: 5, 300 – 307

To link to this Article: DOI: 10.1080/07328300802107437

URL: <http://dx.doi.org/10.1080/07328300802107437>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Montmorillonite K-10 as a Reusable Catalyst for Fischer Type of Glycosylation under Microwave Irradiation

Dipak K. Roy and Manobjyoti Bordoloi

Natural Products Chemistry Division, North East Institute of Science and Technology, Jorhat, Assam, India

Montmorillonite K10 catalyzed Fischer type glycosylation was studied for various monosaccharides with different alcohols under microwave irradiation. The method was found to be efficient, economic, simple and time saving and the catalyst montmorillonite K-10 was reused three times without loss of catalytic activity and anomeric selectivity. With glycerol, the method gave products glycosylated at primary alcohols only.

Keywords Montmorillonite K10, Glycosylation, glycerol glucoside, microwave radiation


Due to immense biological importance of oligosaccharides and glycoconjugates, many glycosylation strategies have been developed.^[1–3] Among all the existing strategies for glycosylation of sugars, Fischer glycosylation procedure is the earliest and the simplest. In the classical Fischer glycosylation methodology, an unprotected carbohydrate is refluxed in an alcohol in presence of catalytic acid to give the corresponding glycoside. In spite of many limitations, Fischer glycosylation is utilized for synthesis of simple glycosides, such as, methyl, allyl, benzyl and p-methoxybenzyl glycosides.^[4–7] In carbohydrate chemistry, methyl, allyl, benzyl and p-methoxybenzyl alcohols are used for temporary protection of the anomeric hydroxyl group, as glycosides of these alcohols can undergo easy deglycosylation.^[8–10] Modification of Fischer glycosylation by replacing soluble acid with cation exchange resins, namely, Permutit Q, Duolite

Received March 20, 2008; accepted April 2, 2008.

The authors wish to thank the Director, NEIST, Jorhat for provision of facilities and DKR is thankful to the CSIR, New Delhi, for providing research fellowship.

Address correspondence to Manobjyoti Bordoloi, Natural Products Chemistry Division, North East Institute of Science and Technology, Jorhat-785006, Assam, India. E-mail: mjb_rrljt@yahoo.co.in

C-3 and C-3 Zeokarb H, has simplified the removal of the catalyst.^[11] In one report, microwave was used for acceleration of Fischer glycosylation process catalyzed by Amberlite resin IRN 120 H⁺.^[12] Recently, sulfuric acid immobilized on silica was reported as a catalyst for Fischer type glycosylation.^[13] In search of an inexpensive and environmentally benign and time saving method as part of our ongoing programme, we have developed an efficient and economic method of Fischer glycosylation catalysed by montmorillonite K-10 under microwave irradiation (Scheme 1).



Scheme 1.

In the initial experiments for testing the feasibility of montmorillonite K10 as catalysts for Fischer glycosylation, a mixture of D-glucose (500 mg) and the clay (250 mg) was refluxed in anhydrous methanol (10 mL). The reaction was monitored by thin layer chromatography. The conversion was found to be almost complete in 5 hours. The reaction mixture was then cooled, filtered and the product was recrystallized from isopropanol. A mixture of both α and β anomer was obtained. To get the best α/β ratio, the reaction was studied in different reaction times (Table 1). Anomeric ratio was derived from the comparison of ¹H NMR peak at δ 4.70 ($J_{1,2} \approx 4.0$ Hz) and δ 4.27 ($J_{1,2} \approx 8.5$ Hz). Improved α/β ratio was obtained in 10 hrs. Increase in reaction time beyond 10 hours improved α/β ratio slightly with increase in decomposition of the product.

We then studied the conversion under microwave radiation. D-glucose (500 mg) and montmorillonite K10 (250 mg) were mixed thoroughly in a mortar and the mixture was placed in a quartz reaction vessel of ProLabo Synthwave Microwave reactor. To it, 5 mL of methyl alcohol was added and

Table 1: Montmorillonite K10 catalyzed glycosylation of methanol with D-glucose under conventional reflux condition

Entry	Reaction time, hours	Yield, ^a %	α/β Ratio ^b
1	5	83	8.6/1
2	6	84	9.2/1
3	8	84	12.4/1
4	10	84	13.3/1
6	12	81	13.6/1
7	14	79	13.8/1
8	24	75	14.5/1

^aisolated yield. ^bThe α/β ratio was determined by 300 MHz ¹H NMR spectroscopy.

Table 2: Montmorillonite K10 catalyzed glycosylation of methanol with D-glucose under microwave radiation

Entry	Reaction time, min	Yield, ^a %	α/β Ratio ^b
1	2	79	4.7/1
2	4	82	6.2/1
3	6	85	8.6/1
4	8	86	11.3/1
6	10	86	14.6/1
7	12	83	14.7/1
8	14	78	14.8/1
9	20	72	15.1/1

^aisolated yield. ^bThe α/β ratio was determined by 300 MHz ¹H NMR spectroscopy.











the mixture was allowed to react under microwave irradiation at a temperature of 90°C for 10 minutes. During the reaction, the temperature was not allowed to rise above 90°C by setting the programmer. The mixture was then allowed to cool, filtered and the product was recrystallized from isopropanol. To get improved α/β ratio, reaction was carried out for different time periods (Table 2). Optimum anomeric selectivity was found in 10 minutes. Further increase in reaction time increased the α/β ratio only slightly with increased decomposition.

After getting good result in microwave-assisted reaction between D-glucose and methanol, we generalized the reaction with a number of monosaccharides and alcohols (Table 3). Very good yields and anomeric selectivity were obtained. After the reaction, the catalyst could be separated by filtration. After washing and activation at 110°C, the recycled catalyst was used again for glycosylation of D-glucose by benzyl alcohol. Over three times of recycle, comparable yields with same anomeric selectivity were obtained (Table 4).

To study the utility of the method for alcohol containing both primary and secondary hydroxyl groups, we treated D-glucopyranose with glycerol in presence of montmorillonite K-10. *In situ* formed product was acetylated with acetic anhydride. Purification of the product by column chromatography gave 1-(2,3,4,6-Tetraacetyl- α -D-glucopyranosyl)-2,3-diacetylglycerol **3a** (yield 45%), 1-(2,3,4,6-Tetraacetyl- β -D-glucopyranosyl)-2,3-diacetylglycerol **3b** (yield 15%) and 1,3-Bis (2,3,4,6-tetraacetyl-D-glucopyranosyl)-2-acetylglycerol **3c** (yield 6%) with mono- and di-glycosylated product in 5:1 ratio (Scheme 2). Anomeric ratio of the monoglycosylated products was determined to be 3:1. However, as in the glucosidase mediated method of glycosylation of glycerol^[14] or similar to the glycerol glucoside isolated from marine blue-green alga,^[15] and *Lilium japonicum*^[16] no product glycosylated at secondary hydroxyl of glycerol was found to form in our reaction.

Thus, the use of montmorillonite K-10 for this Fischer type glycosylation method, has simplified its work up procedure. Only filtration was required to remove the catalyst. In addition, montmorillonite K-10 is stable, inexpensive,

Table 3: Microwave-assisted glycosylation catalyzed by Montmorillonite K10.

Entry	Sugar	Alcohol	Reaction time (min.)	Yields of Products ^a (%)	α/β Ratio ^b
01		CH ₃ OH	10	86	15/1
02		n-C ₄ H ₉ OH	10	82	10/1
03		PhCH ₂ OH	10	81	9/1
04			10	84	10/1
05		C ₂ H ₅ OH	10	77	13/1
06		n-C ₈ H ₁₇ OH	10	74	10/1
07		CH ₃ OH	10	84	11/1
08		PhCH ₂ OH	10	81	8/1
09			10	83	9/1
10		n-C ₄ H ₉ OH	10	80	10/1
11		CH ₃ OH	10	85	14/1
12		PhCH ₂ OH	10	82	9/1
13			10	84	9/1
14		C ₂ H ₅ OH	10	78	12/1
15		CH ₃ OH	10	87	14/1
16			10	85	11/1
17		PhCH ₂ OH	10	83	9/1
18		CH ₃ OH	10	85	15/1
19			10	85	12/1
20		PhCH ₂ OH	10	80	10/1

a. Isolated yield.


b. The α/β ratio was determined by 300 MHz ¹H NMR spectroscopy.**Scheme 2.**

Table 4: Microwave assisted Glycosylation of benzyl alcohol with D-glucose catalysed by recycled montmorillonite K-10

Number of recycle Ratio ^b (%)	Yield, ^a	α/β
1	79	9/1
2	78	9/1
3	76	9/1

^aisolated yield. ^bThe α/β ratio was determined by 300 MHz ¹H NMR spectroscopy.

eco-friendly and found as reusable catalyst. Another advantage of the method is that it is very fast (only 10 minutes is required). As mentioned above, methyl, allyl and benzyl glycosides are very useful precursor for the synthesis of oligosaccharides and other glycoconjugates.

In conclusion, we have developed an ecofriendly, inexpensive and fast method of Fischer glycosylation catalyzed by montmorillonite K-10 under microwave radiation. One notable advantage of the method is that the catalyst is reusable.

EXPERIMENTAL

General Methods

Melting points were recorded on a Buchi melting point apparatus with open capillary method and are uncorrected. Specific rotations were recorded on a Perkin Elmer polarimeter model 343 at sodium light of wavelength 589 nm. IR spectra were recorded on a Perkin Elmer system 2000 FT-IR Spectrometer. All NMR spectra were recorded on a Bruker Avance DPX300 operated at 300 MHz. Chemical shifts are given in parts per million (ppm) and referenced to internal 0.03% TMS. Positive Electro-Spray Ionization mass spectra [(+)-ESIMS] were recorded on Bruker Esquire 3000 Iontrap Mass spectrometer. Elemental analysis was done on a Perkin Elmer Series II 2400 CHNS/O Micro Elemental Analyzer. Prolabo Synthwave Microwave reactor 402 with temperature programmer was used for reactions under microwave irradiation. Buchi rotary evaporator was used for distillation under reduced pressure. Silica gel G was used for TLC and silica gel (60–120 mesh) was used for column chromatography. All solvents used were distilled prior to use.

General Procedure for Glycosylation under Microwave Irradiation

Monosaccharide (500 mg) and Montmorillonite K10 (250 mg) were mixed thoroughly in a mortar. The mixture was then placed in a quartz reaction vessel of Prolabo Synthwave Microwave reactor 402. To it, 5 mL of an alcohol was added and the mixture was allowed to react under microwave irradiation at a

temperature of 90°C for 10 minutes. During the reaction the temperature was not allowed to rise above 90°C by setting the programmer. The reaction mixture was then cooled and filtered. The residue was washed with methanol. The combined filtrate and washings were distilled under reduced pressure with a rotary evaporator to get the product. Product was crystallized from isopropanol and characterized by ¹H NMR (300 MHz), ¹³C NMR (75 MHz), ESIMS, IR spectroscopy and elemental analysis and direct comparison with that of authenticated compound.

Procedure for Reaction between D-Glucopyranose and Glycerol

D-glucopyranose (180 mg, 1 mmol), glycerol (920 mg, 10 mmol) and montmorillonite K-10 (90 mg) were mixed thoroughly in a mortar and then the mixture was placed in a quartz reaction vessel of Prolabo Synthwave Microwave reactor 402. Then the mixture was allowed to react under microwave irradiation at a temperature of 90°C for 10 minutes. During the reaction, the temperature was not allowed to rise above 90°C by setting the programmer. After cooling the reaction mixture to room temperature, 5 mL of acetic anhydride was added to it and stirred for 3 hours. Chloroform (100 mL) was then added to it and filtered. Solid was washed with chloroform (10 mL). The combined filtrate and washings were washed with water (3 × 10 mL) and dried over anhydrous sodium sulfate. Solvent was distilled under reduced pressure to obtain a crude, which was chromatographed over a column of silica gel using ethyl acetate-hexane (1:3) to get two major product **3a** (227 mg, 45%) and **3b** (75 mg, 15%) and a minor product **3c** (48 mg, 6%).

*Spectral data of 1-(2,3,4,6-Tetraacetyl- α -D-glucopyranosyl)-2,3-diacetyl glycerol **3a***

Gummy, $[\alpha]_D^{32} + 169.05^\circ$ (CHCl₃, c 0.5); IR (CHCl₃)_v cm⁻¹: 2958, 2850, 1747, 1435, 1371, 1226, 1167, 1040, 957, 889, 768, 601; ¹H NMR (300 MHz) in CDCl₃: δ 5.43 (dt, 1H, J_{1'b,2'} = J_{2'3'a} = 9.6 Hz, J_{1'a,2'} = 1.2 Hz, H-2'), 5.17 (dd, 1H, J_{3,4} = 10 Hz, J_{4,5} = 8.4 Hz, H-4), 5.09 (d, 1H, J_{1,2} = 3.2 Hz, H-1), 5.04 (t, 1H, J_{2,3} = J_{3,4} = 10 Hz, H-3), 4.84 (dd, 1H, J_{2,3} = 10.0 Hz, J_{1,2} = 3.2 Hz, H-2) 4.32 (m, 1H, H-3'.a), 4.25 (m, 1H, H-3'.b), 4.22 (dd, 1H, J_{6a,6b} = 11 Hz, J_{5,6a} = 2.4 Hz, H-6a), 4.12 (dd, 1H, J_{6a,6b} = 11 Hz, J_{5,6b} = 4.8 Hz, H-6b), 4.02 (m, 1H, H-1'a), 3.81 (m, 1H, H-1'b), 3.62 (ddd, 1H, J_{4,5} = 8.4 Hz, J_{5,6a} = 2.4 Hz, J_{5,6b} = 4.8 Hz, H-5) 2.09 (s, 3H, CH₃-CO), 2.08 (s, 3H, CH₃-CO), 2.07 (s, 3H, CH₃-CO), 2.06 (s, 3H, CH₃-CO), 2.03 (s, 3H, CH₃-CO), 2.01 (s, 3H, CH₃-CO); ¹³C NMR (75 MHz) in CDCl₃: δ 169.96 (C=O), 169.95 (C=O), 169.92 (C=O), 169.81 (C=O), 169.72 (C=O), 169.59 (C=O), 94.95 (C-1), 70.53 (C-2'), 69.24 (C-2), 68.61 (C-1'), 67.25 (C-3'), 65.85 (C-3), 65.73 (C-5), 62.59 (C-4), 61.82 (C-6), 21.80 (CH₃-CO), 20.75 (CH₃-CO), 20.73 (CH₃-CO), 20.65 (CH₃-CO), 20.55 (CH₃-CO), 21.41 (CH₃-CO); (+)-ESIMS for C₂₁H₃₀O₁₄ (M, 506): m/z 529.13 [M+Na]⁺, 507.16 [M+H]⁺, Anal, calcd. for C₂₁H₃₀O₁₄: C, 49.80%; H, 5.97%. Found C, 49.70%; H, 5.84%.

Spectral data of 1-(2,3,4,6-Tetraacetyl-β-D-glucopyranosyl)-2,3-diacetylglycerol 3b

mp 106–107°C (reported^[16] 107–108°C); $[\alpha]_D^{32}$ –35.1° (CHCl₃, c 0.5); IR (CHCl₃) $\nu_{\text{cm}^{-1}}$: 2952, 2856, 1741, 1435, 1366, 1226, 1161, 1045, 955, 888, 757, 607; ¹H NMR (300 MHz) in CDCl₃: δ 5.18 (m, 1H, H-2'), 5.17 (dd, 1H, $J_{2,3}$ = 9.9 Hz, $J_{3,4}$ = 9.6 Hz, H-3), 5.07 (dd, 1H, $J_{2,3}$ = 9.8 Hz, $J_{1,2}$ = 7.2 Hz, H-2), 4.98 (d, 1H, $J_{4,5}$ = 9.2 Hz, $J_{3,4}$ = 9.6 Hz, H-4), 4.52 (d, 1H, J = 7.2 Hz, H-1), 4.32 (m, 2H, H-3'), 4.25 (dd, 1H, $J_{6a,6b}$ = 12.3 Hz, $J_{5,6a}$ = 4.6 Hz, H-6a), 4.12 (dd, 1H, $J_{6a,6b}$ = 12.2 Hz, $J_{5,6b}$ = 2.6 Hz, H-6b), 3.92 (ddd, 1H, $J_{4,5}$ = 9.2 Hz, $J_{5,6a}$ = 4.7 Hz, $J_{5,6b}$ = 2.6 Hz, H-5), 3.68 (m, 2H, H-1'), 2.10 (s, 3H, CH₃-CO), 2.09 (s, 3H, CH₃-CO), 2.085 (s, 3H, CH₃-CO), 2.08 (s, 3H, CH₃-CO), 2.03 (s, 3H, CH₃-CO), 2.01 (s, 3H, CH₃-CO). ¹³C NMR (75 MHz) in CDCl₃: δ 170.12 (C=O), 170.0 (C=O), 169.90 (C=O), 169.83 (C=O), 169.75 (C=O), 169.59 (CO), 95.57 (C-1), 72.96 (C-2), 71.88 (C-1'), 70.53 (C-3'), 70.52 (C-2'), 68.61 (C-4), 65.73 (C-3), 62.59 (C-5), 61.82 (C-6), 21.41 (CH₃-CO), 20.80 (CH₃-CO), 20.78 (CH₃-CO), 20.74 (CH₃-CO), 20.67 (CH₃-CO), 20.52 (CH₃-CO); (+)-ESIMS for C₂₁H₃₀O₁₄ (M, 506): m/z 531.15 [M+2H+Na]⁺, 507.16 [M+H]⁺, Anal, calcd. for C₂₁H₃₀O₁₄: C, 49.80%; H, 5.97%. Found C, 49.67%; H, 5.81%.

Spectral data of 1,3-Bis (2,3,4,6-tetraacetyl-D-glucopyranosyl)-2-acetylglycerol 3c

Gummy; $[\alpha]_D^{32}$ +17.1° (CHCl₃, c 0.5); IR (CHCl₃) $\nu_{\text{cm}^{-1}}$: 2980, 2850, 1715, 1410, 1250; ¹H NMR (300 MHz) in CDCl₃: δ 5.46 (m, 1H, H-2'), 5.30 (m, 1H), 5.22–5.01 (m, 5H), 4.33–4.08 (m, 8H), 3.85–3.54 (m, 4H), 2.13 (s, 3H, CH₃-CO), 2.085 (s, 3H, CH₃-CO), 2.10 (s, 3H, CH₃-CO), 2.096 (s, 3H, CH₃-CO), 2.08 (s, 3H, CH₃-CO), 2.06 (s, 3H, CH₃-CO), 2.03 (s, 3H, CH₃-CO), 2.02 (s, 3H, CH₃-CO), 2.01 (s, 3H, CH₃-CO); ¹³C NMR (75 MHz) in CDCl₃: δ 170, 169.95, 169.33, 169.50 (9 C=O), 100.10 (C-1, C-1'), 71.01 (C-1'' and C-3''), 70.80 (C-2''), 69.63, 69.15, 68.33, 68.12 (6>CHO-Ac), 64.07 (C-5, C-5'), 61.40 (2-CH₂OAc, C-6, C-6'), 21.41, 20.85, 20.78, 20.71, 20.53 (9 CH₃-CO); (+)-ESIMS for C₃₃H₄₆O₂₂ (M, 794): m/z 817.24 [M+Na]⁺, 795.26 [M+H]⁺; Anal, calcd for C₃₃H₄₆O₂₂: C, 49.87%; H, 5.83%. Found : C, 49.99%; H, 5.70%.

REFERENCES

- Schmidt, R.R. New Methods for the Synthesis of Glycosides and Oligosaccharides—Are There Alternatives to the Koenigs-Knorr Method? *Angew. Chem., Int. Ed.* **1986**, *25*, 212.
- Schmidt, R.R. *Comprehensive Organic Synthesis*. Pergamon press, **1991**, *6*, 33–64.
- El Ashry, E.S.H.; Awad, L.F.; Atta, A.I. Synthesis and role of glycosylthio heterocycles in carbohydrate chemistry. *Tetrahedron*, **2006** *62*, 2943–2998.
- Capon, B. Mechanism in carbohydrate chemistry. *Chem. Rev.* **1969**, *69*, 407–498.

5. Ferrier, R.J.; Hatton, L.R. Studies with radioactive sugars : Part I. Aspects of the alcoholysis of D-xylose and D-glucose; the role of the acyclic acetals. *Carbohydr. Res.* **1968**, *6*, 75–86.
6. Heard, D.D.; Barker, R. Investigation of the role of dimethyl acetals in the formation of methyl glycosides. *J. Org. Chem.* **1968**, *33*, 740–746.
7. Kappe, C.O. Controlled microwave heating in modern organic synthesis. *Angew. Chem., Int. Ed.* **2004**, *43*, 6250–6284.
8. Mukhopadhyay, B.; Field, R.A. Synthesis of L-arabinose-containing fragments of the oat root saponin Avenacin A-I. *Carbohydr. Res.* **2004**, *339*, 1285–1291.
9. Kim, I.J.; Park, T.K.; Hu, S.; Abrampah, K.; Zhang, S.; Livingston, P.; Danishefsky, S.J. Defining the molecular recognition of MBrl (human breast cancer) antigen by the MBrl antibody through probe structures prepared by total synthesis. *J. Org. Chem.* **1995**, *60*, 7716–7717.
10. Sarabia, F.; Martí n-Ortiz, L.; López-Herrera, F.J. Synthetic studies towards the tunicamycins and analogues based on diazo chemistry. Total synthesis of tunicaminy uracil. *Org. Biomol. Chem.* **2003**, *21*, 3716–3725.
11. Dean, G.R.; Pyle, R.E. Preparation of methyl glucoside, U.S. Patent 2,606,186 (1952).
12. Bornaghi, L.F.; Poulsen, S.T. Microwave-accelerated Fischer glycosylation. *Tetrahedron Lett.* **2005**, *46*, 3485–3488.
13. Roy, B.; Mukhopadhyay, B. Sulfuric acid immobilized on silica: an excellent catalyst for Fischer type glycosylation. *Tetrahedron Lett.* **2007**, *48*, 3783–3487
14. Matheus de Roode, B.; Peters, S.W.P.G.; Franssen, M.C.R.; van der Padt, A.; de Groot, A.; Boom, R.M. Optimization of production and downstream processing of the almond (3-glucosidase-mediated glycosylation of glycerol. *Biotech & Bioeng.* **2001**, *72*(5), 568–572.
15. Choi, J.S.; Lee, D.; Lee, W.K.; Kim, D.; Choi, H.D.; Son, B.W. 2-O-(α -D-Glucopyranosyl)glycerol, A New Glycerol Glycoside from the Marine Blue-green Alga *Oscillatoria* sp. *Bull. Korean Chem. Soc.* **1999**, *20*(10), 1121–1122.
16. Kaneda, M.; Kobayashi, K.; Nishida, K.; Katsuta, S. Lilioides D and E, two glycerol glucosides from *Lilium japonicum*. *Phytochemistry*, **1984**, *23*(4), 795–798.