

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

First Synthesis of Thienopyrazole Thioglycosides

Galal H. Elgemeie^a; Wafaa A. Zaghary^b; Tamer M. Nasr^b; Kamelia M. Amin^c

^a Faculty of Science, Department of Chemistry, Helwan University, Cairo, Egypt ^b Faculty of Pharmacy, Pharmaceutical Chemistry Department, Helwan University, Cairo, Egypt ^c Faculty of Pharmacy, Pharmaceutical Chemistry Department, Cairo University, Cairo, Egypt

To cite this Article Elgemeie, Galal H. , Zaghary, Wafaa A. , Nasr, Tamer M. and Amin, Kamelia M.(2008) 'First Synthesis of Thienopyrazole Thioglycosides', *Journal of Carbohydrate Chemistry*, 27: 6, 345 — 356

To link to this Article: DOI: 10.1080/07328300802236178

URL: <http://dx.doi.org/10.1080/07328300802236178>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

First Synthesis of Thienopyrazole Thioglycosides

Galal H. Elgemeie,¹ Wafaa A. Zaghary,² Tamer M. Nasr,² and Kamelia M. Amin³

¹Faculty of Science, Department of Chemistry, Helwan University, Cairo, Egypt

²Faculty of Pharmacy, Pharmaceutical Chemistry Department, Helwan University, Cairo, Egypt

³Faculty of Pharmacy, Pharmaceutical Chemistry Department, Cairo University, Cairo, Egypt

Reported is the first method to prepare a new class of thienopyrazole thioglycosides via a one-pot reaction of the sodium thienopyrazolthiolate salts with 2,3,4,6-tetra-*O*-acetyl- α -D-gluco- and galactopyranosyl bromides. The sodium thienopyrazolthiolate salts are prepared using pyrazoldithioic acids and their corresponding mono- and dithiolate salts.

Keywords Thienopyrazole thioglycosides, Sodium thienopyrazolthiolate salts, Pyrazoldithioic acids


There is a great interest in the synthesis of nucleosides due to their incorporation into DNA, RNA, protein, and all intracellular essential nucleotides. Chemical modification of nucleosides is an important approach to fight various infections and cancer. In recent reports from our laboratory, we described the preparation of different novel functionalized pyrimidine and pyridine thioglycosides, which revealed antagonistic activity against human carcinoma cells and HIV.^[1–3] Earlier, we had reported the use of dihydropyridinethione glycosides as P-glycoprotein (Pgp) substrates or inhibitors of protein glycosylation.^[4] These common features encouraged us to develop a new straightforward route for the synthesis of these compounds. Here, we describe the synthesis of thienopyrazole thioglycosides through reaction of sodium thienopyrazolthiolates with α -halogeno sugars. To our knowledge, this is the first thienopyrazole thioglycoside ring system to be reported. The

Received March 20, 2008; accepted May 29, 2008.

Address correspondence to Galal H. Elgemeie, Faculty of Science, Department of Chemistry, Helwan University, Ain-Helwan, Cairo, Egypt. E-mail: elgemeie@yahoo.com

sequence of reactions followed in the preparation of the compounds is summarized in Charts 1 through 3. Thus, it has been found that pyrazolin-5-one **1** reacted with carbon disulfide in the presence of sodium ethoxide to afford the sodium dithiolate salts **2**. Compounds **2** were readily monoalkylated to give the stable sodium salts of monoalkylthio derivatives. Thus, one equivalent of phenacyl bromide or methyl iodide gave the corresponding sodium salts of monoalkylated products **3** or **4**, respectively. When neutralized with hydrochloric acid **3** and **4** gave the novel thiol products **8** and **9**. Compounds **3** and **4** reacted with 2,3,4,6-tetra-*O*-acetyl- α -D-gluco- or galactopyranosyl bromides **5a,b** in ethanol at rt to give the corresponding *S*-glycosides **6** and **7** in high yields. These compounds were also prepared by the reaction of the thiols **8** and **9** with 2,3,4,6-tetra-*O*-acetyl- α -D-gluco- or galacto-pyranosyl bromides **5** in KOH-acetone at rt for 12 h. The IR spectrum of **6a** revealed the presence of NH band at 3300 cm^{-1} , the mass spectrum showed a molecular formula $\text{C}_{27}\text{H}_{30}\text{N}_2\text{O}_{11}\text{S}_2$ ($M^+ = 622$), and the ^1H NMR spectrum indicated the anomeric proton as a doublet at δ 6.05 ppm with spin-spin coupling constant of 10.11 Hz, while the other six glucose protons resonated at δ 3.95 to 5.50 ppm. The four acetyl groups appeared as four singlets at δ 1.95 to 2.00 ppm. Furthermore, the ^{13}C NMR spectrum of **6a** represented a signal at δ 80.00 ppm corresponding to the C-1' atom of the β -configuration.


When compounds **2a,b** (Chart 2) were subjected to monoalkylation with 2,3,4,6-tetra-*O*-acetyl- α -D-gluco- and galactopyranosyl bromides **5** they yielded compounds **8** that, upon neutralization to ph 7, yielded the glycoside thiols **10**. These compounds could also be prepared by alkylation of the dithiol derivatives **9** (obtained by neutralization of compound **2**) with one equivalent of **5**. The structures of compounds **10** were established on the basis of their elemental analyses, MS, IR, and ^1H NMR spectroscopies. The mass spectrum of **10b** was compatible with the molecular formula $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_{10}\text{S}_2$ ($M^+ = 580$). The ^1H NMR spectrum indicated the anomeric proton as a doublet at $\delta = 5.95$ ppm with spin-spin coupling constant of 10.11 Hz, while the other six galactose protons resonated at $\delta = 4.00$ to 5.40 ppm. The acetyl groups appeared as four singlets at $\delta = 1.90$ to 2.15 ppm. The synthetic pathways adopted for the preparation of our target thienopyrazole thioglycosides are illustrated in Chart 3. Compound **3** was cyclized in refluxing sodium ethoxide to give the corresponding sodium thieno[3,4-*c*]pyrazolethiolate **11** that upon alkylation with halosugars **5** yielded the corresponding 2-(glycopyranosylthio)thienopyrazole derivatives **12**. When compound **3** was cyclized in refluxing sodium ethoxide followed by acidification it afforded the thienopyrazole-4-thiol **13**, which was coupled with halosugars **5** in KOH-acetone to yield also compounds **12**. The structures of the reaction products **12** were established and confirmed on the basis of their elemental analyses and spectral data (MS, IR, $^1\text{HNMR}$, $^{13}\text{CNMR}$). Thus, the mass spectrum of **12a** revealed a molecular formula of $\text{C}_{27}\text{H}_{28}\text{N}_2\text{O}_{10}\text{S}_2$ ($M^+ = 604$). The



1-9 X = H

<u>6,7</u>	<u>X</u>	<u>R₁</u>	<u>R₂</u>
a,	H	H	OAc
b,	H	OAc	H

Chart 1




<u>2,9</u>	<u>X</u>
a,	H
b,	Ph

<u>8,10</u>	<u>X</u>	<u>R₁</u>	<u>R₂</u>
a,	H	H	OAc
b,	H	OAc	H
c,	Ph	H	OAc
d,	Ph	OAc	H

Chart 2

¹H NMR spectrum for **12b** showed the anomeric proton as a doublet at δ 5.90 ppm with a spin-spin coupling constant of 9.8 Hz, while the other six galactose protons resonated at δ 4.05 to 5.40 ppm and the four acetyl groups appeared as four singlets at δ 1.80 to 2.00 ppm.



<u>12</u>	<u>X</u>	<u>R₁</u>	<u>R₂</u>
a,	H	H	OAc
b,	H	OAc	H

Chart 3

EXPERIMENTAL

All melting points were uncorrected on a Gallenkamp melting point apparatus. The IR spectra were recorded (KBr disk) on a Perkin Elmer 1650 FT-IR instrument. The ^1H NMR spectra were measured on a Varian 400 MHz spectrometer for solution $(\text{CD}_3)_2\text{SO}$ using $\text{Si}(\text{CH}_3)_4$ as an internal standard. Mass spectra were recorded on a Varian MAT 112 spectrometer. Elemental analyses were obtained from the Microanalytical Data Center at Cairo University, Egypt.

Progress of the reactions was monitored by TLC using aluminum sheets coated with silica gel F254 (Merck). Viewing under a short-wavelength UV lamp effected detection. All evaporations were carried out under reduced pressure at 40°C.

Sodium 5-methylpyrazol-3-one-4-methylenedithiolates (2a,b)

General Procedure

A solution of pyrazolones **1a,b** (0.01 mol) and sodium ethoxide (0.46 g, 0.02 mol) in absolute ethanol (20 mL) was refluxed for 20 min. The reaction mixture was cooled to rt; carbon disulphide (0.01 mol) was added. The reaction mixture was stirred at rt for 30 min, and then the solution was evaporated and the formed solid product was collected by filtration.

Sodium 5-methylpyrazol-3-one-4-((2-oxo-2-phenylethyl)thio)methyl-enethiolates (3a,b)

General Procedure

A solution of compounds **2a,b** (0.01 mol) and phenacyl bromide (1.98 g, 0.01 mol) in ethanol (20 mL) was stirred at rt for 2 h, the solution was evaporated, and the formed solid product was collected and recrystallized from methanol.

Sodium 5-methylpyrazol-3-one-4-(methylthio)methylenethiolates (4a,b)

General Procedure

A solution of compounds **2a,b** (0.01 mol) and methyl iodide (0.01 mol) in ethanol (20 mL) was stirred at rt for 2 h, the solution was evaporated, and the formed solid product was collected by filtration.

(4Z)-4-(mercapto((2-oxo-2-phenylethyl)thio)methylene)-5-methyl-pyrazol-3-ones (8a,b)

General Procedure

A solution of compounds **3a,b** in ethanol (20 mL) was poured on cold water and treated with hydrochloric acid until just acidic and the formed solid products **8a,b** were collected by filtration and recrystallized from ethanol.

4-(Mercapto(methylthio)methylene)-5-methylpyrazol-3-ones (8a,b)

General Procedure

A solution of compounds **4a,b** in ethanol (20 mL) was poured on cold water and treated with hydrochloric acid until just acidic and the formed solid products **9a,b** were collected by filtration and recrystallized from ethanol.

(4E)-5-methyl-4-[(2-oxo-2-phenylethylthio)(2',3',4',6'-tetra-O-acetyl-β-D-glucopyranosylthio)methylene]-2,4-dihydro-3H-pyrazol-3-ones (6a,b)

General Procedure

Method A

To a solution of compound **3a** (3.14 g, 0.01 mol) in ethanol (30 mL), a solution of 2,3,4,6-tetra-O-acetyl-α-D-glucopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (20 mL) was added. The reaction mixture was stirred at rt until completion (TLC, 16 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the sodium bromide formed. The resulting product was recrystallized from ethanol.

Method B

A solution of compound **8a** (2.92 g, 0.01 mol) in aqueous potassium hydroxide [0.56 g (0.01 mol) in distilled water (6 mL)] was added to a solution of 2,3,4,6-tetra-O-acetyl-α-D-glucopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (30 mL). The reaction mixture was stirred at rt until completion (TLC, 16 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the formed potassium bromide salt. The resulting product was recrystallized from ethanol.

6a: brown, m.p. 213°C (from EtOH), yield (65% method A, 75% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3300 (NH), 1750 (CO), 1690 (CO). ^1H NMR (DMSO) δ 1.95–2.00 (4s, 12H, 4 × CH₃CO), 2.40 (s, 3H, CH₃), 2.90 (s, 1H, OH), 3.95–4.15 (m, 3H, 6'-H₂, 5'-H), 4.25 (t, 1H, 4'-H), 4.90 (s, 2H, CH₂), 5.10 (t, 1H, 3'-H), 5.50 (t, 1H, 2'-H), 6.05 (d, 1H, 1'-H), 7.50–8.10 (m, 5H, C₆H₅). ^{13}C NMR (DMSO) δ 14.93 (Ar-CH₃), 20.72–20.87 (4 × CH₃), 42.35 (SCH₂), 61.83 (CH₂, C-6'), 66.42 (C-4'), 68.26 (C-2'), 71.05 (C-3'), 71.52 (C-5'), 72.47 (C-1'), 129.00–156.00 (aromatic carbons), 160.00 (=C-S), 165.00–185.00 (6 × CO). C₂₇H₃₀N₂O₁₁S₂. Calcd: C, 52.07; H, 4.86; N, 4.50. Found: C, 52.29; H, 4.83; N, 4.85%.

6b: Brown, m.p. 230°C (from EtOH), yield (68% method A, 76% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3200 (NH), 1751 (CO), 1681 (CO). ^1H NMR (DMSO) δ 1.90–2.20 (4s, 12H, 4 × CH₃CO), 2.45 (s, 3H, CH₃), 4.00–4.10 (m, 3H,

6'-H₂, 5'-H), 4.45 (t, 1H, 4'-H), 4.95 (s, 2H, CH₂), 5.25–5.45 (m, 2H, 3'-H, 2'-H), 6.00 (d, 1H, 1'-H), 7.40–8.10 (m, 5H, C₆H₅). ¹³C NMR (DMSO) δ 14.93 (Ar-CH₃), 20.70–20.87 (4 × CH₃), 42.36 (SCH₂), 61.41 (CH₂, C-6'), 67.65 (C-4'), 68.60 (C-2'), 70.63 (C-3'), 70.81 (C-5'), 80.00 (C-1'), 128.00–146.00 (aromatic carbons), 160.15 (=C-S), 165.21–183.00 (6 × CO). C₂₇H₃₀N₂O₁₁S₂, Calcd: C, 52.07; H, 4.86; N, 4.50. Found: C, 51.85; H, 5.03; N, 4.51%.

5-Methyl-4-((methylthio)(2',3',4',6'-tetra-O-acetyl-β-D-glucopyranosylthio)methylene)-2,4-dihydro-3H-pyrazol-3-ones (7a,b)

General Procedure

Method A

To a solution of compound **4a** (2.10 g, 0.01 mol) in ethanol (30 mL), a solution of 2,3,4,6-tetra-O-acetyl-α-D-glucopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (20 mL) was added. The reaction mixture was stirred at rt until completion (TLC, 13 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the sodium bromide formed. The resulting product was recrystallized from ethanol.

Method B

A solution of compounds **9a** (1.88 g, 0.01 mol) in aqueous potassium hydroxide [0.56 g (0.01 mol) in distilled water (6 mL)] was added to a solution of 2,3,4,6-tetra-O-acetyl-α-D-glucopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (30 mL). The reaction mixture was stirred at rt until completion (TLC, 13 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the formed potassium bromide. The resulting product was recrystallized from ethanol.

7a: yellow, m.p. 192°C (from EtOH), yield (70% method A, 78% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3371 (NH), 1751 (CO). ¹H NMR (DMSO) δ 1.90–2.10 (4s, 12H, 4 × CH₃CO), 2.45 (s, 1H, OH), 2.60 (s, 3H, CH₃), 2.65 (s, 3H, SCH₃), 3.90–4.05 (m, 3H, 6'-H₂, 5'-H), 4.20 (t, 1H, 4'-H), 5.05 (t, 1H, 3'-H), 5.45 (t, 1H, 2'-H), 6.00 (d, 1H, 1'-H). ¹³C NMR (DMSO) δ 13.92 (Ar-CH₃), 15.66 (S-CH₃), 18.83–20.84 (4 × CH₃), 59.91 (CH₂, C-6'), 68.48 (C-4'), 71.10 (C-2'), 71.48 (C-3'), 73.01 (C-5'), 74.53 (C-1'), 113.00 (C-4), 143.74 (C-5), 144.68 (C-3 enol form), 160.15 (=C-S), 169.21–169.52 (4 × CO). C₂₀H₂₆N₂O₁₀S₂, Calcd: C, 46.31; H, 5.06; N, 5.40. Found: C, 46.70; H, 5.14; N, 5.42%.

7b: Yellow, m.p. 145°C (from EtOH), yield (69% method A, 77% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3340 (NH), 1751 (CO). ¹H NMR (DMSO) δ 1.90–2.15 (4s, 12H, 4 × CH₃CO), 2.45 (s, 1H, OH), 2.60 (s, 3H, CH₃), 2.65 (s, 3H, SCH₃), 4.05–4.15 (m, 3H, 6'-H₂, 5'-H), 4.45 (t, 1H, 4'-H), 5.30–5.45 (m, 2H, 3'-H, 2'-H), 5.90 (d, 1H, 1'-H). ¹³C NMR (DMSO) δ 14.52 (Ar-CH₃), 18.05 (S-CH₃), 18.87–

20.85 (4 × CH₃), 61.39 (CH₂, C-6'), 64.89 (C-4'), 67.42 (C-2'), 68.87 (C-3'), 70.97 (C-5'), 71.08 (C-1'), 113.81 (C-4), 144.63 (C-5), 160.00 (=C-S), 165.00 (NHCO), 169.24–170.32 (4 × CO). C₂₀H₂₆N₂O₁₀S₂, Calcd: C, 46.31; H, 5.06; N, 5.40. Found: C, 46.38; H, 5.31; N, 5.85%.

Sodium 5-methyl-3-one-pyrazole-4-(2',3',4',6'-tetra-O-acetyl-β-D-gluco or galactopyranosylthio)methylenethiolates (8a–d)

General Procedure

To a solution of compounds **2a,b** (0.01 mol) in ethanol (30 mL), a solution of 2,3,4,6-tetra-O-acetyl-α-D-gluco or galactopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (20 mL) was added. The reaction mixture was stirred at rt until completion (TLC, 12 h) and then evaporated under reduced pressure, and the product was separated by filtration.

8a: yellow, m.p. >300°C (from EtOH), yield (74%). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3417 (NH), 1751 (CO), 1596 (C=C). C₁₉H₂₃N₂O₁₀S₂Na.

8b: yellow, m.p. >300°C (from EtOH), yield (73%). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3417 (NH), 1743 (CO), 1596 (C=C). C₁₉H₂₃N₂O₁₀S₂Na.

8c: yellow, m.p. >300°C (from EtOH), yield (82%). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 1751 (CO), 1612 (C=C). C₂₅H₂₇N₂O₁₀S₂Na.

8d: yellow, m.p. >300°C (from EtOH), yield (81%). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 1743 (CO), 1620 (C=C). C₂₅H₂₇N₂O₁₀S₂Na.

4-(Dimercaptomethylene)-5-methyl-pyrazol-3-ones (9a,b)

General Procedure

A solution of compound **2a,b** in ethanol (20 mL) was poured on cold water and treated with hydrochloric acid until just acidic and the formed solid product **9a,b** was collected by filtration and recrystallized from ethanol.

4-(Mercapto(2',3',4',6'-tetra-O-acetyl-β-D-gluco or galactopyrano-sylthio)methylene)-5-methyl-pyrazol-3-ones (10a–d)

General procedures

Method A

A solution of compounds **8a–d** (0.01 mol) in ethanol (30 mL) was poured on ice-cooled water and the medium was adjusted to pH 7 using dilute acetic acid,

and the formed solid product **10a–d** was collected by filtration and recrystallized from ethanol.

Method B

A solution of compounds **9a,b** (0.01 mol) in aqueous potassium hydroxide [0.56 g (0.01 mol) in distilled water (6 mL)] was added to a solution of 2,3,4,6-tetra-*O*-acetyl- α -D-gluco- or galactopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (30 mL). The reaction mixture was stirred at rt until completion (TLC, 15 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the formed potassium bromide. The resulting product was recrystallized from ethanol.

10a: yellow, m.p. 110°C (from EtOH), yield (70% method A, 77% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3200 (NH), 2507 (SH), 1751 (CO). ^1H NMR (DMSO) δ 1.95–2.00 (4s, 12H, 4 \times CH₃CO), 2.45 (s, 3H, CH₃), 4.00–4.20 (m, 3H, 6'-H₂, 5'-H), 4.95 (t, 1H, 4'-H), 5.20 (t, 1H, 3'-H), 5.40 (t, 1H, 2'-H), 5.95 (d, 1H, 1'-H). ^{13}C NMR (DMSO) δ 15.92 (Ar-CH₃), 20.65–20.87 (4 \times CH₃), 61.97 (CH₂, C-6'), 68.37 (C-4'), 68.81 (C-2'), 73.92 (C-3'), 75.25 (C-5'), 81.91 (C-1'), 116.00 (C-4), 147.00 (C-5), 160.10 (C-S), 165.00 (NHCO), 169.45–170.29 (4 \times CO). C₁₉H₂₄N₂O₁₀S₂, Calcd: C, 45.22; H, 4.80; N, 5.55. Found: C, 45.00; H, 4.76; N, 5.35%.

10b: Yellow, m.p. 155°C (from EtOH), yield (73% method A, 80% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3209 (NH), 2550 (SH), 1751 (CO). ^1H NMR (DMSO) δ 1.90–2.15 (4s, 12H, 4 \times CH₃CO), 2.45 (s, 3H, CH₃), 4.00–4.10 (m, 3H, 6'-H₂, 5'-H), 4.30 (t, 1H, 4'-H), 5.30–5.40 (m, 2H, 3'-H, 2'-H), 5.95 (d, 1H, 1'-H). ^{13}C NMR (DMSO) δ 15.91 (Ar-CH₃), 20.69–20.89 (4 \times CH₃), 61.94 (CH₂, C-6'), 66.22 (C-4'), 68.01 (C-2'), 71.73 (C-3'), 74.33 (C-5'), 82.28 (C-1'), 115.00 (C-4), 145.00 (C-5), 160.15 (C-S), 165.00 (NHCO), 169.66–170.33 (4 \times CO). C₁₉H₂₄N₂O₁₀S₂, Calcd: C, 45.22; H, 4.80; N, 5.55. Found: C, 45.21; H, 4.59; N, 5.51%.

10c: Yellow, m.p. 165°C (from EtOH), yield (82% method A, 88% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3020 (CH, aromatic), 2939 (CH, aliphatic), 2507 (SH), 1751 (CO). ^1H NMR (DMSO) δ 1.90–2.00 (4s, 12H, 4 \times CH₃CO), 2.55 (s, 3H, CH₃), 4.00 (s, 2H, 6'-H), 4.15 (m, 1H, 5'-H), 4.95 (t, 1H, 4'-H), 5.20 (t, 1H, 3'-H), 5.35 (t, 1H, 2'-H), 6.10 (d, 1H, 1'-H), 7.20–7.80 (m, 5H, C₆H₅). ^{13}C NMR (DMSO) δ 18.21 (Ar-CH₃), 20.67–20.89 (4 \times CH₃), 62.04 (CH₂, C-6'), 68.49 (C-4'), 68.92 (C-2'), 74.27 (C-3'), 75.14 (C-5'), 81.30 (C-1'), 120.12–150.00 (aromatic and heteroaromatic carbons), 159.00 (C-S), 162.00 (NHCO), 169.38–169.65 (4 \times CO). C₂₅H₂₈N₂O₁₀S₂, Calcd: C, 51.70; H, 4.82; N, 4.87. Found: C, 51.99; H, 5.26; N, 4.47%.

10d: Yellow, m.p. 108°C (from EtOH), yield (81% method A, 87% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3020 (CH, aromatic), 2962 (CH, aliphatic), 2507 (SH), 1751 (CO). ^1H NMR (DMSO) δ 1.90–2.10 (4s, 12H, 4 \times CH₃CO), 2.60 (s, 3H, CH₃),

4.00 (s, 2H, 6'-H), 4.30 (m, 1H, 5'-H), 5.00 (t, 1H, 4'-H), 5.35 (m, 2H, 3'-H, 2'-H), 6.05 (d, 1H, 1'-H), 7.20–7.80 (m, 5H, C₆H₅). C₂₅H₂₈N₂O₁₀S₂, Calcd: C, 51.70; H, 4.82; N, 4.87. Found: C, 51.08; H, 5.46; N, 4.85%.

Sodium 3-methyl-1*H*-thieno(3,4-*c*)pyrazol-6-yl-(phenyl)methanone-4-thiolate (11)

General Procedure

A solution of compound **3** (0.01 mol) and sodium ethoxide (0.01 mol) in ethanol (20 mL) was refluxed for 3 h, the solution was evaporated, and the formed solid product was collected by filtration.

(4-Mercapto-3-methyl-1*H*-thieno(3,4-*c*)pyrazol-6-yl)(phenyl)methan-one (13)

General Procedure

A solution of compound **3** in (20 mL) ethanol was poured on cooled water acidified with hydrochloric acid, and the formed solid product **13** was collected by filtration and recrystallized from ethanol.

(3-Methyl-4-(2',3',4',6'-tetra-*O*-acetyl-β-*D*-gluco- and galacto-pyranosyl-thio)-1*H*-thieno(3,4-*c*)pyrazol-6-yl)(phenyl)methanones (12a,b)

General Procedures

Method A

To a solution of compound **1** (0.01 mol) in ethanol (30 mL), a solution of 2,3,4,6-tetra-*O*-acetyl-α-*D*-gluco- or galactopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone 20 mL was added. The reaction mixture was stirred at rt until completion (TLC, 14 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the sodium bromide formed. The resulting product was recrystallized from ethanol.

Method B

A solution of compounds **13** (0.01 mol) in aqueous potassium hydroxide [0.56 g (0.01 mol) in distilled water (6 mL)] was added to a solution of 2,3,4,6-tetra-*O*-acetyl-α-*D*-gluco- or galactopyranosyl bromide **5a,b** (4.10 g, 0.01 mol) in acetone (30 mL). The reaction mixture was stirred at rt until completion (TLC, 14 h) and then evaporated under reduced pressure, and the residue was washed with distilled water to remove the formed potassium bromide. The resulting product was crystallized from ethanol.

12a: brown, m.p. 160°C (from EtOH), yield (75% method A, 80% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3201 (NH), 1751 (CO). ^1H NMR (DMSO) δ 1.90–2.00 (4s, 12H, 4 \times CH₃CO), 2.10 (s, 3H, CH₃), 3.90 (m, 3H, 6'-H₂, 5'-H), 4.20 (t, 1H, 4'-H), 4.95 (t, 1H, 3'-H), 5.50 (t, 1H, 2'-H), 5.90 (d, 1H, 1'-H), 7.10–8.10 (m, 5H, C₅H₄). C₂₇H₂₈N₂O₁₀S₂, Calcd: C, 53.62; H, 4.67; N, 4.63. Found: C, 53.12; H, 4.84; N, 4.95%.

12b: Brown, m.p. 205°C (from EtOH), yield (77% method A, 85% method B). IR (KBr) $\nu_{\max}/\text{cm}^{-1}$ 3193 (NH), 1751 (CO). ^1H NMR (DMSO) δ 1.80–2.00 (4s, 12H, 4 \times CH₃CO), 2.10 (s, 3H, CH₃), 4.05 (s, 2H, 6'-H), 4.60 (m, 1H, 5'-H), 4.95 (t, 1H, 4'-H), 5.20–5.40 (m, 2H, 3'-H, 2'-H), 5.95 (d, 1H, 1'-H), 7.00–8.10 (m, 5H, C₆H₅). C₂₇H₂₈N₂O₁₀S₂, Calcd: C, 53.62; H, 4.67; N, 4.63. Found: C, 53.57; H, 4.22; N, 4.70%.

REFERENCES

- [1] Elgemeie, G.H.; Mansour, O.A.; Metwally, N.H. Synthesis and anti-HIV activity of different novel nonclassical nucleosides. *Nucleosides Nucleotides* **1999**, *18*, 113–123.
- [2] Elgemeie, G.H.; Attia, A.M.; Hussain, B.A. A synthetic strategy to a new class of cycloalkane ring-fused pyridine nucleosides as potential anti-HIV agents. *Nucleosides Nucleotides* **1998**, *17*, 855–868.
- [3] Elgemeie, G.H.; Attia, A.M.; Al-Kaabi, S.S. Nucleic acid components and their analogues: new synthesis of bicyclic thiopyrimidine nucleosides. *Nucleosides Nucleotides* **2000**, *19*, 723–733.
- [4] Scala, S.; Akhmed, N.; Rao, U.S.; Paull, K.; Lan, L.; Dickstein, B.; Lee, J.; Elgemeie, G.E.H.; Stein, W.D.; Bates, S.E. P-Glycoprotein substrates and antagonists cluster into two distinct groups. *Mol. Pharmacol.* **1977**, *51*, 1024–1033.