

Kinetics and Catalysis, Vol. IV, Issue 1,  
pages 24-34 (Jan.-Feb. 1963)

**Free Radical Recombination Processes on Semiconductor Surfaces and Their Role in Luminescence**

By F. F. VOLKENSHTEIN, A. N. GORBAN,  
AND V. A. SOKOLOV

*Tomsk Polytechnical Institute,  
Institute of Physical Chemistry  
of the Academy of Sciences of  
U.S.S.R.*

The mechanism of luminescence in luminofors was studied as possibly occurring by the radical recombination route under the influence of chemically active gases and by "candor"-luminescence. The intensity of the radical-recombination luminescence was calculated in terms of pressure and the levels of Fermi state. The effect of an outside electric field on the intensity of "candor"-luminescence was determined experimentally. The results qualitatively agree with the theory proposed.

**Possible Explanation of the Compensating Effect in Some Chemical Reactions**

By G. E. LEEHTENSHTEIN

*The Institute of Chemical Physics  
of the Academy of Sciences of  
U.S.S.R.*

The compensating effect can be rationalized by assuming that (1) the experimentally-determined effective rate of a reaction studied is the sum of the local rates which are independent of each other and (2) the free activation energies follow the Gaussian frequency distribution.

**Decomposition of Molecular Ions Formed in Photoionization of Hydrazine and of Some of Its Alkyl Derivatives**

By M. E. AKOPYAN AND F. I. VEELESOV

*Institute of Physics of the  
Leningrad State University*

Summarizes the results of studies of the mass spectra and of the ion formation potentials in photoionization of hydrazine; methyl hydrazine; 1,1-dimethyl hydrazine; and 1-methyl 1-normal butyl hydrazine. Gives calculated values of heats of formation of the ions, ionization potentials of the radicals, and dissociation energies of some of the bonds.

**Formation of Free Radicals in the RH + O<sub>2</sub> System: I. Tetralin, Cyclohexanol, Cyclohexanone**

By E. T. DIENESOV

*The Institute of Chemical Physics  
of the Academy of Sciences of  
U.S.S.R.*

The mechanism of free radical formation (chain initiation) in an RH + O<sub>2</sub> liquid phase system was studied by use of an inhibition technique. Based on the results for tetralin and cyclohexanol, show that the radicals are formed according to the reaction, 2RH + O<sub>2</sub> → 2R· + H<sub>2</sub>O<sub>2</sub>. List values of the reaction rate constants for tetralin, cyclohexanol and cyclohexanone.

**Determination of Absolute Rates of Chain Initiation and Growth in Anionic Polymerization of Styrene**

By H. S. BAGDASARYAN AND  
N. I. NIEPOMNYASHCHII

*The L. Ya. Karpov  
Physico-Chemical Institute*

The kinetics of chain growth was studied in a single-pass continuous unit by reacting the "live" polymers of styrene with the monomer in a tetrahydrofuran solution. The chain growth rate constant is  $k_p = (1.00 \times 10^6) \cdot \exp(-4,670/RT)$  l/mol-sec.

An analysis of the kinetic data for interaction of sodium diphenyl with styrene shows that conversion of the ion-radicals of styrene into the dianionic dimers is the limiting step in the "live" polymer formation process. The rate constant for this reaction is  $k_2 = (5.6 \times 10^{11}) \cdot \exp(-14,200/RT)$  l/mol-sec. The effects of solvation and interaction of the ions upon the activation energies and entropies of the reactions studied were also examined.

**Mechanisms of Energy Transfer in Luminescence and Chemical Radiation**

By V. A. KRONGAUZ AND I. N. VASEELIYEV

*The L. Ya. Karpov  
Physico-Chemical Institute*

Radiolysis and luminescence of three component solutions: toluene, benzoyl peroxide, and