

### Polarographic Investigation of Mechanism and Kinetics of Protolytic Reactions in Nitro-Aci-Tautomeric Conversion of Nitro-Cyclohexane

By P. M. ZAITZEV, YA. I. TOOR'YAN,  
AND Z. V. ZAITZEVA

*Leesechansk Affiliate of State Institute of Atomic Industry, Yaroslavl Scientific Research Institute for Monomers*

The nitro-aci dissociation and aci-nitro recombination rate constants were determined by polarographic technique in tautomeric conversion of nitrocyclohexane in presence of various proton donors and acceptors. To describe the recombination reaction kinetics in a 3 to 7 pH range requires a consideration of all possible protolytic reactions and of the aci-form molecule-anion equilibrium. At pH values of 10 to 12, the dissociation rate is primarily a kinetic function of the protolytic reaction involving participation of hydroxyl ions.

The transformations of acidic protons into aci-form anions (recombination) and of nitro-form protons into bases (dissociation) are the controlling steps in these processes.

In general, the relationship between the protolytic reaction rate constants and the dissociation constants of the corresponding acids can be expressed by Brönsted equations.

### Photochemical Decomposition of Alcohols at Low Temperatures: Decomposition Kinetics of Methyl Alcohol

By B. N. SHELEMOV, N. V. FOK,  
AND V. V. VOYEVODSKII

*Chemistry Department, M. V. Lomonosov Moscow State University, Institute of Chemical Physics of the Academy of Sciences of USSR*

The accumulation, disappearance, and mutual transformation kinetics of the radicals formed in ultra-violet irradiation of dilute solutions of  $\text{H}_2\text{O}_2$  and  $\text{C}_6\text{H}_6$  in  $\text{CH}_3\text{OH}$  at  $77^\circ$  were studied, using a direct-type resonating EPR spectrometer. Formation of the  $\cdot\text{CH}_3$ ,  $\cdot\text{HCO}$ , and  $\cdot\text{CH}_2\text{OH}$  radicals was uncovered. The ultra-violet light effectively decomposes the  $\cdot\text{CH}_2\text{OH}$  radical into  $\cdot\text{CH}_3$  and  $\cdot\text{HCO}$ . The concentrations of  $\text{CO}$ ,  $\text{CH}_4$  and  $\text{H}_2$  in the gaseous end-products are 26, 17, and 57 per cent, respectively. At the conditions employed, the  $\text{H}_2\text{O}_2$ - and  $\text{C}_6\text{H}_6$ -initiated photochemical decomposition of methyl alcohol is a radical-forming process. Multiplication of free valences is one of its basic characteristics.

### A Quanto-Mechanical Basis to Classify Thermal Decomposition Reactions of Solids

By V. V. BOLDIREV AND A. A. MEDVEENSKII

*Scientific Research Institute of Nuclear Physics, Electronics and Automation of the S. M. Karpov Polytechnical Institute of the City of Tomsk*

The quanto-mechanical concepts can serve as a basis to classify thermal decomposition reactions of solids (ionic salts). In this classification, the Group 2 reactions involve electron migration to produce conductive zones and those of Group 1, involve transition of electrons into excited states.

### Formation of Radicals in Radiolysis of Organic Solids: Yields of Radicals from Benzene and Diphenyl Derivatives

By YU. I. MOLEEN, J. J. CHKHE'EEDZÉ,  
E. P. KAPLAN, N. YA. BOOBEN,  
AND V. V. VOYEVODSKII

*Institute of Chemical Physics of the Academy of Sciences of USSR. Institute of Chemical Kinetics and Combustion of Siberian Division of the Academy of Sciences of USSR. Institute of Organic Chemistry of the Academy of Sciences of USSR*

The irradiation yields of some derivatives of benzene and diphenyl were determined by the EPR method. For the compounds investigated, the experimental  $G_R$  values are 1.5-3.0 times greater than that of benzene ( $G_R = 0.2$ ) but significantly smaller than the values based on equation,  $G_{ADD} = \gamma_A G_A + \gamma_D G_D$ , where  $\gamma_A$  and  $\gamma_D$  and  $G_A$  and  $G_D$  are electron fractions and yields, respectively, of the aromatic (A) and saturated (D) groups. Probability of energy transfer from the substitute to the aromatic groups was evaluated on the basis of these experimental data. The values of energy transfer coefficients of a majority of these compounds are close to unity.

### Entropies of Activated Complexes in Hydrogenation of Ethylene Over Metals

By G. I. GOLODIETZ

*L. V. Peesarijevskii Institute of Physical Chemistry of the Academy of Sciences of Ukrainian SSR*

The absolute entropy values of activated complexes produced in hydrogenation of ethylene over various metals were calculated from experimental data. The Jenkins-Rideal mechanism is not applicable to the cases of palladium and platinum. With nickel and a number of other metals, only a small portion of the surface participates in the catalysis.