

Heterogeneous Catalytic Condensation of Olefins in the Presence of Hydrogen, Using Oxygen as the Reaction Initiator

N. I. YERSHOV, YA. T. EIDOOS,
AND V. R. YEROKHEENA

N. D. Z'yeleensky Institute of Organic Chemistry of the Academy of Sciences of USSR

Hydrogenation of pentene-1 by hydrogen over a cobalt/clay catalyst at a temperature of 100° results in its condensation to form C₆, C₇, C₈, C₉, and higher hydrocarbons. The yield of the condensed products is about 15%, based on the olefin charge. Addition of oxygen in amounts of 0.7–1.2% increases the product yield by the factor of 1.5–1.7; on the other hand, increasing the reaction temperature to 180°, decreases the yield by 50%.

Oxidation in Gas Phase of Partially Hydrogenated Derivatives of Benzene

A. N. SHAREPOV AND T. M. HANNANOV

Scientific Research Institute for Petrochemical Industries

A study of oxidation of 4,5-dimethyl-1,2,3,6-tetrahydrophthalic anhydride, 1,2,4,5-tetramethylcyclohexane, 4-methyl-1,2,3,6-tetrahydrophthalic anhydride, and of 2,6-dimethyldecalin was carried out in the gaseous phase over a vanadium catalyst. The oxidation of 4,5-dimethyl-1,2,3,6-tetrahydrophthalic anhydride and of 1,2,4,5-tetramethylcyclohexane resulted in formation of pyromellitic anhydride in amounts of 35–45% and the oxidation of 2,6-dimethyldecalin produced up to 12% of phthalic anhydride. The processes investigated demonstrate that dehydrogenation of naphthenes and of partially hydrogenated aromatics—to form alkyl benzenes—occurs ahead of the oxidation reaction. A hypothesized mechanism is described for oxidation in the gaseous phase of the subject hydrocarbons.

Valence State of Chromium in Active Centers of a Potassia-Chromia-Alumina Dehydrogenation Catalyst

K. I. SLOV'YETZKAYA AND
A. M. ROOBEENSHEIN

N. D. Z'yeleensky Institute of Organic Chemistry of the Academy of Sciences of USSR

The results of this study demonstrate that the active centers, responsible for chemisorption of hydrocarbons on a reduced potassia-chromia-alumina catalyst, contain Cr³⁺ ions.

Decomposition of Isopropyl Alcohol Over Titanium Dioxide With an Octahedral (Anatase) Structure

T. P. KHOKHLOVA, I. S. SAZONOVA,
AND I. P. KAY'YER

Institute of Catalysis of Siberian Division of the Academy of Sciences of USSR

Decomposition of isopropyl alcohol was investigated, using octahedral (anatase) TiO₂ and the TiO₂ + 0.5 mol % WO₃ catalysts. The added WO₃ has substantially no effect on catalytic properties of octahedral TiO₂. The experimentally-determined catalytic activities, electroconductance, and the electron work function values of the two catalysts are compared with the like performance characteristics of another set of two catalysts of identical composition but using the TiO₂ component with rutile structure.

Chemisorption of Nitrogen on Precipitated Iron Catalysts Used in Ammonia Synthesis

R. V. CHESNOKOVA, A. I. GORBOONOV,
S. S. LACHEENOV, G. K. MOORAVSKAYA,
AND G. A. ERDEDEE

Scientific Research and Planning Institute for Industrial Nitrogen and Synthetic Organic Products

Chemisorption of nitrogen was studied at a temperature of 475° on the precipitated iron catalysts promoted with Al₂O₃ (1–37.5% by wt), K₂O (~2%, by wt), and by Al₂O₃ + K₂O. The unpromoted iron catalyst is used in commercial production of ammonia.

Addition of individual promoters—particularly, of K₂O—to the precipitated iron catalyst increases its capacity to chemisorb nitrogen; on the other hand, simultaneous addition of the two promoters results in co-inhibition of their individual ability to promote the chemisorption.

The fused and precipitated iron catalysts with near-identical compositions have similar capacities for chemisorption of nitrogen.

Certain discrepancy was found to exist between capacity of the promoted iron catalysts to chemisorb nitrogen and their catalytic activities in the ammonia synthesis at atmospheric pressure.

Phase Analysis of Ferric Oxide-Molybdena Catalysts

A. S. KOOZ'YETZOVA, L. M. SKR'YABKOVA,
AND B. S. KHREESTOFOROV

Institute of Catalysis of Siberian Division of the Academy of Sciences of USSR

The results show that a 0.3 M solution of hydrazine in hydrochloric acid selectively dis-