

is stopped by injection of α -naphthylamine. This fact indicates that these products are formed via a chain mechanism.

Acetophenone is formed in recombination of the peroxide radicals and in decomposition of the oxycumene radicals. Basically, dimethylphenylcarbinol is produced in decomposition of the hydroperoxide. The rate of cumene hydroperoxide decomposition into radicals is not affected by the addition of dimethylphenylcarbinol and acetophenone.

In the presence of benzoic acid, the hydroperoxide decomposition rate, v , is described by the equation:

$$v = k[\text{C}_6\text{H}_5\text{C}(\text{CH}_3)_2\text{OHH}][\text{C}_6\text{H}_5\text{COOH}].$$

With chlorobenzene as the solvent, the value of k is $7.1 \times 10^6 \exp(-20600/\text{RT})$ liter/mol-sec.

Kinetics of Reaction of Divalent Ions of Iron with Diphenylpicrylhydrazyl

R. I. ZOOSMAN, YU. I. NAB'YEROOKHEEN,
YU. N. MOLEEN, AND N. V. VOYEVOFSKY

*Institute of Chemical Kinetics and
Combustion of Siberian Division of
the Academy of Sciences of USSR*

A spectrophotometric study was made of the general kinetic relationships governing reactions of Fe^{2+} ions with diphenylpicrylhydrazyl (DPhPH). In a mixture with a 74% alcohol—26% water solution, the reaction rate is expressed by the equation: $W = kRM$, where R is concentration of the radical and M is concentration of Fe^{2+} ions. The reaction products are Fe^{3+} and diphenylpicrylhydrazyl only, which is in agreement with the published data. The effective activation energy and the preexponential factor values are 8.1 ± 0.5 kcal/mol and 1.2×10^8 liter/mol \times sec, respectively.

Qualitative data were obtained of the effects of pH of the media and the nature of the solvents and the ligands on the reaction rate.

Interaction of Atomic Oxygen, Produced in the Gas Phase, with Solid Hydrocarbons: Reactions of the Atomic Oxygen with Solid Propylene at Temperatures of 77°–67°K

V. M. ORLOV AND A. N. PONAMAR'YOV

*Institute of Chemical Physics of
the Academy of Sciences of USSR*

The effects of the temperature of a solid hydrocarbon and of the degree of atomic oxygen conversion on the reaction rate of the oxygen were determined for the case of solid propylene at 77°–67°K. The results show that the yields of

the principal reaction products are: propylene oxide, 54%; propion aldehyde, 43%; and acetone, 3%. The product distribution remains unchanged at the conditions covered in this study.

Alteration by Additives of Physico-Chemical Properties of Solids: Effect of Admixed Heterophase Semiconductors on Thermal Stability of Silver Azide

YU. A. ZAKHAROV, E. S. KOOROCKHEEN,
G. G. SAV'YEL'YEV, AND YU. N. ROOFOV

*S. M. Keerov Polytechnical
Institute in the City of Tomsk*

A study was made of the effect of admixed heterophase semiconductors: CdO , ZnO , NiO , Ni_2O_3 , Bi_2O_3 , CdS , CoO , Co_2O_3 , and Co_3O_4 — on the rate and activation energy of thermal decomposition of AgN_3 . In this reaction, catalytic activity of the semiconductors for production of the respective compounds increases with increasing electron work function of the semiconductors at the decomposition conditions employed.

The changes in activity of the semiconductors were determined following the irradiation and partial thermolysis of the AgN_3 — semiconductor mixtures. The effect of the admixed semiconductors on the AgN_3 decomposition rate is compared with the effect of static electric field on the rate of decomposition of "pure" AgN_3 . The results obtained are discussed in the light of the concepts advanced by the authors regarding the decisive role played in catalytic thermolysis of solids by the contact phenomena and by the processes on the border of separation of solid-semiconductor mixtures onto the phases.

The nature of the elementary limiting stage of the AgN_3 decomposition process is considered.

Effect of Admixtures on Radiolysis of Potassium Nitrate

V. V. BOLDIR'YEV, V. M. LIKHEEN,
A. N. OBLEEVANTZEV, AND K. M. SALEEKHOV

*Institute of Chemical Kinetics and
Combustion of Siberian Division of
the Academy of Sciences of USSR
Scientific Research Institute of
Nuclear Physics of the Tomsk
Polytechnical Institute*

The effect of admixtures of Ti^{2+} , Sr^{2+} , Pb^{2+} , and SO_4^{2-} was determined in radiolysis of potassium nitrate by the γ -rays of Co^{60} . The results show that in the radiation-chemical conversion of KNO_3 , the yield of the end product, potassium nitrite, is increased by small dosages of the Sr^{2+} and Pb^{2+} , whereas in large dosages these additives decrease the KNO_2 yield. In addition, the Ti^{2+} in