

### Kinetics of the Alkaline Hydrolysis of $[\text{F}_3\text{BONH}_2]^-$ and of the Acidic Hydrolysis of $\text{F}_3\text{BONH}_3$

I. G. RISS, S. L. EEDEL'S,  
AND M. B. BOOCHEENSKAYA

*The Dn'yeprop'yetrovsk Engineering  
Institute for Railroad Industry*

$[\text{F}_3\text{BONH}_2]^-$ , the anion formed in the coordination reaction of  $\text{BF}_3$  and hydroxylamine, is hydrolyzed in aqueous alkaline solutions via the solvolytic dissociation mechanism; the rate of split-off of the first  $\text{F}^-$  is the limiting step of the reaction. The activation energy and entropy of the process are 17.10 kcal/mol and  $-20.5$  entropy units.

Hydrolysis of  $\text{F}_3\text{BONH}_3$  is catalyzed by the  $\text{H}_3\text{O}^+$  cation due to the reversible formation in small amounts of a protonated complex, followed by rapid solvolytic dissociation of the B—O bond of the complex.

The kinetic data for the hydrolysis of  $[\text{F}_3\text{BONH}_2]^-$  and  $\text{F}_3\text{BONH}_3$  are compared with the known kinetic data for formation of  $\text{BF}_4^-$  and  $\text{F}_3\text{BON}(\text{CH}_3)_3$ , and for the combination reactions of  $\text{BF}_3$  with amines.

### Stereospecific Mutarotation of $\alpha$ -L-Rhamnose by the Catalytically Active Optical Isomers and Racemate of $\alpha$ -Phenylethylamine

V. A. PAVLOV, E. I. KLABONOVSKY,  
AND A. A. BALANDEEN

*N. D. Z'yeleensky Institute of Organic Chemistry  
of the Academy of Sciences of USSR*

A study was made of the effect of stereospecificity of  $\alpha$ -phenylethylamine catalyst on the kinetics of conversion of the L-substrate. The results show that the optical isomers and the racemate of the catalyst affect differently the kinetics of the isomers transition and that the observed difference is entirely due to the stereospecificity of the catalyst.

### Effect of Electronic Structure on Catalytic Properties of Iron-Nickel Catalysts

R. F. LAVR'YENTOVEECH, M. T. ROOSOV,  
AND YU. N. ART'YUKH

*L. V. Peesariyevsky Institute of Physical Chemistry  
of the Academy of Sciences of USSR*

The data presented cover catalytic properties of iron-nickel alloys in decomposition of ammonia. The listed properties include the values of magnetic saturation moment and electron work function of the catalysts. The authors conclude that activity of the catalysts increases with increasing number of the unpaired  $d$ -electrons and

with increasing value of the electron work function.

### Effect of Alkali Metals on Properties of Platinum Catalysts in Dehydrocyclization of $n$ -Hexane

N. R. BOORSEAN, S. N. KOGAN,  
G. M. OSMOLOVSKY, AND Z. A. DAVIDOVA

*The All-Union Scientific Research Institute  
of Petrochemical Processes*

The promoting effect of Li, Na, and Cs additives on activity of Pt/ $\text{Al}_2\text{O}_3$  and Pt/ $\text{SiO}_2$  catalysts was evaluated in dehydrocyclization of  $n$ -hexane. X-ray analyses of the Pt/ $\text{SiO}_2$  structure show that the alkali metals retard crystallization of the Pt-component and that effectiveness of the alkali metals as the promoters decreases in the following order: Cs > Na > Li. The difference in the effectiveness of these metals is attributed to the differences in the size of their ionic radii.

### Catalytic Activity of Aluminum Fluoride

T. V. ANTEEPOVA AND N. D. V'YERSHEENEENA

*Chemistry Department of M. V. Lomonosov  
State University in the City of Moscow*

Aluminum fluoride is an active and stable catalyst in dehydration of ethyl alcohol and in cracking reactions. Kinetics of these reactions over the  $\text{AlF}_3$  were determined.

### pH of Silica-Alumina Cracking Catalysts at the Isoelectric Point

L. G. MAIDANOVSKAYA AND  
T. I. SHEER'YAYEVA

*V. V. Kooibishev State University  
in the City of Tomsk*

The pH values of aqueous solutions of potassium chloride were determined before and after adsorption on various silica-alumina catalysts. Based on the experimental data, the pH values of a series of mixtures of silica-alumina catalysts of different composition were determined at the isoelectric point. At this point, the samples containing 30%  $\text{Al}_2\text{O}_3$  and 70%  $\text{SiO}_2$  had the lowest pH.

### Electron Work Functions of the Samples of Germanium of Different Types of Conductivity in Catalytic Reactions

V. M. FROLOV

*Institute of Chemical Physics of  
the Academy of Sciences of USSR*

In catalytic dehydration of ethanol, the electron work function values of germanium catalysts