

to chain termination on the surface of the channel black, a reaction which of substantially higher than first order in terms of RO_2^* .

Effect of Small Additions of Platinum on the Formation and Growth of New-Phase Nuclei in V_2O_5 Reduction by H_2

N. I. ILCHENKO

The Pizarzhevskiy Institute of Physical Chemistry, Academy of Sciences, Ukr. SSR

Small (0.001–0.01 wt %) concentrations of Pt not only accelerate the V_2O_5 reduction by H_2 (higher k_{specific}), but also alter the process of the metal nuclei. Below the maximum formation rate, the rate of formation of the nuclei in the absence of Pt obeys an exponential function with a constant exponent; however, if Pt is present, the rate is an exponential function with a variable exponent.

Kinetics of the Catalytic Hydrolysis of Allyl Chloride in the Presence of Metals of Group VIII

R. M. FLEED AND YU. M. TREGER

Hydrolysis of allyl chloride in an alkaline medium is catalyzed Ni, Pt, Pd, and Rh. The k_{specific} , as well as the activation energies decrease in the order $\text{Rh} > \text{Ni} \approx \text{Pd} > \text{Pt}$. This series correlates with the corresponding work functions and ionization potentials. It appears that the donor-acceptor bond plays a substantial role in the activation of allyl chloride. It also appears that the transformation of the intermediate π -allyl complex is the controlling step of the reaction.

Alkylation, Conjugated Alkylation, and Disproportionation of Ethylisopropylbenzenes

E. P. BABIN, N. L. ZOTOVA, M. S. BEGININA,
AND V. P. MARSHTUPA

The Donets Section of the All-Union Research Institute of Reagents and Superpure Compounds

Ethylisopropylbenzenes and ethyldiisopropylbenzenes were disproportionated at various temperatures on AlCl_3 . The $K_{\text{eq. dispr.}}$ for ethylisopropylbenzene is 28. This compound can be obtained by conjugated alkylation of ethylbenzene by ethyldi- or ethyltriisopropylbenzenes over AlCl_3 (yields of 75 and 44.5%, respectively). However, alkylation with propylene at 60°C in the presence of AlCl_3 is accompanied by disproportionation, and the maximum yield of ethylbenzene does not exceed 11–12%.

The Energies of the Bonds between Some Oxide Catalysts and Oxygen, Hydrogen, and Carbon Atoms

S. M. RESHETNIKOV AND A. M. SOKOLSKAYA

The S. M. Kirov Kazakh State University

The kinetic technique for calculating the energies of bonds between an oxide catalyst and the oxygen of a reacting molecule does not hold in the case of isotopic oxygen exchange, as well as in oxidation of hydrogen and propylene. These reactions cannot be encompassed by the multiplet theory of catalysis.

Study of the Reaction of Germanium with Oxygen by Mass Spectrometry

G. F. ROMANOVA, I. I. STEPKO,
AND V. I. PEREPELITSA

Semiconductor Institute, Academy of Sciences, Ukr. SSR

Mass spectrometry has shown that it is possible to regenerate an oxidized germanium surface by heating in high vacuum, whereby oxygen is removed above 520°C .

In a steady stream of oxygen, the oxidation of germanium below 500°C is accompanied by vaporization, but above 600°C oxidation slows down. The most probable reason for this slow down is formation of a protective film of GeO_2 .

Chemisorption of *n*-Butane on a Chromia-Alumina-Potassium Dehydrogenation Catalyst

K. I. SLOVETSKAYA, T. P. BRUYEVA,
R. V. DMITRIYEV, AND
A. M. RUBINSHTEYN

The Zelinskiy Institute of Organic Chemistry, Academy of Sciences, USSR

Between 100° and 150°C . *n*-butane absorbs only physically on the above catalyst. Between 170° and 240°C , there is chemisorption without decomposition of the butane. Above 250°C , the chemisorbed *n*-butane is dehydrogenated to butylenes, which in some complex consecutive reactions transform to benzene and toluene.

Chemisorption of Organotin Compounds on $\gamma\text{-Al}_2\text{O}_3$

A. N. KARASEV, YU. A. KOLBANOVSKIY,
L. S. POLAK, AND E. B. SHLIKHTEV

The Topchiev Institute of Petrochemistry, Academy of Sciences, USSR

Using $(\text{CH}_3)_4\text{Sn}$ as a type compound, the authors examined the structure of the Mössbauer