

to chain termination on the surface of the channel black, a reaction which of substantially higher than first order in terms of  $\text{RO}_2^*$ .

### Effect of Small Additions of Platinum on the Formation and Growth of New-Phase Nuclei in $\text{V}_2\text{O}_5$ Reduction by $\text{H}_2$

N. I. ILCHENKO

*The Pizarzhevskiy Institute of Physical Chemistry, Academy of Sciences, Ukr. SSR*

Small (0.001–0.01 wt %) concentrations of Pt not only accelerate the  $\text{V}_2\text{O}_5$  reduction by  $\text{H}_2$  (higher  $k_{\text{specific}}$ ), but also alter the process of the metal nuclei. Below the maximum formation rate, the rate of formation of the nuclei in the absence of Pt obeys an exponential function with a constant exponent; however, if Pt is present, the rate is an exponential function with a variable exponent.

### Kinetics of the Catalytic Hydrolysis of Allyl Chloride in the Presence of Metals of Group VIII

R. M. FLEED AND YU. M. TREGER

Hydrolysis of allyl chloride in an alkaline medium is catalyzed Ni, Pt, Pd, and Rh. The  $k_{\text{specific}}$ , as well as the activation energies decrease in the order  $\text{Rh} > \text{Ni} \approx \text{Pd} > \text{Pt}$ . This series correlates with the corresponding work functions and ionization potentials. It appears that the donor-acceptor bond plays a substantial role in the activation of allyl chloride. It also appears that the transformation of the intermediate  $\pi$ -allyl complex is the controlling step of the reaction.

### Alkylation, Conjugated Alkylation, and Disproportionation of Ethylisopropylbenzenes

E. P. BABIN, N. L. ZOTOVA, M. S. BEGININA,  
AND V. P. MARSHTUPA

*The Donets Section of the All-Union Research Institute of Reagents and Superpure Compounds*

Ethylisopropylbenzenes and ethyldiisopropylbenzenes were disproportionated at various temperatures on  $\text{AlCl}_3$ . The  $K_{\text{eq. dispr.}}$  for ethylisopropylbenzene is 28. This compound can be obtained by conjugated alkylation of ethylbenzene by ethyldi- or ethyltriisopropylbenzenes over  $\text{AlCl}_3$  (yields of 75 and 44.5%, respectively). However, alkylation with propylene at  $60^\circ\text{C}$  in the presence of  $\text{AlCl}_3$  is accompanied by disproportionation, and the maximum yield of ethylbenzene does not exceed 11–12%.

### The Energies of the Bonds between Some Oxide Catalysts and Oxygen, Hydrogen, and Carbon Atoms

S. M. RESHETNIKOV AND A. M. SOKOLSKAYA

*The S. M. Kirov Kazakh State University*

The kinetic technique for calculating the energies of bonds between an oxide catalyst and the oxygen of a reacting molecule does not hold in the case of isotopic oxygen exchange, as well as in oxidation of hydrogen and propylene. These reactions cannot be encompassed by the multiplet theory of catalysis.

### Study of the Reaction of Germanium with Oxygen by Mass Spectrometry

G. F. ROMANOVA, I. I. STEPKO,  
AND V. I. PEREPELITSA

*Semiconductor Institute, Academy of Sciences, Ukr. SSR*

Mass spectrometry has shown that it is possible to regenerate an oxidized germanium surface by heating in high vacuum, whereby oxygen is removed above  $520^\circ\text{C}$ .

In a steady stream of oxygen, the oxidation of germanium below  $500^\circ\text{C}$  is accompanied by vaporization, but above  $600^\circ\text{C}$  oxidation slows down. The most probable reason for this slow down is formation of a protective film of  $\text{GeO}_2$ .

### Chemisorption of *n*-Butane on a Chromia-Alumina-Potassium Dehydrogenation Catalyst

K. I. SLOVETSKAYA, T. P. BRUYEVA,  
R. V. DMITRIYEV, AND  
A. M. RUBINSHTEYN

*The Zelinskiy Institute of Organic Chemistry, Academy of Sciences, USSR*

Between  $100^\circ$  and  $150^\circ\text{C}$ . *n*-butane absorbs only physically on the above catalyst. Between  $170^\circ$  and  $240^\circ\text{C}$ , there is chemisorption without decomposition of the butane. Above  $250^\circ\text{C}$ , the chemisorbed *n*-butane is dehydrogenated to butylenes, which in some complex consecutive reactions transform to benzene and toluene.

### Chemisorption of Organotin Compounds on $\gamma\text{-Al}_2\text{O}_3$

A. N. KARASEV, YU. A. KOLBANOVSKIY,  
L. S. POLAK, AND E. B. SHLIKHTEV

*The Topchiev Institute of Petrochemistry, Academy of Sciences, USSR*

Using  $(\text{CH}_3)_4\text{Sn}$  as a type compound, the authors examined the structure of the Mössbauer