

Salvia miltiorrhiza Inhibits Intimal Hyperplasia and Monocyte Chemotactic Protein-1 Expression After Balloon Injury in Cholesterol-Fed Rabbits

Yuh-Lien Chen,¹ Sung-Pao Yang,² Ming-Shi Shiao,³ Jaw-Wen Chen,^{4,5} and Shing-Jong Lin^{4,5,6*}

¹Institute of Anatomy and Cell Biology, National Yang-Ming University, Taipei, Taiwan

²Institute of Traditional Medicine, National Yang-Ming University, Taipei, Taiwan

³Department of Education and Research, Taipei Veterans General Hospital, Taipei, Taiwan

⁴Division of Cardiology, Taipei Veterans General Hospital, Taipei, Taiwan

⁵Cardiovascular Research Center, National Yang-Ming University, Taipei, Taiwan

⁶Institute of Clinical Medicine, National Yang-Ming University, Taipei, Taiwan

Abstract Antioxidants that prevent low density lipoproteins (LDL) from oxidation may inhibit atherosclerosis and post-angioplasty restenosis. *Salvia miltiorrhiza* (SM) has been shown to inhibit LDL oxidation and reduce atherosclerosis in cholesterol-fed rabbits. The effects of SM on neointimal hyperplasia and monocyte chemotactic protein-1 (MCP-1) expression after balloon injury were studied. Male New Zealand white rabbits were fed a 2% cholesterol diet together with daily SM (4.8 gm/kg body wt.) treatment (SM; n=10) or without SM as a control (C; n=9) for 6 weeks. Probucol-treated (0.6 gm/kg body wt.) rabbits (P; n=9) were used as a positive control group. A balloon injury of the abdominal aorta was performed at the end of the third week. Aortas were harvested at the end of 6 weeks. The plasma cholesterol levels were lowered in SM group. The neointimal hyperplasia in abdominal aortas was significantly inhibited in SM group [neointima/media area ratio: 0.63 ± 0.05 (SM) versus 0.78 ± 0.05 (C); $P < 0.05$] and in P group [0.45 ± 0.02 (P) versus 0.78 ± 0.05 (C); $P < 0.05$] when compared with C group. SM treatment significantly reduced MCP-1 mRNA and protein expression in balloon-injured abdominal aorta. These inhibitory effects on intimal response after balloon injury might be attributed to antioxidant capacity and cholesterol lowering effect of SM. SM treatment may offer some protection against post-angioplasty restenosis. *J. Cell. Biochem.* 83: 484–493, 2001. © 2001 Wiley-Liss, Inc.

Key words: angioplasty; atherosclerosis; cholesterol; cytokines; restenosis; *Salvia miltiorrhiza*

Restenosis after successful coronary intervention remains a major limitation to the long-term outcome of this procedure [Leimgruber et al., 1986]. Intimal hyperplasia, elastic recoil, and vessel wall remodeling are some major mechanisms responsible for post-angioplasty restenosis [Post et al., 1994]. The formation of intimal hyperplasia is mainly due to the phenotypic conversion and proliferation of

smooth muscle cells as well as the accumulation of activated macrophages and foam cells [Hanke et al., 1994; Chen et al., 1997]. Activated macrophages and smooth muscle cells may express a number of cytokines, growth factors, and chemoattractant proteins, which involved in the vessel wall remodeling, and could further exacerbate the progression of restenotic lesions [Schwartz et al., 1995]. The monocyte chemotactic protein-1 (MCP-1) is an important mediator of monocyte recruitment into the vascular wall at sites of active inflammation or injury. Increased levels of MCP-1 have been demonstrated in human atherosclerotic arteries [Yla-Herttuala et al., 1991], experimental atherosclerotic lesions [Yla-Herttuala et al., 1991; Yu et al., 1992; Wysocki et al., 1996], and some animal models of post-angioplasty restenosis [Wysocki et al.,

Grant sponsor: National Science Council; Grant numbers: NSC 86-2314-B-010-029, NSC 87-2314-B-010-052; Grant sponsor: Taipei Veterans General Hospital; Grant number: VGH-305; Grant sponsor: Yen Tjing-Ling Medical Foundation; Grant number: CI-86-18.

*Correspondence to: Shing-Jong Lin, Division of Cardiology, Taipei Veterans General Hospital, 201, Section 2, Shih-Pai Rd, Taipei, Taiwan. E-mail: sjlin@vghtpe.gov.tw

Received 30 March 2001; Accepted 25 May 2001

© 2001 Wiley-Liss, Inc.
DOI 10.1002/jcb.1233

1996; Merritt et al., 1997]. These findings suggested that MCP-1 may play an important role in the pathogenesis of vascular lesion formation.

An evolving concept in recent years is the role of oxygen free radicals and oxidative stress in the development of atherosclerosis and post-angioplasty restenosis [Halliwell, 1989]. There is considerable evidence supporting the notion that oxidized low density lipoprotein (oxLDL) contributes to foam cell formation and subsequently, the progression of atherosclerotic and restenotic lesions [Steinberg et al., 1989]. Products of low density lipoprotein (LDL) oxidation have been reported to be chemotactic for vascular smooth muscle cells [Autio et al., 1990] and monocytes [Berliner et al., 1990]. It is plausible that antioxidants, which prevent LDL from oxidative modification, may interrupt the progression of atherosclerosis and restenosis [Lusis and Navab, 1993]. Probuocol, a potent lipophilic antioxidant, has been shown to inhibit atherosclerosis [Daugherty et al., 1991; Sasahara et al., 1994] and reduce restenosis after angioplasty [Ferns et al., 1992; Schneider et al., 1993; Tardif et al., 1997]. Several studies also suggest that water-soluble antioxidants, such as vitamin C [Jialal and Grundy, 1991], glabridin [Fuhrman et al., 1997], and flavonoids [Hertog et al., 1993], are effective in inhibiting LDL oxidation and reducing the risk of coronary heart disease [Hertog et al., 1993]. Our previous study also showed that red wine with antioxidant capacity inhibited MCP-1 expression and reduced intimal hyperplasia in a hyperlipidemic rabbit restenosis model [Feng et al., 1999]. *Salvia miltiorrhiza* (SM) has been used as a blood-quickening, stasis-dispelling agent in traditional Chinese medicine [Yagi et al., 1989; Wu et al., 1998]. Salvianolic acid B (Sal B), a potent water-soluble polyphenolic antioxidant extracted from the roots of SM, has been shown to scavenge 1,1-diphenyl-2-picrylhydrazyl radicals and inhibit LDL oxidation more effectively than probuocol [Wu et al., 1998]. In a recent study, we demonstrated an increase of vitamin E content in LDL and reduction of atherosclerosis in cholesterol-fed rabbits by a water-soluble antioxidant-rich fraction of SM [Wu et al., 1998]. The purposes of the present study were to examine the effects of SM treatment on intimal response and MCP-1 expression in the aorta after balloon injury in cholesterol-fed rabbits.

MATERIALS AND METHODS

Agents

Probuocol was obtained from Marion Merrell Dow, USA. Sudan IV and hematoxylin were purchased from Sigma Chemical Co., USA. Sal B was a gift from the Institute of Materia Medica, Chinese Academy of Medical Sciences at Beijing, China.

Preparation of SM and Determination of Sal B Content

The dry roots of SM were extracted with a mixture of water and ethanol (4:1, v/v) at room temperature for 24 h. After evaporation of solvent under reduced pressure, the SM extract was stored under nitrogen at 4°C before use. The content of Sal B in SM was determined by reversed-phase HPLC.

Animal Experiments

Twenty-eight male New Zealand white rabbits, 3 months of age and weighing 2.5–3.0 kg, were housed in the animal center of the Taipei Veterans General Hospital, Taiwan. After 1 week on a commercial rabbit chow diet (Scientific Diet Services, Essex, UK), the animals were then fed a 2% high cholesterol diet (Purina Mills, Inc., USA) and randomly divided into three groups: (1) without drug treatment as the control group (group C; n=9), (2) probuocol treatment with oral ingestion of probuocol (0.6 gm/kg body wt. per day) as the positive control group (group P; n=9), and (3) SM treatment with oral ingestion of SM extract (4.8 gm/kg body wt. per day) as the experimental group (group SM; n=10). The investigation conforms with the *Guide for the Care and Use of Laboratory Animals* published by the US National Institutes of Health (NIH Publication No. 85–23, revised 1996). The protocol was approved by the Animal Care and Use Committee of National Yang-Ming University, Taipei, Taiwan. The choice of SM dose was based on its antioxidant activity comparable to that of probuocol in Cu²⁺-induced LDL oxidation [Wu et al., 1998]. Water was allowed ad libitum. Animals were bled periodically for measurement of plasma cholesterol, liver function, and renal function. At the end of the third week of high cholesterol diet feeding with or without drug treatment, the animals in each group were fasted for 12 h and anesthetized with an intramuscular injection of xylazine (5 mg/kg)

and ketamine hydrochloride (35 mg/kg). The surgery was performed under sterile conditions as previously described [Feng et al., 1999]. The right femoral artery was exposed and a 3 Fr. (1 French=0.33 mm) Fogarty arterial embolectomy balloon catheter (Baxter Healthcare, Buckinghamshire, U.K.) was introduced retrogradely into the femoral artery and the lower abdominal aorta. Denudation was then performed by inflating the balloon with normal saline. The surgical wound was closed and the animals were continued on the same diets and treatments. At the end of the sixth week of the experiment, the rabbits were sacrificed. The abdominal aorta and right iliac artery were harvested for 16 cm by measuring from the previous arteriotomy site. The abdominal aorta was cut into six segments. A small part of each arterial segment was taken, immersion-fixed with 4% buffered paraformaldehyde and paraffin-embedded, then cross-sectioned for morphometry, *in situ* hybridization, and immunohistochemistry. The remaining larger portion of each arterial segment was immediately frozen in liquid nitrogen for RNA isolation.

Blood Sampling

Blood samples were collected from each animal before the start of experiment, and at the times of balloon denudation (third week), and sacrifice (sixth week). Samples were separated by centrifugation, and plasma was stored at -20°C prior to analysis. The levels of total cholesterol and triglycerides were measured with Kodak Ektachem DT 60 and DTSC analyzer.

Analysis of Extent of Atherosclerosis

The thoracic aortas (from aortic arch to the diaphragm) were also collected and cleaned of adhering tissue when animals were sacrificed. Thoracic aortas were stained with a solution of Sudan VI to visualize the atherosclerotic lesion area. Sudanophilic areas were quantitatively measured by computer-assisted planimetry. The extent of the lesions was expressed as the proportion of the total area with sudanophilic areas (the surface area of lesions/the surface area of the whole thoracic aorta).

Hematoxylin-Eosin Stain and Morphometry

One paraffin-embedded cross-section, at a thickness of 5 μm , was obtained from each segment of the abdominal aorta and stained

with hematoxylin and eosin. Morphometric analysis of the six arterial sections per animal was performed by the use of a LV-2 Image Analyzer (Winhow Instruments, Taipei, Taiwan). For each arterial cross-section specimen, the intimal and medial thickness were measured at eight separate points, the intima/media thickness ratio was calculated for each point, and the overall intima/media ratio for each section was expressed as a mean of these eight values. The intimal and medial areas for each arterial cross-section specimen were measured and the intima/media area ratio was determined.

RNA Isolation and Northern Blot Analysis

The MCP-1 expression in the arterial specimen was examined at the RNA level by using Northern blot analysis. Total RNA was obtained as described previously [Chirgwin et al., 1979; Feng et al., 1999]. RNA was transferred onto a nylon membrane (Nytran 0.45 μm , Schleicher & Schuell, Inc., USA) and fixed by ultraviolet irradiation. After hybridizing with the ^{32}P -labeled MCP-1 cDNA, the membrane was washed with 1X standard saline citrate containing 1% SDS at room temperature for 15 min and then exposed to X-ray films (X-OMAT, Kodak, Rochester, NY) at -70°C for autoradiography. The intensity of each hybridized band on XAR film was determined using optical densitometry (Molecular Dynamics, USA). The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA levels served as the internal standard to normalize the MCP-1 signals [Alwine et al., 1977]. The following DNA fragments were used as probes: a 0.7 kb EcoRI and BamHI fragment containing full-length MCP-1 cDNA and a 1.3 kb PstI fragment containing GAPDH cDNA (received from American Type Culture Collection, Rockville, MD).

In Situ Hybridization and Immunohistochemistry Analysis

To examine the cellular expression and localization of the MCP-1 gene and protein, *in situ* hybridization and immunohistochemistry were performed on serial sections of the aorta. The first tissue section was hybridized with digoxigenin-MCP-1 cDNA. The second and the third sections were incubated with smooth muscle cell-specific and macrophage-specific antibodies to identify smooth muscle cells and macrophages, respectively. The last section was used to detect the expression of MCP-1 protein.

In situ hybridization. MCP-1 cDNA was labeled with digoxigenin (DIG)-dUTP according to the manufacturer's instructions (DIG labeling and detection kit, Boehringer Mannheim Biochemica, Mannheim, Germany) and was used as a probe for in situ hybridization, which was performed according to a previously published method [Pang et al., 1996]. Paraffin-embedded arterial sections (5- μ m thickness) were placed on poly-L-lysine-coated slides, deparaffinized, treated with proteinase K (1 μ g/ml) for 15 min at 37°C, and acetylated (0.25% acetic anhydride in 0.1 M triethanolamine and 0.9% NaCl) for 10 min. Sections were then washed with 2X SSC, dehydrated, air dried for 30 min and stored at -70°C before use. Before hybridization, each section was prehybridized in a humid chamber with 100 μ l prehybridization solution (5X SSC, 5X Denhardt's solution, 50% deionized formamide, 250 μ g/ml yeast t-RNA, 250 μ g denatured salmon sperm DNA and 4 mmol/L EDTA) for 3 h. Hybridization was performed at 50°C for 16–24 h in a humid chamber with a 25 μ l/section prehybridization solution containing 10 ng/ μ l DNA probe. Following the hybridization, sections were washed with SSC. Sections were then blocked for 30 min, incubated with alkaline phosphatase-conjugated anti-DIG antibody (30 min) and detected with a color solution containing 337.5 μ g/ml nitroblue tetrazolium salt and 175 μ g/ml 5-bromo-4-chloro-3-indolylphosphate (10–30 min) according to the manufacturer's instructions. In some experiments, tissue sections were hybridized with DIG-labeled probes plus 50-fold unlabeled cDNA, which abolished the signal as controls.

Immunohistochemistry. For immunohistochemistry, the second, the third, and the last serial paraffin-embedded sections were used. The tissue sections were deparaffinized, rehydrated, washed with PBS, and treated with 5mg/ml serum albumin in PBS for 1 h to block non-specific binding. The last section was

incubated with goat anti-human MCP-1 primary antibody (1:15 dilutions, R&D systems, USA). Antibodies were localized by an indirect immunoperoxidase technique (avidine-biotin-horseradish peroxidase complex) employing diaminobenzidine as a chromogen (Vector Lab, USA). The other two sections were incubated with mouse anti- α -smooth muscle actin (IA4, Sigma Chemical Co.) and mouse anti-rabbit macrophage (Ram II, Dako Corp., USA) to detect smooth muscle cells and macrophages, respectively. These two sections were then incubated with FITC-conjugated goat anti-mouse secondary antibody (Sigma Chemical Co.) and observed by fluorescent microscopy. Negative control was performed by omitting the incubation of the primary antibodies in the tissue sections.

Statistics

Values were expressed as mean \pm SEM. Data for the three groups of animals were compared using one-way ANOVA. When ANOVA indicated a significant difference between groups, data for individual groups were compared by the Dunnett test. A probability of $P < 0.05$ was considered statistically significant.

RESULTS

The content of Sal B in SM extract was 8.61%. Each group of animals was matched for age and weight at the start of the experiment. There was no difference in body weight at the end of the 6-week feeding period.

Blood Biochemistry

The plasma cholesterol concentrations in the C group were apparently increased after 3 weeks of feeding a 2% high cholesterol diet, and cholesterol levels increased further after 6 weeks of feeding, up to 53.92 \pm 4.10 mmole/L (Table I). The plasma cholesterol concentrations in the SM group at 3-week and 6-week

TABLE I. Plasma Cholesterol Levels at Baseline and at 3, 6 Weeks After High-Cholesterol Feeding in Different Groups of Rabbits

	C	P	SM
Baseline	1.27 \pm 0.14	0.84 \pm 0.20	1.21 \pm 0.17
3 weeks	39.33 \pm 7.22	30.04 \pm 7.00	20.16 \pm 3.56
6 weeks	53.92 \pm 4.10	38.94 \pm 6.19	35.30 \pm 3.79*

Values are means \pm SEM. *: $P < 0.05$ compared with the C group at 6 weeks. C: control, P: probucol, SM: *Salvia Miltiorrhiza*, Cholesterol: mmole/L.

were 20.16 ± 3.56 mmole/L and 35.30 ± 3.79 mmole/L, respectively. The SM group showed significantly lower cholesterol levels than the control group at the 6-week point ($P < 0.05$). Probucol treatment showed a trend of lower cholesterol levels but was not significant when compared to that of the control group. Plasma triglyceride levels were not increased significantly during the experimental period in these three groups.

SM and Probucol Treatment Inhibit the Formation of Atherosclerotic Lesions

The extent of atherosclerotic lesions, expressed as atherosclerotic area/total surface area ratio, in the thoracic aortas of the control group was 0.232 ± 0.040 . In the probucol and SM groups, the corresponding values were 0.038 ± 0.023 and 0.106 ± 0.041 , respectively. Probucol treatment resulted in an 84% decrease and SM treatment caused a 47% decrease in atherosclerotic areas. Both decreases were statistically significant compared with the control group ($P < 0.05$).

SM and Probucol Treatment Reduce Neointimal Formation After Balloon Injury

Histopathological sections of abdominal aortas following balloon injury were shown in Figure 1. The control group showed markedly thickened neointima with foam cells accumulation (Fig. 1A). Both probucol (Fig. 1B) and SM (Fig. 1C) treatment significantly reduced the neointimal hyperplasia. The thickness ratios of the neointima to the media in the probucol (0.61 ± 0.03) and the SM (0.75 ± 0.06) groups were significantly lower than that in the control group (1.07 ± 0.06), as shown in Figure 2A. Similarly, the area ratios of the neointima to the media in the probucol (0.45 ± 0.02) and the SM (0.63 ± 0.05) groups were significantly less when compared to the control group (0.78 ± 0.05), as shown in Figure 2B.

MCP-1 mRNA Expression by Northern Blot Analysis

The cholesterol-fed control group exhibited a significant expression of MCP-1 mRNA in the abdominal aorta after balloon injury (Fig. 3). The expression of MCP-1 mRNA was significantly inhibited in the probucol treatment group (0.69-fold lower) and in the SM




Fig. 1. Hematoxylin and eosin-stained sections of abdominal aortas for 3 weeks after balloon injury from the control (A), probucol (B), and SM (C) groups. The lumen is uppermost in all sections. The neointima is indicated between two arrows. The neointima is less thickened in both the P and SM groups compared with the C group.

treatment group (0.23-fold lower) than in the control group. The SM group showed significantly the lower MCP-1 expression than the P group.




Fig. 2. The mean of neointima/media thickness ratio (A) and the mean of neointima/media area ratio (B) of abdominal aortas at 3 weeks after balloon injury from the control (C), probucol (P), and *Salvia miltiorrhiza* (SM) groups. Values represent means \pm SEM. In the SM and P groups, the neointima/media thickness ratio and the neointima/media area ratio were significantly reduced compared with the control group (* $P < 0.05$).

In Situ Hybridization Combined With Immunohistochemistry for the Expression of MCP-1 mRNA and Protein

The cellular localization and specific cell type associated with the MCP-1 gene and protein expression were examined in the abdominal aorta after balloon injury (Figs. 4–6). In the cholesterol-fed control group, the MCP-1 mRNA was strongly expressed in the thickened neointima (Fig. 4A) where abundant smooth muscle cells (Fig. 4B) and a few macrophages (Fig. 4C) were present. The MCP-1 protein was strongly expressed in the thickened intima (Fig. 4D) in this group of rabbits. The probucol-treated group also showed the expression of MCP-1 mRNA (Fig. 5A) and protein (Fig. 5D) in the intima. Significantly less MCP-1 mRNA was

detected in the aortic wall of the SM-treated (Fig. 6A) group. A weaker MCP-1 protein reaction was seen in the SM (Fig. 6D) groups. In the probucol and the SM-treated groups the less thickened intima was composed of smooth muscle cells mainly (Figs. 5B, 6B) and few macrophages (Figs. 5C, 6C).

DISCUSSION

In the present investigation, a moderate dietary-induced hypercholesterolemia combining balloon injury of abdominal aortas was used as an experimental restenosis animal model. The major findings of this study were that a water-soluble fraction of SM, which contained potent antioxidants including Sal B, limited the intimal response to balloon injury in cholesterol-fed rabbits by inhibiting intimal hyperplasia and reducing MCP-1 expression in abdominal aortas. In addition, SM treatment also markedly decreased atherosclerotic lesion areas in thoracic aortas without endothelial denudation. In this study, plasma cholesterol levels were significantly lowered after six weeks of daily SM treatment when compared with cholesterol-fed control rabbits. This cholesterol-lowering effect may partly explain the less atherosclerotic lesion formation in thoracic aorta, and a decreased intimal response in balloon-injured abdominal aorta of SM-treated rabbits.

A number of lipophilic antioxidants, such as α -tocopherol, probucol, or butylated hydroxytoluene, can block the oxidation of lipoproteins in vitro [Morel et al., 1983; Steinbrecher et al., 1984] and in vivo [Morel and Chisolm, 1989; Lafont et al., 1995]. Probucol has been shown to significantly reduce the neointimal formation in response to balloon injury in the hypercholesterolemic rabbit model [Ferns et al., 1992] and in a swine model of coronary artery balloon injury [Schneider et al., 1993]. In the present study, probucol treatment was used as a positive control for comparison. We found that probucol not only significantly reduced atherosclerotic lesions in thoracic aortas but also inhibited intimal hyperplasia and MCP-1 expression in abdominal aortas after balloon injury. Several studies also suggest that water-soluble antioxidants, such as vitamin C, are effective in inhibiting LDL oxidation by the preservation of endogenous antioxidants in LDL [Jialal and Grundy, 1991]. Flavonoids, a





Fig. 3. Northern blot analysis. The MCP-1 mRNA expression was significantly reduced in the SM and P groups compared with the control group ($*P < 0.05$). The MCP-1 mRNA expression of the SM group was also significantly reduced compared with the P group ($*P < 0.05$).

class of water-soluble antioxidants, are useful in reducing the risk of coronary heart disease [Hertog et al., 1993]. Glabridin, a polyphenolic compound from licorice, protects LDL against lipid peroxidation in humans and reduces atherosclerotic lesion areas in apoE-deficient mice [Fuhrman et al., 1997]. Our recent study demonstrated that red wine with abundant flavonoids inhibits MCP-1 expression and modestly reduces neointimal hyperplasia after balloon injury in cholesterol-fed rabbits [Feng et al., 1999]. In this study, SM was chosen as treatment drug because it has usually been used as a blood-quickening, stasis-dispelling agent for treatment of cardiovascular disorders in traditional Chinese medicine. Its non-polar extracts contain tanshinones, which can inhibit platelet aggregation and protect myocardium against ischemia-induced derangement [Yagi et al., 1989]. Sal B is a potent antioxidant in the water-soluble extract of SM. In a previous study, we observed the increase of vitamin E

content in LDL, inhibition of LDL oxidation, protection of endothelial damage, and reduction of atherosclerosis in cholesterol-fed rabbits by a water-soluble antioxidant-rich fraction of SM [Wu et al., 1998]. In the present study, we further demonstrated that this fraction of SM was able to significantly reduce intimal hyperplasia after balloon injury in hypercholesterolemic animals.

Monocyte infiltration into the neointima after balloon injury depends on their ability to attach to the injured endothelium and their migration into the subendothelial space. The activation state of circulating monocytes and their expression of adhesion molecules, and the secretion of chemotactic factors by the vascular cells of the damaged vessels are involved in these processes, which may be inhibited by SM treatment. Monocyte chemotactic protein (MCP-1) is a chemotactic factor for monocytes with great potency [Valente et al., 1988]. MCP-1 is secreted by stimulated human lymphocytes, endothelial



Figs. 4–6. Detection of MCP-1 mRNA and protein in serial sections of abdominal aortas from the control (Fig. 4), probucol (Fig. 5), and SM (Fig. 6) groups. The lumen is uppermost in all sections. The internal elastic lamina is indicated by arrows. **A:** MCP-1 mRNA by in situ hybridization (purple reaction products) was observed in control (Fig. 4A) group, probucol (Fig. 5A), and SM (Fig. 6A) groups. **B:** Immunohistochemical staining for smooth muscle cell-specific antibody. Staining was

obvious in thickened intima of control (Fig. 4B), probucol (Fig. 5B), and SM (Fig. 6B) groups. **C:** Immunohistochemical staining for macrophage-specific antibody. Staining was observed in a few cells in the thickened intima of control (Fig. 4C), probucol (Fig. 5C), and SM (Fig. 6C) groups. **D:** MCP-1 protein by immunohistochemical staining (brown reaction products) was observed in control (Fig. 4D) group, probucol (Fig. 5D), and SM (Fig. 6D) groups.

cells, fibroblasts, monocytes, and smooth muscle cells in vitro [Kaczmarek et al., 1985; Valente et al., 1988; Strieter et al., 1989; Yoshimura et al., 1989; Rollins et al., 1990]. Strong expression of MCP-1 mRNA has been observed in *Cynomolgus* monkey fed with 2% cholesterol and 10% fat diet for 6, 11 and 18 months, and in New Zealand white rabbits fed with a 2% cholesterol diet for 10 weeks [Yla-Herttuala et al., 1991; Yu et al., 1992]. In addition, the elevation of MCP-1 expression was also demonstrated in balloon-injured aortas of pigs [Wysocki et al., 1996] or rabbits

[Merritt et al., 1997]. The MCP-1 mRNA expression in SM-treated or probucol-treated rabbits was lower than that in the control group in the present study. These findings implicated that antioxidants might inhibit the expression of MCP-1 and thereby reduced the intimal response after balloon injury of aortas in cholesterol-fed rabbits.

Immunostaining with specific antibody to MCP-1 revealed strong reactivity in the thickened intima of the control groups. The serial sections with specific antibodies to smooth muscle cells and macrophages demonstrated

that both cells are responsible for the expression of MCP-1 during vascular response after balloon injury. The production of MCP-1 by either smooth muscle cells or macrophages in the neointima in this study may implicate a recruitment of more monocytes/macrophages to the injured vessel sites and the progression of neointimal hyperplasia. SM treatment obviously decreased MCP-1 mRNA and protein expression, thereby resulted in the reduction of intimal response in this hyperlipidemic restenosis rabbit model.

In conclusion, this study provides experimental evidence that SM treatment attenuates intimal thickening in the balloon-injured abdominal aorta of cholesterol-fed rabbits. The reduction of intimal response may be attributed to the suppression of MCP-1 release from activated macrophages and smooth muscle cells, cholesterol-lowering effect, and antioxidant capacity of SM. It is suggested that SM may be of potential benefit in preventing restenosis after coronary angioplasty, particularly in patients with hypercholesterolemia.

REFERENCES

- Alwine JC, Kemp DJ, Stark GR. 1977. Method for detection of specific RNAs in agarose gels by transfer to diazobenzyloxymethyl-paper and hybridization with DNA probes. *Proc Natl Acad Sci USA* 74:5350-5354.
- Autio I, Jaakkola O, Solakivi T, Nikkari T. 1990. Oxidized low-density lipoprotein is chemotactic for arterial smooth muscle cells in culture. *FEBS Lett* 277:247-249.
- Berliner JA, Territo MC, Sevanian A, Ramin S, Kim JA, Bamshad B, Esterson M, Fogelman AM. 1990. Minimally modified low density lipoprotein stimulates monocyte endothelial interactions. *J Clin Invest* 85:1260-1266.
- Chen YH, Chen YL, Lin SJ, Chou CY, Mar GY, Chang MS, Wang SP. 1997. Electron microscopic studies of phenotypic modulation of smooth muscle cells in coronary arteries of patients with unstable angina pectoris and postangioplasty restenosis. *Circulation* 95:1169-1175.
- Chirgwin JM, Przybyla AE, MacDonald RJ, Rutter WJ. 1979. Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry* 18:5294-5299.
- Daugherty A, Zweifel BS, Schonfeld G. 1991. The effects of probucol on the progression of atherosclerosis in mature Watanabe heritable hyperlipidaemic rabbits. *Br J Pharmacol* 103:1013-1018.
- Feng AN, Chen YL, Chen YT, Ding YZ, Lin SJ. 1999. Red wine inhibits monocyte chemotactic protein-1 expression and modestly reduces neointimal hyperplasia after balloon injury in cholesterol-fed rabbits. *Circulation* 100:2254-2259.
- Ferns GA, Forster L, Stewart-Lee A, Konneh M, Nourooz-Zadeh J, Anggard EE. 1992. Probucol inhibits neointimal thickening and macrophage accumulation after balloon injury in the cholesterol-fed rabbit. *Proc Natl Acad Sci USA* 89:11312-11316.
- Fuhrman B, Buch S, Vaya J, Belinky PA, Coleman R, Hayek T, Aviram M. 1997. Licorice extract and its major polyphenol glabridin protect low-density lipoprotein against lipid peroxidation: in vitro and ex vivo studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *Am J Clin Nutr* 66:267-275.
- Halliwell B. 1989. Free radicals, reactive oxygen species and human disease: a critical evaluation with special reference to atherosclerosis. *Br J Exp Pathol* 70:737-757.
- Hanke H, Hassenstein S, Ulmer A, Kamenz J, Oberhoff M, Haase KK, Baumbach A. 1994. Accumulation of macrophages in the arterial vessel wall following experimental balloon angioplasty. *Eur Heart J* 15:691-698.
- Hertog MG, Feskens EJ, Hollman PC, Katan MB, Kromhout D. 1993. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen elderly study. *Lancet* 342:1007-1011.
- Jialal I, Grundy SM. 1991. Preservation of the endogenous antioxidants in low density lipoprotein by ascorbate but not probucol during oxidative modification. *J Clin Invest* 87:597-601.
- Kaczmarek L, Calabretta B, Baserga R. 1985. Expression of cell-cycle-dependent genes in phytohemagglutinin-stimulated human lymphocytes. *Proc Natl Acad Sci USA* 82:5375-5379.
- Lafont AM, Chai YC, Cornhill JF, Whitlow PL, Howe PH, Chisolm GM. 1995. Effect of alpha-tocopherol on restenosis after angioplasty in a model of experimental atherosclerosis. *J Clin Invest* 95:1018-1025.
- Leimgruber PP, Roubin GS, Hollman J, Cotsonis GA, Meier B, Douglas JS, King SB Jr., Gruentzig AR. 1986. Restenosis after successful coronary angioplasty in patients with single-vessel disease. *Circulation* 73:710-717.
- Lusis AJ, Navab M. 1993. Lipoprotein oxidation and gene expression in the artery wall. New opportunities for pharmacologic intervention in atherosclerosis. *Biochem Pharmacol* 46:2119-2126.
- Merritt R, Guruge BL, Miller DD, Chaitman BR, Bora PS. 1997. Moderate alcohol feeding attenuates postinjury vascular cell proliferation in rabbit angioplasty model. *J Cardiovasc Pharm* 30:19-25.
- Morel DW, Chisolm GM. 1989. Antioxidant treatment of diabetic rats inhibits lipoprotein oxidation and cytotoxicity. *J Lipid Res* 30:1827-1834.
- Morel DW, Hessler JR, Chisolm GM. 1983. Low density lipoprotein cytotoxicity induced by free radical peroxidation of lipid. *J Lipid Res* 24:1070-1076.
- Pang JH, Jiang MJ, Chen YL, Wang FW, Wang DL, Chu SH, Chau LY. 1996. Increased ferritin gene expression in atherosclerotic lesions. *J Clin Invest* 97:2204-2212.
- Post MJ, Borst C, Kuntz RE. 1994. The relative importance of arterial remodeling compared with intimal hyperplasia in lumen renarrowing after balloon angioplasty. A study in the normal rabbit and the hypercholesterolemic Yucatan micropig. *Circulation* 89:2816-2821.
- Rollins BJ, Yoshimura T, Leonard EJ, Pober JS. 1990. Cytokine-activated human endothelial cells synthesize and secrete a monocyte chemoattractant, MCP-1/JE. *Am J Pathol* 136:1229-1233.
- Sasahara M, Raines EW, Chait A, Carew TE, Steinberg D, Wahl PW, Ross R. 1994. Inhibition of hypercholester-

- olemia-induced atherosclerosis in the nonhuman primate by probucol. I. Is the extent of atherosclerosis related to resistance of LDL to oxidation? *J Clin Invest* 94:155–164.
- Schneider JE, Berk BC, Gravanis MB, Santoian EC, Cipolla GD, Tarazona N, Lassegue B, King SB 3rd. 1993. Probucol decreases neointimal formation in a swine model of coronary artery balloon injury. A possible role for antioxidants in restenosis. *Circulation* 88:628–637.
- Schwartz SM, Reidy MA, O'Brien ER. 1995. Assessment of factors important in atherosclerotic occlusion and restenosis. *Thromb Haemostasis* 74:541–551.
- Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witztum JL. 1989. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med* 320:915–924.
- Steinbrecher UP, Parthasarathy S, Leake DS, Witztum JL, Steinberg D. 1984. Modification of low density lipoprotein by endothelial cells involves lipid peroxidation and degradation of low density lipoprotein phospholipids. *Proc Natl Acad Sci USA* 81:3883–3887.
- Strieter RM, Wiggins R, Phan SH, Wharram BL, Showell HJ, Remick DG, Chensue SW, Kunkel SL. 1989. Monocyte chemotactic protein gene expression by cytokine-treated human fibroblasts and endothelial cells. *Biochem Biophys Res Co* 162:694–700.
- Tardif JC, Cote G, Lesperance J, Bourassa M, Lambert J, Doucet S, Bilodeau L, Nattel S, de Guise P. 1997. Probucol and multivitamins in the prevention of restenosis after coronary angioplasty. *N Engl J Med* 337:365–372.
- Valente AJ, Graves DT, Vialle-Valentin CE, Delgado R, Schwartz CJ. 1988. Purification of a monocyte chemotactic factor secreted by nonhuman primate vascular cells in culture. *Biochemistry* 27:4162–4168.
- Wu YJ, Hong CY, Lin SJ, Wu P, Shiao MS. 1998. Increase of vitamin E content in LDL and reduction of atherosclerosis in cholesterol-fed rabbits by a water-soluble antioxidant-rich fraction of *Salvia miltiorrhiza*. *Arterioscler Thromb Vasc Biol* 18:481–486.
- Wysocki SJ, Zheng MH, Smith A, Lamawansa MD, Iacopetta BJ, Robertson TA, Papadimitriou JM, House AK, Norman PE. 1996. Monocyte chemoattractant protein-1 gene expression in injured pig artery coincides with early appearance of infiltrating monocyte/macrophages. *J Cell Biochem* 62:303–313.
- Yagi A, Fujimoto K, Tanonaka K, Hirai K, Takeo S. 1989. Possible active components of tan-shen (*Salvia miltiorrhiza*) for protection of the myocardium against ischemia-induced derangements. *Planta Med* 55:51–54.
- Yla-Herttuala S, Lipton BA, Rosenfeld ME, Sarkioja T, Yoshimura T, Leonard EJ, Witztum JL, Steinberg D. 1991. Expression of monocyte chemoattractant protein 1 in macrophage-rich areas of human and rabbit atherosclerotic lesions. *Proc Natl Acad Sci USA* 88:5252–5256.
- Yoshimura T, Yuhki N, Moore SK, Appella E, Lerman MI, Leonard EJ. 1989. Human monocyte chemoattractant protein-1 (MCP-1). Full-length cDNA cloning, expression in mitogen-stimulated blood mononuclear leukocytes, and sequence similarity to mouse competence gene JE. *FEBS Lett* 244:487–493.
- Yu X, Dluz S, Graves DT, Zhang L, Antoniadis HN, Hollander W, Prusty S, Valente AJ, Schwartz CJ, Sonenshein GE. 1992. Elevated expression of monocyte chemoattractant protein 1 by vascular smooth muscle cells in hypercholesterolemic primates. *Proc Natl Acad Sci USA* 89:6953–6957.