

Molecular Analysis of RANKL-Independent Cell Fusion of Osteoclast-Like Cells Induced by TNF- α , Lipopolysaccharide, or Peptidoglycan

Hitoshi Hotokezaka,^{1*} Eiko Sakai,² Naoya Ohara,³ Yuka Hotokezaka,⁴ Carmen Gonzales,¹ Ken-ichiro Matsuo,¹ Yuji Fujimura,¹ Noriaki Yoshida,¹ and Koji Nakayama³

¹Department of Developmental and Reconstructive Medicine, Division of Orthodontics and Biomedical Engineering, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki 852-8588, Japan

²Department of Developmental and Reconstructive Medicine, Division of Oral Molecular Pharmacology, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki 852-8588, Japan

³Department of Developmental and Reconstructive Medicine, Division of Microbiology and Oral Infection, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki 852-8588, Japan

⁴Department of Developmental and Reconstructive Medicine, Division of Oral Radiology and Cancer Biology, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki 852-8588, Japan

Abstract Focusing on the final step of osteoclastogenesis, we studied cell fusion from tartrate-resistant acid phosphatase (TRAP)-positive mononuclear cells into multinuclear cells. TRAP-positive mononuclear cells before generation of multinuclear cells by cell fusion were differentiated from RAW264.7 cells by treatment with receptor activator of nuclear factor kappa B ligand (RANKL), and then the cells were treated with lipopolysaccharide (LPS), followed by culturing for further 12 h. LPS-induced cell fusion even in the absence of RANKL. Similarly, tumor necrosis factor (TNF)- α and peptidoglycan (PGN) induced cell fusion, but M-CSF did not. The cell fusion induced by RANKL, TNF- α , and LPS was specifically blocked by osteoprotegerin (OPG), anti-TNF- α antibody, and polymyxin B, respectively. LPS- and PGN-induced cell fusion was partly inhibited by anti-TNF- α antibody but not by OPG. When TRAP-positive mononuclear cells fused to yield multinuclear cells, phosphorylation of Akt, Src, extracellular signal-regulated kinase (ERK), p38MAPK (p38), and c-Jun NH2-terminal kinase (JNK) was observed. The specific chemical inhibitors LY294002 (PI3K), PP2 (Src), U0126 (MAPK-ERK kinase (MEK)/ERK), and SP600125 (JNK) effectively suppressed cell fusion, although SB203580 (p38) did not. mRNA of nuclear factor of activated T-cells c1 (NFATc1) and dendritic cell-specific transmembrane protein (DC-STAMP) during the cell fusion was quantified, however, there was no obvious difference among the TRAP-positive mononuclear cells treated with or without M-CSF, RANKL, TNF- α , LPS, or PGN. Collectively, RANKL, TNF- α , LPS, and PGN induced cell fusion of osteoclasts through their own receptors. Subsequent activation of signaling pathways involving PI3K, Src, ERK, and JNK molecules was required for the cell fusion. Although DC-STAMP is considered to be a requisite for cell fusion of osteoclasts, cell fusion-inducing factors other than DC-STAMP might be necessary for the cell fusion. *J. Cell. Biochem.* 101: 122–134, 2007. © 2006 Wiley-Liss, Inc.

Key words: TNF- α ; lipopolysaccharide; peptidoglycan; cell fusion; osteoclasts

Abbreviations used: Akt, PKB, protein kinase B; c-Fos, cellular homolog of v-fos; c-Src, cellular homolog of v-src; DC-STAMP, dendritic cell-specific transmembrane protein; ERK, extracellular signal-regulated kinase; FCS, fetal calf serum; HRP, horse radish peroxidase; IL-1 β , interleukine-1 β ; IL-6, interleukine-6; JNK, c-Jun NH2-terminal kinase; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; MEK, MAPK-ERK kinase; MyD88, myeloid differentiation factor 88; NFATc1, nuclear factor of activated T-cells c1; NF- κ B, nuclear factor kappa B; OPG, osteoprotegerin ligand; P-, phosphorylated; PBS, phosphate-buffered saline; PGN, peptidoglycan; p38, p38MAPK; p65, 65 kD subunit of NF- κ B (RelA); RANKL, receptor activator of nuclear factor kappa B ligand; SDS, sodium dodecyl sulfate; TBST, 150 mM NaCl and 0.1% Tween-20 in 25 mM Tris/HCl, pH 7.6; TLR, Toll-like receptor; TNF, tumor

necrosis factor; TRAF, tumor necrosis factor receptor-associated factor; TRAP, tartrate-resistant acid phosphatase.

Grant sponsor: Ministry of Education, Science, Sports, and Culture of Japan; Grant numbers: 15592169, DC2-16-6032.

*Correspondence to: Hitoshi Hotokezaka, Department of Developmental and Reconstructive Medicine, Division of Orthodontics and Biomedical Engineering, Graduate School of Biomedical Sciences, Nagasaki University, Sakamoto 1-7-1, Nagasaki City, Nagasaki 852-8588, Japan.
E-mail: hotoke@nagasaki-u.ac.jp

Received 21 June 2006; Accepted 12 September 2006

DOI 10.1002/jcb.21167

Bone is continuously remodeled by bone formation and resorption, and the cooperative bone metabolism is precisely regulated to maintain homeostasis. Osteoclasts, which are responsible for bone resorption in bone metabolism, are multinucleated cells originating from hematopoietic precursor cells of the monocyte/macrophage lineage. One of the key factors for osteoclastogenesis, which is induced by osteoblasts, is receptor activator of nuclear factor kappa B ligand (RANKL), a member of the tumor necrosis factor (TNF) family [Anderson et al., 1997; Wong et al., 1997; Lacey et al., 1998; Yasuda et al., 1998]. RANKL was found to be expressed by T-cells [Wong et al., 1997; Josien et al., 1999; Kong et al., 1999] and B-cells [Li et al., 2000] as well as osteoblastic/stromal cells and to be essential for osteoclast differentiation. Receptor activator of nuclear factor kappa B (RANK), which is one of TNF receptor family members, is expressed in osteoclasts and their precursor cells as the receptor of RANKL [Josien et al., 1999; Li et al., 2000; Choi et al., 2001]. Downstream signaling through RANK is essential for osteoclastogenesis.

Osteoclasts are differentiated through multiple steps that include cell fusion at the latest step of differentiation, yielding multi-nuclear cells. Dendritic cell-specific transmembrane protein (DC-STAMP) was recently found to play a role in the cell fusion of osteoclasts, which seems to support the activity of bone resorption by osteoclasts [Kukita et al., 2004; Yagi et al., 2005]; however, the factors involved in and the precise mechanism of the cell fusion are unknown.

Deviation from the normal conditions of bone resorption results in bone diseases such as osteopetrosis, osteoporosis, and bone resorptive infectious disease. Periodontal disease is the most frequent bone resorptive infectious disease and is thought to be caused mainly by infection with Gram-negative bacteria. Such an infectious and pathological condition induces inflammation, resulting in bone resorption. A major bacterial factor for inflammation has been believed to be lipopolysaccharide (LPS), a main component of the cell surface of Gram-negative bacteria. LPS has the ability to induce proinflammatory cytokines such as TNF- α , interleukine-1 β (IL-1 β), and interleukine-6 (IL-6) in various kinds of cells [Wang and Ohura, 2002]. Bacterial components such as LPS or various cytokines elicited in infectious

lesions may modulate physiological osteoclastogenesis, leading to a pathological bone resorptive condition.

In the present study, we investigated what bacterial components or cytokines affect osteoclastogenesis at the stage of cell fusion, what receptor molecules are involved in the cell fusion, and what signaling pathways are necessary for the cell fusion.

MATERIALS AND METHODS

Reagents

Anti-nuclear factor of activated T-cells c1 (NFATc1) (7A6), anti-phospho extracellular signal-regulated kinase (ERK) (E-4), anti-TRAF6 (H-274) and anti-cellular homolog of v-src (c-Src) were purchased from Santa Cruz (Santa Cruz, CA). TNF- α , anti-TNF- α , osteoprotegrin ligand (OPG), and transforming growth factor- β (TGF- β) were from R&D Systems Inc. (Mineapolis, MN). Anti-ERK, anti-p38 MAPK (p38) (5F11), anti-phospho-p38 (28B10), anti-Akt, anti-phospho-Akt, and anti-phospho-c-Jun NH2-terminal kinase (JNK) antibodies were from Cell Signaling Technology (Beverly, MA). RANKL was from Peprotech EC Ltd (London, United Kingdom). LY294002, PD169316, SB203580, SP600125, PP2, U0126, and PD98059 were purchased from Calbiochem Corp. (La Jolla, CA). M-CSF was from Kyowa Hakko Kogyo Co., Ltd (Tokyo, Japan). RANKL was from Peprotech (Rocky Hill, NJ). cDNA of mouse kidney was from Clontech. Minimal essential medium α modification (α -MEM) and all other reagents including LPS, PGN, IL-1 β , and IL-6 were obtained from Sigma (St. Louis, MO).

Cell Culture

The murine monocytic cell line RAW264.7 (ATCC, Rockville, MD) was cultured in a humidified incubator with an atmosphere of 5% CO₂ in air at 37°C and maintained on uncoated plastic dishes of 9 cm in diameter in α -MEM containing 10% (v/v) heat-inactivated fetal calf serum (FCS) with 2 mM glutamine, 100 U/ml penicillin, and 100 μ g/ml streptomycin. For subculture, the cells were resuspended with 10 ml of fresh medium by mild pipetting and transferred to a 15 ml conical tube. After sitting for 5 min, the upper 14.5 ml fraction was removed, and aggregated cells were collected from the bottom and seeded into

a fresh dish containing 12 ml of fresh medium. For osteoclastogenesis experiments, the indicated number of cells was seeded on tissue culture plates in the presence or absence of RANKL and chemical reagents.

Tartrate-Resistant Acid Phosphatase (TRAP) Staining

Cells were fixed with 4% paraformaldehyde in phosphate-buffered saline (PBS) at 4°C for more than 60 min. The cells were then treated with 0.2% Triton X-100 in PBS at room temperature for 5 min, followed by rinsing twice with PBS. Finally, the fixed cells were stained with 0.01% naphthol AS-MX phosphate (Sigma) and 0.05% fast red violet LB salt (Sigma) in the presence of 50 mM sodium tartrate and 90 mM sodium acetate (pH 5.0) for 15–60 min and then rinsed twice with PBS.

Measurement of TRAP Intensity

Following TRAP staining, the plates were scanned by a transparent light scanner, and the red color image was extracted from the scanned image using the Photoshop (Adobe Systems Inc., San Jose, CA) computer program. The intensity of the red color image was measured using National Institutes of Health (NIH) Image computer program and is represented as TRAP intensity in this paper.

Cell Proliferation Assays

Cell proliferation was measured using a Cell Counting Kit (Dojindo, Kumamoto, Japan) according to the manufacturer's instructions. Similar to the MTT [3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl 2H-tetrazolium bromide] assay, this kit measures intracellular mitochondrial dehydrogenase activity in living but not in dead cells by forming water-soluble formazan dye with the tetrazolium compound WST-1, a sodium salt of 4-[3-(4-iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazolium]-1,3-benzene disulfonate.

Immunoblotting

Cells were seeded at 5×10^6 cells/well on a 24-well plate and kept in a humidified incubator with an atmosphere of 5% CO₂ in air at 37°C for 4 h. The cells were stimulated with RANKL or/and other chemical reagents and incubated for the indicated time. After incubation, the cells in the 24-well plate were rinsed twice with ice-cold PBS, followed by addition of 50 µl of sodium

dodecyl sulfate (SDS)-sample buffer [1% SDS, 100 mM dithiothreitol (DTT), 10% glycerol, 0.0025% bromophenol blue, 62.5 mM Tris/HCl, and pH 6.8] containing 1 mM sodium orthovanadate (protein tyrosine phosphatase inhibitor), 1 mM phenylmethylsulfonyl fluoride, 1 µg/ml aprotinin, 1 µg/ml pepstatin, 1 µg/ml chymostatin, and 1 µg/ml leupeptin. The whole cell lysate was then treated in ice-cold water by sonication (two 15-s pulses with a 60-s interval) (Bioruptor UDC-200T, Cosmo Bio, Tokyo, Japan). After boiling for 5 min, 4–7 µl of the lysate (20 µg of protein) was subjected to SDS-polyacrylamide gel electrophoresis (PAGE), and the proteins in the gel were transferred to a polyvinylidene difluoride (PVDF) membrane. The membrane was incubated in 5% skim milk, 25 mM Tris/HCl (pH 7.6), 150 mM NaCl, and 0.1% Tween-20 (TBST) at room temperature for 1 h, washed twice with TBST for 5 min, and then incubated with an antibody at 2000–4000 × dilution in TBST at 4°C for 16 h. The membrane was washed three times with TBST for 10 min, incubated with a horse radish peroxidase (HRP)-conjugated second antibody at 4000–8000 × dilution in TBST at room temperature for 1 h, washed vigorously five times for 10 min, and subjected to chemiluminescence (ECL-plus, Amersham Pharmacia Biotech, Piscataway, NJ) to visualize HRP. In some experiments for reprobing, the membrane was stripped of antibody with Reblot Plus Strong (Chemicon, CA) according to the manufacturer's instructions.

Real-Time PCR

Total RNA was prepared using an RNeasy Mini Spin Column (Quiagen) according to the manufacturer's instructions, and the cDNA was reverse-transcribed by SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, CA). Using an M × 3005PTM Real-time PCR System and Brilliant SYBR Green QPCR Mastermix (Stratagene), the reverse-transcribed cDNA was amplified and quantified with specific primers according to the manufacturer's instructions. The primer sequences used for amplification were as follows: mouse GAPDH, 5'-ACCCAGAAGACTGTG-GATGG-3' and 5'-CACATTGGGGGTAGGAA-CAC-3'; NF-ATc1, 5'-TCATCCTGTCCAACA-CCAAA-3' and 5'-TCACCCTGGTGTTCCTCC-TC-3'; DC-STA-MP, 5'-GGGCACCAGTATTTT-CCTGA-3' and 5'-TGGCAGGATCCAGTAAAA-

GG-3'; and TRAP, 5'-CAGCAGCCAAGGAG-GACTAC-3' and 5'-ACATAGCCCACACCGT-TCTC-3'. The relative amounts of each mRNA were normalized by the GAPDH expression.

RESULTS

LPS-Mediated Inhibition in the Initial Step of Osteoclast Differentiation

LPS, a major component of the cell wall of Gram-negative bacteria, has been reported to strongly induce inflammation, which is thought to cause bone destruction. LPS induces bone resorption when administered *in vivo*; however, it suppresses osteoclast formation from bone marrow macrophages *in vitro* cell cultures. In

order to determine what step of osteoclastogenesis is influenced by LPS, we examined whether LPS affected RANKL-induced osteoclastogenesis when it was present in RAW264.7 cell culture at different time periods after treatment with RANKL. In a previous study, we found that 1–3 μM U0126, a MAPK-ERK kinase (MEK)/ERK inhibitor, accelerated osteoclastogenesis of RAW264.7 cells in the presence of RANKL but that 10 μM of U0126 suppressed osteoclastogenesis [Hotokezaka et al., 2002]. In culture conditions with or without 2 μM U0126, cells were treated with LPS for different periods of 12 h during the total 48-h culture. In both culture conditions, LPS suppressed osteoclastogenesis when added for the first 12 h, and the




Fig. 1. Effect of LPS on differentiation of RAW264.7 cells into osteoclast-like cells. RAW264.7 cells in 250 μl of α -MEM were cultured in a 96-well tissue culture plate in the presence of 25 ng/ml RANKL with (upper; 16,000 cells) or without (lower; 8,000 cells) 2 μM U0126. The cells were stimulated with 25 ng/ml LPS during the indicated period in the absence of RANKL. TRAP activity was visualized by TRAP staining. **A:** culture plate scanned. **B:** measured TRAP intensity, the mean of three different determinations was plotted. **C:** cell shapes under an optical microscope. Bars indicate 50 μm . [Color figure can be viewed in the online issue, which is available at www.interscience.wiley.com.]

suppression was markedly weaker when LPS was added at a later 12-h period (Fig. 1). Similar to the previous reports [Takami et al., 2002; Zou and Bar-Shavit, 2002; Zou et al., 2002; Hayashi et al., 2003], LPS inhibits osteoclastogenesis only in the early step of differentiation suggests that effects of cytokines or Toll-like receptor (TLR) ligand molecules such as LPS on osteoclastogenesis may vary among steps of differentiation.

Effects of Cytokines or TLR Ligands on the Latest Step of Osteoclastogenesis

In this study, we used an enhanced culture condition in which the cells were cultured in the presence of both RANKL and U0126 since TRAP-positive multinuclear cells were generated in the culture condition within 48 h (Fig. 1). Effects of various cytokines or TLR ligands on the latest step of osteoclastogenesis and fusion of TRAP-positive mononuclear cells were investigated. First, TRAP-positive mononuclear cells were generated from RAW264.7 cells by incubation with RANKL and U0126 for 36 h. The cells were then incubated in culture medium with various cytokines or TLR ligand molecules for 12 h in the absence of RANKL. Similar to the previous reports [Lam et al., 2000; Zou and Bar-Shavit, 2002; Zou et al., 2002], LPS, PGN, and TNF- α induced cell fusion as well as RANKL (Fig. 2). Although M-CSF failed to induce cell fusion, it increased the number of TRAP-positive mononuclear cells. IL-1 β had no ability to induce cell fusion in this condition. In addition, we examined IL-6 (0.1–50 ng/ml), TGF- β (0.1–50 ng/ml), phorbol myristic acetate (PMA; 0.1–10 μ M), concanavalin A (0.1–10 μ g/ml), amphotericin B (AmB; 0.1–10 μ g/ml), and oxidized low-density lipoprotein (LDL; 0.1–10 μ g/ml). None of them induced cell fusion (data not shown) although PMA, AmB, and oxidized LDL are known as inducers of proinflammatory cytokines in macrophages [Stuart and Hamilton, 1980; Cleary et al., 1992; Pollaud-Cherion et al., 1998]. These

results suggested that cell fusion in the latest step of osteoclastogenesis might be induced by treatment with RANKL, TNF- α , LPS, and PGN but not by treatment with others.

It is known that IL-1 β -induced cell signaling is similar to that of LPS and PGN with respect to involvement of MyD88, an adaptor molecule of IL1R family members [Subramaniam et al., 2004]. We investigated whether the IL-1 β receptor IL1RI was expressed in RAW264.7 cells. Real-time RT-PCR analysis revealed that IL1RI was not expressed in RAW264.7 cells, whereas RANK, c-fms, TLR2, and TLR4, receptors for RANKL, M-CSF, PGN, and LPS, respectively, were expressed in the cells (Fig. 2F). Absence of the IL-1 β receptor in the cell line may account for the disability of IL-1 β to induce cell fusion.

Ability of LPS and PGN to Induce TNF- α -Independent Cell Fusion

LPS and PGN are known to induce TNF- α in RAW264.7 and other cells [Remick et al., 1988; Gupta et al., 1995]. Furthermore, TNF- α was reported to induce osteoclastogenesis in macrophages exposed to permissive levels of RANKL [Lam et al., 2000]. In order to determine whether cytokines or TLR ligands directly or indirectly induced cell fusion, OPG—a neutralizing antibody against TNF- α (anti-TNF- α), and polymyxin B were used to block RANKL, TNF- α , and LPS, respectively (Fig. 3). OPG, anti-TNF- α , and polymyxin B (each 0.3 μ g/ml) specifically inhibited the osteoclast cell fusion induced by RANKL, TNF- α , and LPS, respectively. Anti-TNF- α strongly inhibited TNF- α -induced cell fusion and also had some inhibitory effects at 1 μ g/ml on LPS- and PGN-induced cell fusion (Fig. 3A). Anti-TNF- α (1 μ g/ml) completely suppressed cell fusion induced by TNF- α (10 ng/ml) and the secretion of TNF- α induced by LPS or PGN was less than 1.2 ng/ml, suggesting that LPS and PGN had the ability to induce TNF- α -independent cell fusion (Fig. 3B).

Fig. 2. Effects of factors on cell fusion of osteoclasts. RAW264.7 cells (1.6×10^3) were cultured for 36 h in 250 μ l of α -MEM containing 10% FCS, 25 ng/ml RANKL, and 2 μ M U0126 in a 96-well tissue culture plate. Then the medium was replaced with 250 μ l of α -MEM containing 10% FCS and indicated factors. Concentrations used were 50 ng/ml for M-CSF, 10 ng/ml RANKL, 10 ng/ml LPS, 100 ng/ml PGN, 10 ng/ml TNF- α , and 50 ng/ml IL-1 β . **A:** scanned culture plate. **B:** area % of multinuclear cells.

C: number of mononuclear cells. **D:** TRAP intensity. **E:** cell shapes under an optical microscope. **F:** mRNA quantified by real-time RT-PCR. Total RNAs were prepared from RAW264.7 cells treated with RANKL and U0126 for 36 h and kidney cells of mice, and they were subjected to real-time RT-PCR using primers for RANK, c-fms, TLR4, and IL-1R type 1. Bar indicates 100 μ m. [Color figure can be viewed in the online issue, which is available at www.interscience.wiley.com.]




Fig. 2.

Immunoblot Analysis During the Cell Fusion Process

An intracellular domain of RANK interacts with TNF receptor-associated factor (TRAF) 2 and TRAF 6 [Galibert et al., 1998; Wong et al., 1998, 1999; Darnay et al., 1999; Kim et al., 1999], which appear to be involved in the activation of downstream signaling molecules such as nuclear factor kappa B (NF- κ B), Src [Horne et al., 1992; Lowe et al., 1993], phosphatidylinositol 3 (PI3), protein kinase B (PKB/Akt), and mitogen-activated protein kinases (MAPKs) including p38, ERK, and JNK [Xia et al., 1995; Wong et al., 1998, 1999; Matsumoto et al., 2000; Chang and Karin, 2001; Lee et al., 2002; Wei et al., 2002]. In order to study cell signaling during the cell fusion process, we determined the amounts of several transcription factors and phosphorylation of various signaling molecules by immunoblot analysis. In our culture system, TRAP-positive mononuclear cells began to fuse at 8 h, and they were fully fused at 16 h after stimulation. As shown in Figure 2, RANKL, TNF- α , and LPS equally induced fusion of TRAP-positive mononuclear cells, but M-CSF did not. Phosphorylation of Akt, Src, ERK, JNK, and p38 was observed at 480 or 960 min in cells stimulated with RANKL, TNF- α , and LPS (Fig. 4). p38 showed the most prominent phosphorylation by stimulation with RANKL, TNF- α , and LPS throughout the incubation period. Continuous phosphorylation of Akt was observed in cells treated with M-CSF, in contrast to the transient phosphorylation in cells treated with RANKL, TNF- α , or LPS at 8 and 16 h. The continuous or transient phosphorylation of Akt may be involved in cell fate, growth, or fusion. Since phosphorylation of signaling molecules is thought to be involved in the subsequent cellular responses, we determined the importance of the signaling molecules in the cell fusion process by using specific inhibitory compounds. As expected, LY294002 (PI3K inhibitor), PP2 (Src inhibitor), U0126 (ERK inhibitor), and SP600125 (JNK inhibitor) efficiently inhibited the RANKL-induced cell fusion, but SB203580 (p38 inhibitor) had no inhibitory effect (Fig. 5). Another p38 inhibitor, PD169316, also had no effect (data not shown). In order to study whether p38 is involved in an activity of osteoclast, resorption pit formation was measured. However, resorption pit forma-


tion induced by RANKL, TNF- α , or LPS was not inhibited by the p38 inhibitor SB203580. The role of p38 during the cell fusion process in osteoclastogenesis remains unclear.

Real-Time PCR Analysis of Expression of NFATc1, DC-STAMP, and TRAP

It was recently found that DC-STAMP participates in the cell fusion of osteoclasts [Kukita et al., 2004; Yagi et al., 2005]. First, we determined levels of mRNA of NFATc1, DC-STAMP, and TRAP in cells treated with RANKL at several time points. The mRNA level of NFATc1 increased at an earlier time point than did the mRNA levels of DC-STAMP and TRAP. DC-STAMP and TRAP mRNAs increased immediately before cell fusion (Fig. 6). We then investigated the expression of NFATc1, DC-STAMP, and TRAP in the latest step of osteoclastogenesis. RAW264.7 cells that had been treated with RANKL for 36 h were then treated with RANKL, M-CSF, LPS, or TNF- α for 6 h. At this time point, the cells began to fuse. Then the expression of NFATc1, DC-STAMP, and TRAP in the cells was determined by real-time PCR analysis. Expression of DC-STAMP in RANKL- or LPS-treated cells was not significantly different from that in M-CSF-treated cells or non-treated cells. The M-CSF-treated cells and non-treated cells showed no cell fusion, whereas RANKL-, LPS-, and TNF- α -treated cells showed cell fusion. We could not find any significant difference in expression of NFATc1, DC-STAMP, and TRAP between the fused cells and non-fused cells. These results suggest that an essential factor other than DC-STAMP may determine whether TRAP-positive mononuclear cells are fused or not.

DISCUSSION

In an *in vitro* culture system using osteoclast precursor cells purified from various tissues such as bone marrow, spleen, and liver, it is difficult to avoid contamination with other cell lineages such as T-cells and mesenchymal stromal cells. In this study, we used RAW264.7—a murine macrophage cell line that can differentiate into osteoclast-like cells in the presence of RANKL [Hsu et al., 1999]. There are some characteristic differences between RAW264.7 cells and macrophages; for example, RAW264.7 cells do not respond to IL-1 β , and



the osteoclast-like cells differentiated from RAW264.7 cells form smaller and shallower resorption pits on dentin slices than do osteoclasts derived from bone marrow macrophages. However, the cell line is useful to analyze the detailed mechanisms of osteoclast differentiation.

In this study, we focused on the cell fusion step of osteoclastogenesis. Several investigators reported the later stage of osteoclastogenesis [Lam et al., 2000; Zou and Bar-Shavit, 2002; Zou et al., 2002]. However, this process is not characterized well since it is not easy to isolate cells at each step during the process of osteoclast differentiation. In this study, we used an enhanced culture condition for osteoclastogenesis of RAW264.7 cells. By culturing the cell line in the presence of RANKL and the MEK/ERK inhibitor U0126, the culture period for osteoclastogenesis was shortened to 48 h, less than half of the standard culture period, and almost all the cells are fused to one another at the end of the culture period. Therefore, the TRAP-positive mononuclear cells (preosteoclasts) prepared shortly before cell fusion in the enhanced culture condition appear to be almost homogeneous in terms of differentiation.

In the present study, LPS suppressed RANKL-induced osteoclast formation at the initial step; in contrast, at the latest step it induced cell fusion without RANKL, resulting in osteoclast formation. The suppressive effect of LPS on osteoclastogenesis, which was observed in the present study when LPS and RANKL were simultaneously added at the initial step, has previously been reported for bone marrow macrophages by Takami et al. [2002]. Moreover, it has been reported that LPS promotes the survival of mature osteoclasts via TLR4 [Itoh et al., 2003]. Taken together, the results indicate that the effects of LPS on

Fig. 3. RANKL, TNF- α , LPS, and PGN induced cell fusion specifically through their receptors. RAW264.7 cells (1.6×10^5) were cultured for 36 h in 250 μ l of α -MEM containing 10% FCS, 25 ng/ml RANKL, and 2 μ M U0126 in a 96-well tissue culture plate. Then the medium was replaced with 250 μ l of α -MEM containing 10% FCS, indicated stimulation factors (RANKL, TNF- α , LPS, and PGN), and blocking reagents (OPG, PMXB, and Anti-TNF- α ; indicated number μ g/ml were used). The fused cell area was measured as percent of total plate area. Concentrations used were 10 ng/ml RANKL, 10 ng/ml TNF- α , 10 ng/ml LPS, and 100 ng/ml PGN (A). Amounts of TNF- α in the culture media stimulated with RANKL, LPS, and PGN were measured by ELISA (B). The mean of three different determinations was plotted.







Fig. 4. Western blotting analysis during cell fusion process. The cells were treated with 25 ng/ml RANKL, 25 ng/ml TNF- α , 25 ng/ml LPS, or 100 ng/ml PGN for the indicated time and then subjected to Western blotting analysis probing with antibodies against c-Fos, NFATc1, phosphorylated (P-) Akt, P-Src, P-ERK, P-p38, P-p65 (RelA, 65 kD subunit of NF- κ B), Akt, TRAF6, tubulin, and P-JNK.


osteoclastogenesis vary among steps of osteoclast differentiation. In this context, it is notable that LPS induces bone resorption when injected into bone surfaces of mice [Umezumi et al., 1989]. Since osteoclast precursor cells of various

steps of differentiation exist together in an *in vivo* condition, osteoclast precursor cells of the pre-fusion step around bones might differentiate into mature osteoclasts in response to LPS, resulting in bone resorption.

RANKL	-	+	+	+	+	+	+	+
	-	-	LY294002	PP2	PP1 analog	U0126	SB203580	SP600125



(-)	RANKL	TNF α	LPS
	SB203580	SB203580	SB203580
	LY294002	LY294002	LY294002



Cell fusion is seen in muscle, nerves, bone in their development, and in the liver in its repair and regeneration [Ogle et al., 2005]. TRAP-positive multinuclear osteoclasts appear in bone. TRAP-positive mononuclear cells fuse to one another because the cell fusion increases cell size and enables the cells to resorb bone to a larger extent [Vignery, 2005]. This explanation is supported by the finding that a DC-STAMP knockout mouse in which TRAP-positive mononuclear cells do not fuse to one another shows an increase in bone density due to a decrease in bone resorption by osteoclasts [Yagi et al., 2005]. Osteoclasts seem to have the same origin in cell lineage as that of macrophages. Macrophages sometimes fuse with one another during infection and tissue repair. The resulting multinucleated macrophages effectively phagocytise pathogens and repair tissues. The mechanism by which macrophages repair tissues seems to be similar to bone resorption of osteoclasts. In this study, inflammatory factors such as TNF- α , LPS, and PGN induced cell fusion, which may be one of mechanisms to repair the inflammatory circumstance of bone, although the inflammation results in an unfavorable bone loss.

Cell fusion-inducing factors include several groups. One is known as fusogens that can directly induce cell fusion. And other groups that include receptors, signaling proteins, transcription factors, and proteins organizing cytoskeleton and membrane, indirectly induce cell fusion [Ogle et al., 2005]. Although EFF-1 of *Caenorhabditis elegans* might be the only one fusogen that have been found in higher eukaryotes so far as we know [Kontani and Rothman, 2005], no fusogens have been found in mammalian cells. Some molecules such as meltrin- α [Harris et al., 1997; Inoue et al., 1998], CD47, MFR (also reported as SIRP,

Fig. 5. Effects of chemical inhibitors of signal transduction on cell fusion and pit formation. RAW264.7 cells (1.6×10^5) were cultured for 36 h in 250 μ l of α -MEM containing 10% FCS, 25 ng/ml RANKL, and 2 μ M U0126 in a 96-well tissue culture plate. Then the medium was replaced with 250 μ l of α -MEM containing 10% FCS, 25 ng/ml RANKL, TNF- α , and 10 of chemical inhibitors, and the cells were cultured for a further 12 h. The fused cell area was measured as percent of total plate area (A). The TRAP intensity and cell viability were also measured (B and C). D: Pit formation was measured on a plate coated with calcium phosphate (BD BioCoat Osteologic Bone Cell Culture System, Nippon BD, Tokyo, Japan) stimulated with 25 ng/ml RANKL, TNF- α , or LPS in the presence or absence of 10 μ M SB203580 or LY294002.




Fig. 6. Real-time PCR for NFATc1, DC-STAMP, and TRAP. RAW264.7 cells (1.6×10^5) were cultured for 36 h in 250 μ l of α -MEM containing 10% FCS, 25 ng/ml RANKL, and 2 μ M U0126 in a 96-well tissue culture plate (A). Then the medium was replaced with 250 μ l of α -MEM containing 10% FCS, 25 ng/ml RANKL, or TNF- α , and the cells were cultured for further 6 h (B), followed by RNA preparation and real-time RT-PCR. Each value was normalized by that of GAPDH. The mean of three different determinations was plotted.

SHPS-1, BIT, and MyD-1) [Vignery, 2005], and DC-STAMP have been suggested as cell fusion-inducing factors in osteoclasts. In the bones of mice lacking DC-STAMP multinuclear osteoclasts were completely absent, although development of mononuclear osteoclasts was normal. The DC-STAMP-deficient mice suffer from mild osteopetrosis probably because mononuclear osteoclasts can still resorb bones [Yagi et al., 2005]. The expression of DC-STAMP was not significantly different between fused and non-fused cells in this study, suggesting the presence of other cell fusion-inducing factor(s) that directly function at the latest stage of osteoclastogenesis. Further study is needed to understand the molecular mechanisms of cell fusion of osteoclasts.

ACKNOWLEDGMENTS

We thank the members of Division of Microbiology and Oral Infection for helpful comments and discussions. This work was supported by Grants-in-Aid from the Ministry of Education,

Science, Sports, and Culture of Japan to Hotokezaka (#15592169) and Matsuo (#DC2-16-6032).

REFERENCES

- Anderson DM, Maraskovsky E, Billingsley WL, Dougall WC, Tometsko ME, Roux ER, Teepe MC, DuBose RF, Cosman D, Galibert L. 1997. A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature* 390:175–179.
- Chang L, Karin M. 2001. Mammalian MAP kinase signalling cascades. *Nature* 410:37–40.
- Choi Y, Woo KM, Ko SH, Lee YJ, Park SJ, Kim HM, Kwon BS. 2001. Osteoclastogenesis is enhanced by activated B cells but suppressed by activated CD8(+) T cells. *Eur J Immunol* 31:2179–2188.
- Cleary JD, Chapman SW, Nolan RL. 1992. Pharmacologic modulation of interleukin-1 expression by amphotericin B-stimulated human mononuclear cells. *Antimicrob Agents Chemother* 36:977–981.
- Darnay BG, Ni J, Moore PA, Aggarwal BB. 1999. Activation of NF-kappaB by RANK requires tumor necrosis factor receptor-associated factor (TRAF) 6 and NF-kappaB-inducing kinase. Identification of a novel TRAF6 interaction motif. *J Biol Chem* 274:7724–7731.
- Galibert L, Tometsko ME, Anderson DM, Cosman D, Dougall WC. 1998. The involvement of multiple tumor

- necrosis factor receptor (TNFR)-associated factors in the signaling mechanisms of receptor activator of NF- κ B, a member of the TNFR superfamily. *J Biol Chem* 273:34120–34127.
- Gupta D, Jin YP, Dziarski R. 1995. Peptidoglycan induces transcription and secretion of TNF- α and activation of lyn, extracellular signal-regulated kinase, and rsk signal transduction proteins in mouse macrophages. *J Immunol* 155:2620–2630.
- Harris HA, Murrills RJ, Komm BS. 1997. Expression of meltrin- α mRNA is not restricted to fusagenic cells. *J Cell Biochem* 67:136–142.
- Hayashi S, Yamada T, Tsuneto M, Yamane T, Takahashi M, Shultz LD, Yamazaki H. 2003. Distinct osteoclast precursors in the bone marrow and extramedullary organs characterized by responsiveness to Toll-like receptor ligands and TNF- α . *J Immunol* 171:5130–5139.
- Horne WC, Neff L, Chatterjee D, Lomri A, Levy JB, Baron R. 1992. Osteoclasts express high levels of pp60c-src in association with intracellular membranes. *J Cell Biol* 119:1003–1013.
- Hotokezaka H, Sakai E, Kanaoka K, Saito K, Matsuo K, Kitaura H, Yoshida N, Nakayama K. 2002. U0126 and PD98059, specific inhibitors of MEK, accelerate differentiation of RAW264.7 cells into osteoclast-like cells. *J Biol Chem* 277:47366–47372.
- Hsu H, Lacey DL, Dunstan CR, Solovyev I, Colombero A, Timms E, Tan HL, Elliott G, Kelley MJ, Sarosi I, Wang L, Xia XZ, Elliott R, Chiu L, Black T, Scully S, Capparelli C, Morony S, Shimamoto G, Bass MB, Boyle WJ. 1999. Tumor necrosis factor receptor family member RANK mediates osteoclast differentiation and activation induced by osteoprotegerin ligand. *Proc Natl Acad Sci USA* 96:3540–3545.
- Inoue D, Reid M, Lum L, Kratzschmar J, Weskamp G, Myung YM, Baron R, Blobel CP. 1998. Cloning and initial characterization of mouse meltrin β and analysis of the expression of four metalloprotease-disintegrins in bone cells. *J Biol Chem* 273:4180–4187.
- Itoh K, Udagawa N, Kobayashi K, Suda K, Li X, Takami M, Okahashi N, Nishihara T, Takahashi N. 2003. Lipopolysaccharide promotes the survival of osteoclasts via Toll-like receptor 4, but cytokine production of osteoclasts in response to lipopolysaccharide is different from that of macrophages. *J Immunol* 170:3688–3695.
- Josien R, Wong BR, Li HL, Steinman RM, Choi Y. 1999. TRANCE, a TNF family member, is differentially expressed on T cell subsets and induces cytokine production in dendritic cells. *J Immunol* 162:2562–2568.
- Kim HH, Lee DE, Shin JN, Lee YS, Jeon YM, Chung CH, Ni J, Kwon BS, Lee ZH. 1999. Receptor activator of NF- κ B recruits multiple TRAF family adaptors and activates c-Jun N-terminal kinase. *FEBS Lett* 443:297–302.
- Kong YY, Feige U, Sarosi I, Bolon B, Tafuri A, Morony S, Capparelli C, Li J, Elliott R, McCabe S, Wong T, Campagnuolo G, Moran E, Bogoch ER, Van G, Nguyen LT, Ohashi PS, Lacey DL, Fish E, Boyle WJ, Penninger JM. 1999. Activated T cells regulate bone loss and joint destruction in adjuvant arthritis through osteoprotegerin ligand. *Nature* 402:304–309.
- Kontani K, Rothman JH. 2005. Cell fusion: EFF is enough. *Curr Biol* 15:R252–R254.
- Kukita T, Wada N, Kukita A, Kakimoto T, Sandra F, Toh K, Nagata K, Iijima T, Horiuchi M, Matsusaki H, Hieshima K, Yoshie O, Nomiyama H. 2004. RANKL-induced DC-STAMP is essential for osteoclastogenesis. *J Exp Med* 200:941–946.
- Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ. 1998. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 93:165–176.
- Lam J, Takeshita S, Barker JE, Kanagawa O, Ross FP, Teitelbaum SL. 2000. TNF- α induces osteoclastogenesis by direct stimulation of macrophages exposed to permissive levels of RANK ligand. *J Clin Invest* 106:1481–1488.
- Lee SE, Woo KM, Kim SY, Kim HM, Kwack K, Lee ZH, Kim HH. 2002. The phosphatidylinositol 3-kinase, p38, and extracellular signal-regulated kinase pathways are involved in osteoclast differentiation. *Bone* 30:71–77.
- Li J, Sarosi I, Yan XQ, Morony S, Capparelli C, Tan HL, McCabe S, Elliott R, Scully S, Van G, Kaufman S, Juan SC, Sun Y, Tarpley J, Martin L, Christensen K, McCabe J, Kostenuik P, Hsu H, Fletcher F, Dunstan CR, Lacey DL, Boyle WJ. 2000. RANK is the intrinsic hematopoietic cell surface receptor that controls osteoclastogenesis and regulation of bone mass and calcium metabolism. *Proc Natl Acad Sci USA* 97:1566–1571.
- Lowe C, Yoneda T, Boyce BF, Chen H, Mundy GR, Soriano P. 1993. Osteopetrosis in Src-deficient mice is due to an autonomous defect of osteoclasts. *Proc Natl Acad Sci USA* 90:4485–4489.
- Matsumoto M, Sudo T, Saito T, Osada H, Tsujimoto M. 2000. Involvement of p38 mitogen-activated protein kinase signaling pathway in osteoclastogenesis mediated by receptor activator of NF- κ B ligand (RANKL). *J Biol Chem* 275:31155–31161.
- Ogle BM, Cascalho M, Platt JL. 2005. Biological implications of cell fusion. *Nat Rev Mol Cell Biol* 6:567–575.
- Pollaud-Cherion C, Vandaele J, Quartulli F, Seguelas MH, Decerprit J, Pipy B. 1998. Involvement of calcium and arachidonate metabolism in acetylated-low-density-lipoprotein-stimulated tumor-necrosis-factor- α production by rat peritoneal macrophages. *Eur J Biochem* 253:345–353.
- Remick DG, Scales WE, May MA, Spengler M, Nguyen D, Kunkel SL. 1988. In situ hybridization analysis of macrophage-derived tumor necrosis factor and interleukin-1 mRNA. *Lab Invest* 59:809–816.
- Stuart RK, Hamilton JA. 1980. Tumor-promoting phorbol esters stimulate hematopoietic colony formation in vitro. *Science* 208:402–404.
- Subramaniam S, Stansberg C, Cunningham C. 2004. The interleukin 1 receptor family. *Dev Comp Immunol* 28:415–428.
- Takami M, Kim N, Rho J, Choi Y. 2002. Stimulation by toll-like receptors inhibits osteoclast differentiation. *J Immunol* 169:1516–1523.
- Umezawa A, Kaneko N, Toyama Y, Watanabe Y, Itoh H. 1989. Appearance of osteoclasts by injections of lipopolysaccharides in rat periodontal tissue. *J Periodontol Res* 24:378–383.
- Vignery A. 2005. Macrophage fusion: Are somatic and cancer cells possible partners? *Trends Cell Biol* 15:188–193.

- Wang PL, Ohura K. 2002. Porphyromonas gingivalis lipopolysaccharide signaling in gingival fibroblasts-CD14 and Toll-like receptors. *Crit Rev Oral Biol Med* 13:132–142.
- Wei S, Wang MW, Teitelbaum SL, Ross FP. 2002. Interleukin-4 reversibly inhibits osteoclastogenesis via inhibition of NF-kappa B and mitogen-activated protein kinase signaling. *J Biol Chem* 277:6622–6630.
- Wong BR, Rho J, Arron J, Robinson E, Orlinick J, Chao M, Kalachikov S, Cayani E, Bartlett FS, 3rd, Frankel WN, Lee SY, Choi Y. 1997. TRANCE is a novel ligand of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. *J Biol Chem* 272:25190–25194.
- Wong BR, Josien R, Lee SY, Vologodskaja M, Steinman RM, Choi Y. 1998. The TRAF family of signal transducers mediates NF-kappaB activation by the TRANCE receptor. *J Biol Chem* 273:28355–28359.
- Wong BR, Besser D, Kim N, Arron JR, Vologodskaja M, Hanafusa H, Choi Y. 1999. TRANCE, a TNF family member, activates Akt/PKB through a signaling complex involving TRAF6 and c-Src. *Mol Cell* 4:1041–1049.
- Xia Z, Dickens M, Raingeaud J, Davis RJ, Greenberg ME. 1995. Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 270:1326–1331.
- Yagi M, Miyamoto T, Sawatani Y, Iwamoto K, Hosogane N, Fujita N, Morita K, Ninomiya K, Suzuki T, Miyamoto K, Oike Y, Takeya M, Toyama Y, Suda T. 2005. DC-STAMP is essential for cell-cell fusion in osteoclasts and foreign body giant cells. *J Exp Med* 202: 345–351.
- Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinosaki M, Mochizuki S, Tomoyasu A, Yano K, Goto M, Murakami A, Tsuda E, Morinaga T, Higashio K, Udagawa N, Takahashi N, Suda T. 1998. Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci USA* 95:3597–3602.
- Zou W, Bar-Shavit Z. 2002. Dual modulation of osteoclast differentiation by lipopolysaccharide. *J Bone Miner Res* 17:1211–1218.
- Zou W, Schwartz H, Endres S, Hartmann G, Bar-Shavit Z. 2002. CpG oligonucleotides: Novel regulators of osteoclast differentiation. *FASEB J* 16:274–282.