


Scheme I



for 24 h. Then water was added and extracted several times with chloroform. After evaporation of the solvent the residue was acetylated with pyridine-acetic anhydride; yield 7.5 g; mp 132–134 °C (acetone-water); IR V_{max} 1730 cm^{-1} ; 1240 cm^{-1} (CH_3CO); 1610 cm^{-1} ($\text{C}=\text{C}$).

3-β-Acetoxy-5-pregnen-20,21-dione-1'-phenyl-5'-(2-pyridinyl)pyrazoline (III). To a solution of II (4.7 g) in ethanol (100 mL) containing hydrochloric acid (3 mL), phenylhydrazine (1.2 g) is added. The resultant solution is refluxed for 24 h. Then water was added and neutralized with ammonium hydroxide. The precipitate was collected by filtration, dried, and acetylated with pyridine-acetic anhydride at room temperature. Then the usual work up and column chromatography from silica gel (eluent chloroform) gave 4.0 g of III. Recrystallization from methanol gives mp 115–117 °C; IR V_{max} 1730 cm^{-1} ; 1250 cm^{-1} (CH_3CO).

3-Oxo-4-pregnen-20,21-dione-1'-phenyl-5'-(2-pyridinyl)pyrazole (IV). Pyrazoline (III) (1.5 g) was dissolved in 200 mL of methanol containing 2 g of sodium hydroxide. The mixture was refluxed for 1.5 h. The reaction mixture was poured into water and the precipitate, collected by filtration, was washed several times with water and dried to yield III β (1.3 g).

To a solution of III β (0.5 g) in 7 mL of cyclohexanone, 20 mL of dry dioxane and 20 mL of dry toluene was distilled slowly

as a solution of aluminum isopropylate (0.8 g) in dry toluene (6 mL) was added. Distillation was continued for 2 h as 15 mL of toluene was added and 30 mL of distillate was collected. Then the mixture was refluxed for 4 h and left to stand at room temperature overnight. The mixture was filtered to remove the precipitate containing the aluminum. The filtrate was distilled, extracted with chloroform, and evaporated. The volatile constituents were removed by steam distillation. The residue was extracted with chloroform and evaporated. Column chromatography from silica gel (eluent chloroform:methanol, 98:2) gave pyrazole IV (1.15 g), which was recrystallized from chloroform-methanol; mp 249–251 °C; IR V_{max} 1670 cm^{-1} ; 1615 cm^{-1} ($\text{C}=\text{C}$).

Registry No. I, 1778-02-5; (E)-II, 110027-50-4; (Z)-II, 110027-56-0; (E)-II-AcOH, 110027-51-5; (Z)-II-AcOH, 110027-57-1; IIIa, 110027-52-6; IIIa-AcOH, 110027-53-7; IIIb, 110027-54-8; IV, 110027-55-9; pyridine-2-carboxaldehyde, 1121-60-4.

Literature Cited

- (1) Catsoulacos, P.; Stassinopoulou, C. I. *J. Heterocycl. Chem.* **1978**, *15*, 313.
- (2) Elderfield, R. C. *Heterocyclic Compounds*, Wiley: New York, 1957; Vol. 5, pp 48, 49.

Received for review February 13, 1986. Accepted May 26, 1987.

2-Pyrazinyl-2-arylalkanenitriles

Alan R. Katritzky,* Konstantina Yannakopoulou, Julie Thomson, Franciszek Saczewski, and Boguslaw Pilarski†

Department of Chemistry, University of Florida, Gainesville, Florida 32611

A series of new 2-alkyl-2-(3,4-dialkylphenyl)-2-pyrazineacetonitriles (3) was prepared from readily available starting materials by a simple, efficient, two-step sequence. The products are potential herbicides.

Compounds of the type 2 and 3 have been reported in the patent literature (1, 2) to possess sedative and anticonvulsive

*Department of Chemistry, Medical Academy, 80-416 Gdansk, Poland.

activity. Pyrazineacetonitriles (3) (Scheme I) were prepared in our lab for testing as herbicides.


2-Substituted acetonitriles were reacted at room temperature with 2-chloropyrazine under basic conditions (NaOH) in dimethyl sulfoxide (DMSO) (3, 4) to afford 2 in high yields. Syntheses of these compounds previously reported (1) used drastic conditions (NaNH_2 /toluene, reflux or liquor ammonia/dioxane) and the yields were lower (27–80%) than ours (70–90%). Compounds 2 with base (see Table I) and the appropriate alkyl halide gave 3 in fair to very good yields, usually by simple workup. Thus, quenching of the reaction mixture with water

Table I. Physical Properties and NMR Data of Compounds 3

compd	yield, %	mp, °C, or bp, °C/mmHg	recryst solvent	¹ H NMR chem shifts ^a for R ³ group ^b
3a	82 ^c	155–157/0.6		2.6 (m, 2 H), 1.4 (m, 4 H), 0.8 (m, 3 H)
3b	72 ^c	77–78	MeOH ⁿ	4.1 (CH ₂ -S, dd, 2H, <i>J</i> = 15 Hz)
3c	78 ^c	94–96	MeOH	7.0 (ap. s, 4 H), 3.8 (dd, 2 H, CH ₂), 2.2 (s, CH ₃)
3d	52 ^c	178–179	DMF/H ₂ O	8.2 (d, 2 H), ^r 7.2 (d, 2 H), ^r 4.1 (ap. s, 2 H, CH ₂) (in DMSO- <i>d</i> ₆)
3e	49 ^c	192–193	DMF/H ₂ O	6.7 (d, enolic H, <i>J</i> = 1.5 Hz)
3f	80 ^d	87–89	MeOH/hexane	2.8 (m, 4 H)
3g	62 ^e	76–79	<i>f, g</i>	3.7 (dd, 2 H, <i>J</i> = 17 Hz)
3h	24 ^h	oil	<i>f, m</i>	5.7 (dd, 2 H, <i>J</i> = 15 Hz)
3i	63 ⁱ	97–99	MeOH ⁱ	4.1 (dd, 2 H, <i>J</i> = 14 Hz)
3j	34 ^j	91–93	MeOH/CH ₂ Cl ₂	2.7 (m, 4 H)
3k	70 ⁱ	95–97	MeOH	6.9 (AA'BB', 4 H, <i>J</i> = 4 Hz), 3.8 (dd, 2 H, <i>J</i> = 14 Hz), 3.8 (s, 3 H)
3l	77 ^c	108–109	MeOH	3.8 (dd, 2 H, <i>J</i> = 14 Hz)
3m	66 ^c	138–139	MeOH	3.8 (dd, 2 H, <i>J</i> = 14 Hz), 2.3 (s, 6 H)
3n	69 ^c	113–115	MeOH	3.8 (dd, 2 H, <i>J</i> = 13 Hz)
3o	69 ^c	86–87	MeOH	4.0–3.4 (m, 3 H), 2.3 (s, 3 H)
3p	61 ^k	89–91	MeOH	2.2 (s, 3 H)
3q	76 ^k	~130/0.1	oil	6.5–5.2 (m, CH=CH ₂), 3.4 (t, CH ₂)
3r	79	111–113	MeOH	3.7 (2 H, CH ₂ -CO ₂ Et), 4.2 (q, CH ₂ CH ₃), 1.2 (t, CH ₂ CH ₃)
3s	74 ^c	205–207/0.4	oil	3.8 (CH ₂ -Ph, covered by adjacent peaks)
3t	77	172–174	MeOH	9.9 (br s, 1 H), 3.9 (q, <i>J</i> = 16 Hz)
3u	61	l		10.9 (br s, 1 H), 3.3–2.3 (m, 4 H)

^aThe spectra were recorded on a Varian EM 360 L (60 MHz) instrument, for solutions in deuteriochloroform (tetramethylsilane as the internal standard) unless otherwise stated. ^bPeaks due to aromatic protons are not included, since they always come together with all other aromatic peaks, making assignment impossible. In all cases, however, integration showed the expected number of aromatic protons. ^cConditions: 50% aqueous NaOH (5 mL), DMSO (10 mL), **2** (20 mmol), at 60–70 °C for 3–4 h. ^dConditions: *t*-BuOH (4 mL), 30% NaOH (1 drop) in *t*-BuOH, **2** (2 mmol), at 60 °C for 6 h. ^eIn anhydrous DMSO (4 mL) and powdered NaOH (see ref 3). ^fPurified by column chromatography on silica gel. ^gThe obtained oil crystallized upon standing. ^hIn NaH (60% oil dispersion)/toluene, reflux, incomplete reaction after 2 days. ⁱIn NaH (60% oil dispersion)/DMF, overnight stirring (see ref 5). ^jIn excess Na/absolute EtOH, reflux, 3 days. ^kConditions: 50% aqueous NaOH (4 mL), DMSO (20 mL), **2** (20 mmol), at 60–70 °C for 4–5 h. ^lThe solid was extremely hydroscopic and was characterized only by its NMR spectra (see Tables I and II). ^mThe necessary alkyl halide, chloromethyl-1-benzotriazole, was prepared as described in ref 6. ⁿFor the preparation of the necessary alkyl halide, see ref 7. ^oProduct was first distilled (bp 180–185 °C/0.4 mmHg) and then it crystallized on standing. ^pConditions: **2d** (50 mmol) in DMSO (150 mL)/50% aqueous NaOH (2 mL). ^qThe product was obtained on hydrolysis (NaOH) of **3r** in water (20 min). ^r*J* = 8 Hz. ^ss, 3 H.

Table II. ¹³C NMR Chemical Shifts for Compounds 3^c




compd	pyrazinyl group		R ¹	R ²	C(CN	R ³
3b	152.2, 143.8, 143.1						53.8 119.6 134.8, 131.4, 128.6, 128.5, 44.2
3c	153.8, 144.2, 143.7, 143.3						53.7 120.2 137.8, 131.4, 130.2, 44.2 (CH ₂), 20.8 (CH ₃)
3d ^b	144.7, 144.3, 143.2 ^b						53.7 b 147.0, 131.8, 123.2 ^b
3f	151.9, 144.1, 143.8, 143.3						50.9 119.0 ^c 117.6 (-CN), ^c 33.9, 13.6
3g	150.9, 144.8, 144.4, 143.3						49.2 118.4 115.5 (-CN), 28.6
3h	151.5, 144.6, 143.2						52.6 118.9 145.4, 133.5, 127.4, 123.8, 119.8, 109.4, 54.4
3i	153.7, 143.9, 143.5, 143.2						52.1 120.2 155.0, 149.0, 135.9, 124.2, 121.9, 46.0
3j	153.6, 143.6, 143.4						52.1 120.3 40.4, ^c 31.7
3k	153.6, 144.1, 143.7, 143.3						53.8 120.2 158.7, 131.3, 126.4, 113.4, 54.9, 43.7
3l	153.9, 144.1, 143.6, 143.2		20.7				53.3 120.2 134.8, 129.4, 127.1, 126.7, 44.5 (CH ₂)
3m	154.0, 144.1, 143.5, 143.2		20.7				53.4 120.3 136.7, 131.6, 129.4, 126.8, 44.2 (CH ₂), 20.7
3o	154.2, 144.3, 143.6, 143.3		55.3				53.2 120.5 136.9, 131.7, 130.0, 128.2, 44.5, 20.9
3p	155.0, 143.4, 143.2, 142.9		149.2, 148.9, 131.0, 118.4, 111.1, 109.5	55.8 55.7			46.6 121.8 26.4
3q	153.7, 144.5, 143.6, 143.4		149.1, 148.8, 129.5, 118.3, 111.1, 109.7	55.8 55.7	51.7 119.8		131.25, 120.6, 42.8
3r	153.3, 144.0, 143.7, 143.0		149.3, 149.1, 129.3, 118.6, 111.3, 109.4	55.9 55.8	48.0 120.3		168.4 (-CO ₂ -), 61.0, 43.0 (-CH ₂), 13.8
3u	153.8, 143.8, 143.5, 143.4		149.4, 149.2, 128.9, 118.9, 111.4, 109.5	55.8 55.8	51.1 120.1		176.7 (-CO ₂ H), 38.4, 30.2

^aThe spectra were recorded on a JEOL FX-100 (25 MHz, FT mode) spectrometer. Deuteriochloroform ($\delta = 77.0$) was the solvent, unless otherwise stated. The chemical shifts were expressed in parts per million (ppm) relative to the deuterium signal of the solvent. ^bThe solubility of this compound in DMSO-*d*₆ ($\delta = 39.5$) is very low; therefore some of the peaks are missing in the noise. ^cAssignments can be reversed. ^dSignals at 128.9, 128.1, 126.2, and 126.0 correspond to both aromatic nuclei; some peaks overlap.

most often precipitated the product; otherwise extraction with chloroform was used. Column chromatography was necessary in only the cases of **3g**, **3h**, and **3l**. The products were characterized by ¹H (60 MHz) and ¹³C (25 MHz) NMR spectroscopy. Satisfactory elemental analyses and/or high-resolution mass spectra were obtained for all compounds, except for **3u**, which,

due to its extreme hydroscopicity, was characterized only by its ¹H and ¹³C NMR spectra. The proton NMR spectra of all compounds showed a doublet ($\delta \sim 9.0$, 1 H, *J* = 1 Hz), and a multiplet ($\delta \sim 8.8$, 2 H) due to the pyrazinyl group, and also a multiplet ($\delta \sim 7.0$ – 8.0) due to the protons of the aromatic nucleus bearing R¹ and R². These latter signals frequently over-

Scheme I



lapped with peaks due to aromatic protons of R³ (Table I). In all cases where R¹ = CH₃, the protons appeared at $\delta \sim 2.2$,

whereas the methoxy protons, for R¹ or R² = OMe, resonated at 4.0 ppm.

Registry No. 2a, 1080-87-1; 2b, 1082-48-0; 2c, 1084-83-9; 2d, 1088-67-1; 3a, 109929-53-5; 3b, 109929-54-6; 3c, 109929-55-7; 3d, 109929-56-8; 3e, 109929-57-9; 3f, 4422-44-0; 3g, 4190-80-1; 3h, 109929-58-0; 3i, 109929-59-1; 3j, 109929-60-4; 3k, 109929-61-5; 3l, 109929-62-6; 3m, 109929-63-7; 3n, 109929-64-8; 3o, 109929-65-9; 3p, 109929-66-0; 3q, 109929-67-1; 3r, 109929-68-2; 3s, 109929-69-3; 3t, 109929-70-6; 3u, 109929-71-7; Me(CH₂)₃Br, 109-65-9; Br(CH₂)SPh, 35572-08-8; 4-MeC₆H₄CH₂Br, 104-81-4; 4-NO₂C₆H₄CH₂Br, 100-11-8; BrCH₂COPh, 70-11-1; Br(CH₂)₂CN, 2417-90-5; BrCH₂CN, 590-17-0; Br-(CH₂)₂Ph, 103-63-9; 4-MeOC₆H₄CH₂Br, 2746-25-0; PhCH₂Br, 100-39-0; MeBr, 74-83-9; CH₂=CHCH₂Br, 106-95-6; MeCO₂Et, 105-36-2; Br-(C-H₂)₂CO₂H, 590-92-1; 1-(chloromethyl)benzotriazole, 54187-96-1; 2-(bromomethyl)pyridine, 55401-97-3.

Literature Cited

- (1) Akkerman, A. M.; Kofman, H.; de Vries, G. Neth. Patent 105 432, 1963; *Chem. Abstr.* **1965**, *62*, 6495c.
- (2) Akkerman, A. M.; Kofman, H. de Vries, G. Ger. Patent 1 101 425, 1961; *Chem. Abstr.* **1962**, *57*, 842e.
- (3) Johnstone, R. A. W.; Rose, M. E. *Tetrahedron* **1979**, *35*, 2169.
- (4) Pliarski, B.; Foks, H. Polish Patent, Application P-251771, 1986.
- (5) Dyke, S. F.; Tilley, E. P.; White, A. W. C.; Gale, D. P. *Tetrahedron* **1975**, *31*, 1219.
- (6) Burckhalter, J. H.; Stephens, V. C.; Hall, L. A. R. *J. Am. Chem. Soc.* **1952**, *74*, 3868.
- (7) Trost, B. M.; Kunz, R. A. *J. Org. Chem.* **1974**, *39*, 2648.

Received for review February 11, 1987. Accepted May 18, 1987.

Heterocycles. 14. Synthesis of 5H-Indenopyrimidines

Nizar El-Rayyes,* Sheekha Al-Qatami, and Mustafa Edun

Department of Chemistry, Kuwait University, Kuwait

1-Indanone (I) was reacted with aryl aldehydes (II) to give the corresponding 2-arylidene-1-indanones (III). Condensation of the chalcones III with guanidine revealed the formation of the corresponding 5H-2-amino-4-arylideneopyrimidines (V). The structures of all products were substantiated by chemical and spectroscopic methods.

Various aromatic and heterocyclic aldehydes (IIa-k) were reacted with 1-indenone to give the corresponding 2-arylidene-1-indanones (IIIa-k) (Scheme I). The structures of these products were evident from the infrared (1, 2), electronic (3, 4-7), NMR spectra (8), and chemical analyses (Tables I and II). The infrared spectra of IIIa show absorption bands at 1625 and 1695 cm⁻¹ attributed to $\nu_{C=C}$ and C=O, respectively. The chalcones (III a-k) were condensed with guanidine to yield the corresponding 5H-2-amino-4-arylideneopyrimidines (Va-k) (3, 9, 10) (Scheme I). The structures of these products were substantiated by spectroscopy (3, 8) (Tables I and II). The IR spectrum of Va shows absorptions at 1600 cm⁻¹ (C=C), 1640 cm⁻¹ (C=N), and 3140, 3280, and 3460 cm⁻¹ (NH₂).

Chemical evidence can be also adduced in favor of the structure of compounds V. Thus acetylation of Vf,h leads to the formation of the corresponding acetylamido derivatives (VI f,h) (Scheme I) (3, 9). Their structures were established from their infrared (11), electronic (12), and mass spectra (Table I). The IR spectrum of VI f shows absorptions at 1720 cm⁻¹ (C=O) and 3400 cm⁻¹ (NH).

Treatment of the 2-aminopyrimidines (Vf,h,j) with nitrous acid revealed the formation of the corresponding 2(1H)-pyrimidinones (VII f,h,j) (Scheme I). The lactam form of these products was inferred from their infrared (9), electronic and mass spectra (Table I). The IR spectrum of VII f revealed absorptions at 1640 cm⁻¹ (C=O) and 2990 and 3080 cm⁻¹ (NH).

Experimental Section

Melting points were measured using a Bock-Monoscope (thermal microscope) and are uncorrected. Electronic and infrared spectra were run on Cary 17 and Perkin-Elmer 580B, respectively. The ¹H NMR and the mass spectra were measured with Varian T60A and Varian MAT 311A, respectively. Microanalyses were determined by A. Bernhardt Microanalytical Laboratory, GFR.