

Henry's Law Constants and Infinite Dilution Activity Coefficients of Propane, Propene, Butane, 2-Methylpropane, 1-Butene, 2-Methylpropene, *trans*-2-Butene, *cis*-2-Butene, 1,3-Butadiene, Dimethyl Ether, Chloroethane, and 1,1-Difluoroethane in 2-Methyl-3-buten-2-ol and 3-Methyl-3-buten-1-ol

Yoshimori Miyano,* Osamu Mitsuoka, Hirofumi Ikeda, Akimasa Ouchi, Tomoki Ono, Kayoko Tsuchida, and Yoko Tateishi

Department of Chemistry and Bioscience, Kurashiki University of Science and the Arts, 2640 Nishinoura, Tsurajimacho, Kurashiki 712-8505, Japan

Henry's law constants and infinite dilution activity coefficients of propane, propene, butane, 2-methylpropane, 1-butene, 2-methylpropene, *trans*-2-butene, *cis*-2-butene, 1,3-butadiene, dimethyl ether, chloroethane, and 1,1-difluoroethane in 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol in the temperature range of (250 to 330) K were measured by a gas stripping method, and partial molar excess enthalpies were calculated from the activity coefficients. The estimated uncertainties are about 2 % for the Henry's law constants and 3 % for the infinite dilution activity coefficients. In general the Henry's law constants followed the order of increasing Henry's law constant with decreases in the normal boiling point temperature of the liquefied gas except for 1,3-butadiene, and the partial molar excess enthalpies of gases followed the order of increasing partial molar excess enthalpy with decreases of the polarity of the gases except for chloroethane and 1,1-difluoroethane.

Introduction

This work is a continuation of a systematic study on the Henry's law constants and the infinite dilution activity coefficients of gases in various alcohols.^{1–7} The gas stripping method originally proposed by Leroi et al.⁸ has been used to measure the activity coefficients at infinite dilution of solutes in nonvolatile solvents when the vapor pressures of solutes are negligibly small. In previous work,^{1–7} the Henry's law constants for propane, propene, butane, 2-methylpropane, 1-butene, 2-methylpropene, *trans*-2-butene, and 1,3-butadiene in methanol, propanol (propan-1-ol, propan-2-ol), butanol (butan-1-ol, butan-2-ol, 2-methylbutan-1-ol, 2-methylbutan-2-ol), and pentanol (pentan-1-ol, pentan-2-ol, pentan-3-ol, 2-methylbutan-1-ol, 3-methylbutan-1-ol, 2-methylbutan-2-ol) were measured with this method. For these highly volatile solutes and solvents, a rigorous expression was derived for data reduction.

In this work, the Henry's law constants for propane, propene, butane, 2-methylpropane, 1-butene, 2-methylpropene, *trans*-2-butene, *cis*-2-butene, 1,3-butadiene, dimethyl ether, chloroethane, and 1,1-difluoroethane in 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol are measured by the gas stripping method, and the infinite dilution activity coefficients of solutes and the partial molar excess enthalpies are evaluated. These data are not available in the literature.

Theory

The gas stripping method, originally proposed by Leroi et al.,⁸ is based on the variation of the vapor-phase composition when the highly diluted solute in a liquid

mixture in an equilibrium cell is stripped from the solution by the flow of inert gas (helium). The composition of the gas leaving the cell is periodically sampled and analyzed by means of gas chromatography. The peak area (S) of the solute decreases exponentially with the volume of inert gas flowing out of the cell by the following rigorous relation:¹

$$\ln \frac{S}{S_0} = \left(\frac{(H_g/\varphi_g^V)}{(1 + \alpha)P_s^{\text{sat}} - \alpha(H_g/\varphi_g^V)} - 1 \right) \times \ln \left(1 - \frac{(1 + \alpha)P_s^{\text{sat}} - \alpha(H_g/\varphi_g^V)}{ZRTn_{s,0}^L + V_{\text{GP},0}(H_g/\varphi_g^V)} V \right) \quad (1)$$

where S_0 is the peak area of the solute at time $t = 0$, P_s^{sat} is the saturated vapor pressure of the pure solvent (s), H_g is the Henry's law constant of the solute (g), Z is the compressibility factor of the saturated vapor in the cell, and φ_g^V represents the fugacity coefficient of the solute in the vapor phase. Superscripts V and L denote vapor and liquid phases, respectively. The $n_{s,0}^L$ is the number of moles of solvent in the liquid phase in the cell at $t = 0$. V is the volume of the saturated gas flowing out of the cell. $V_{\text{GP},0}$ is the initial volume of the vapor phase in the cell. The volume of the vapor phase in the cell at time t (V_{GP}) slightly depends on the gas volume withdrawn from the cell, and it is expressed as

$$V_{\text{GP}} = V_{\text{GP},0} + \alpha V \quad \alpha = \frac{P_s^{\text{sat}} v_s^{\text{L,sat}}}{ZRT} \quad (2)$$

where $v_s^{\text{L,sat}}$ is the liquid molar volume of the solvent at saturation, R is the gas constant, and T is the absolute temperature.

* Corresponding author. E-mail: miyano@chem.kusa.ac.jp. Fax: +81-86-440-1062.

From eq 1, the Henry's law constant of the solute can be obtained from the slope of the line in a log–log plot as follows. The initial value of the Henry's law constant was set equal to zero, and the logarithm on the right-hand side in eq 1 was calculated. By a process of successive iterations, the value of the Henry's law constant was determined.

The infinite dilution activity coefficient of the solute (γ^∞) can be obtained from the following relationships:

$$H_g \equiv \lim_{x_g \rightarrow 0} \frac{f_g^V}{x_g} = f_g^{L,0} \gamma^\infty = f_g^{\text{sat}} \lambda \gamma^\infty = \varphi_g^{\text{sat}} P_g^{\text{sat}} \lambda \gamma^\infty \quad (3)$$

$$\gamma^\infty = \frac{H_g}{\varphi_g^{\text{sat}} P_g^{\text{sat}} \lambda} \quad (4)$$

where x_g is the mole fraction of the solute in the liquid phase, f_g^V is the fugacity of the solute in the vapor phase, $f_g^{L,0}$ is the fugacity of the pure solute in the reference state (pure liquid at system temperature and pressure), f_g^{sat} and φ_g^{sat} are respectively the fugacity and the fugacity coefficient of the solute at saturation, P_g^{sat} is the saturated vapor pressure of the pure solute. λ is the Poynting correction factor and is well-approximated as

$$\lambda \approx \exp\left(\frac{(P - P_g^{\text{sat}})v_g^{L,\text{sat}}}{RT}\right) \quad (5)$$

where P is the system pressure. The vapor pressures and the saturated liquid densities for pure substances were taken from ref 9. The thermodynamic properties appearing in the above equations (φ_g^V , φ_g^{sat} , Z , etc.) were calculated from the virial equation of state truncated after the second virial coefficient. For propane and propene, however, the virial equation of state truncated after the third virial coefficient was used for this purpose because of their high volatility. The details of the calculation methods were described in the previous papers.^{1,2}

According to the Gibbs–Helmholtz equation, the value for the partial molar excess enthalpy of the solute at infinite dilution ($H^{E,\infty}$) can be obtained from the infinite dilution activity coefficients:¹⁰

$$\frac{\partial \ln \gamma^\infty}{\partial(1/T)} = \frac{H^{E,\infty}}{R} \quad (6)$$

Experimental Section

Details of the experimental apparatus were presented in our earlier paper.^{1,4} About 36 cm³ of the solvent was introduced into the equilibrium cell, of volume about 40 cm³, and the accurate quantity was determined by mass with the uncertainty of 1 mg. Then the equilibrium cell was immersed in a constant-temperature bath filled with (ethylene glycol + water) and connected to a supply of helium. The temperature was controlled to within ± 0.02 K and measured with a quartz thermometer (Hewlett-Packard Co., model 2804A) with an uncertainty of 0.01 K. About 1 cm³ of the solute gas was introduced into the equilibrium cell by a syringe. The amount of the solute gas introduced into the cell was adjusted to keep the mole fraction lower than 10^{-4} in solution. Then helium began to flow into the equilibrium cell at a flow rate of about 2 cm³·min⁻¹, which was measured with a mass flowmeter (Kofloc Co., model 3300) with a maximum flow rate of 2 cm³·min⁻¹ and an uncertainty of 1 %. The flow rate was

controlled with a fine metering valve (double needles, Swagelok Co., SS2-D). The gas flowing out of the equilibrium cell was kept at a higher temperature than the others to avoid any condensation and was introduced into a gas chromatograph (Hitachi Ltd., model G-3000, with double FID detectors) to measure the solute peak area (S). Sampling for the gas chromatography (GC) was performed every 60 min and continued for about 20 h. The pressure in the cell, the temperatures of the bath, and the amount of helium (n_{He}), which could be obtained from the integration of the flow rate, were also measured every 60 min. The pressure in the cell was approximately equal to atmospheric pressure measured with a pressure transducer (Paroscientific, Inc., Digiquartz pressure transducer model 215A and Tsukasa Sokken Co. Ltd., Digiquartz Pressure Computer model 600S) to an uncertainty of 10 Pa (0.01 %).

The uncertainty of this measurement for the Henry's law constants may be considered to be within 2 %, and it mostly depends on the uncertainty of the mass flowmeter (1 %). The total uncertainty of the other measured variables such as the temperature, pressure, and mass of the solvent was estimated to be not larger than 1 %.

Materials. Butane, 2-methylpropane, 1-butene, and 2-methylpropene were supplied by Takachiho Kagaku Kogyo with specified minimum mass fraction purities of 0.998, 0.99, 0.99, and 0.99, respectively. Propane and propene were supplied by Japan Fine Products at specified minimum mass fraction purities of 0.995 and 0.995, respectively. *cis*-2-Butene, *trans*-2-butene, chloroethane, 1,1-difluoroethane, and 1,3-butadiene were supplied by Aldrich Chemicals with specified minimum purities by GC of 99+ %, 99+ %, 99.7+ %, 98+ %, and 99+ %, respectively. 2-Methyl-3-buten-2-ol, 3-methyl-3-buten-1-ol, and dimethyl ether were supplied by Tokyo Kasei Kogyo Co. with specified minimum purities of 98+ %, 98+ %, and 99+ % by GC, respectively. All materials were used without further purification.

Results and Discussion

The Henry's law constants, the infinite dilution activity coefficients, and the partial molar excess enthalpies measured in this work are numerically indicated in Tables 1 and 2. All experiments were conducted under atmospheric pressure, and the estimated fugacity coefficients of the solute in the vapor phase and the compressibility factors of the vapor were around unity ($\varphi_g^V = 1.00$, $Z = 1.00$) for all systems. However, for the evaluation of the infinite dilution activity coefficients, the nonideality of gases at saturation is not negligible. For highly volatile solutes such as propene, the vapor pressure at $T = 330$ K is about $P = 2400$ kPa, and the nonideality (φ_g^{sat}) should be evaluated. The estimated uncertainty for the vapor phase fugacity coefficient at saturation was about 1 % as discussed in the previous paper¹ (the difference between the calculated values from the virial equation and the Soave equation of state¹¹ was less than 1 %). Therefore, the uncertainty of the obtained infinite dilution activity coefficients may be considered to be no greater than 3 %.

Figures 1 and 2 show the temperature dependence of the Henry's law constants of the 12 gases in 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol, respectively. The Henry's law constants of nonpolar gases in any solvents generally follow the order of increasing Henry's law constant with decreases in the normal boiling point temperature (T_b) of the liquefied gas. The order of the Henry's law constants

Table 1. Henry's Law Constants and Infinite Dilution Activity Coefficients of Solutes in 2-Methyl-3-buten-2-ol

T K	H_g kPa	γ^∞	P_g^{sat} kPa	φ_g^{sat}	$H^{E,\infty}$ kJ·mol ⁻¹	T K	H_g kPa	γ^∞	P_g^{sat} kPa	φ_g^{sat}	$H^{E,\infty}$ kJ·mol ⁻¹
Propane											
249.98	630	3.08	218.4	0.94	0.56	309.99	2810	2.70	1274.4	0.85	2.23
269.95	1160	2.97	430.9	0.92	1.20	330.00	3840	2.54	1985.6	0.82	2.65
289.99	1900	2.86	770.8	0.89	1.75						
Propene											
249.98	630	2.46	272.7	0.94	0.42	309.99	2650	2.17	1531.0	0.84	2.22
269.95	1130	2.38	529.5	0.91	1.11	330.00	3600	2.05	2367.1	0.80	2.68
289.99	1820	2.30	935.1	0.87	1.70						
Butane											
249.98	127	3.30	39.2	0.98	0.00	309.99	900	2.87	347.1	0.92	2.98
269.95	285	3.22	91.6	0.96	1.14	330.00	1370	2.65	592.1	0.89	3.73
289.99	540	3.08	187.8	0.94	2.12						
2-Methylpropane											
249.98	218	3.56	63.0	0.97	0.46	309.99	1310	3.06	485.6	0.90	2.81
269.95	452	3.42	139.2	0.95	1.36	330.00	1920	2.83	802.5	0.87	3.40
289.99	820	3.27	272.8	0.93	2.13						
1-Butene											
249.98	133	2.74	49.8	0.98	0.43	309.99	890	2.37	419.7	0.91	2.72
269.95	290	2.65	114.0	0.96	1.31	330.00	1340	2.20	707.4	0.88	3.29
289.99	540	2.52	230.2	0.94	2.06						
2-Methylpropene											
249.98	135	2.72	50.6	0.98	0.73	309.99	890	2.31	426.4	0.91	2.77
269.95	289	2.61	115.8	0.96	1.51	330.00	1330	2.15	719.1	0.88	3.29
289.99	540	2.47	233.7	0.94	2.19						
<i>trans</i> -2-Butene											
249.98	98	2.73	36.6	0.98	0.62	309.99	720	2.35	332.5	0.92	2.56
269.95	219	2.62	86.6	0.97	1.36	330.00	1100	2.20	568.4	0.90	3.05
289.99	424	2.51	179.0	0.95	2.00						
<i>cis</i> -2-Butene											
250.00	80	2.55	31.9	0.98	0.19	309.99	630	2.23	307.4	0.93	2.83
269.96	186	2.48	77.2	0.97	1.20	330.00	970	2.05	533.6	0.90	3.49
289.99	364	2.36	162.7	0.95	2.07						
1,3-Butadiene											
249.98	110	2.43	46.1	0.98	0.26	309.99	740	2.05	399.7	0.92	3.37
269.95	243	2.36	106.8	0.96	1.45	330.00	1110	1.88	678.2	0.89	4.16
289.99	454	2.22	217.6	0.94	2.48						
Dimethyl Ether											
250.01	65	0.62	108.9	0.97	-2.25	309.99	500	0.72	812.0	0.89	-1.02
269.96	151	0.67	237.7	0.95	-1.78	330.00	810	0.74	1330.9	0.86	-0.71
289.99	294	0.70	460.7	0.92	-1.37						
Chloroethane											
250.01	48	2.28	21.4	0.99	1.30	309.99	375	1.69	235.3	0.95	4.68
269.96	115	2.16	54.5	0.98	2.59	330.00	600	1.56	420.8	0.92	5.53
289.99	227	1.96	120.1	0.97	3.70						
1,1-Difluoroethane											
250.00	495	4.68	109.8	0.96	4.00	309.99	2100	2.89	846.0	0.88	6.34
269.96	910	3.99	242.9	0.94	4.90	330.00	2800	2.47	1398.3	0.85	6.93
289.99	1450	3.39	475.7	0.91	5.67						

of the 12 gases in 3-methyl-3-buten-1-ol at $T = 290$ K is as follows:

propane ($T_b = 231.1$ K, $\mu = 0$ D) >
 propene (225.5 K, 0.4 D) >
 1,1-difluoroethane (248.2 K, 2.3 D) >
 2-methylpropane (261.4 K, 0.1 D) >
 butane (272.7 K, 0 D) > 1-butene (266.9 K, 0.3 D) >
 2-methylpropene (266.2 K, 0.5 D) >
trans-2-butene (274.0 K, 0 D) >
 1,3-butadiene (268.7 K, 0 D) >
 dimethyl ether (248.3 K, 1.3 D) >
cis-2-butene (276.9 K, 0.3 D) >
 chloroethane (285.5 K, 2.0 D)

where μ is the dipole moment (1 D = 3.162×10^{-25} J^{1/2}·m^{3/2})

and the values were cited from ref 12. For nonpolar gases ($\mu = 0$ D), the Henry's law constants in 3-methyl-3-buten-1-ol follow the above simple relationship except for 1,3-butadiene. The molecular size of 1,3-butadiene is smaller than the others because it has two double bonds, which will make it easier to insert the molecule into a liquid. As the result, under the condition that the solutes have almost the same boiling point temperatures, the Henry's law constant of the smaller molecule becomes smaller than that of larger molecules. This may be the reason that 1,3-butadiene does not follow the simple relationship.

In a polar solvents such as 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol, on the other hand, the Henry's law constants of polar gases decreases in general because of the dipole-pair intermolecular interaction. As the result, the Henry's law constant of propene becomes smaller than that of propane, the Henry's law constant of 2-methylpro-

Table 2. Henry's Law Constants and Infinite Dilution Activity Coefficients of Solutes in 3-Methyl-3-buten-1-ol

T	H_g	P_g^{sat}	φ_g^{sat}	$H^{E,\infty}$	T	H_g	P_g^{sat}	φ_g^{sat}	$H^{E,\infty}$		
K	kPa	kPa		$\text{kJ}\cdot\text{mol}^{-1}$	K	kPa	kPa		$\text{kJ}\cdot\text{mol}^{-1}$		
Propane											
249.92	890	4.36	217.9	0.94	0.58	309.99	4020	3.86	1274.4	0.85	1.88
270.04	1640	4.19	432.1	0.92	1.08	330.00	5590	3.70	1985.6	0.82	2.20
289.99	2730	4.10	770.8	0.89	1.51						
Propene											
249.92	780	3.07	272.2	0.94	-0.13	309.99	3560	2.92	1531.0	0.84	1.28
270.04	1440	3.04	531.0	0.91	0.41	330.00	4920	2.80	2367.1	0.80	1.64
289.99	2370	2.99	935.1	0.87	0.88						
Butane											
249.92	188	4.89	39.1	0.98	1.41	309.99	1320	4.18	347.1	0.92	1.92
270.04	408	4.61	91.9	0.96	1.61	330.00	2060	3.99	592.1	0.89	2.05
289.99	780	4.41	187.8	0.94	1.77						
2-Methylpropane											
249.92	322	5.27	62.8	0.97	1.37	309.99	1950	4.54	485.6	0.90	1.93
270.04	660	4.98	139.6	0.95	1.59	330.00	2900	4.29	802.5	0.87	2.07
289.99	1190	4.73	272.8	0.93	1.77						
1-Butene											
249.92	174	3.57	49.6	0.98	0.30	309.99	1240	3.28	419.7	0.91	1.55
270.04	381	3.47	114.4	0.96	0.78	330.00	1910	3.13	707.4	0.88	1.86
289.99	730	3.40	230.2	0.94	1.19						
2-Methylpropene											
249.92	172	3.49	50.4	0.98	0.17	309.99	1230	3.21	426.4	0.91	1.71
270.04	380	3.41	116.2	0.96	0.76	330.00	1880	3.04	719.1	0.88	2.10
289.99	720	3.32	233.7	0.94	1.27						
<i>trans</i> -2-Butene											
249.92	128	3.55	36.5	0.98	0.59	309.99	1000	3.27	332.5	0.92	1.30
270.04	287	3.42	86.9	0.97	0.86	330.00	1560	3.12	568.4	0.90	1.48
289.99	570	3.35	179.0	0.95	1.09						
<i>cis</i> -2-Butene											
249.95	103	3.26	31.9	0.98	0.24	309.99	820	2.90	307.4	0.93	2.15
269.95	240	3.20	77.1	0.97	0.97	330.00	1310	2.77	533.6	0.90	2.63
289.90	473	3.08	162.2	0.95	1.60						
1,3-Butadiene											
249.92	124	2.76	45.9	0.98	0.19	309.99	930	2.56	399.7	0.92	1.44
270.04	277	2.69	107.1	0.96	0.67	330.00	1450	2.45	678.2	0.89	1.75
289.99	540	2.65	217.6	0.94	1.08						
Dimethyl Ether											
249.95	135	1.28	108.7	0.97	1.00	309.99	830	1.17	812.0	0.89	0.81
269.95	276	1.23	237.6	0.95	0.93	330.00	1260	1.15	1330.9	0.86	0.76
289.90	500	1.20	459.4	0.92	0.87						
Chloroethane											
249.94	53	2.49	21.3	0.99	0.06	309.99	495	2.23	235.3	0.95	2.65
269.95	125	2.35	54.5	0.98	1.05	330.00	770	1.99	420.8	0.92	3.30
289.90	268	2.32	119.7	0.97	1.91						
1,1-Difluoroethane											
249.95	520	4.94	109.6	0.96	2.48	309.99	2720	3.74	846.0	0.88	3.33
269.95	1030	4.53	242.8	0.94	2.80	330.00	3980	3.50	1398.3	0.85	3.54
289.90	1760	4.12	474.4	0.91	3.08						

pene becomes smaller than that of butane and 1-butene, and so on.

The boiling point temperature of dimethyl ether is almost the same as to that of 1,1-difluoroethane, but the Henry's law constants of dimethyl ether are significantly smaller than those of 1,1-difluoroethane. This means that dimethyl ether should be considered to be a strong polar gas in this solvent. Similar phenomena can be observed for 2-methyl-3-buten-2-ol systems.

The infinite dilution activity coefficients for C_4 gases in methanol, 1-propanol, 2-propanol, 1-butanol, and 2-butanol could be classified into three groups. The first group is the alkane (butane and its isomers), the second group is the alkene (1-butene and its isomers), and the last group is the dialkene (1,3-butadiene). Each group showed very similar values of the infinite dilution activity coefficients, and the order of the infinite dilution activity coefficients

was alkane > alkene > dialkene for C_4 gases. Similar phenomena can be observed for the 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol systems as show in Figures 3 and 4.

As observed in the previous paper,¹³ the infinite dilution activity coefficients for asymmetric mixtures (mixture of nonpolar + strong polar substances) strongly depend on the temperature, and the partial molar excess enthalpies for these asymmetric mixtures have large values in general. On the other hand, for symmetric mixtures such as (nonpolar + nonpolar) or (strong polar + strong polar), the partial molar excess enthalpies have small values in general and the activity coefficients have values near to unity. As 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol are polar solvents, the values of the partial molar excess enthalpies of nonpolar gases such as propane or butane are large, but the values of dimethyl ether, which is a




Figure 1. Henry's law constants of 12 gases in 2-methyl-3-buten-2-ol: ●, propane; ■, propene; ▲, butane; □, 2-methylpropane; ○, 1-butene; ×, 2-methylpropene; △, *trans*-2-butene; +, *cis*-2-butene; ◇, 1,3-butadiene; ⊙, dimethyl ether; ◆, chloroethane; ▽, 1,1-difluoroethane.




Figure 2. Henry's law constants of 12 gases in 3-methyl-3-buten-1-ol: ■, propane; ■, propene; ▲, butane; □, 2-methylpropane; ○, 1-butene; ×, 2-methylpropene; △, *trans*-2-butene; +, *cis*-2-butene; ◇, 1,3-butadiene; ⊙, dimethyl ether; ◆, chloroethane; ▽, 1,1-difluoroethane.

strong polar solute, are small as expected. While 1,1-difluoroethane and chloroethane are polar gases as well as dimethyl ether, the partial molar excess enthalpies of these two gases are very large in these polar solvents. From the viewpoint of a molecular simulation of the Henry's law constant, it is considered that the volume change of the solvent may cause a drastic change of the probability of the insertion of a solute molecule into the solvent. When the molecular size of the solute is sufficiently larger than the void size in the solvent, all insertions will be rejected. When the molecular size of the solute is sufficiently smaller than the void size in the solvent, all insertions will be accepted. In the both cases the probabilities of the insertion will be stable. On the other hand, when the molecular size of the solute fits well with the void size in the solvent, the probability of the insertion will strongly depend on the void size of the solvent and therefore on the volume change of the solvent. As a result the partial molar excess enthalpies




Figure 3. Infinite dilution activity coefficients of gases in 2-methyl-3-buten-2-ol: ▲, butane; □, 2-methylpropane; ○, 1-butene; ×, 2-methylpropene; △, *trans*-2-butene; +, *cis*-2-butene; ◇, 1,3-butadiene; ⊙, dimethyl ether; ◆, chloroethane; ▽, 1,1-difluoroethane.




Figure 4. Infinite dilution activity coefficients of gases in 3-methyl-3-buten-1-ol: ▲, butane; □, 2-methylpropane; ○, 1-butene; ×, 2-methylpropene; △, *trans*-2-butene; +, *cis*-2-butene; ◇, 1,3-butadiene; ⊙, dimethyl ether; ◆, chloroethane; ▽, 1,1-difluoroethane.

of the solute become large. The molecular sizes of 1,1-difluoroethane and chloroethane are smaller than the other 10 solutes, and these sizes may fit well to the void in these solvents.

Conclusions

Henry's law constants, the infinite dilution activity coefficients, and the partial molar excess enthalpies of 12 gases in 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol at $T = (250 \text{ to } 330) \text{ K}$ have been obtained from gas stripping measurements. The Henry's law constant did not depend on the nonideality (the fugacity coefficient) of the solute and the solvent for the systems studied in this work. However, the nonideality of the solute at the reference state should be considered in order to obtain infinite dilution activity coefficients precisely.

The Henry's law constants of the nonpolar gases in the 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol have fol-

lowed the order of increasing Henry's law constant with decreases in the normal boiling point temperature of the liquefied gas except for 1,3-butadiene. The Henry's law constants of polar gases in 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol, however, have become smaller than that expected from the simple relationship.

The partial molar excess enthalpies of gases in 2-methyl-3-buten-2-ol and 3-methyl-3-buten-1-ol generally follow the order of increasing partial molar excess enthalpy with decreases of the polarity of the gases except for 1,1-difluoroethane and chloroethane. This exception may be due to the volumetric balance between the volume of the solute molecule and the void in the solvents.

Literature Cited

- (1) Miyano, Y.; Nakanishi, K.; Fukuchi, K. Henry's constants of butane, isobutane, 1-butene, and isobutene in methanol at 255–320 K. *Fluid Phase Equilib.* **2003**, *208*, 223–238.
- (2) Miyano, Y. Henry's constants and infinite dilution activity coefficients of propane, propene, butane, isobutane, 1-butene, isobutene, *trans*-2-butene and 1,3-butadiene in 1-propanol at $T = (260 \text{ to } 340) \text{ K}$. *J. Chem. Thermodyn.* **2004**, *36*, 101–106.
- (3) Miyano, Y. Henry's constants and infinite dilution activity coefficients of propane, propene, butane, isobutane, 1-butene, isobutene, *trans*-2-butene, and 1,3-butadiene in 2-propanol at 250–330 K. *J. Chem. Eng. Data* **2004**, *49*, 368–371.
- (4) Miyano, Y. Henry's constants and infinite dilution activity coefficients of propane, propene, butane, isobutane, 1-butene, isobutene, *trans*-2-butene, and 1,3-butadiene in 1-butanol and 2-butanol from (250 to 330) K. *J. Chem. Eng. Data* **2004**, *49*, 1285–1289.
- (5) Miyano, Y. Henry's constants and infinite dilution activity coefficients of propane, propene, butane, isobutane, 1-butene, isobutene, *trans*-2-butene, and 1,3-butadiene in isobutanol and *tert*-butanol. *J. Chem. Thermodyn.* **2004**, *36*, 865–869.
- (6) Miyano, Y. Henry's law constants and infinite dilution activity coefficients of propane, propene, butane, isobutane, 1-butene, isobutene, *trans*-2-butene, and 1,3-butadiene in 2-methyl-1-butanol, 3-methyl-1-butanol, and 2-methyl-2-butanol. *J. Chem. Eng. Data* **2005**, *50*, 211–215.
- (7) Miyano, Y. Henry's law constants and infinite dilution activity coefficients of propane, propene, butane, isobutane, 1-butene, isobutene, *trans*-2-butene, and 1,3-butadiene in 1-pentanol, 2-pentanol, and 3-pentanol. *J. Chem. Thermodyn.* **2005**, *37*, 459–465.
- (8) Leroi, J.-C.; Masson, J.-C.; Renon, H.; Fabries, J.-F.; Sannier, H. Accurate measurement of activity coefficients at infinite dilution by inert gas stripping and gas chromatography. *Ind. Eng. Chem. Process Des. Dev.* **1977**, *16*, 139–144.
- (9) Smith, B. D.; Srivastava, R. *Thermodynamic Data for Pure Compounds*, Parts A and B; Elsevier: Amsterdam, 1986.
- (10) Heintz, A.; Kulikov, D. V.; Verevkin, S. P. Thermodynamic properties of mixtures containing ionic liquids. 1. Activity coefficients at infinite dilution of alkanes, alkenes, and alkylbenzenes in 4-methyl-*n*-butylpyridinium tetrafluoroborate using gas–liquid chromatography. *J. Chem. Eng. Data* **2001**, *46*, 1526–1529.
- (11) Soave, G. Equilibrium constants from a modified Redlich–Kwong equation of state. *Chem. Eng. Sci.* **1972**, *27*, 1197–1203.
- (12) Reid, R. C.; Prausnitz, J. M.; Poling, B. E. *The Properties of Gases and Liquids*; McGraw-Hill: New York, 1987.
- (13) Miyano, Y.; Kobashi, T.; Shinjo, H.; Kumada, S.; Watanabe, Y.; Niya, W.; Tateishi, Y. Henry's law constants and infinite dilution activity coefficients of *cis*-2-butene, dimethyl ether, chloroethane, and 1,1-difluoroethane in methanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, isobutanol, *tert*-butanol, 1-pentanol, 2-pentanol, 3-pentanol, 2-methyl-1-butanol, 3-methyl-1-butanol, and 2-methyl-2-butanol. *J. Chem. Thermodyn.* (in press).

Received for review August 23, 2005. Accepted September 22, 2005. This paper reports part of the work supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (16560663), which is gratefully acknowledged.

JE050343I