

Liquid–Liquid Equilibria for Monoethylene Glycol + Water + Alkane Systems in the Range (273 to 313) K and Atmospheric Pressure. Philippe Sentenac, Charles Berro, Evelyne Rauzy,* Ilham Mokbel, and Jacques Jose, *J. Chem. Eng. Data* **2004**, *131*, 1577–1580.

The authors of “Liquid–Liquid Equilibria for Monoethylene Glycol + Water + Alkane Systems in the Range (273 to 313) K and Atmospheric Pressure” (*J. Chem. Eng. Data* **2004**, *131*, 1577–1580) declare that the experimental solubilities published there are completely erroneous. The reader is invited to consult another article (DOI: 10.1021/je900295u) that reports new data over the temperature range (283 to 323) K for two of the eight ternary systems appearing in *J. Chem. Eng. Data* **2004**, *131*, 1577–1580, namely, monoethylene glycol + water + *n*-hexane/2,2,4-trimethylpentane. In addition, the article reports data for four binary systems, namely, monoethylene glycol + *n*-hexane/2,2,4-trimethylpentane and water + *n*-hexane/2,2,4-trimethylpentane.

JE100122S

10.1021/je100122s

Published on Web 03/12/2010

Isothermal Binary Vapor–Liquid Equilibrium for 2-Methylpropane and *n*-Butane with 1,2-Ethanedithiol and 2-Methyl-2-propanethiol. Anne Penttilä,* Petri Uusi-Kyyny, Juha-Pekka Pokki, Minna Pakkanen, and Ville Alopaeus, *J. Chem. Eng. Data* **2010**, *55*, 291–296.

The temperature of the refractive index of 1,2-ethanedithiol and 2-methyl-2-propanethiol in Table 1 in the above paper was not reported. The temperature at which the refractive index was measured has been added to the heading of the table.

Table 1. Materials, Their Purities, and Refractive Indexes Measured at $T = 298.15$ K

component	supplier	GC analysis (mass fraction)	refractive index		
			this work	ref 4	ref 5
2-methylpropane	Oy Aga Ab	0.9995			
<i>n</i> -butane	Oy Aga Ab	0.9995			
1,2-ethanedithiol	Fluka	0.9976	1.5562	1.5562	
2-methyl-2-propanethiol	Aldrich	0.9987	1.4200	1.42004	1.4200

JE100086J

10.1021/je100086j

Published on Web 02/25/2010