


Synthesis and Tautomerization of 2-Nitro-1-nitrosoethylbenzene in Acetone†

Ahmad Shaabani,* Majid Ameri and Hamid Reza Bijanzadeh

Chemistry Department, Shahid Beheshti University, P.O. Box 19395-4716, Tehran, Iran

2-Nitro-1-nitrosoethylbenzene has been synthesized in fairly high yield and its tautomerization studied by ^1H NMR spectroscopy.

The reaction of dinitrogen trioxide with olefins affords 1-nitro-2-nitroso derivatives, commonly referred to as pseudo-nitrosites **1**.¹ These adducts can be converted into the more soluble isomers, the corresponding 1,2-nitroximes **2**.^{1,2} Reaction of styrene with arsenic³ and concentrated nitric acid or dinitrogen trioxide^{1b} produces 2-nitro-1-nitrosoethylbenzene **3**. Compound **3** is transformed easily into the more stable isomer 2-isonitroso-1-nitro-2-phenylethane **4** by boiling in ethanol.⁴ The structure of compounds **3** and **4** were earlier deduced only from their elemental analyses and for compound **4** ^1H NMR data were also reported.⁵



2-Nitro-1-nitrosoethylbenzene **3** and 2-isonitroso-1-nitro-2-phenylethane **4** are frequently used as useful products⁶ and as versatile intermediates in organic synthesis.⁷ In addition **4** is used in analytical chemistry for indirect determination of styrene.⁸ Various approaches can lead to these compounds.⁹ However, much demand still exists for their preparation in high yields and free of compound **4** under mild and safe conditions.

We now report a new method for the preparation of compound **3** in high yields under easy and safe conditions and deduce the essential structure of compounds **3** and **4** from their IR, ^1H , ^{13}C NMR and mass spectra. In addition, the tautomerization of **3** to **4** was studied by using ^1H NMR spectroscopy in acetone at $25\text{--}40^\circ\text{C}$.

For the tautomerization study a fresh sample of compound **3** was dissolved in $(\text{CD}_3)_2\text{CO}$ [0.01 g in 0.4 ml $(\text{CD}_3)_2\text{CO}$ in a 5 mm NMR tube] and equilibrated at the required temperature. The progress of the tautomerization was monitored by recording the appearance and disappearance of the methylene signals of **3** and **4** (Fig. 1). Integration of the area under the methylene signal of **4** with respect to that of the methylene signals of **3** gave the concentration of the species present. The rate of the reaction is given by $\text{rate} = k[\mathbf{3}]^x$ or $k[\mathbf{4}]^y$, where x or y is the reaction order. The integrated rate equation for a first order reaction is $\ln[\mathbf{4}]_t = -kt + \ln[\mathbf{3}]_0$.

The concentration of compound **4** at time t was found by setting the peak area for **3** equal to one and measuring the peak area of **4** relative to it. Typical data are presented in




Fig. 1

Table 1. A plot of $\ln[\mathbf{4}]_t$ versus time should yield a straight line if the reaction is first order [see Fig. 2(a)]. The slope of this line is equal to k . The rate constants were obtained at several temperatures (298, 303 and 313 K) [Fig. 2(a)–(d), Table 2].

By taking the natural logarithm of the Arrhenius equation, the activation energy, $E_a = 65.63 \text{ kJ mol}^{-1}$, was obtained from a plot $\ln k$ versus $1/T$. Activation parameters, $\Delta H^\ddagger = 57.85 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -89.58 \text{ J K}^{-1} \text{ mol}^{-1}$, were also determined from a plot of $\ln(k/T)$ versus $1/T$. The relatively low enthalpy of activation for this tautomerization suggests a mechanism which is compatible with a cyclic activated complex, in which bond making accompanies bond breaking. In addition, a negative entropy of activation, due to the loss of rotational degrees of freedom associated with the highly ordered transition state, supports this complex formation.




Fig. 2

*To receive any correspondence.

†This is a **Short Paper** as defined in the Instructions for Authors, Section 5.0 [see *J. Chem. Research (S)*, 1998, Issue 1]; there is therefore no corresponding material in *J. Chem. Research (M)*.

Table 1 Integrated area of compound **4** with respect to that of **3** at different temperatures

$T/^\circ\text{C}$	t/min	Integrated area (A_t)	$\ln A_t$
25	0	2.2	0.8
	15	3.0	1.1
	34	4.0	1.4
	45	5.0	1.6
	60	6.0	1.8
	94	9.0	2.2
30	0	2.0	0.7
	17	3.0	1.1
	30	4.0	1.4
	46	6.0	1.8
	59	8.0	2.1
35	0	2.7	1.0
	14	5.0	1.6
	29	9.0	2.2
	46	13.5	2.6
40	0	2.0	0.7
	10	4.0	1.4
	21	7.4	2.0
	31	12.2	2.5
	45	20.1	3.0

Experimental

NMR spectra were recorded on JEOL EX 90-MHz spectrometer using tetramethylsilane as the internal standard. Temperature was calibrated by the shift difference in methanol. The temperature range was 25 to 40 °C. Infrared spectra were taken on a Shimadzu IR-470 spectrophotometer, mass spectra on a Finnigan-Matt 8430 mass spectrometer. Elemental analyses were performed with a CHN Heracus-O-Rapid analyzer.

Preparation of 2-Nitro-1-nitrosoethylbenzene 3.—Sodium nitrite (34.5 g, 0.5 mol) and styrene (20.8 g, 0.2 mol) were added to chloroform (150 ml) in a 250 cm³ two-necked round-bottomed flask equipped with condenser and dropping funnel. To this stirred mixture was added phosphoric acid (57.6 g, 85 wt.% solution in water; 0.5 mol) from a dropping funnel over a period of 20 min. After complete addition of phosphoric acid, the mixture was stirred for 4 h at 50 °C, then neutralized with saturated sodium bicarbonate solution. The precipitate was filtered off and washed with water (100 ml) and then *n*-hexane (40 ml). Compound **3** was obtained as a white solid (25.2 g, yield 70%), mp 129 °C; δ_{H} [90 MHz, (CD₃)₂CO] 7.60 (5 H, m, aromatic), 6.90 (1 H, dd, ³*J* 9.5, ³*J* 2.9, CH), 5.41 (1 H, dd, ²*J* 14.0, ³*J* 9.5, CH₂), 4.97 (1 H, dd, ²*J* 14.0, ³*J* 2.9 Hz, CH₂); δ_{C} [22.5 MHz, (CD₃)₂CO], 69.2 (CNO₂), 74.7 (CNO), 127.4,

Table 2 Temperature dependence of rate constants for the tautomerization of **3** → **4** in acetone

T/K	$10^2k/\text{min}^{-1}$	$\ln(k/T)$
298	1.485	-9.907
303	2.381	-9.451
308	3.509	-9.100
313	5.097	-8.800

130.0, 130.9, 135.7 (C₆H₅); $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 2925, 1559, 1373; M⁺ at m/z 180, C₈H₈N₂O₃ requires 180 (Found: C, 53.6; H, 4.5; N, 15.3. C₈H₈N₂O₃ requires C, 53.3; H, 4.4; N, 15.5%).

Preparation of 2-Isonitroso-1-nitro-2-phenylethane 4.—Compound **3** was completely transformed into **4** in acetone at room temperature after about 24 h. Mp 95–96 °C; δ_{H} (CDCl₃) 5.6 (2 H, s, CH₂) 7.5 (5 H, m, C₆H₅), 9.3 (1 H, s, NOH exchange with D₂O); δ_{C} (CDCl₃) 69.9 (CNO₂), 128.0, 130.6, 131.6, 135.7 (C₆H₅), 149.3 (C=NOH); $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3279, 1559, 1373.

We gratefully acknowledge financial support from the research of Shahid Beheshti University.

Received, 13th February 1998; Accepted, 28th May 1998
Paper E/8/01282B

References

- (a) H. Wieland, *Justus Liebigs Ann. Chem.*, 1920, **424**, 71; (b) H. Wieland, *Ber.*, 1903, **36**, 2558; (c) D. Klamann, W. Koser, P. Weyerstahl and M. Fligge, *Chem. Ber.*, 1965, **98**, 1831; (d) L. Scheinbaum, *Am. Chem. Soc., Div. Pet. Chem. Prepr.*, 1968, **13**, 193.
- M. L. Scheinbaum, *J. Org. Chem.*, 1970, **35**, 2785.
- E. A. Sommer, *Ber. Bunsenges. Phys. Chem.*, 1895, **28**, 1329.
- H. Wieland, *Ann. N.Y. Acad. Sci.*, 1903, **329**, 225.
- M. L. Scheinbaum, *J. Org. Chem.*, 1970, **35**, 2790.
- Ger. Offen.*, 2036681, 1971 (*Chem. Abstr.*, 1971, **74**, 99646c).
- (a) *U.S.S.R. Pat.*, 536179, 1976 (*Chem. Abstr.*, 1977, **86**, 121319v); (b) A. Kunai, T. Doi, T. Kishimoto and K. Sasaki, *Chem. Express*, 1990, **5**, 245.
- (a) G. R. Bond, *Anal. Chem.*, 1947, **19**, 390; (b) V. Sedivec and J. Flek, *Collect. Czech. Chem. Commun.*, 1969 **25**, 1293.
- (a) V. Novak and J. Seidl, *Chem. Prum.*, 1978, **28**, 186; (b) C. D. Hard and J. Patterson, *J. Am. Chem. Soc.*, 1953, **75**, 285.