

Ionic liquid-accelerated N-arylation of 5-arylidene-2,4-thiazolidinediones with diaryliodonium salts

De-Hong Yang^{a*}, Hong-Liang Gao^a, Ben-Yong Yang^a and Zhen-Chu Chen^b

^aDepartment of Materials and Chemistry, Zhongyuan University of Technology, Zhengzhou, 450007, P.R. China

^bDepartment of Chemistry, Zhejiang University, Hangzhou, 310028, P.R. China

An efficient method for the synthesis of N-aryl-5-arylidene-2,4-thiazolidinediones has been developed involving the N-arylation of 5-arylidene-2,4-thiazolidinediones derivatives with diaryliodonium salts in the ionic liquid [bmim]PF₆ (1-butyl-3-methylimidazolium hexafluorophosphate).

Keywords: ionic liquid, N-arylation, 2,4-thiazolidinediones derivatives, diaryliodonium salts


Thiazolidinone derivatives are a bio-active group of heterocyclic compounds.^{1,2} They have again attracted interest because of their antidiabetic activities and pioglitazone and rosiglitazone have been launched for type 2 diabetes mellitus.^{3,4} The common method of synthesising N-aryl-thiazolidinedione moieties utilised the condensation of chloroacetic acid with the aryl-substituted thiourea followed by acid hydrolysis. Another method utilised the reaction of aryl isocyanates with ethyl thioglycolate in the presence of sodium metal.⁵ These preparations are unsatisfactory because of their long time reaction, harmful solvents and low yields.

The applications of hypervalent iodine compounds in organic synthesis are interesting.^{6,7} Recently we showed that diaryliodonium salts are efficient electrophilic arylating reagents, and ionic liquids can accelerate the arylation reactions rates of the diaryliodonium salts.⁸ Relative to conventional molecular solvents, ionic liquids have many advantages.^{9,10} We considered that the preparation of N-aryl-5-arylidene-2,4-thiazolidinediones in ionic liquids, coupled with the high electrophilic reactivity of diaryliodonium salts, would be useful.

First, we examined the efficiency of different solvents for the N-phenylation of 2,4-thiazolidinedione with diphenyliodonium in the presence of KOH (see Scheme 1). The results, summarised in Table 1, show that the ionic liquid [bmim]PF₆ gave the best results in terms of yields and reaction times. As can be seen from Table 1, the ionic liquids compared with classical molecular solvents, have the advantages of accelerating rate and increasing yield. For example, the preparation of N-phenyl-2,4-thiazolidinedione (**3a**) needed refluxing for 12 long hours in the classical molecular solvents dimethylformamide or acetonitrile. Yet, the same reaction was successful in ionic liquid [bmim]PF₆ at only 80 °C within 2 h, and gave higher yield (80%). Furthermore, the temperature of the reaction in the ionic liquid, 80 °C, is lower than that using the solvent dimethylformamide (b.p. 158 °C), and is approximately equivalent to that using acetonitrile (b.p. 81 °C).

The ionic liquid can be typically recovered by extracting the product, and then washing with water followed by vacuum drying. No obvious decrease of yields was observed while reusing the recovered solvent. The results are summarised in Table 2.

We investigated the reactions of 5-arylidene-2,4-thiazolidinediones (**1**) with diaryliodonium tetrafluoroborate (**2**) at 80 °C in the ionic liquid [bmim]PF₆ in the presence of KOH (see Scheme 2). The results summarised in Table 3, show that 5-arylidene-2,4-thiazolidinediones (Ar = 4-CH₃-Ph, 4-CH₃O-Ph, 3,4-(CH₃O)₂-Ph, 4-NO₂-Ph, Ph, Ph-CH=CH, 2-furyl) reacted smoothly with diphenyliodonium tetrafluoroborates and gave adequate yields (50%–85%) after 2–4 h at 80 °C in the ionic liquid [bmim]PF₆ in the presence of KOH. The reactions



Scheme 1

Table 1 Reactions of 2,4-thiazolidinedione (**1a**) with diphenyliodonium salts (**2**) to form 3-phenyl-2,4-thiazolidinedione (**3a**) in different solvents in the presence of KOH

Entry ^a	Solvent	Temp/°C	Time/h	Yield/% ^b
1	DMF	Reflux	12	46
2	CH ₃ CN	Reflux	12	39
3	[bmim]PF ₆	80	2	80
4	[bmim]BF ₄	80	6	72
5	[bpy]BF ₄	80	6	60

^aAll reactions were run with 2,4-thiazolidinedione (**1a**, 0.5 mmol) and diphenyliodonium salts (**2**, 0.55 mmol) in solvents (5 ml) in the presence of KOH (0.5 mmol).

^bIsolated yields based on 2,4-thiazolidinedione (**1a**).

Table 2 Results obtained using recycled ionic liquid

Entry ^a	Product	Cycle	Yield/% ^b
1	3a	1	80
2	3a	2	78
3	3a	3	80
4	3a	4	81

^aAll reactions were run with 2,4-thiazolidinedione (0.5 mmol, 0.078 g) and diphenyliodonium tetrafluoroborate (0.55 mmol, 0.202 g) in [bmim]PF₆ (5 ml) in the presence of KOH (0.5 mmol) at 80 °C for 2 h.

^bIsolated yields based on 2,4-thiazolidinedione.

also proceeded successfully for the substituted (Ar' = 4-CH₃-Ph, 4-CH₃O-Ph) diaryliodonium tetrafluoroborates without variation in the yield. All the products were characterised by ¹H NMR, IR, and the melting points which were consistent with the literature data^{11–13}. Elemental analyses for the new compound also matches the calculated value.

As shown in Schemes 1 and 2, the iodoarene was another product as well as the N-aryl-2,4-thiazolidinediones in the reaction. It could be converted easily to the respective diaryliodonium salts in adequate yields as described in our previous work.¹⁴

In conclusion, an efficient, clean, and novel method for the synthesis of N-aryl-5-arylidene-2,4-thiazolidinediones has been developed using a non toxic arylation agent (diaryliodonium salts) which can be regenerated in adequate yield to replace aryl halides. Employment of ionic liquid rather than classical molecular solvents leads to mild reaction conditions. The ionic liquid can be recycled.

* Correspondent. E-mail: janeydh@hotmail.com

