

Copper(II), cobalt(II), nickel(II) and zinc(II) complexes of Schiff base derived from benzil-2,4-dinitrophenylhydrazone with aniline

N RAMAN*, S RAVICHANDRAN and C THANGARAJA

Department of Chemistry, VHNSN College, Virudhunagar 626 001, India
e-mail: drn_raman@yahoo.co.in

MS received 13 August 2003; revised 1 June 2004

Abstract. New Schiff base chelates of Cu(II), Co(II), Ni(II) and Zn(II) derived from benzil-2,4-dinitrophenylhydrazone with aniline have been synthesised. Microanalytical data, molar conductance, and magnetic susceptibility values have been obtained, and IR, ^1H NMR, ^{13}C NMR, UV-Vis, CV and EPR spectral studies have been carried out to suggest tentative structures for the complexes.

Keywords. Benzil-2,4-dinitrophenylhydrazone; Schiff base complexes; EPR; cyclic voltammetry.

1. Introduction

Metal complexes of Schiff bases have played a central role in the development of coordination chemistry. From the survey of existing literature, it appears that benzil monophenylhydrazone and its related compounds have been extensively used as biologically active complexing agents and analytical reagents.^{1–6} Keeping the above facts in mind and in continuation of our research work^{7–9} on transition metal (II) complexes with Schiff bases, in the present paper we report the synthesis and characterization of Cu(II), Co(II), Ni(II) and Zn(II) complexes of Schiff base derived from the condensation of benzil-2,4-dinitrophenylhydrazone with aniline. The proposed structure of the complexes is shown in chart 1.

2. Experimental

All the chemicals used were of AnalaR grade. Aniline was distilled prior to use. The solvents were dried and distilled before use according to standard procedures. The supporting electrolyte, NaClO_4 used in the voltammetric experiment was purchased from Sigma. IR spectra were recorded in KBr medium on a Perkin–Elmer 783 spectrophotometer. The ^1H and ^{13}C NMR spectra were recorded at Madurai Kamaraj University, on a Bruker instrument using TMS as internal standard. The UV-Vis spectra of all the

complexes were recorded in acetone on a Shimadzu UV-1601 spectrophotometer. Microanalytical data were performed at the Regional Sophisticated Instrumentation Centre, Central Drug Research Institute, Lucknow. Magnetic susceptibility measurements of the complexes in the solid state were determined by Gouy balance using CuSO_4 as the calibrant. Molar conductances of the complexes were measured in acetone at room temperature using systronic conductivity bridge type 305. Electrochemical measurements were carried out in Electrochemical Analyser model BAS-50 voltammograph. The three-electrode cell contained a reference Ag/AgCl electrode, Pt wire auxiliary electrode and glassy carbon




Chart 1.

*For correspondence

working electrode. ESR spectra of the copper complex were recorded on a Varian E112 X-band spectrometer at the Regional Sophisticated Instrumentation Centre, Indian Institute of Technology, Mumbai using tetracyanoethylene (TCNE) as the internal standard.

2.1 Synthesis of Schiff base

The Schiff base was synthesised by the condensation of benzil-2,4-dinitrophenylhydrazone and aniline (1 : 1 molar ratio), dissolved in ethanol. The resulting reaction mixture was refluxed for ≈ 1 h. The yellow solid precipitate of Schiff base obtained was filtered, washed with distilled water dried, recrystallised from ethanol and finally preserved in a desiccator. Yield: 58% (m.p.: 138°C).

2.2 Synthesis of complexes

An ethanolic (10 ml) solution of Schiff base (10 mM) was mixed with metal (II) chloride (5 mM) in ethanol (10 ml) solution keeping ligand-metal ratio 2 : 1 followed by few drops of acetic acid (pH = 6). The mixture was then refluxed for ≈ 1 h on a water bath till the complex precipitated out. Precipitate obtained was dried over KOH pellets. The solid product obtained was filtered, washed with distilled water and dried *in vacuo*.

3. Results and discussion

The analytical data along with some physical properties of the complexes are summarised in table 1. The ligand L, on interaction with Cu(II), Co(II), Ni(II) and Zn(II) chlorides, yields complexes corresponding to the general formula $[\text{ML}_2(\text{H}_2\text{O})_2]$. The analytical data show that the metal to ligand ratio is 1 : 2. They are soluble in common organic solvents. The low molar conductance values of the complexes reveal their non-electrolytic nature.¹⁰

3.1 IR spectra

In order to study the binding mode of Schiff base to metal in the complexes, IR spectrum of the free ligand was compared with the spectra of the metal complexes. Benzil-2,4-dinitrophenylhydrazone shows its characteristics absorption bands in the 3100, 1645, 1620 and 920 cm^{-1} regions, assignable to N–H, C=O, C=N and N–N vibrations respectively. Infrared spe-

tra of the Schiff base ligand, benzil-2,4-dinitrophenylhydrazone with aniline has the important absorption bands in the 3100, 1605–1630 and 920 cm^{-1} regions assignable¹¹ to N–H, C=N and N–N respectively. The band(s) at 1605–1630 cm^{-1} due to the azomethine group of the Schiff base underwent a shift to lower frequency (1580–1590 cm^{-1}) after complexation, indicating the coordination of azomethine nitrogen to metal atom and this can be explained by the donation of electrons from nitrogen to the empty *d*-orbitals of the metal atom. In the spectra of all complexes, the broad band at ≈ 3400 cm^{-1} , together with new band at ≈ 890 cm^{-1} indicating the presence of coordinated water. The nature of metal–ligand bonding is confirmed by the newly formed band at ≈ 440 cm^{-1} in the spectra of the complexes which is tentatively assigned to M–N vibration.

3.2 ¹H and ¹³C NMR spectra

The ¹H-NMR spectrum of the Schiff base ligand exhibits a multiplet signal at 7.3–7.8 *d* (*m*, Ar-H) and 10.1 *d* (*s*, N–H). In Zn(II) complex, the N–H proton is shifted slightly downfield at 10.3 *d* which reveals the bonding of the azomethine nitrogen to Zn(II) ion. The ¹³C-NMR spectra of Zn(II) complex was recorded by employing TMS as internal reference and CDCl₃ as solvent at ambient temperature. The number of signals of sharp peaks represent the number of carbons of the compound which are chemically non-equivalent. The spectra exhibit the azomethine, C=N carbon at 178 ppm. The chemical shift of aromatic carbons appear at 138, 128, 126 and 121 ppm.

3.3 Electronic absorption spectra

The electronic absorption spectra are often very helpful in the evaluation of results furnished by other methods of structural investigation. The electronic spectral measurements were used for assigning the stereochemistries of metal ions in the complexes based on the positions and number of *d–d* transition peaks. The electronic absorption spectra of the Schiff base and its Cu(II), Co(II), Ni(II) and Zn(II) complexes were recorded at room temperature using acetone as solvent. Only one broad band is observed at 16,638 cm^{-1} in the electronic spectrum of the Cu(II) complex assigned to ${}^2E_g \rightarrow {}^2T_{2g}$ transition which is in conformity with octahedral geometry¹². Though three transitions are expected in this case, they are very close in energy and often appear in the

Table 1. Physical characteristics and analytical data of the complexes.

Compound	Analysis, found (calcd) (%)				Yield (%)	m_{eff} (BM)	I_m (mho cm ⁻² mol ⁻¹)
	M	C	H	N			
Ligand	–	66.4 (67.1)	3.7 (4.1)	14.9 (15.1)	60	–	–
[CuL ₂ (H ₂ O) ₂]	5.0 (6.2)	58.9 (61.5)	3.8 (4.1)	13.1 (13.6)	56	1.9	12.6
[CoL ₂ (H ₂ O) ₂]	5.9 (5.8)	59.4 (61.8)	4.0 (4.1)	13.2 (13.6)	59	4.8	11.5
[NiL ₂ (H ₂ O) ₂]	5.7 (5.8)	60.7 (61.9)	3.7 (4.1)	12.4 (13.7)	55	2.8	11.2
[ZnL ₂ (H ₂ O) ₂]	5.4 (6.3)	60.3 (62.8)	3.4 (4.2)	12.9 (13.6)	58	–	10.7

L = Schiff base ligand

form of one broad band envelope. The electronic spectra of Co(II) complex showed two spin-allowed transitions at 17856 and 21734 cm⁻¹ assignable to ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$ and ${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$ transitions respectively, are in conformity with octahedral arrangements for Co(II) ion.¹³ The appearance of a band at 19240 cm⁻¹ due to ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(P)$ transition favours an octahedral geometry¹⁴ for the Ni(II) complex. The absence of any band below 10,000 cm⁻¹ eliminates the possibility of a tetrahedral environment in this complex. The electronic spectrum of the Zn(II) complex shows an absorption band at 23,455 cm⁻¹ attributed to the LMCT transition, which is compatible with this complex having an octahedral structure.^{15,16}

3.4 Magnetic susceptibility measurements

The magnetic moment value of Cu(II) complex is 1.90 BM which suggests a distorted octahedral geometry^{17,18} around the metal ion. The magnetic moment of Co(II) complex is 4.81 BM which suggests the high spin six-coordinated octahedral arrangement^{19,20} of ligand molecules around the metal ion. The Ni(II) complex has magnetic moment value of 2.8 BM indicating a spin-free octahedral^{21,22} configuration. The Zn(II) complex is found to be diamagnetic as expected for d^{10} configuration.

3.5 ESR spectra

The ESR spectra of copper complex provide information of importance in studying the metal ion environment. The X-band ESR spectra of the Cu(II)

complex, recorded in DMSO at liquid nitrogen temperature and at room temperature (300 K), are shown in figures 1 and 2. The spectrum of the copper complex at room temperature shows one intense absorption band in the high field and is isotropic due to the tumbling motion of the molecules. However, this complex in the frozen state shows four well resolved peaks with low field region. The copper complex exhibits the g_{\parallel} value of 2.31 and g_{\perp} value of 2.16. These values indicate that the ground state²³ of Cu(II) is predominantly $d_{x^2-y^2}$. The spin-orbit coupling constant, I value (-486 cm⁻¹) calculated using the relations, $g_{\text{av}} = 1/3[g_{\parallel} + 2g_{\perp}]$ and $g_{\text{av}} = 2(1-2I/10Dq)$, is less than the free Cu(II) ion (-832 cm⁻¹) which also supports covalent character²⁴ of M–L bond in the complex. The G values of 3.82 indicates negligible exchange interaction of Cu–Cu in the complex. The covalency parameter a^2 is calculated ($a^2 = 0.82$) using the following equation:

$$a^2_{\text{Cu}} = -(A_{\parallel}/0.036) + (g_{\parallel} - 2.0023) + 3/7(g_{\perp} - 2.0023) + 0.04.$$

If the value of $a^2 = 0.5$, it indicates complete covalent bonding, while the value of $a^2 = 1.0$ suggests complete ionic bonding. The observed value of a^2 (0.82) of the complex is less than unity, which indicates that the complex has some covalent character in the ligand environment.²⁵

3.6 Cyclic voltammetric study

The cyclic voltammogram of Cu(II) complex (0.01 M) in MeCN (1.0 to -1.2 V potential range)

shows a well-defined redox process corresponding to the formation of the Cu(II)/Cu(I) couple at $E_{p_a} = 0.22$ V and the associated cathodic peak at $E_{p_c} = 0.18$ V. This couple is found to be reversible with $\Delta E_p \approx 0.04$ V and the ratio of anodic to cathodic peak currents ($I_{p_c}/I_{p_a} = 1$) corresponding to a simple one-electron process.²⁶ The complex also shows a quasi-reversible peak in the negative region, characteristic of the Cu(II) \rightarrow Cu(I) couple at $E_{p_c} = -0.66$ V, with associated anodic peak at $E_{p_a} = -0.47$ V for Cu(I) \rightarrow Cu(II) oxidation.

The cyclic voltammogram of Zn(II) complex (0.01 M) in MeCN solution in the absence of molecular oxygen at room temperature in 1.0 to -1.0 V potential range at scan rate 50 mVs^{-1} indicating quasi-reversible one-electron process. A noteworthy feature has been observed in the cyclic voltammogram of Zn(II) complex. During the forward scan it shows two cathodic reduction peaks, one at $+0.45$ V

and another at -0.61 V which are attributed to reduction of Zn(II) \rightarrow Zn(I) and Zn(I) \rightarrow Zn(0) respectively. During the reverse scan it shows two anodic oxidation peaks, one at -0.26 V and another at $+0.62$ V which are attributed to oxidation of Zn(0) \rightarrow Zn(I) and Zn(I) \rightarrow Zn(II) respectively.

4. Conclusions

In this paper coordination chemistry of a Schiff base ligand, obtained from the reaction of benzil-2,4-dinitrophenylhydrazone and aniline, is described. Cu(II), Co(II), Ni(II) and Zn(II) complexes have been synthesized using the above Schiff base ligand and characterized on the basis of analytical, magnetic and spectral data. The Schiff base coordinates through its azomethine nitrogens to the metal ion and acts as a neutral bidentate ligand. All the complexes exhibit octahedral geometry.

Acknowledgements

We thank the Regional Sophisticated Instrumentation Centre, Central Drug Research Institute, Lucknow and Regional Sophisticated Instrumentation Centre, Indian Institute of Technology, Mumbai for use of instrumentation facilities. SR and CT thank the Tamilnadu Government, Chennai for financial support.

References

1. Orget L E and Dunit J D 1957 *Nature (London)* **179** 462
2. Ito K and Ito J 1958 *Aust. J. Chem.* **2** 406
3. Goodgame D M L and Cotton F A 1961 *J. Am. Chem. Soc.* **83** 4161
4. Singh A K and Sharma U N 2002 *Asian J. Chem.* **14** 1221
5. Sacconi L and Ciampolini M 1964 *J. Chem. Soc.* 276
6. Idcher A D 1963 *J. Phys. Chem.* **67** 1214
7. Raman N, Selvan A and Shunmugasundaram A 1997 *Asian J. Chem.* **9** 183
8. Raman N, Kulandaisamy A, Thangaraja C and Jeyasubramanian K 2003 *Transition Met. Chem.* **28** 29
9. Raman N, Ravichandran S and Kulandaisamy A 2002 *Asian J. Chem.* **14** 1261
10. Geary W J 1971 *Coord. Chem. Rev.* **7** 81
11. Silverstein R M, Bassler G C and Morvil T C 1981 *Spectroscopic identification of organic compounds* 4th edn (New York: Wiley)
12. Dunn T M 1960 *The visible and ultraviolet spectra of complex compounds in modern coordination chemistry* (New York: Interscience)




Figure 1. ESR spectra of Cu(II) complex at 77 K in DMSO solution.




Figure 2. ESR spectra of Cu(II) complex at 300 K in DMSO solution.

13. Krishna C H, Mahapatra C M and Dush K C 1977 *J. Inorg. Nucl. Chem.* **39** 1253
14. Kasim A N M, Venkappayya D and Prabhu G V 1999 *J. Indian Chem. Soc.* **76** 67
15. Chohan Z H and Parvez H H 1993 *Synth. React. Inorg. Met.-Org. Chem.* **23** 1061
16. Sekerci M and Tas E 2000 *Heteroatom Chem.* **11** 254
17. Figgis B N and Lewis J 1960 *Modern coordination chemistry* (New York: Wiley Interscience)
18. Sabastiyam A and Venkappayya D 1990 *J. Indian Chem. Soc.* **67** 584
19. Konig E 1971 *Structure and bonding* (Berlin: Springer Verlag) p. 175
20. Rana A K and Shah J R 1981 *Indian J. Chem.* **A20** 142
21. Cotton F A and Wilkinson G 1962 *Advanced inorganic chemistry* (New York: Wiley Interscience)
22. Figgis B N 1976 *An introduction to ligand fields* (New Delhi: Wiley Eastern)
23. Shirin Z and Mukherjee R M 1992 *Polyhedron* **11** 2625
24. Hathway B J and Billing D E 1961 *Coord. Chem. Rev.* **5** 143
25. Kevelson D and Niedman R J 1961 *Chem. Phys.* **35** 149
26. Shirin Z and Mukherjee R M 1992 *Polyhedron* **11** 2625