

The detection of guanidine compounds on paper chromatograms

CONN AND DAVIS^{1,2} found that ninhydrin (triketohydrindene hydrate) in alkaline media forms highly fluorescent products with guanidine, and with monosubstituted and N,N-disubstituted guanidines. The above authors developed a method for the fluorimetric estimation of guanidinium compounds.

It has now been found that the above method could be applied to the detection of certain guanidinium compounds on paper chromatograms. The method used was as follows:

The chromatogram was dried and sprayed with a solution of ninhydrin (0.25 % ethanolic solution). After drying the paper was sprayed with ethanolic sodium hydroxide (2 % ethanolic solution). On exposure to ultraviolet light, guanidinium compounds appeared as green fluorescent spots, which faded slowly.

The spray was tested successfully on the following compounds: guanidine, guanidine hydrochloride, guanidine nitrate, creatine, creatinine, glycoamine, and arginine. Streptidine and diphenyl guanidine could not be detected by this method. Urea also did not give the fluorescent compound.

Although CONN AND DAVIS said that N,N'-disubstituted guanidines did not give the fluorescent compound, it was found that creatinine was detectable on paper chromatograms by this method.

Thanks are due to the Department of Scientific and Industrial Research for a Research Scholarship (to T.W.T.).

*The Chemistry Department,
The University of Birmingham (Great Britain)*

A. S. JONES
T. W. THOMPSON*

¹ R. B. CONN, JR. AND R. B. DAVIS, *Nature*, 183 (1959) 1053.

² R. B. CONN, JR., *Clin. Chem.*, 6 (1960) 547.

Received August 7th, 1962

* Present address: Division of Biology, California Institute of Technology, Pasadena, Calif., U.S.A.

J. Chromatog., 10 (1963) 248

Colour photography of polycyclic compounds on alumina columns

Most polycyclic compounds are highly fluorescent in ultraviolet light hence their chromatographic development on alumina columns can be readily followed. However, the colour of various compounds and clarity of separation can be permanently recorded by modification of techniques recently developed for medical and biological photographic illustrations¹.

The equipment is illustrated in Fig. 1. A Witrona, 180 Joule Electronic Flash Unit at half power was used as the ultraviolet source. A Woods Glass Filter which

J. Chromatog., 10 (1963) 248-249