

- HAUBACH, W. J., KNOBLER, C. M., KATORSKY, A. AND WHITE, D.: The low-temperature chromatographic separation of the isotopic hydrogens at 27° and 55° K. *J. Phys. Chem.*, 71 (1967) 1398-1402 — retention time of the heteronuclear HT is smaller than that of homonuclear D₂ of the same mass.
- SWELL, L.: Gas-liquid radiochromatography of intact labeled cholesterol esters. *Proc. Soc. Exptl. Biol. Med.*, 121 (1966) 1290-1294.

35. MISCELLANEOUS COMPOUNDS AND COMPLEX MIXTURES

- BAILEY, R. L.: Method for detecting volatile organic contaminants in reusable containers. *U.S. Pat.*, 3,266,292 (Aug. 16, 1966).
- MACFARLANE, C.: The determination of trace volatiles in whisky by gas chromatography. *Proc. Soc. Anal. Chem.*, 4 (1967) 99-100.
- MELKONYAN, S. A. AND VARTANYAN, S. D.: (Gas-liquid chromatography of high-boiling vinyl-acetylene compounds). *Zh. Analit. Khim.*, 22 (1967) 930-934 — retention data of 15 compounds (alcohols, ketoalcohols, chlorides) on silicone at 100°, 120° and 140°.
- WUCHERPENNIG, K. AND BRETHAUER, G.: Vergleich einiger gaschromatographischer Untersuchungsmethoden für Aromastoffe von Getränken sowie Beschreibung einer "Säulenextraktion" für Aromastoffe aus Getränken. *Z. Anal. Chem.*, 228 (1967) 342-360.
- YOUNG, F. AND HULCHER, F. H.: Cholesterol esters in myelin and the component fatty acids. *Proc. Soc. Exptl. Biol. Med.*, 123 (1966) 385-387.

J. Chromatog., 32 (1968) 205-220

Erratum

J. Chromatog., 28 (1967) 82-88

On the following pages the length of the column should be "50 cm" instead of "30 cm":

- Page 82, line 24
- Page 83, lines 31 and 33
- Page 85, line 2 of the legend to Fig. 1
- Page 86, line 2 of the legend to Fig. 2
- Page 87, line 10.

J. Chromatog., 31 (1967) 551-556

Page 552, line 28: For "at 110° 1 for h" read "at 110° for 1 h".

J. Chromatog., 32 (1968) 220