

Author Index

- Ackman, R. G., see Banerjee, A. K. 215
- Åkerblom, M.
Simple precolumn sample enrichment in high-performance liquid chromatography for determination of phenoxy acid herbicides in water samples from exposure studies 427
- Alexander, R., see Wajon, J. E. 187
- Aries, R. E.
—, Gutteridge, C. S. and Macrae, R.
Pyrolysis-mass spectrometry investigations of reversed-phase high-performance liquid chromatographic stationary phases 285
- Banerjee, A. K.
—, Ratnayake, W. M. N. and Ackman, R. G.
Enhanced response of more highly unsaturated lipids on thin-layer chromatography-laser-scanned flame ionization detection after exposing the developed Chromarods to iodine vapour 215
- Baranowski, J. D.
High-performance liquid chromatographic separation of pungency components of ginger 471
- Bergenheim, N.
—, Carlsson, U. and Klasson, K.
Rapid ion-exchange chromatography for preparative separation of proteins. Application to porcine and bovine carbonic anhydrases 59
- Bertrand, C., see Felix, G. 432
- Binsard, A. M., see Paphassarang, S. 412
- Blitterswijk, H. van, see Jansen, E. H. J. M. 436
- Booth, C., see Styring, M. G. 115
- Both-Miedema, R., see Jansen, E. H. J. M. 436
- Boukerche, H., see Clezardin, P. 67
- Burford, R. P., see Vimalasiri, P. A. D. T. 121
- Carlsson, U., see Bergenheim, N. 59
- Chang, J.-P.
—, El Rassi, Z. and Horváth, Cs.
Silica-bound polyethyleneglycol as stationary phase for separation of proteins by high-performance liquid chromatography 396
- Chooi, W. Y., see Stark, B. C. 367
- Clezardin, P.
—, McGregor, J. L., Manach, M., Boukerche, H. and Dechavanne, M.
One-step procedure for the rapid isolation of mouse monoclonal antibodies and their antigen binding fragments by fast protein liquid chromatography of a mono Q anion-exchange column 67
- Clift, R. G., see Morreal, C. E. 153
- Corradini, D.
— and Horváth, Cs.
Chromatography of Astra Blue dyes with eluents containing potassium thiocyanate in organic solvents 400
- Cotts, P. M., see Mumby, S. J. 241
- Crews, C., see Gilbert, J. 376
- Curran, N. M., see Lovering, E. G. 235
- Dechavanne, M., see Clezardin, P. 67
- De Jong, J.
—, Schouten, J. P., Muusze, R. G. and Tjaden, Y. R.
Analysis of β -carbolines by reversed-phase ion-pair partition chromatography with fluorometric detection 23
- Dellacherie, E., see Mathis, R. 43
- Dreyer, R. N., see Siegel, S. A. 351
- Dunn, D. L., see Scott, B. S. 419
- Edqvist, L.-E., see Hallin, P. 195
- El Rassi, Z., see Chang, J.-P. 396
- Felix, G.
— and Bertrand, C.
Separation of polyaromatic hydrocarbons on caffeine-bonded silica gel 432
- Fisher, D., see Nazareth, A. 382
- Freeman, C. J., see Musial, S. P. 467
- Fukunaga, K., see Shoji, S. 359
- Funakoshi, T., see Shoji, S. 359
- Funazo, K.
—, Wu, H.-L., Morita, K., Tanaka, M. and Shono, T.
Butylation of inorganic anions for simultaneous determination by gas chromatography 143
- Giese, R. W., see Nazareth, A. 382
- Gilbert, J.
—, Startin, J. R. and Crews, C.
Optimisation of conditions for the trimethylsilylation of trichothecene mycotoxins 376
- Godeau, R. P., see Paphassarang, S. 412
- Greenhut, J.
— and Rudolph, F. B.
Preparative separation of nucleotides by high-performance ion-exchange liquid chromatography using a volatile buffer system 461
- Gulyás, H.
Bestimmung von Aflatoxinen B₁, B₂, G₁, G₂ und M₁ durch Überdruck-Schicht-Chromatographie 105
- Gustine, D. L.
Determination of S-methyl cysteine sulfoxide in *Brassica* extracts by high-performance liquid chromatography 450

- Gutteridge, C. S., see Aries, R. E. 285
- Haas, M. J., see Hicks, K. B. 159
- Haken, J. K.
- and Korhonen, I. O. O.
Gas chromatography of homologous esters. XXVII. Retention increments of C₁-C₁₈ primary alkanols and their 2-chloropropanoyl and 3-chloropropanoyl derivatives on SE-30 and OV-351 capillary columns 131
- , see Vimalasiri, P. A. D. T. 121
- Hallin, P.
- , Madej, A. and Edqvist, L.-E.
Subunits of bovine lutropin. Hormonal and immunological activity after separation with reversed-phase high-performance liquid chromatography 195
- Hamacher, K.
- , Schmid, G., Sahn, H. and Wandrey, Ch.
Structural heterogeneity of cellooligomers homogeneous according to high-resolution size-exclusion chromatography 311
- Harvan, D. J., see Parker, C. E. 273
- Harvey, T. G.
- , Matheson, T. W., Pratt, K. C. and Stanborough, M. S.
Determination of carbonyl compounds in an Australian (Rundle) shale oil 230
- Hass, J. R., see Parker, C. E. 273
- Hawkins, B. L., see Miller, M. L. 9
- Hicks, K. B.
- , Lim, P. C. and Haas, M. J.
Analysis of uronic and aldonic acids, their lactones, and related compounds by high-performance liquid chromatography on cation-exchange resins 159
- Hirota, K.
- and Shimamura, M.
Sephacrose derivatives containing citric acid as affinity ligand. Purification of fumarase 173
- Hlaváč, C., see Kovář, J. 341
- Hori, K., see Matsushashi, S. 79
- Horowitz, P. M.
Rapid fluorescamine based protein assay usable in the presence of interfering substances 446
- Horváth, Cs., see Chang, J.-P. 396
- , see Corradini, D. 400
- Hubert, P., see Mathis, R. 43
- Hülm, M., see Pötter, H. 440
- Hunter, A. T.
Separation of anthraquinone sulphonic acids by ion-pair liquid chromatography 319
- Jansen, E. H. J. M.
- , Zoontjes, P. W., Van Blitterswijk, H., Both-Miedema, R. and Stephany, R. W.
Fast high-performance liquid chromatographic screening method for the presence of trenbolone and its major metabolite in urine of slaughter cattle 436
- Jones, R. S.
- and Stutte, C. A.
Chromatographic analysis of choline and acetylcholine by UV visualization 454
- Jong, J. de, see De Jong, J. 23
- Joppich, M., see Nazareth, A. 382
- Kagi, R. I., see Wajon, J. E. 187
- Kahle, V., see Kovář, J. 341
- Kamata, K.
- and Motohashi, N.
Separation of methyl-substituted benz[c]acridines by gas-liquid chromatography and high-performance liquid chromatography 331
- Kissa, E.
Determination of 2H-hexafluoropropanol-2 in air 407
- Klasson, K., see Bergenheimer, N. 59
- Kobayashi, S.
- , Schwartz, S. J. and Lineback, D. R.
Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 205
- Korhonen, I. O. O., see Haken, J. K. 131
- Kovář, J.
- , Turánek, J., Hlaváč, C., Vala, V. and Kahle, V.
Liquid chromatographic separations of dimers of nicotinamide adenine dinucleotide and nicotinamide adenine dinucleotide phosphate 341
- Krainhöfner, A., see Schade, W. 475
- Kubota, Y., see Shoji, S. 359
- Lascu, I., see Porumb, T. 218
- Lim, P. C., see Hicks, K. B. 159
- Lineback, D. R., see Kobayashi, S. 205
- Linton, R. W., see Miller, M. L. 9
- Lockwood, F. E., see Sprissler, B. 222
- Lovering, E. G.
- and Curran, N. M.
High-performance liquid chromatographic determination of cimetidine and related compounds (raw material and tablets) 235
- McGregor, J. L., see Clezardin, P. 67
- Maciel, G. E., see Miller, M. L. 9
- MacKenzie, S. L.
- and Tenaschuk, D.
Analysis of γ -carboxyglutamic acid by gas-liquid chromatography 404
- Macrae, R., see Aries, R. E. 285

- Madej, A., see Hallin, P. 195
- Manach, M., see Clezardin, P. 67
- Marsden, N. V. B.
Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.*-butanol 247
- Matheson, T. W., see Harvey, T. G. 230
- Mathis, R.
—, Hubert, P. and Dellacherie, E.
Sephacrose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 43
- Matsuhashi, S.
—, Mukai, T. and Hori, K.
Determination of protein spots separated by two-dimensional polyacrylamide gel electrophoresis 79
- Meagher, M. M., see Nikolov, Z. L. 51
- Miller, M. L.
—, Linton, R. W., Maciel, G. E. and Hawkins, B. L.
Characterization of silanol reactivity and acidity on octadecyl-bonded chromatographic supports by ²⁹Si solid-state nuclear magnetic resonance spectroscopy and surface titration 9
- Miyamoto, E., see Shoji, S. 359
- Morita, K., see Funazo, K. 143
- Morreal, C. E.
— and Clift, R. G.
Identification of phenols of 7,12-dimethylbenz[*a*]anthracene by conversion into their methyl ethers 153
- Motohashi, N., see Kamata, K. 331
- Mukai, T., see Matsuhashi, S. 79
- Mumby, S. J.
—, Cotts, P. M. and Russell, T. P.
Preparative fractionation of fluorescent-dye-labeled poly(ethylene oxide) by aqueous size-exclusion chromatography 241
- Musial, S. P.
—, C. J. Freeman and Sinsheimer, J. E.
Mitotane (*o,p'*-DDD) emulsion and tablet analysis by high-performance liquid chromatography 467
- Muusze, R. G., see De Jong, J. 23
- Nakatani, K., see Yoneda, H. 35
- Nazareth, A.
—, Joppich, M., Panthani, A., Fisher, D. and Giese, R. W.
Alkylation with dialkylsulfate diisopropylethylamine 382
- Nikolov, Z. L.
—, Maegher, M. M. and Reilly, P. J.
High-performance liquid chromatography of disaccharides on amine-bonded silica columns 51
- Ohnishi, J., see Shoji, S. 359
- Panthani, A., see Nazareth, A. 382
- Paphassarang, S.
—, Raynaud, J., Godeau, R. P. and Binsard, A. M.
Analyses qualitatives et quantitatives de la hyoscyamine-atropine et de la scopolamine dans les teintures mères de Solanacées par chromatographie liquide haute performance 412
- Parker, C. E.
—, Yamaguchi, K., Harvan, D. J., Smith, R. W. and Hass, J. R.
Liquid chromatography-chloride-attachment negative chemical ionization mass spectrometry 273
- Pirkle, W. H.
—, Tsiouras, A. and Sowin, T. J.
Preparative separation of enantiomers by flash chromatography 392
- Porumb, H., see Porumb, T. 218
- Porumb, T.
—, Porumb, H., Lascu, I. and Proinov, I.
Cibacron Blue 3G-A-substituted cross-linked agar beads: an inexpensive fast-flow affinity medium for large scale applications 218
- Pötter, H.
—, Hülm, M. and Schumann, S.
Verteilungschromatographie von Mutterkornalkaloiden auf Formamidgesättigten Kieselgel-Säulen 440
- Pratt, K. C., see Harvey, T. G. 230
- Price, C., see Styring, M. G. 115
- Proinov, I., see Porumb, T. 218
- Pyysalo, H., see Sjöberg, A.-M. 90
- Rassi, Z. El, see Chang, J.-P. 396
- Ratnayake, W. M. N., see Banerjee, A. K. 215
- Raynaud, J., see Paphassarang, S. 412
- Reilly, P. J., see Nikolov, Z. L. 51
- Rudolph, F. B., see Greenhut, J. 461
- Russell, T. P., see Mumby, S. J. 241
- Sabljić, A.
Calculation of retention indices by molecular topology. Chlorinated benzenes 1
- Sahm, H., see Hamacher, K. 311
- Schade, W.
— and Krainhöfner, A.
Die Bestimmung von 9 α -Hydroxy-androst-4-en-3,17-dion in Fermentationslösungen mittels Dünnschichtchromatographie 475
- Schieffer, G. W.
Limitation of assessing high-performance liquid chromatographic peak purity with photodiode array detectors 387
- Schmid, G., see Hamacher, K. 311
- Schouten, J. P., see De Jong, J. 23
- Schumann, S., see Pötter, H. 440

- Schwartz, S. J., see Kobayashi, S. 205
- Scott, B. S.
— and Dunn, D. L.
High-performance liquid chromatographic analysis of epinephrine enantiomers using a UV detector in series with an optical activity detector 419
- Shimamura, M., see Hirota, K. 173
- Shoji, S.
—, Ohnishi, J., Funakoshi, T., Kubota, Y., Fukunaga, K., Miyamoto, E. and Ueki, H.
Simple and rapid identification of phosphorylated peptides from bovine brain myelin basic protein by reversed-phase high-performance liquid chromatography 359
- Shono, T., see Funazo, K. 143
- Siegel, S. A.
—, Woods, K. L. and Dreyer, R. N.
Pyrimidine nucleoside phosphorylase assay by automated high-performance liquid chromatography 351
- Sinsheimer, J. E., see Musial, S. P. 467
- Sjöberg, A.-M.
— and Pyysalo, H.
Identification of food thickeners by monitoring of their pyrolytic products 90
- Smith, R. W., see Parker, C. E. 273
- Sowin, T. J., see Pirkle, W. H. 392
- Sprissler, B.
— and Lockwood, F. E.
Analysis of mechanical lubricant contaminants in aluminium cold rolling oil by reversed-phase liquid chromatography 222
- Stanborough, M. S., see Harvey, T. G. 230
- Stark, B. C.
— and Chooi, W. Y.
Affinity chromatography of *Drosophila melanogaster* ribosomal proteins to 5S rRNA 367
- Startin, J. R., see Gilbert, J. 376
- Stephany, R. W., see Jansen, E. H. J. M. 436
- Stutte, C. A., see Jones, R. S. 454
- Styring, M. G.
—, Price, C. and Booth, C.
The effect of solute size and shape on the elution volume in size-exclusion chromatography 115
- Tanaka, M., see Funazo, K. 143
- Tenaschuk, D., see MacKenzie, S. L. 404
- Thyssen, K.
Determination of chloroacetanilides and chloroanilines in sulphanilamides and waste water by high-performance liquid chromatography 99
- Tjaden, U. R., see De Jong, J. 23
- Tsipouras, A., see Pirkle, W. H. 392
- Turánek, J., see Kovář, J. 341
- Ueki, H., see Shoji, S. 359
- Vala, V., see Kovář, J. 341
- Van Blitterswijk, H., see Jansen, E. H. J. M. 436
- Vimalasiri, P. A. D. T.
—, Haken, J. K. and Burford, R. P.
Chromatographic analysis of polyether based polyurethanes after alkali fusion 121
- Wajon, J. E.
—, Alexander, R. and Kagi, R. I.
Determination of trace levels of dimethyl polysulphides by capillary gas chromatography 187
- Wandrey, Ch., see Hamacher, K. 311
- Wells, G.
Design and performance of a mass-flow-modulated detector for gas chromatography 263
- Woods, K. L., see Siegel, S. A. 351
- Wu, H.-L., see Funazo, K. 143
- Yamagishi, A.
Clay column chromatography for optical resolution. Resolution of aromatic compounds on a *A*-Ru(1,10-phenanthroline)₃²⁺-montmorillonite column 299
- Yamaguchi, K., see Parker, C. E. 273
- Yamazaki, S., Yoneda, H. 35
- Yoneda, H.
—, Nakatani, K. and Yamazaki, S.
Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 35
- Zoontjes, P. W., see Jansen, E. H. J. M. 436