

Author Index

- Aerts, M. M. L., see Keukens, H. J. 445
 Aguilar, M. I., see Hearn, M. T. W. 35
 Araki, M., see Tanaka, N. 307
 Beek, W. M. J., see Keukens, H. J. 445
 Berg, J. H. M. van den, see Glöckner, G. 511
 Beynen, J. H., see Lingeman, H. 261
 Billiet, H. A. H., see Drouen, A. C. J. H. 127
 Brown, P. R., see Yi, C. 221
 Burda, J., see Kříž, J. 427
 Chase, D. B., see Köhler, J. 275
 Coddens, M. E., see Poole, C. F. 407
 Dawkins, J. V.
 —, Lloyd, L. L. and Warner, F. P.
 Chromatographic characteristics of polymer-based high-performance liquid chromatography packings 157
 De Galan, L., see Drouen, A. C. J. H. 127
 Drouen, A. C. J. H.
 —, Billiet, H. A. H. and De Galan, L.
 Practical procedure for the optimization of reversed-phase separations with quaternary mobile phase mixtures 127
 Farlee, R. D., see Köhler, J. 275
 Fasching, J. L., see Yi, C. 221
 Fazio, S. D., see Kaliszian, R. 141
 Frei, R. W., see Kok, W. Th. 27
 Furton, K. G., see Poole, C. F. 407
 Galan, L. de, see Drouen, A. C. J. H. 127
 Garteiz, D., see McCown, S. M. 465, 483
 Glöckner, G.
 — and Van den Berg, J. H. M.
 Precipitation and adsorption phenomena in polymer chromatography 511
 Goworek, J.
 —, Nooitgedacht, F., Rijkhof, M. and Poppe, H.
 Determination of silanol groups on packings for high-performance liquid chromatography by means of isotopic exchange 399
 Green, J. S.
 — and Jorgenson, J. W.
 Variable-wavelength on-column fluorescence detector for open-tubular zone electrophoresis 337
 Grego, B.
 —, Nice, E. C. and Simpson, R. J.
 Use of scanning diode array detector with reversed-phase microbore columns for the real-time spectral analysis of aromatic amino acids in peptides and proteins at the submicrogram level. Applications to peptide and protein microsequencing 359
 Halvax, J. J., see Kok, W. Th. 27
 Hartwick, R. A., see Kaliszian, R. 141
 Hearn, M. T. W.
 — and Aguilar, M. I.
 High-performance liquid chromatography of amino acids, peptides and proteins. LXVII. Evaluation of bandwidth relationships of peptides related to human β -endorphin, separated by gradient-elution reversed-phase high-performance liquid chromatography 35
 Halfman, G. W., see Spittler, T. D. 439
 Ho, S. S. J., see Poole, C. F. 407
 Honda, K., see Miyaguchi, K. 255
 Horiguchi, M., see Takahagi, H. 369
 Hsu, S.-H., see Kaliszian, R. 141
 Hulshoff, A., see Lingeman, H. 261
 Imai, K., see Miyaguchi, K. 255
 Inoue, K., see Takahagi, H. 369
 Jackson, M. A., see Stout, R. W. 381
 Jandera, P.
 Method for characterization of selectivity in reversed-phase liquid chromatography. I. Derivation of the method and verification of the assumptions 91
 —
 Method for characterization of selectivity in reversed-phase liquid chromatography. II. Possibilities for the prediction of retention data 111
 Jorgenson, J. W., see Green, J. S. 337
 Kaliszian, R.
 —, Osmialowski, K., Tomellini, S. A., Hsu, S.-H., Fazio, S. D. and Hartwick, R. A.
 Quantitative retention relationships as a function of mobile and C_{18} stationary phase composition for non-cogeneric solutes 141
 Katz, E. D.
 —, Ogan, K. and Scott, R. P. W.
 Distribution of a solute between two phases. The basic theory and its application to the prediction of chromatographic retention 67
 Kaur, B., see Knox, J. H. 3
 Kersten, B.R., see Poole, C. F. 407
 Keukens, H. J.
 —, Beek, W. M. J. and Aerts, M. M. L.
 High-performance liquid chromatographic screening and confirmation methods for chloramphenicol residues in meat with off-line cartridge sample clean-up and on-line diode array UV-VIS detection 445
 Kimata, K., see Tanaka, N. 307
 Kinkel, J. N., see Lork, K. D. 199

- Kinoshita, T., see Nimura, N. 169
 Kirkland, J. J., see Köhler, J. 275
 Köhler, J.
 —, Chase, D. B., Farlee, R. D., Vega, A. J. and Kirkland, J. J.
 Comprehensive characterization of some silica-based stationary phases for high-performance liquid chromatography 275
- Kok, W. Th.
 —, Halvax, J. J. and Frei, R. W.
 Comparison of electrochemical detection methods in liquid chromatography for the determination of N-acetylcysteine in plasma 27
- Kouřilová, D., see Šlais, K. 179
- Knox, J. H.
 —, Kaur, B. and Millward, G. R.
 Structure and performance of porous graphitic carbon in liquid chromatography 3
- Kraak, J. C., see Veening, H. 345
- , Van Rooij, H. H. and Thus, J. L. G.
 Reversed-phase ion-pair systems for the prediction of *n*-octanol-water partition coefficients of basic compounds by high-performance liquid chromatography 455
- Krejčí, M., see Šlais, K. 179
- Kříž, J.
 —, Vodička, L. and Burda, J.
 High-performance liquid chromatography of methyl esters of diamantane and adamantine carboxylic acids 427
- Lawson, A. M., see Meijers, J. C. M. 231
- Lim, C. K., see Meijers, J. C. M. 231
- Lingeman, H.
 —, Van Munster, H. A., Beynen, J. H., Udenberg, W. J. M. and Hulshoff, A.
 High-performance liquid chromatographic analysis of basic compounds on non-modified silica gel and aluminium oxide with aqueous solvent mixtures 261
- Lloyd, L. L., see Dawkins, J. V. 157
- Lork, K. D.
 —, Unger, K. K. and Kinkel, J. N.
 Role of the functional group in *n*-octyldimethylsilanes in the synthesis of C₈ reversed-phase packings for high-performance liquid chromatography 199
- McCown, S. M.
 —, Southern, D., Morrison, B. E. and Garteiz, D.
 Solvent properties and their effects on gradient elution high-performance liquid chromatography. II. Temperature gradients in high-performance liquid chromatographic columns 483
 —, Southern, D. and Morrison, B. E.
 Solvent properties and their effects on gradient elution high-performance liquid chromatography. III. Experimental findings for water and acetonitrile 493
- Maeda, M.
 — and Tsuji, A.
 Chemiluminescence with lucigenin as post-column reagent in high-performance liquid chromatography of corticosteroids and *p*-nitrophenacyl esters 213
- Marafioti, R. A., see Spittler, T. D. 439
- Meijers, J. C. M.
 —, Lim, C. K., Lawson, A. M. and Peters, T. J.
 Analysis of tumour-localizing haematoxophyrin derivative by high-performance liquid chromatography and fast-atom bombardment mass spectrometry 231
- Millward, G. R., see Knox, J. H. 3
- Miyaguchi, K.
 —, Honda, K., Toyo'oka, T. and Imai, K.
 Application of a microbore high-performance liquid chromatography-chemiluminescence detection system to the N-terminal amino acid of bradykinin 255
- Morrison, B. E., see McCown, S. M. 465, 483, 493
- Morse, R. A., see Spittler, T. D. 439
- Munster, H. A. van, see Lingeman, H. 261
- Nambara, T., see Shimada, K. 329
- Nice, E. C., see Grego, B. 359
- Nimura, N.
 — and Kinoshita, T.
o-Phthalaldehyde-N-acetyl-L-cysteine as a chiral derivatization reagent for liquid chromatographic optical resolution of amino acid enantiomers and its application to conventional amino acid analysis 169
- Nooitgedacht, F., see Goworek, J. 399
- Odiorne, T. J., see Stout, R. W. 381
- Ogan, K., see Katz, E. D. 67
- Orii, S., see Shimada, K. 329
- Osmialowski, K., see Kaliszan, R. 141
- Palmer, H. C., see Stout, R. W. 381

- Patience, R. L.
— and Rees, L. H.
Comparison of reversed-phase and anion-exchange high-performance liquid chromatography for the analysis of human growth hormones 241
- Peters, T. J., see Meijers, J. C. M. 231
- Poole, C. F.
—, Kersten, B. R., Ho, S. S. J., Coddens, M. E. and Furton, K. G.
Organic salts, liquid at room temperature, as mobile phases in liquid chromatography 407
- Poppe, H., see Goworek, K. J. 399
—, see Veening, H. 345
- Rees, L. H., see Patience, R. L. 241
- Ricker, R. D., see Stout, R. W. 381
- Rijkhof, M., see Goworek, J. 399
- Rooij, H. H. van, see Kraak, J. C. 455
- Schoenmakers, P. J.
— and Verhoeven, F. C. C. J. G.
Effect of pressure on retention in supercritical-fluid chromatography with packed columns 315
- Scott, R. P. W., see Katz, E. D. 67
- Shimada, K.
—, Orii, S., Tanaka, M. and Nambara, T.
New ferrocene reagents for derivatization of alcohols in high-performance liquid chromatography with electrochemical detection 329
- Simpson, R. J., see Grego, B. 359
- Sivakoff, S. I., see Stout, R. W. 381
- Šlais, K.
—, Krejčí, M. and Kouřilová, D.
Trace analysis by injection-generated gradient ion-pair chromatography with micro-bore column high-performance liquid chromatography 179
- Southern, D., see McCown, S. M. 465, 483, 493
- Spittler, T. D.
—, Marafioti, R. A., Helfman, G. W. and Morse, R. A.
Determination of carbaryl in honeybees and pollen by high-performance liquid chromatography 439
- Stout, R. W.
—, Sivakoff, S. I., Ricker, R. D., Palmer, H. C., Jackson, M. A. and Odiorne, T. J.
New ion-exchange packings based on zirconium oxide surface-stabilized, diol-bonded, silica substrates 381
- Takahagi, H.
—, Inoue, K. and Horiguchi, M.
Drug monitoring by a fully automated high-performance liquid chromatographic technique, involving direct injection of plasma 369
- Tanaka, M., see Shimada, K. 329
- Tanaka, N.
—, Araki, M. and Kimata, K.
Separation of oxygen isotopic compounds by reversed-phase liquid chromatography 307
- Thus, J. L. G., see Kraak, J. C. 455
- Tock, P. P. H., see Veening, H. 345
- Tomellini, S. A., see Kalisz, R. 141
- Toyo'oka, T., see Miyaguchi, K. 255
- Tsuji, A., see Maeda, M. 213
- Underberg, W. J. M., see Lingeman, H. 261
- Unger, K. K., see Lork, K. D. 199
- Van den Berg, J. H. M., see Glöckner, G. 511
- Van der Wal, Sj.
Comparison of some high-performance liquid chromatographic fluorescence detector flow cells 351
- Van Munster, H. A., see Lingeman, H. 261
- Van Rooij, H. H., see Kraak, J. C. 455
- Veening, H.
—, Tock, P. P. H., Kraak, J. C. and Poppe, H.
Microbore high-performance liquid chromatography with a moving-wire flame ionization detector 345
- Vega, A. J., see Köhler, J. 275
- Verhoeven, F. C. C. J. G., see Schoenmakers, P. J. 315
- Vodička, L., see Kříž, J. 427
- Wal, Sj. van der, see Van der Wal, Sj. 351
- Warner, F. P., see Dawkins, J. V. 157
- Yi, C.
—, Fasching, J. L. and Brown, P. R.
Temperature, methanol, and structure effects in the reversed-phase liquid chromatography of guanine and hypoxanthine and their ribonucleosides and ribonucleotides 221