

Author Index

- Aburatani, R., see Okamoto, Y. 454
- Adachi, S., see Hashimoto, K. 241
- Al-Khalidi, A. H. E., see Girard, J. E. 355
- Altmayer, B., see Schwenk, S. 424
- Amato, A., see Gagliardi, L. 296
- Andreolini, F., see Flurer, C. L. 73
- Arai, T.
- , Koike, H., Hirata, K. and Ōzumi, H.
Separation of pyridone carboxylic acid enantiomers by high-performance liquid chromatography using copper(II)-L-amino acid as the eluent 439
- Araki, M., see Tanaka, N. 95
- Armstrong, D. W.
- , He, F.-Y. and Han, S. M.
Planar chromatographic separation of enantiomers and diastereomers with cyclodextrin mobile phase additives 345
- Beasley, V. R., see Harada, K.-I. 275
- Beger, J., see Voelkel, A. 219
- Benická, E., see Krupčík, J. 203
- Bezard, J., see Narce, M. 249
- Biedermann, M., see Grob, K. 387
- Borra, C., see Flurer, C. L. 73
- Buell, P. E., see Girard, J. E. 355
- Caponecchi, G., see Polcaro, C. 127
- Carmichael, W. W., see Harada, K.-I. 275
- Caude, M., see Lienne, M. 55
- Cavazzutti, G., see Gagliardi, L. 296
- Chiou, S.-H.
- , and Wang, K.-T.
Simplified protein hydrolysis with methanesulphonic acid at elevated temperature for the complete amino acid analysis of proteins 404
- Cho, S. Y., see Nam, H. S. 445
- Cohen, A. S., see Guttman, A. 41
- Cozzi, R., see Polcaro, C. 127
- Crouch, Jr., F. W.
- , Riley, C. M. and Stobaugh, J. F.
High-performance liquid chromatography of bis[1,2-bis-(diphenylphosphino)ethane]gold(I) chloride, a potential antineoplastic agent 333
- Dahiya, J. S.
- and Rimmer, S. R.
High-performance liquid chromatography of phytoalexins in stem callus tissues of rapeseed 448
- Dahlem, A. M., see Harada, K.-I. 275
- Eichhorn, K. W., see Schwenk, S. 424
- Fiore, M., see Polcaro, C. 127
- Flurer, C. L.
- , Borra, C., Andreolini, F. and Novotny, M.
Separation of proteins by microcolumn liquid chromatography based on the reversed-phase and size-exclusion principles 73
- Fujii, Y.
- , Ikeda, Y. and Yamazaki, M.
Micro high-performance liquid chromatographic determination of cardiac glycosides in β -methyl digoxin and digoxin tablets 157
- Gagliardi, L.
- , Amato, A., Cavazzutti, G., Tonelli, D. and Montanarella, L.
Identification of xanthene dyes in lipsticks by reversed-phase high-performance liquid chromatography 296
- Gál, S., see Kiss, L. 109
- Galensa, R., see Risch, B. 291
- Gangoda, M. E.
- and Gilpin, R. K.
Comparative study of alkyl and fluoroalkyl N-(2-hydroxy-ethoxyethyl)-amides as reversed-phase liquid chromatographic modifiers 365
- Gawdzik, B.
- and Patrykiewicz, A.
Gas chromatography on porous polymers. IV. Influence of the geometric structure of porous copolymers of 1,4-di(methacryloyloxymethyl)-naphthalene with 1,4-divinylbenzene on their chromatographic behaviour 233
- Gilpin, R. K., see Gangoda, M. E. 365
- Girard, J. E.
- , Rebbani, N., Buell, P. E. and Al-Khalidi, A. H. E.
Carbohydrate-borate eluents for anion chromatography 355
- Gorbunov, A. A.
- , Solovyova, L. Ya. and Pasechnik, V. A.
Fundamentals of the theory and practice of polymer gel-permeation chromatography as a method of chromatographic porosimetry 307
- Greenaway, W.
- , Wollenweber, E., Scaysbrook, T. and Whatley, F. R.
Novel isoferulate esters identified by gas chromatography-mass spectrometry in bud exudate of *Populus nigra* 284
- Gref, R.
Gas chromatographic analysis of underivatized resin acids 428

- Gresti, J., see Narce, M. 249
 Grinberg, N., see Guttman, A. 41
 Grob, K.
 —, Biedermann, M. and Li, Z.
 Checking the capacity of a splitless injector —
 a simple test 387
 Günther, K.
 Thin-layer chromatographic enantiomeric res-
 olution via ligand exchange 11
 Guttman, A.
 —, Paulus, A., Cohen, A. S., Grinberg, N. and
 Karger, B. L.
 Use of complexing agents for selective separa-
 tion in high-performance capillary electropho-
 resis. Chiral resolution via cyclodextrins in-
 corporated within polyacrylamide gel
 columns 41
 Hagman, A.
 — and Jacobsson, S.
 Trapping efficiency of capillary cold
 traps 117
 Han, S. M., see Armstrong, D. W. 345
 Harada, K.-I.
 —, Matsuura, K., Suzuki, M., Oka, H., Wata-
 nabe, M. F., Oishi, S., Dahlem, A. M., Beasley,
 V. R. and Carmichael, W. W.
 Analysis and purification of toxic peptides
 from cyanobacteria by reversed-phase high-
 performance liquid chromatography 275
 Hashidzume, K., see Tanaka, N. 95
 Hashimoto, H.
 —, Koyama, Y. and Shimura, T.
 Isolation of *cis-trans* isomers of canthaxan-
 thin by high-performance liquid chromatogra-
 phy using a calcium hydroxide column and
 identification of their configurations by ¹H
 NMR spectroscopy 182
 Hashimoto, K.
 —, Shirai, Y., Adachi, S. and Horie, M.
 Effect of the content of divinylbenzene in ion-
 exchange resins on the chromatographic sep-
 aration of α -cyclodextrin and glucose 241
 Hatada, K., see Okamoto, Y. 454
 He, F.-Y., see Armstrong, D. W. 345
 Heimler, D.
 —, Vidrich, V.
 Thin-layer chromatographic characterization
 of essential oils 301
 Hernádi, F. J., see Kiss, L. 109
 Herrmann, K., see Risch, B. 291
 Hevesi, T., see Krupčik, J. 203
 Hirata, K., see Arai, T. 439
 Hoque, E.
 Isocratic reversed-phase high-performance
 liquid chromatographic analysis of pigments
 in Norway spruce 417
 Horie, M., see Hashimoto, K. 241
 Hostettmann, K., see Slacanin, I. 265
 Ikeda, Y., see Fujii, Y. 157
 Iwaguchi, K., see Tanaka, N. 95
 Jacobsson, S., see Hagman, A. 117
 Jinno, K., see Wahyuni, W. T. 398
 Karger, B. L., see Guttman, A. 41
 Kaufmann, S. H. E., see Schoel, B. 165
 Kiss, L.
 —, Tar, A., Gál, S., Toth-Martinez, B. L. and
 Hernádi, F. J.
 Modified general affinity adsorbent for large-
 scale purification of penicillinases 109
 Koike, H., see Arai, T. 439
 Korpela, T., see Mäkelä, M. 391
 Koyama, Y., see Hashimoto, H. 182
 Krupčik, J.
 —, Repka, D., Benická, E., Hevesi, T., Nolte, J.,
 Paschold, B. and Mayer, H.
 Use of retention temperatures for the identifi-
 cation of phenols separated on OV-1701 capil-
 lary columns using linear temperature-pro-
 grammed gas chromatography 203
 Lawson, D. E. M., see Okano, T. 145
 Lázaro, F.
 —, Luque de Castro, M. D. and Valcárcel, M.
 Fluorimetric determination of aflatoxins in
 foodstuffs by high-performance liquid chro-
 matography with flow injection analysis 173
 Li, Z., see Grob, K. 387
 Lienne, M.
 —, Caude, M., Rosset, R. and Tambute, A.
 Optimization of the direct chiral separation of
 potential cytotoxic α -methylene- γ -butyrolac-
 tones and α -methylene- γ -butyrolactams by
 liquid chromatography 55
 Luque de Castro, M. D., see Lázaro, F. 173
 Mäkelä, M.
 —, Mattsson, P., Pintamo-Kenttälä, K. and
 Korpela, T.
 Cyclodextrin-ligand interaction as a simpli-
 fied model of biospecific affinity chromatogra-
 phy 391
 Marston, A., see Slacanin, I. 265
 Matsuura, K., see Harada, K.-I. 275
 Mattsson, P., see Mäkelä, M. 391
 Mayer, H., see Krupčik, J. 203
 Montanarella, L., see Gagliardi, L. 296
 Morris, M. D., see Peck, K. 193
 Nam, H. S.
 —, Cho, S. Y. and Rhee, J. S.
 High-performance liquid chromatographic
 analysis of major carotenoids from *Rhodotor-
 ula glutinis* 445

- Narce, M.
 —, Gresti, J. and Bezar, J.
 Method for evaluating the bioconversion of radioactive polyunsaturated fatty acids by use of reversed-phase liquid chromatography 249
- Nau, H., see Wittfoht, W. 433
- Netting, A. G., see Rhodes, S. H. 135
- Nicoletti, I., see Polcaro, C. 127
- Nolte, J., see Krupčik, J. 203
- Nomura, A.
 —, Yamada, J. and Tsunoda, K.-I.
 Effect of the pore structure of silica gel on retention behaviour in pressure programming supercritical fluid chromatography 87
- Novotny, M., see Flurer, C. L. 73
- Ohnishi, S., see Tanaka, N. 95
- Oishi, S., see Harada, K.-I. 275
- Ōizumi, H., see Arai, T. 439
- Oka, H., see Harada, K.-I. 275
- Okamoto, Y.
 —, Aburatani, R. and Hatada, K.
 Chromatographic resolution. XXI. Direct optical resolution of abscisic acid by high-performance liquid chromatography on cellulose tris(3,5-dimethylphenylcarbamate) 454
- Okano, T.
 — and Lawson, D. E. M.
 Chick 28 000 *M_v* vitamin D-dependent calcium-binding protein in intestine, kidney and cerebellum. Purification using chromatofocusing 145
- Okuno, A., see Tanaka, N. 95
- Ossicini, L., see Polcaro, C. 127
- Palitti, F., see Polcaro, C. 127
- Paschold, B., see Krupčik, J. 203
- Pasechnik, V. A., see Gorbunov, A. A. 307
- Patrykiewicz, A., see Gawdzik, B. 233
- Paulus, A., see Guttman, A. 41
- Peck, K.
 — and Morris, M. D.
 Optical errors in a liquid chromatography absorbance cell 193
- Pintamo-Kenttälä, see Mäkelä, M. 391
- Polcaro, C.
 —, Nicoletti, I., Ossicini, L., Caponecchi, G., Cozzi, R., Fiore, M. and Palitti, F.
 Chromatographic and cytogenetic analysis of *in vivo* metabolites of fluoranthene 127
- Rebbani, N., see Girard, J. E. 355
- Repka, D., see Krupčik, J. 203
- Rhee, J. S., see Nam, H. S. 445
- Rhodes, S. H.
 — and Netting, A. G.
 Normal-phase high-performance liquid chromatography of triacylglycerols 135
- Rimmer, S. R., see Dahiya, J. S. 448
- Riley, C. M., see Crouch, Jr., F. W. 333
- Risch, B.
 —, Galensa, R. and Herrmann, K.
 Analysis of catechin and epicatechin by high-performance liquid chromatography after benzoylation 291
- Rosset, R., see Lienne, M. 55
- Rüstig, H., see Voelkel, A. 219
- Scaysbrook, T., see Greenaway, W. 284
- Schoel, B.
 — and Kaufmann, S. H. E.
 Isolation of recombinant mycobacterial antigens by an automatic and generally applicable purification method for β -galactosidase fusion proteins 165
- Schwenk, S.
 —, Altmayer, B. and Eichhorn, K. W.
 Simultaneous detection of trichothecenes and rosenonolactone in grape juice and wine by capillary gas chromatography 424
- Scott, W. J., see Wittfoht, W. 433
- Shimamura, T., see Hashimoto, H. 182
- Shirai, Y., see Hashimoto, K. 241
- Siefermann-Harms, D.
 High-performance liquid chromatography of chloroplast pigments. One-step separation of carotene and xanthophyll isomers, chlorophylls and pheophytins 411
- Slacanin, I.
 —, Marston, A. and Hostettmann, K.
 High-performance liquid chromatographic determination of molluscicidal saponins from *Phytolacca dodecandra* (phytolaccaceae) 265
- Solovyova, L. Ya., see Gorbunov, A. A. 307
- Stephens, R. E.
 Separation of tubulin subunits by reversed-phase high-performance liquid chromatography 371
- Stobaugh, J. F., see Crouch, Jr., F. W. 333
- Stromberg, R.
 Statistical optimization of a reversed-phase ion-pair liquid chromatographic method for the analysis of tolmetin sodium in dosage forms 1
- Suzuki, M., see Harada, K.-I. 275
- Szymanowski, J., see Voelkel, A. 219
- Takai, N., see Tanaka, N. 95
- Tambute, A., see Lienne, M. 55
- Tanaka, N.
 —, Hashidzume, K., Araki, M., Tsuchiya, H., Okuno, A., Iwaguchi, K., Ohnishi, S. and Takai, N.
 Internal structures of wide-pore packing materials for high-performance liquid chromatography studied by transmission electron microscopy 95
- Tar, A., see Kiss, L. 109

- Tonelli, D., see Gagliardi, L. 296
Toth-Martinez, B. L., see Kiss, L. 109
Tsuchiya, H., see Tanaka, N. 95
Tsunoda, K.-I., see Nomura, A. 87
Valcárcel, M., see Lázaro, F. 173
Vidrich, V., see Heimler, D. 301
Voelkel, A.
——, Szymanowski, J., Beger, J. and Rüstig, H.
Polarity of some individual derivatives of α,ω -diaminoalcohols as measured by gas chromatography 219
Wahyuni, W. T.
—— and Jinno, K.
Separation of tocopherols on various chemically bonded phases in microcolumn liquid chromatography 398
Wang, K.-T., see Chiou, S.-H. 404
Watanabe, M. F., see Harada, K.-I. 275
Whatley, F. R., see Greenaway, W. 284
Wičar, S.
Flow velocity profiles in rectangular channels containing two liquid layers of different densities and viscosities for sedimentation field flow fractionation 456
Wilson, T. D.
Sample solvent effects in an apparent chiral high-performance liquid chromatographic separation on β -cyclodextrin 31
Wittfoht, W.
——, Scott, W. J. and Nau, H.
Assay of methoxyacetic acid in body fluids and tissues by gas chromatography-mass spectrometry following *tert.*-butyldimethylsilylation 433
Wollenweber, E., see Greenaway, W. 284
Yamada, J., see Nomura, A. 87
Yamazaki, M., see Fujii, Y. 157