

## Short Communication

---

# Identification of vitamin D<sub>2</sub> by thermospray-interface mass spectrometry

KAZUNORI TAKAMURA\* and HIROKO HOSHINO

*Seitoku Junior College of Nutrition, 1–4–6, Nishi-shinkoiwa, Katsushika-ku, Tokyo 124 (Japan)*

NORIAKI HARIMA

*JASCO International Co., Ltd., 2–4–21, Sennin-cho, Hachioji City, Tokyo 193 (Japan)*

TATSUYUKI SUGAHARA

*Kagawa Nutrition College, 3–24–2, Komagome, Toshima-ku, Tokyo 170 (Japan)*

and

HISAO AMANO

*Toho University School of Medicine, 5–21–16, Ohmori-nishi, Ohta-ku, Tokyo 143 (Japan)*

(First received July 16th, 1990; revised manuscript received December 11th, 1990)

---

## ABSTRACT

In order to identify vitamin D<sub>2</sub> contained in shiitake mushroom (*Lentinus edodes*), which is taken routinely in Japan, vitamin D<sub>2</sub> was isolated by thin-layer liquid chromatography and high-performance liquid chromatography and identified by thermospray-interface mass spectrometry; this procedure prevents the decomposition of vitamin by heat, which is a common problem in the gas chromatography–mass spectrometry of vitamin D<sub>2</sub>.

---

## INTRODUCTION

The identification of vitamin D<sub>2</sub> [1] (hereafter called D<sub>2</sub>) and the determination of vitamin D<sub>3</sub> [2] metabolites have usually been conducted by gas chromatography–mass spectrometry (GC–MS). It is well known, however, that the heating process during GC causes ring closure of a steroid ring between the 9- and 10-positions, converting D<sub>2</sub> into pyro D<sub>2</sub> and isopyro D<sub>2</sub> [3] (Fig. 1). In order to identify D<sub>2</sub> contained in shiitake mushroom (*Lentinus edodes*), which is taken routinely in Japan, we isolated D<sub>2</sub> by thin-layer liquid chromatography (TLC) and high-performance liquid chromatography (HPLC) and the isolated substance was identified by thermospray-interface mass spectrometry (TSP–MS). We have subsequently succeeded in obtaining a mass spectrum of D<sub>2</sub> that does not suffer from thermal ring closure.




Fig. 1. Thermal isomerization of vitamin D<sub>2</sub>.

## EXPERIMENTAL

All reagents were of analytical-reagent grade from Wake (Osaka, Japan). Shiitake mushroom (brand name Koshin) produced in Japan from March to May was used as a sample. A 500-g sample was ground with a homogenizer (one fifth at a time), and all of it was placed in a digestion flask and decomposed by heating at 80°C for 30 min after addition of absolute non-aldehyde ethanol (400 ml), pyrogallol (40 g) and 50% potassium hydroxide solution (100 ml). After cooling to room temperature, unsaponifiable matter was extracted with 1000 ml of benzene, followed by washing once each with 500 ml of 1 *M* and 300 ml of 0.5 *M* potassium hydroxide solution and then four times with 100 ml of water [4]. The upper benzene layer was separated by TLC using a Wako-gel B5FM silica gel TLC plate (Wako) with benzene–acetone (95:5) as developing solvent and UV detection (254 nm) so as to obtain a D<sub>2</sub> fraction.

This D<sub>2</sub> fraction was extracted with chloroform and evaporated to dryness below 35°C. To the resulting residue 2 ml of methanol–acetonitrile (1:1) were added and this solution was further purified twice by HPLC on an NSLC Model 100A chromatograph (Nihon Seimitsu Kagaku). In the first step a LiChrosorb RP-18 column (250 × 7.5 mm I.D.) was used with methanol–acetonitrile (1:1) as eluent and in the second step a Nucleosil 100-5 column column (150 × 4.6 mm I.D.) with *n*-hexane containing 0.1% of *n*-amyl alcohol and 0.4% of isopropyl alcohol as eluent.

Finally, the isolated substance was dissolved in 0.5 ml of 0.1 *M* ammonium acetate–methanol (4:6) and directly subjected to identification of D<sub>2</sub> by TSP-MS on a VG Model 12-250 instrument (VG Analytical) without any HPLC column, under the following conditions: solvent flow-rate, 0.4 ml/min; source temperature, 230°C; probe temperature, 250°C; electron energy, 70 eV.

## RESULTS AND DISCUSSION

Fig. 2 is a TSP mass spectrum the purified extract of shiitake mushroom. A major peak was detected at  $m/z$  397,  $[M+H]^+$ , but little fragmentation was observed. Similar results were obtained with standard D<sub>2</sub>. Pyro D<sub>2</sub> and isopyro D<sub>2</sub> were also subjected to TSP-MS under the same conditions as those employed for the




Fig. 2. Mass spectrum of the purified  $D_2$  fraction obtained from shiitake mushroom (concentration  $5 \mu\text{g}/\mu\text{l}$ ). Injection volume,  $3 \mu\text{l}$ .

standard  $D_2$ . No peak at  $m/z$  397 was obtained with these samples, indicating that  $D_2$  suffering no thermal ring closure can be identified by TSP-MS.

The identification of aldosterone and corticosterone, each having a steroid ring similar to that in vitamin  $D_3$ , by LC-MS [5,6] (measurement concentration  $1 \mu\text{g}$  in  $100 \mu\text{l}$ ) and the determination of vitamin  $D_3$  metabolites (1,25-dihydroxy-vitamin  $D_3$  and 24,25-dihydroxy-vitamin  $D_3$ ) by GC-MS [7,8] (determination concentration 0–250 ng/l) have been reported previously. In contrast to dihydroxy-vitamin  $D_3$ , the measurement concentration employed in this study was as high as  $5 \mu\text{g}/\mu\text{l}$ , as  $D_2$  is poorly ionized in TSP-MS as there are few functional groups. Previously  $D_2$  has not been identified by TSP-MS.

#### REFERENCES

- 1 A. Takeuchi, T. Okano, S. Teraoka, Y. Murakami, M. Sayamoto, S. Sawamura and T. Kobayashi, *Bitamin*, 58 (1984) 439.
- 2 J. Fohlman, P. A. Peterson, I. Kamensky, P. Hakansson and B. B. Sundqvist, *Nucl. Instrum. Methods. Phys. Res.*, 198 (1982) 169.
- 3 P. Busse, *Hoppe-Seyler's Z. Physiol. Chem.*, 214 (1933) 211.
- 4 F. J. Mulder, E. J. DeVries and K. J. Keuning, *Pharm. Weekbl.*, 100 (1965) 1457.
- 5 D. Watson, G. W. Taylor, S. Laird and G. P. Vinson, *Biochem. J.*, 242 (1987) 109.
- 6 P. R. Das, B. N. Pramanik, R. D. Malchow and K. J. Ng, *Biomed. Environ. Mass Spectrom.*, 15 (1988) 253.
- 7 I. Bjorkhem, I. Holmberg, T. Kristiansen and J. I. Pedersen, *Clin. Chem.*, 25 (1979) 584.
- 8 R. D. Coldwell, D. J. H. Trafford, H. L. J. Makin, M. J. Varley and D. N. Kirk, *Clin. Chem.*, 30 (1984) 1193.