

## Bibliography Section

---

### Liquid Column Chromatography

#### 1. REVIEWS AND BOOKS

- 2546 Beelaram, A. and Sadana, A.: High-resolution fractionation processes: chromatographic techniques. *Isol. Purif.*, 2 (1994) 45-62; C.A., 122 (1995) 54016r - a review with 33 refs.
- 2547 Diez-Masa, J.C., Crego, A., de Frutos, M. and Polo, M.C.: High performance liquid chromatography with microcolumns. *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, 1992, (1993) 27-32; C.A., 122 (1995) 75621z - a review with 18 refs.
- 2548 Hayashi, Y. and Matsuda, R.: Uncertainty structure information theory, and optimization of quantitative analysis in separation science. *Adv. Chromatogr. (N.Y.)*, 34 (1994) 347-427; C.A., 122 (1995) 150209b - a review with 198 refs.
- 2549 Janos, P.: Complex-forming equilibria in ion chromatography. *J. Chromatogr. A*, 699 (1995) 1-10 - a review with 84 refs.
- 2550 Jellum, E.: Chromatography and capillary electrophoresis in biomedicine. *Spec. Publ.-R. Soc. Chem.*, 154 (Reviews on Analytical Chemistry-Euroanalysis VIII) (1994) 308-316; C.A., 122 (1995) 50306n - a review with 14 refs.
- 2551 Kitazume, E.: (Countercurrent chromatography). *Bunseki*, (1994) 711-717; C.A., 122 (1995) 170873s - a review with 99 refs.
- 2552 Scott, R.P.W.: Liquid chromatography phase systems. *Chromatogr. Sci. Ser.*, 67 (1994) 51-92; C.A., 122 (1995) 143233b - a review with 15 refs.
- 2553 Scott, R.P.W.: Resolution retention and selectivity. *Chromatogr. Sci. Ser.*, 67 (1994) 15-49; C.A., 122 (1995) 143232a - a review with 13 refs.
- 2554 Sewell, P.A.: High-performance liquid chromatography. In: Hamilton, R.J. and Hamilton, S. (Editors), *Lipid Anal.*, IRL, Oxford, 1992, pp. 153-203; C.A., 122 (1995) 127737p - a review with 53 refs.
- 2555 Tachibana, K.: (Overview and basic theory of chromatography). *Bunseki*, (1995) 2-11; C.A., 122 (1995) 143283t - an elementary review with 5 refs.

See also 2556, 2578, 2602, 2630, 2638, 2655, 2656, 2662, 2676, 2695, 2707, 2709, 2717, 2734, 2737, 2742, 2744, 2754, 2755, 2783, 2819, 2914, 2934, 2936, 2953, 2992, 3009, 3022, 3080, 3081, 3123, 3129, 3423, 3491, 3518, 3519, 3520, 3533, 3549, 3623, 3631, 3634, 3675, 3764, 3806, 3809, 3820, 3822, 3832, 3849, 3882.

#### 2. FUNDAMENTALS, THEORY AND GENERAL

##### 2a. General

- 2556 Anonymous: (Ninety years of history of chromatography (1903-1993)). *Zh. Anal. Khim.*, 49 (1994) 1047-1058; C.A., 122 (1995) 80297c - a review with 106 refs.
- 2557 Bakken, G.A. and Kalivas, J.H.: Assessing chromatographic peak purity using condition index and singular value evolving profiles. *Anal. Chim. Acta*, 300 (1995) 173-181.
- 2558 Buergisser, C.S., Scheidegger, A.M., Borkovec, M. and Sticher, H.: Chromatographic charge density determination of materials with low surface area. *Langmuir*, 10 (1994) 855-860; C.A., 120 (1994) 174398x.
- 2559 Dolan, J.W.: Case study - irreproducible retention times. *LC-GC*, 12 (1994) 900-904; C.A., 122 (1995) 100747e.
- 2560 Dolan, J.W.: Obtaining separations. Part IV: Putting it all together. *LC-GC*, 12 (1994) 666-668; C.A., 122 (1995) 100740x.
- 2561 Dolan, J.W.: Problems with water. *LC-GC Int.*, 8, No. 3 (1995) 134-136.
- 2562 Dolan, J.W.: Reader mailbag. *LC-GC Int.*, 8, No. 5 (1995) 250-252.
- 2563 Dolan, J.W.: Reader questions. *LC-GC Int.*, 8, No. 2 (1995) 76-82.
- 2564 Edlen, B.B.: Consistent chromatographic performance: fundamentals in media selection. *Am. Biotechnol. Lab.*, 12 (1994) 40, 42; C.A., 122 (1995) 50389s.
- 2565 Gagnon, P., Grund, E. and Lindbaeck, T.: Large-scale process development for hydrophobic interaction chromatography, part I - gel selection and development of binding conditions. *LC-GC*, 13 (1995) 318-328; C.A., 122 (1995) 234545w.
- 2566 Gennaro, M.C., Abrigo, C., Pobozy, E. and Marengo, E.: Retention dependence on organic modifier and interaction reagent concentration in reversed-phase ion-interaction HPLC. *J. Liq. Chromatogr.*, 18 (1995) 311-330.
- 2567 Glód, B.K., Alexander, P.W., Chen, Z.L. and Haddad, P.R.: Thermodynamic investigation of sample retention mechanism in ion-exclusion chromatography with inclusion compound in the mobile phase. *Anal. Chim. Acta*, 306 (1995) 267-272.
- 2568 Grung, B. and Kvalheim, O.M.: Retention time shift adjustments of two-way chromatograms using Bessel's inequality. *Anal. Chim. Acta*, 304 (1995) 57-66.

- 2569 Hanson, M., Unger, K.K., Denoyel, R. and Rouquerol, J.: Interactions of lysozyme with hydrophilic and hydrophobic polymethacrylate stationary phases in reversed phase chromatography (RPC). *J. Biochem. Biophys. Methods*, 29 (1994) 283-293; C.A., 122 (1995) 208266m.
- 2570 Hawkes, S.J.: Distorted chromatographic peaks. *J. Chem. Educ.*, 71 (1994) 1032-1033; C.A., 122 (1995) 55249f.
- 2571 Hill, H.M., Smith, I., Fairbrother, I., Tennant, K. and Robson, J.: Protein and other interference problems, and some solutions. *Methodol. Surv. Bioanal. Drugs*, 23(Biofluid and Tissue Analysis for Drugs, Including Hypolipidaemics) (1994) 327-334; C.A., 122 (1995) 229994n.
- 2572 Kaiser, E. and Wojtusik, M.J.: Improvements in on-line ion chromatography. *Off. Proc.-Int. Water Conf.*, 54th (1993) 266-274; C.A., 122 (1995) 150268v.
- 2573 Kohn, A.: Automating method validation and system suitability testing in HPLC and CE. *Am. Biotechnol. Lab.*, 12 (1994) 44-47; C.A., 122 (1995) 26972f.
- 2574 Kozulic, B.: On the "donor-corridor" model of gel electrophoresis. II. Developments related to new gels capillary gel electrophoresis. *Appl. Theor. Electrophor.*, 4 (1994) 137-148; C.A., 122 (1995) 234664j.
- 2575 Löre, A., Szász, G. and Budvári-Bárány, Z.: Further investigations in HPLC systems containing silica and polar bonded silica packing dynamically modified by camphorsulfonic acid. *J. Liq. Chromatogr.*, 18 (1995) 287-295.
- 2576 McDowall, R.D.: A suitable system for chromatography? *LC-GC Int.*, 8, No. 4 (1995) 196-200.
- 2577 McDowall, R.D.: Change? I've always worked this way. *LC-GC Int.*, 8, No. 2 (1995) 89-91.
- 2578 Mori, S.: (Size exclusion chromatography). *Bunseki*, (1995) 43-50; C.A., 122 (1995) 143284u - a review with 81 refs.
- 2579 Ouchi, G.I.: Statistical techniques for chromatography data handling. *LC-GC*, 12 (1994) 826-830; C.A., 122 (1995) 101065t.
- 2580 Poppe, H.: The role of miniaturization in the development of analytical fractionation methods. *Analisis*, 22 (1994) M22-M24; C.A., 122 (1995) 177132h.
- 2581 Rowe, R.C., Mulley, V.J., Hughes, J.C., Nabney, I.T. and Debenham, R.M.: Neural networks for chromatographic peak classification - a preliminary study. *LC-GC*, 12 (1994) 690-698; C.A., 122 (1995) 100741y.
- 2582 Rózylo, J.K., Janicka, M. and Siembida, R.: Advantages of TLC as a pilot technique for HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3641-3653.
- 2583 Ryan, T.W.: HPLC method transfer to narrow bore columns: an evaluation. *J. Liq. Chromatogr.*, 18 (1995) 51-62.
- 2584 Seaver, C. and Przybytek, J.: Solvent selection, Part II - physical properties. *LC-GC Int.*, 8, No. 4 (1995) 190-195.
- 2585 Seaver, C. and Sadek, P.: Solvent selection. Part I. UV absorption characteristics. *LC-GC*, 12 (1994) 742; C.A., 122 (1995) 177383r.
- 2586 Sharina, I.G., Cherkasov, A.V. and Nedospasov, A.A.: (Quantitative evaluations of the degree of similarity of chromatograms and electrophoretograms). *Zh. Fiz. Khim.*, 68 (1994) 1752-1757; C.A., 122 (1995) 25434v.
- 2587 Shibukawa, M.: Theoretical interpretation of the retention of system peaks in partition chromatography with a mobile phase containing electrolytes. *J. Chromatogr. A*, 696 (1995) 165-172.
- 2588 Tanimura, M. and Tamura, M.: Chromatographic separation of three or more compositions with different adsorbability by cyclic pseudo-moving bed-like systems. *Jpn. Kokai Tokkyo Koho JP 06,170,112 [94,170,112]* (Cl. B01D15/00), 21 Jun. 1994, Appl. 92/329,516, 09 Dec. 1992; 7 p.; C.A., 122 (1995) 137316q.
- 2589 Uhl, S., Rempfer, K., Egelhaaf, H.-J., Lehr, B. and Oelkrug, D.: Fluorescence characterization of acid-base interaction and mobility at chromatographic interfaces. *Anal. Chim. Acta*, 303 (1995) 17-23.
- 2590 Woodbury, C.P., Jr., Fitzloff, J.F. and Vincent, S.S.: Sample multiplexing for greater throughput in HPLC and related methods. *Anal. Chem.*, 67 (1995) 885-890.

For additional information see C.A.:

122 (1995) 27168k, 30628r, 177436k.

See also 2548, 2549, 2555, 2676, 2753, 3075, 3408.

## 2b. Thermodynamics and theoretical relationships

- 2591 Baumeister, E., Klose, U., Albert, K., Bayer, E. and Guiochon, G.: Determination of the apparent transverse and axial dispersion coefficients in a chromatographic column by pulsed field gradient nuclear magnetic resonance. *J. Chromatogr. A*, 694 (1995) 321-331.
- 2592 Blumberg, L.M.: Limits of resolution and speed of analysis in linear chromatography with and without focusing. *Chromatographia*, 39 (1994) 719-728.
- 2593 Borowko, M.: Isoelutotropic diagrams for ternary mobile phases systems in liquid-solid chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 319-324; C.A., 122 (1995) 143536c.
- 2594 Bosch, E., Bou, P. and Rosés, M.: Linear description of solute retention in reversed-phase liquid chromatography by a new mobile phase polarity parameter. *Anal. Chim. Acta*, 299 (1994) 219-229.
- 2595 Cheong, W.J. and Choi, J.D.: Limitations of the linear solvation energy relationships in reversed phase liquid chromatography. *Bull. Korean Chem. Soc.*, 15 (1994) 868-873; C.A., 122 (1995) 90393v.
- 2596 Foti, G. and sz. Kovats, E.: To the relationship of excess adsorption and retention in liquid/solid chromatography. In: Czempesz, F., Horvorigyi, Z. and Paszli, I. (Editors), *Proc.-Conf. Colloid Chem.*, 6th 1992, Hung. Chem. Soc., Budapest, 1993, pp. 96-99; C.A., 122 (1995) 171021z.
- 2597 Furman, W.B., Layloff, T.P. and Tetzlaff, R.F.: Validation of computerized liquid chromatographic systems. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1314-1318.
- 2598 Glod, B.K.: Retention mechanism of ion exclusion chromatography. *Chem. Anal. (Warsaw)*, 39 (1994) 399-422; C.A., 122 (1995) 171043h.
- 2599 Harlan, J.E., Picot, D., Loll, P.J. and Garavito, R.M.: Calibration of size-exclusion chromatography: use of a double Gaussian distribution function to describe pore sizes. *Anal. Biochem.*, 224 (1995) 557-563.
- 2600 Koo, Y.-M. and Hong, J.: Analysis of a nonlinear, coupled chromatographic system. *Korean J. Chem. Eng.*, 11 (1994) 227-231; C.A., 122 (1995) 177075s.
- 2601 Meyer, V.R.: Quantitation of chromatographic peaks in the 0.1 to 1.0% range. *Chromatographia*, 40 (1995) 15-22.

- 2602 Miyabe, K.: Adsorption equilibrium and rate in reverse-phase liquid chromatography separation system. *Adsorpt. News*, 8 (1994) 16-20; C.A., 122 (1995) 143205u - a review with 8 refs.
- 2603 Nicolas-Simonnot, M.-O., Sardin, M. and Grévillet, G.: Consequences of ion-exchange and sorption effects on pH waves propagating in a strong-base anion-exchange column. *J. Liq. Chromatogr.*, 18 (1995) 463-487.
- 2604 Parriott, P.J. and Lai, Y.-H.: Dynamic secondary equilibria in chromatography columns. *Chem. Aust.*, 61 (1994) 386-389; C.A., 122 (1995) 177345e.
- 2605 Patrickios, C.S. and Patrickios, E.S.: Stoichiometric mass-action ion-exchange model. Explicit isotherms for mono-, di-, tri- and tetrameric ions. *J. Chromatogr. A*, 694 (1995) 480-485.
- 2606 Shatirishvili, Sh.I., Zakalashvili, G.N. and Sirbiladze, A.L.: Evaluation of the minimum separation time of complex systems on high-performance chromatographic columns. *Khim. Prom-st (Moscow)*, (1994) 189-191; C.A., 122 (1995) 203934t.
- 2607 Tan, L.C.: Study of retention mechanism in reversed-phase liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9428946, 1994, 305 p.; C.A., 122 (1995) 177131g.
- 2608 Thompson, R.A., Ge, Z., Grinberg, N., Ellison, D. and Tway, P.: Mechanistic aspects of the stereospecific interaction for aminoindanol with a crown ether column. *Anal. Chem.*, 67 (1995) 1580-1587.
- 2609 Zhu, J. and Guiochon, G.: Production rate of an isotactic train in displacement chromatography. *AIChE J.*, 41 (1995) 45-57; C.A., 122 (1995) 84525y.
- See also 2568, 2581, 2648, 2657, 2688, 2695, 2739, 2761, 3495, 3845.
- 2c. *Relationship between structure and chromatographic behaviour*
- 2610 Gerasimenko, V.A. and Nabivach, V.M.: (Sorption-structural correlations in chromatography of aromatic compounds). *Zh. Fiz. Khim.*, 68 (1994) 1740-1748; C.A., 122 (1995) 55540u.
- 2611 Hahai, T., Hatano, H., Nimura, N. and Kinoshita, T.: Molecular recognition in chromatography aided by computational chemistry. *Supramol. Chem.*, 3 (1994) 243-247; C.A., 122 (1995) 171039m.
- 2612 Kumskov, M.I., Ponomareva, L.A., Smolenskii, E.A., Mitushev, D.F. and Zefirov, N.S.: (Automated method for formation of structure descriptors of organic compounds for quantitative structure-properties relations (QSPR)). *Izv. Akad. Nauk, Ser. Khim.*, (1994) 1391-1393; C.A., 122 (1995) 104998k.
- 2613 Takács-Novák, K., Szász, G., Budvári-Bárány, Z., Józán, M. and Löre, A.: Relationship study between reversed phase HPLC retention and octanol/water partition among amphoteric compounds. *J. Liq. Chromatogr.*, 18 (1995) 807-825.
- 2614 Wilce, M.C.J., Aguilar, M.-I. and Hearn, M.T.W.: Physicochemical basis of amino acid hydrophobicity scales: evaluation of four new scales of amino acid hydrophobicity coefficients derived from RP-HPLC of peptides. *Anal. Chem.*, 67 (1995) 1210-1219.
- See also 3493.
- 2d. *Measurement of physico-chemical and related values*
- 2615 Bodzek, M. and Korus, I.: Characterization of ultrafiltration membranes using gel permeation chromatography. *Acta Chromatogr.*, 2 (1993) 40-55; C.A., 122 (1995) 107741g.
- 2616 Miyabe, K. and Suzuki, M.: Chromatographic study on liquid-phase adsorption on octadecylsilyl-silica gel. *AIChE J.*, 41 (1995) 548-558; C.A., 122 (1995) 191286t.
- 2617 Miyakoda, G., Nakayama, N., Matsui, K. and Yabuuchi, Y.: Mechanisms of the potent and long-lasting antihypertensive action of the new calcium channel blocker pranidipine. Evidence for strong affinity to the calcium channels or membranes. *Arzneim.-Forsch.*, 45 (1995) 471-475.
- See also 2614, 2650, 2681, 2735, 2762, 2874, 3057, 3156, 3446, 3447, 3607, 3610, 3611, 3612, 3620, 3842.
3. GENERAL TECHNIQUES
- 3a. *Apparatus and accessories*
- 2618 Goosens, E.C., de Jong, D., de Jong, G.J. and Brinkman, U.A.T.: Reversed-phase liquid chromatography coupled online with capillary gas chromatography II. Use of a solvent vapor exit to increase introduction volumes and introduction rates into the gas chromatograph. *J. Microcolumn Sep.*, 6 (1994) 207-215. C.A., 122 (1995) 95569s.
- 2619 Karlsson, K. and Hellstroem, K.-G.: Method and apparatus for liquid chromatography. *PCT Int. Appl. WO 94 25,134 (Cl. B01D15/08)*, 10 Nov. 1994, SE Appl. 93/1,395, 26 Apr. 1993; 17 p.; C.A., 122 (1995) 109480v.
- 2620 Rothchild, R.D.: Method and apparatus for moving-bed or simulated moving-bed continuous chromatographic separation. *U.S. US 5,387,347 (Cl. 210-659; B01D15/08)*, 07 Feb. 1995, Appl. 120,777, 15 Sep. 1993; 7 p.; C.A., 122 (1995) 191538b.
- 2621 Stanin, V.V., Petrov, D.B., Isaev, V.B. and Ilyushin, A.A.: Thermostat for a chromatograph. *U.S.S.R. SU 1,807,385 (Cl. G01N30/30)*, 07 Apr. 1993, Appl. 4,933,992, 05 May 1991; C.A., 122 (1995) 95640h.
- 2622 Wenk, M. and Haefeli, W.E.: A user-built automatic solid phase extraction device for parallel sample preparation prior to HPLC or RIA. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A18.
- For additional information see C.A.:  
122 (1995) 155722y, 164202f, 164232r, 177425f.
- See also 3090.
- 3b. *Detectors and detection reagents*
- 2623 Crego, A.L., Diez-Masa, J.C. and Dabrio, M.V.: (Criteria for the modification of a UV-VIS HPLC detector to permit detection in open capillary columns). *Quim. Anal. (Barcelona)*, 12 (1993) 83-87; C.A., 122 (1995) 150524a.
- 2624 Fallick, G. and Romano, J.: Enhanced photodiode array detection: HPLC applications in environmental research, development, and quality assurance using a spectral contrasting technique. *Int. Lab.*, 25, No. 3 (1995) 12C-12H.

- 2625 Glod, B.K., Alexander, P.W., Haddad, P.R. and Chen, Z.L.: Potentiometric detection using a metallic copper electrode in reversed-phase and ion-exclusion chromatography with eluents containing ion-interaction reagents. *J. Chromatogr. A*, 699 (1995) 31-37.
- 2626 Gorenstein, M.V., Li, J.B., Antwerp, J.V. and Chapman, D.: Detecting coeluted impurities by spectral comparison. *LC-GC*, 12 (1994) 768-772; *C.A.*, 122 (1995) 100744b.
- 2627 Jane, Y.S. and Shih, J.S.: Application of crown ether coated piezoelectric crystal as a detector for ion chromatography. *Analyst (Cambridge)*, 120 (1995) 517-522.
- 2628 Kim, J.-S. and Raines, R.T.: Dibromobimane as a fluorescent crosslinking reagent. *Anal. Biochem.*, 225 (1995) 174-176.
- 2629 Krull, I.S., Mhatre, R. and Cunniff, J.: Biopolymer detection, Part II - low-angle laser light scattering. *LC-GC Int.*, 8, No. 2 (1995) 83-88.
- 2630 Laubscher, J.C.: UV-VIS detection in HPLC. *Schweiz. Lab.-Z.*, 51 (1994) 239-243; *C.A.*, 122 (1995) 122195a - a review with 13 refs.
- 2631 Mank, A.J.G., Velthorst, N.H., Brinkman, U.A.T. and Gooijer, C.: Near-infrared laser-induced fluorescence detection in column liquid chromatography. A comparison of various lasers and detection systems. I. Continuous wave lasers. *J. Chromatogr. A*, 695 (1995) 165-174.
- 2632 Mank, A.J.G., Velthorst, N.H., Brinkman, U.A.T. and Gooijer, C.: Near-infrared laser-induced fluorescence detection in column liquid chromatography. A comparison of various lasers and detection systems. II. Pulsed lasers. *J. Chromatogr. A*, 695 (1995) 175-183.
- 2633 Roepke, W.: (Measurement in the femtomol range. The chemiluminescence detector in HPLC). *LaborPraxis*, 18 (1994) 52-55; *C.A.*, 122 (1995) 203943v.
- 2634 Smalley, M.B. and McGown, L.B.: Limits of detection and resolution for on-the-fly fluorescence lifetime detection in HPLC. *Anal. Chem.*, 67 (1995) 1371-1376.
- 2635 Takeuchi, T. and Miwa, T.: Effect of sodium dodecyl sulfate as stationary phase on signal intensities of dansylamino acids in microcolumn liquid chromatography with on-column fluorimetric detection. *J. Chromatogr. A*, 696 (1995) 185-192.
- 2636 Toyooka, T. and Liu, Y.-M.: Development of optically active fluorescent Edman-type reagents. *Analyst (Cambridge)*, 120 (1995) 385-390.
- 2637 Tran, C.D., Huang, G. and Grishko, V.I.: Direct and indirect detection of liquid chromatography by infrared thermal lens spectrometry. *Anal. Chim. Acta*, 299 (1995) 361-369.
- 2638 Wirth, M.J.: Spectroscopic probing of reversed-phase chromatographic retention. *LC-GC*, 12 (1994) 656-664; *C.A.*, 122 (1995) 100689n - a review with 39 refs.
- 2639 Wodecki, Z., Slebioda, M. and Kolodziejczyk, A.M.: Application of polarimetric detector for the high-performance liquid chromatographic determination of the optical purity of 5(4H)-oxazolones. *J. Chromatogr. A*, 696 (1995) 149-152.
- 2640 Yamamoto, A., Matsunaga, A., Mizukami, E., Hayakawa, K., Miyazaki, M., Nishimura, M., Kitaoka, M. and Fujita, T.: Evaluation of split-type flow cell for a polarized spectrophotometric detector. *Analyst (Cambridge)*, 120 (1995) 1137-1139.
- 2641 Yang, L., Janle, E., Huang, T., Gitzen, J., Kissinger, P.T., Vreeke, M. and Heller, A.: Applications of "wired" peroxidase electrodes for peroxide determination in liquid chromatography coupled to oxidase immobilized enzyme reactors. *Anal. Chem.*, 67 (1995) 1326-1331.
- 2642 Zegers, B.N., de Brouwer, J.F.C., Poppema, A., Lingeman, H. and Brinkman, U.A.T.: Electron-capture detection in reversed-phase liquid chromatography using packed-capillary columns. *Anal. Chim. Acta*, 304 (1995) 47-56.
- 2643 Zegers, B.N., Hessels, R., Jagesar, J., Rozenbrand, J., Lingeman, H. and Brinkman, U.A.T.: Photoionization detection in packed-capillary liquid and supercritical-fluid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 413-440.

For additional information see *C.A.*:

122 (1995) 122158r, 177389x, 204235c.

See also 2768, 2820, 2921, 2933, 2966, 2984, 2993, 3007, 3017, 3027, 3034, 3035, 3050, 3052, 3473, 3509, 3541, 3613, 3717, 3737, 3745, 3773, 3872, 3895, 3896, 3899, 3902.

### 3c. Sorbents and columns, packing procedures

- 2644 Afeyan, N.B., Hodjat, H.K. and Varady, L.: Sulphonamide-bonded hydrophilic coatings for preparing chromatography matrix. *U.S. US 5,389,449* (Cl. 428-523; 801D15/08), 14 Feb. 1995, US Appl. 637,482, 04 Jan. 1991; 11 p.; *C.A.*, 122 (1995) 191540w.
- 2645 Alvarez, C., Bertorello, H. and Strumia, M.: Preparation of new synthetic hydrogels containing galactose, applicable in affinity chromatography. *J. Appl. Polym. Sci.*, 54 (1994) 1887-1891; *C.A.*, 122 (1995) 50602f.
- 2646 Anonymous: Chromatographic hydrations of sulphonated cation exchange resins. *Res. Discl.*, 361 (1994)36129; *C.A.*, 122 (1995) 135179y.
- 2647 Anspach, F.B. and Altmann-Haase, G.: Immobilized-metal-chelate regenerable carriers: (II) stability of copper chelate:concanavalin A sandwich affinity-chromatographic sorbents. *Biotechnol. Appl. Biochem.*, 20 (1994) 323-333; *C.A.*, 122 (1995) 50411t.
- 2648 Arenas, R.V. and Foley, J.P.: Polymer coat4d alumina columns in reversed-phase liquid chromatography. *Am. Lab. (Shelton)*, 26 (1994) 32CC, 32DD; *C.A.*, 122 (1995) 150540c.
- 2649 Benmoussa, A., Mikou, M., Lacout, J.L. and Siouffi, A.M.: Lead phosphate hydroxyapatite high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 486-491.
- 2650 Biffis, A., Corain, B., Zecca, M., Corvaja, C. and Jerabek, K.: On the macromolecular structure and molecular accessibility of swollen microporous resins: a combined ESR-ISEC approach. *J. Am. Chem. Soc.*, 117 (1995) 1603-1606; *C.A.*, 122 (1995) 82759d.
- 2651 Bluemel, J.: Reaction of ethoxysilanes with silica: a solid state NMR study. *J. Am. Chem. Soc.*, 117 (1995) 2112-2113; *C.A.*, 122 (1995) 133271y.
- 2652 Buszewski, B., Gilpin, R.K. and Jaroniec, M.: Comparative studies of thermal properties of alkylamide and alkyl bonded phases. *Chem. Anal. (Warsaw)*, 39 (1994) 673-679; *C.A.*, 122 (1995) 80670u.

- 2653 Da, S.-L., Yue, W.-G., Wen, Y.-F., Da, H.-L. and Wang, Z.-H.: Preparation and characterization of bonded stationary phases of nitrogen-containing crown ether for high-performance liquid chromatography. *Anal. Chim. Acta*, 299 (1994) 239-247.
- 2654 Favrou, A. and Roussel, C.: Use of  $\gamma$ -cyclodextrin in chiral chromatography. In: Hedges, A.R. (Editor), *Minutes Int. Symp. Cyclodextrins, 6th*, Ed. Sante, Paris, 1992, pp. 603-606; *C.A.*, 122 (1995) 143530w.
- 2655 Francisco Lorenzano Porras, C., Carr, P.W. and McCormick, A.V.: Pore characterization of SiO<sub>2</sub> chromatographic gels using nuclear magnetic resonance. In: Adair, J.H., Casey, J.A. and Venigalla, S. (Editors), *Handb. Charact. Tech. Solid-Solution Interface*, Am. Ceram. Soc., Westerville, 1993, pp. 273-281; *C.A.*, 122 (1995) 197686v - a review with 22 refs.
- 2656 Giddings, J., Ratanathanawongs, S.K., Barman, B.N., Moon, M.H., Liu, G., Tjelta, B.L. and Hansen, M.E.: Characterization of colloidal and particulate silica by field-flow fractionation. *Adv. Chem. Ser.*, 234 (1994) 309-340; *C.A.*, 122 (1995) 197692u - a review with 23 refs.
- 2657 Göhlin, K., Buschke, A. and Larsson, M.: Kinetic performance of open-tubular and packed columns in LC using the same stationary phase: immobilized polymethyloctadecylsiloxane. *Chromatographia*, 39 (1994) 729-739.
- 2658 He, B., Yan, H., Yu, K., Cheng, X. and Ni, A.: Synthesis of chiral ligand exchange resin. *Chin. Sci. Bull.*, 38 (1993) 1451-1453; *C.A.*, 122 (1995) 134561e.
- 2659 Honda, F., Honda, H. and Koishi, M.: Utilization of the dry impact blending method to prepare irregularly shaped particles for high-performance liquid chromatographic column packings. *J. Chromatogr. A*, 696 (1995) 19-30.
- 2660 Kaminski, M.: Reasons for non-uniformity of bed structure in wet packed preparative liquid chromatographic columns. *Isol. Purif.*, 2 (1994) 1-17; *C.A.*, 122 (1995) 30712p.
- 2661 Kitagawa, S. and Tsuda, T.: New packing method for liquid chromatographic columns by the aid of electrostatic field (II): application for packing of silica gels in short period. *Kuromatogurafi*, 15 (1994) 137-140; *C.A.*, 122 (1995) 171044j.
- 2662 Levison, P.R.: Cellulosics as ion-exchange materials. In: Kennedy, J.F., Phillips, G.O. and Williams, P.A. (Editors), *Cellulosics*, Horwood, Chichester, 1993, pp. 25-36; *C.A.*, 122 (1995) 131038d - a review with 25 refs.
- 2663 Lodkowski, R., Kruszynska, E., Supryniewicz, Z. and Buszewski, B.: Narrow bore HPLC column with chemically bonded phase on porous glass. *Chem. Anal. (Warsaw)*, 39 (1994) 543-554; *C.A.*, 122 (1995) 171061n.
- 2664 Majors, R.E.: New chromatography columns and accessories at the 1995 Pittsburgh Conference, Part I. *LC-GC Int.*, 8, No. 5 (1995) 266-283.
- 2665 Masłowska, J. and Gasinska, E.: (Chelexes as a polymeric column packing in the chromatographic separation of amino acids). *Zesz. Nauk.-Politech. Lodz., Technol. Chem. Spozyw.*, 589 (1993) 41-55; *C.A.*, 122 (1995) 150502s.
- 2666 Miyabe, K. and Suzuki, M.: Adsorption characteristics of reversed-phase liquid chromatography with various alkyl bonded phases. *J. Chem. Eng. Jpn.*, 27 (1994) 785-789; *C.A.*, 122 (1995) 115973z.
- 2667 Myers, T.A., Healy, M.G., Murphy, W.R. and Al-Duri, B.: The assessment of chitosan beads as support materials for use in dye-ligand affinity chromatography. In: *ICHEME Res. Event, Two-Day Symp.*, 1, Inst. Chem. Eng., Rugby, 1994, pp. 210-213; *C.A.*, 122 (1995) 26974h.
- 2668 Nagae, N.: (Packed column for reversed-phase liquid chromatography: "Develosil ODS-UG"). *Kuromatogurafi*, 15 (1994) 175-179; *C.A.*, 122 (1995) 171045k.
- 2669 Noel, R., Sanderson, A. and Spark, L.: A monolithic ion-exchange material suitable for downstream processing of bio-products. In: Kennedy, J.F., Phillips, G.O. and Williams, P.A. (Editors), *Cellulosics*, Horwood, Chichester, 1993, pp. 17-24; *C.A.*, 122 (1995) 131047f.
- 2670 Ogino, K., Sato, H., Aihara, Y., Suzuki, H. and Moriguchi, S.: Preparation and characterization of monodisperse oligo(ethylene glycol) dimethacrylate polymer beads for aqueous high-performance liquid chromatography. *J. Chromatogr. A*, 699 (1995) 67-72.
- 2671 Ogino, K., Sato, H., Tsuchiya, K., Suzuki, H. and Moriguchi, S.: Synthesis of monodisperse macroreticular styrene-divinylbenzene gel particles by a single-step swelling and polymerization method. *J. Chromatogr. A*, 699 (1995) 59-66.
- 2672 Pesek, J.J. and Tang, V.H.: The modification of alumina, zirconia, thoria and titania for potential use as HPLC stationary phases through a silanization/hydrosilation reaction scheme. *Chromatographia*, 39 (1994) 649-654.
- 2673 Pesek, J.J., Matyska, M.T., Soczewinski, E. and Christensen, P.: Spectroscopic studies of butylphenyl, mono-ol and perfluorinated bonded phases. *Chromatographia*, 39 (1994) 520-528.
- 2674 Schmid, J., Albert, K. and Bayer, E.: Synthesis and characterization of unsaturated bonded phases for high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 333-345.
- 2675 Schurig, F.V., Schleimer, M. and Mayer, S.: Preparation and of chiral, cyclodextrin-containing polymers immobilized on supports. *Ger. Offen DE 4,324,636 (Cl. C08B37/16)*, 11 May 1994, Appl. 22 Jul. 1993; 5 p.; *C.A.*, 122 (1995) 84129x.
- 2676 Scott, R.P.W.: The liquid chromatography column. *Chromatogr. Sci. Ser.*, 67 (1994) 93-122; *C.A.*, 122 (1995) 39513g - a review with 14 refs.
- 2677 Shalliker, R.A., Kavanagh, P.E. and Russell, I.M.: Reversed phase behaviour of high molecular mass polystyrenes in ethyl acetate solvent mixture. *Chromatographia*, 39 (1994) 663-667.
- 2678 Terabe, S., Ozaki, H., Tanaka, Y. and Okada, Y.: (Pseudo-stationary phases for electrokinetic chromatography). *Kuromatogurafi*, 15 (1994) 114-115; *C.A.*, 122 (1995) 155375n.
- 2679 Tong, D., Bartle, K.D. and Clifford, A.A.: Preparation and evaluation of supercritical carbon dioxide-packed capillary columns for HPLC and SFC. *J. Microcolumn Sep.*, 6 (1994) 249-255; *C.A.*, 122 (1995) 95583s.
- 2680 Wang, Q.: Desing preparation and characterization of novel polymeric separation media for high-performance liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9501304, 1994, 172 p.; *C.A.*, 122 (1995) 162792z.
- 2681 Yamamoto, F. and Rokushika, S.: (Characteristics of stationary phase on fluorocarbon bonded silica column). *Kuromatogurafi*, 15 (1994) 100-101; *C.A.*, 122 (1995) 171103c.
- 2682 Zusman, R. and Zusman, I.: Gel fiberglass as a new support for affinity chromatography. *Biotechnol. Appl. Biochem.*, 21 (1995) 161-172; *C.A.*, 122 (1995) 234537v.

For additional information see C.A.:

122 (1995) 50735b, 76031u, 82648s, 83166v, 90358n,  
101130k, 122270w, 134188p, 161725t, 177435j,  
177437m.

See also 2591, 2615, 2635, 2699, 2708, 2711, 2712, 2726,  
2728, 2730, 2731, 2733, 2738, 2739, 2740, 2742,  
2743, 2749, 2750, 2751, 2771, 2830, 2981, 3033,  
3112, 3116, 3120, 3124, 3126, 3144, 3158, 3193,  
3215, 3234, 3322, 3614, 3715, 3776, 3873.

### 3d. Quantitative analysis

2683 Meyer, V.R.: Errors in the area determination of incompletely resolved chromatographic peaks. *J. Chromatogr. Sci.*, 33 (1995) 26-33.

For additional information see C.A.:

122 (1995) 177349j.

See also 2548, 2779, 2781, 2805, 2809, 2980, 3624, 3667,  
3752.

### 3e. Preparative scale chromatography

2684 Tharakan, J.P. and Belizaire, M.: Protein band dispersion in axial and radial flow chromatography. *J. Liq. Chromatogr.*, 18 (1995) 39-49.

See also 2620, 2746, 2842, 2928, 2981, 2983, 3026, 3173,  
3193, 3481, 3499, 3563.

### 3f. Programmed temperature, pressure, vapors, gradients

2685 Treiber, L.R.: Normal-phase high-performance liquid chromatography with relay gradient elution. I. Description of the method. *J. Chromatogr. A*, 696 (1995) 193-199.

See also 2698, 3606.

## 4. SPECIAL TECHNIQUES

### 4a. Automation

2686 Olivè, J., Mánuel, A., Rodríguez, P. and Grimalt, J.O.: A low cost, open and flexible on-line data acquisition system for instrumental chromatography. *Anal. Chim. Acta*, 306 (1995) 285-295.

2687 Pullen, F.S. and Richards, D.S.: Automated liquid chromatography/mass spectrometry for chromatographers. *Rapid Commun. Mass Spectrom.*, 9 (1995) 188-190; C.A., 122 (1995) 142700q.

For additional information see C.A.:

122 (1995) 204188q, 234488e.

See also 2573, 3017, 3078, 3463, 3579, 3587, 3658, 3821.

### 4b. Computerization and modelling

2688 Carta, G.: Linear driving force approximation for intraparticle diffusion and convection in permeable supports. *Chem. Eng. Sci.*, 50 (1995) 887-889; C.A., 122 (1995) 191295v.

2689 De Aguiar, P.F., Burquignon, B., Khots, M.S., Penninckx, W. and Massart, D.L.: The use of logistic transformation in HPLC optimization. *Quim. Anal. (Barcelona)*, 12 (1993) 177-183; C.A., 122 (1995) 150546j.

2690 Galushko, S.V., Kamenchuk, A.A. and Pit, G.L.: Software for method development in reversed-phase liquid chromatography. *Int. Lab.*, 25, No. 2 (1995) 14-17.

2691 Hanai, T., Hatano, H., Nimura, N. and Kinoshita, T.: Computational chemical analysis of the chiral recognition of binuclear copper(II) of N-salicylidene (R)-2-amino-1,2-bis(2-butoxy-5-tert-butylphenyl)-3-phenyl-1-propanol in liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4327-4334.

2692 Kevra, S.A., Bergman, D.L. and Maloy, J.T.: A computational introduction to chromatographic bandshape analysis. *J. Chem. Educ.*, 71 (1994) 1023-1028; C.A., 122 (1995) 55248e.

2693 Levy, D. and Miller, C.: Implementing client/server computing in the chromatography laboratory. *Int. Lab.*, 25, No. 2 (1995) 29-31.

2694 Mew, C.D.: A three dimensional surface viewer for Microsoft Windows. *Lab. Microcomput.*, 13 (1994) 17-24; C.A., 122 (1995) 204182h.

2695 Nabivach, V.M.: (Topological models and chromatographic retention of aromatic and heterocyclic compounds. Review. 1. Characteristics of design calculations and comparative characteristics of topological indexes). *Koks Khim.*, (1994) 16-22; C.A., 122 (1995) 150509z - a review with 32 refs.

2696 Naef, O.: (Analysis of a chromatogram). *Chimia*, 48 (1994) 409-412; C.A., 122 (1995) 204169j.

2697 Popov, A.A., Sergeev, S.K. and Prokhorova, E.K.: (Computerized chromatographic system for air quality monitoring). *Ekol. i Bezopas. Truda v Prom.-sti: Mater. Semin. Tsent. Ros. Dom Znanii, M.*, (1991) 43-47; C.A., 120 (1994) 306103j.

2698 Soczewinski, E. and Markowski, W.: Application of stepwise gradient elution chromatography to preliminary separation in short columns. *Chem. Anal. (Warsaw)*, 39 (1994) 467-474; C.A., 122 (1995) 177127k.

2699 Van Straten, M.A., Lynch, R.J. and Claessens, H.A.: The role of stationary phases in reversed-phase liquid chromatography in the application of solvent optimization procedures. *LC-GC Int.*, 8, No. 3 (1995) 143-150.

2700 Villioth, O.: (Processing of chromatographic data under Windows NT). *LaborPraxis*, 18 (1994) 22-27; C.A., 122 (1995) 204171d.

See also 2597, 2598, 2611, 2686, 2718, 2752, 2754, 2761,  
3062, 3122.

### 4c. Combination with other physico-chemical techniques (MS, IR etc.)

2701 Debets, A.J.J., Mekes, T.J.L., Ritburg, A. and Jacobs, P.L.: Column liquid chromatography coupled on-line with mass spectrometry using post-column phosphate suppression. *J. High Resolut. Chromatogr.*, 18 (1995) 45-48.

- 2702 Eaton, A.N. and Rao, R.: Mass detectors: analyte-specific detection for HPLC. *Int. Lab.*, 25, No. 2 (1995) 18-21.
- 2703 Gagne, J.P.: Systematic approach to the development and optimization of mass spectrometry-liquid chromatography systems for the analysis of organic substances. Avail. NLC Order No. DANN88921, 1993, 268 p.; C.A., 122 (1995) 177404y.
- 2704 Joos, P.E.: Electrospray and APCL-mass spectrometry. *LC-GC Int.*, 8, No. 2 (1995) 92-95.
- 2705 Kaltashov, I.A., Yu, X. and Fenselau, C.: Simple interface for microbore LC and electrospray ionization mass spectrometry and analysis of melphalan-alkylation sites in metallothionein. *J. Pharm. Biomed. Anal.*, 13 (1995) 279-284.
- 2706 Manne, R.: On the resolution problem in hyphenated chromatography. *Chemom. Intell. Lab. Syst.*, 27 (1995) 89-94; C.A., 122 (1995) 150308h.
- 2707 Murakoshi, R.: (Recent development of liquid chromatography (LC)-mass spectrometry (MS)). *Idemitsu Giho*, 37 (1994) 732-736; C.A., 122 (1995) 150511u - a review with 9 refs.
- See also 2687, 2776, 2876, 2925, 2936, 2946, 2952, 2960, 2962, 2968, 2976, 2977, 2985, 3002, 3003, 3092, 3108, 3162, 3243, 3341, 3372, 3461, 3467, 3487, 3488, 3489, 3490, 3492, 3495, 3500, 3538, 3570, 3576, 3581, 3582, 3583, 3587, 3603, 3615, 3641, 3646, 3647, 3701, 3704, 3714, 3723, 3746, 3758, 3779, 3786, 3792, 3803, 3824, 3827, 3837, 3840, 3850, 3852, 3855, 3858, 3859, 3864, 3871, 3883, 3884.
- 4d. *Affinity chromatography (advances)*
- 2708 Brena, B.M., Ryden, L.G. and Porath, J.: Immobilization of  $\beta$ -galactosidase on metal-chelate-substituted gels. *Biotechnol. Appl. Biochem.*, 19 (1994) 217-231; C.A., 122 (1995) 208657q.
- 2709 De Frutos, M.: Chromatography-immunology coupling, a powerful tool for environmental analysis. *TrAC*, 14 (1995) 133-140 - a review with 49 refs.
- 2710 Galaev, I.Yu. and Mattiasson, B.: Shielding affinity chromatography. *BioTechnology*, 12 (1994) 1086; C.A., 122 (1995) 26983k.
- 2711 Glatz, Z., Psotova, J., Janiczek, O. and Novotny, M.V.: Immobilized lectins as a means of reversible immobilization of ligands for affinity chromatography. *Int. J. Bio-Chromatogr.*, 1 (1994) 143-149; C.A., 122 (1995) 75706f.
- 2712 Guo, W., Shang, Z.H., Yu, Y.N. and Zhou, L.M.: Cellulose membrane used as stationary phase of membrane affinity chromatography. *Chin. Chem. Lett.*, 5 (1994) 869-872; C.A., 122 (1995) 75691x.
- 2713 Heywood, S.P. and Hooper, N.M.: Purification of GPI anchors by immunoaffinity chromatography. *Biochem. Soc. Trans.*, 23 (1995) 1015; C.A., 122 (1995) 155591e.
- 2714 Jennissen, H.P. and Botzet, G.: The binding of phosphorylase kinase to immobilized calmodulin. *J. Mol. Recognit.*, 6 (1993) 117-130; C.A., 122 (1995) 26386t.
- 2715 Kasai, K.-L., Sakurai, R. and Ishii, S.-I.: Frontal affinity chromatography using microcolumns as investigative tools for molecular recognition. *Int. J. Bio-Chromatogr.*, 1 (1994) 83-91; C.A., 122 (1995) 50420v.
- 2716 Pang, I.-H., Smrcka, A.V. and Sternweis, P.C.: Synthesis and applications of affinity matrix containing immobilized  $\beta$  subunits of G proteins. *Methods Enzymol.*, 237 (Heterotrimeric G Proteins) (1994) 164-174; C.A., 122 (1995) 26970d.
- 2717 Pepper, D.S.: Some alternative coupling chemistries for affinity chromatography. *Mol. Biotechnol.*, 2 (1994) 157-178; C.A., 122 (1995) 26909r - a review with 11 refs.
- 2718 Sridhar, P. and Sastri, N.V.S.: Modeling of bioseparations in an affinity percolation column packed with porous beads. *Indian Chem. Eng., Sect. A*, 36 (1994) 72-78; C.A., 122 (1995) 50630p.
- 2719 Yoshihara, Y. and Kuroda, M.: Use of an SDS-gel-separated protein band as a ligand for affinity chromatography: procedure and application to the purification of domain-specific antibodies against  $\alpha$ -actinin. *J. Biochem. (Tokyo)*, 117 (1995) 443-446.
- For additional information see C.A.:  
122 (1995) 27255m, 155421z, 212170t.
- See also 2645, 2647, 2667, 2682, 2848, 2959, 3029, 3061, 3153, 3154, 3158, 3164, 3169, 3175, 3234, 3257, 3334, 3349, 3362, 3397, 3846, 3847, 3848, 3851.
- 4e. *Functional analysis*
- See 3475, 3609.
- 4f. *Trace analysis and pre-separation techniques*
- 2720 Buttler, T., Jaroskog, H., Gorton, L., Marko-Varga, G. and Ramnemark, L.: The use of microdialysis for sampling in column liquid chromatography. *Curr. Top. Membr.*, 41 (Cell Biology and Membrane Transport Processes) (1994) 45-69; C.A., 122 (1995) 127783a.
- 2721 Gy, P.M.: Sampling - diving into the unknown. *LC-GC Int.*, 8, No. 2 (1995) 70-74.
- 2722 Majors, R.E.: New approaches to sample preparation. *LC-GC Int.*, 8, No. 3 (1995) 128-133.
- 2723 Mguyen, V.S. and Nguyen, M.S.: Neutron activation analysis for determination of trace impurities of light rare earth elements in high purity  $Y_2O_3$ . *Tap. Chi Hoa Hoc*, 32 (1994) 72-73; C.A., 122 (1995) 122009t.
- See also 2792, 2797, 3451, 3576, 3632, 3823, 3825, 3826, 3873, 3882, 3884.
- 4g. *Enantiomers, separation*
- 2724 Bielejewska, A., Kozbial, M., Nowakowski, R., Duszczyk, K. and Sybilska, D.: Studies on the behaviour of  $\alpha$ -,  $\beta$ - and  $\gamma$ -cyclodextrins and some derivatives under reversed-phase liquid chromatographic. *Anal. Chim. Acta*, 300 (1995) 201-206.
- 2725 Buntun, B.M. and Walker, T.A.: Chiral separation of MDL 73,005EF enantiomers using an  $\alpha_1$ -acid glycoprotein column. *J. Chromatogr. A*, 699 (1995) 389-394.
- 2726 Chankvetadze, B., Yashima, E. and Okamoto, Y.: Dimethyl-, dichloro- and chloromethylphenylcarbamates of amylose as chiral stationary phases for high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 101-109.

- 2727 Cleveland, T.: Pirkle-concept chiral stationary phases for the HPLC separation of pharmaceutical racemates. *J. Liq. Chromatogr.*, 18 (1995) 649-671.
- 2728 Cortes, H.J. and Nicholson, L.W.: Chiral separations using packed capillary liquid chromatography. *J. Microcolumn Sep.*, 6 (1994) 257-263; *C.A.*, 122 (1995) 122211c.
- 2729 Gasparrini, F., Lunazzi, L., Misiti, D. and Villani, C.: Organic stereochemistry and conformational analysis from enantioselective chromatography and dynamic nuclear magnetic resonance measurements. *Acc. Chem. Res.*, 28 (1995) 163-170; *C.A.*, 122 (1995) 213225b.
- 2730 Haginaka, J. and Kanasugi, N.: Enantioselectivity of bovine serum albumin-bonded columns produced with isolated protein fragments. *J. Chromatogr. A*, 694 (1995) 71-80.
- 2731 Hyun, M.H., Cho, Y.J., Ryoo, J.-J., Jyung, K.K. and Heo, G.S.: Preparation and application of an (S)-naproxen chiral stationary phase. *J. Chromatogr. A*, 696 (1995) 173-183.
- 2732 Hyun, M.H., Min, C.S. and Cho, Y.J.: Examples of liquid chromatographic resolution of  $\pi$ -acidic racemates on a  $\pi$ -acidic chiral stationary phase. *J. High Resolut. Chromatogr.*, 18 (1995) 63-65.
- 2733 Ihara, T., Sugimoto, Y., Asada, M., Nakagama, T. and Hobo, T.: Influence of the method of preparation of chiral stationary phases on enantiomer separations in high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 49-56.
- 2734 Kempe, M. and Mosbach, K.: Molecular imprinting used for chiral separations. *J. Chromatogr. A*, 694 (1995) 3-13 - a review with 88 refs.
- 2735 Levin, S. and Abu-Lafi, S.: Experimental studies of competition between enantiomers in chiral liquid chromatography through their system peak. *Chirality*, 6 (1994) 148-155; *C.A.*, 122 (1995) 150539j.
- 2736 Li, S.: Liquid chromatographic separation of enantiomers and structurally-related compounds on  $\beta$ -cyclodextrin stationary phases. Avail. *NCL Order No. DANN74572*, 1992, 205 pp.; *C.A.*, 120 (1994) 280434r.
- 2737 Lipkowitz, K.B.: Theoretical studies of type II-V chiral stationary phases. *J. Chromatogr. A*, 694 (1995) 14-37 - a review with 56 refs.
- 2738 Maier, N.M., Uray, G., Kleidernigg, O.P. and Lindner, W.: Diphenylethanediamine (DPEDA) derivatives as chiral selectors: IV. A comparison of 3,5-dinitrobenzoylated with Pirkle's standard(R)-phenylglycine-derived phase in normal phase HPLC. *Chirality*, 6 (1994) 116-128; *C.A.*, 122 (1995) 95579v.
- 2739 Matlin, S.A., Tiritan, E.M., Crawford, A.J., Cass, Q.B. and Boyd, D.: HPLC with carbohydrate carbamate chiral phases: influence of chiral phase structure on enantioselectivity. *Chirality*, 6 (1994) 135-140; *C.A.*, 122 (1995) 150538h.
- 2740 Nakamura, K., Fujima, H., Kitagawa, H., Wada, H. and Makino, K.: Preparation and chromatographic characteristics of a chiral-recognizing perphenylated cyclodextrin column. *J. Chromatogr. A*, 694 (1995) 111-118.
- 2741 Oda, Y., Mano, N., Asakawa, N., Yoshida, Y. and Sato, T.: Investigation of retention behavior for racemate drugs on avidin- and modified avidin-column. *J. Liq. Chromatogr.*, 17 (1994) 3393-3409.
- 2742 Oguni, K., Oda, H. and Ichida, A.: Development of chiral stationary phases consisting of polysaccharide derivatives. *J. Chromatogr. A*, 694 (1995) 91-100 - a review with 18 refs.
- 2743 Ôi, N., Kitahara, H. and Aoki, F.: Direct enantiomer separations by high-performance liquid chromatography with chiral urea derivatives as stationary phases. *J. Chromatogr. A*, 694 (1995) 129-134.
- 2744 Perrin, S.R.: Fast liquid chromatography for the analysis of enantiomers. *Prog. Pharm. Biomed. Anal.*, 1 (1994) 91-124; *C.A.*, 122 (1995) 234469z - a review with refs.
- 2745 Rustum, A.M.: Determination of chiral purity of ethyl nipeccate using a Chiralcel-OG column. *J. Chromatogr. A*, 696 (1995) 75-81.
- 2746 Sato, Y., Shinomiya, K. and Kabasawa, Y.: (Aqueous two-phase partitioning method by using rotation locular countercurrent chromatograph. An application to enantiomeric separation). *Nippon Kagaku Kaishi*, (1994) 1067-1071; *C.A.*, 122 (1995) 75692y.
- 2747 Schurig, V., Jung, M., Mayer, S., Fluck, M., Negura, S. and Jakubetz, H.: Unified enantioselective capillary chromatography on a Chiralcel-DEX stationary phase. Advantages of column miniaturization. *J. Chromatogr. A*, 694 (1995) 119-128.
- 2748 Shibukawa, A., Kadohara, M., He, J.-y., Nishimura, M., Naito, S. and Nakagawa, T.: Study of the enantioselective binding between BOF-4272 and serum albumins by means of high-performance frontal analysis. *J. Chromatogr. A*, 694 (1995) 81-89.
- 2749 Stalcup, A.M. and Williams, K.L.: Comparison 1-(1-naphthyl)ethylcarbamate derivatives of a carbohydrate bonded chiral stationary phase. *J. Chromatogr. A*, 695 (1995) 185-193.
- 2750 Veigl, E., Böhs, B., Mandl, A., Krametter, D. and Lindner, W.: Comparison of (S)-N-(3,5-dinitrobenzoyl)tyrosine derivatives as chiral selectors for high-performance liquid chromatographic enantioseparations. *J. Chromatogr. A*, 694 (1995) 135-150.
- 2751 Veigl, E., Böhs, B., Mandl, A., Krametter, D. and Lindner, W.: Evaluation of silica gel-based brush type chiral cation exchangers with (S)-N-(3,5-dinitrobenzoyl)tyrosine as chiral selector: attempt to interpret the discouraging results. *J. Chromatogr. A*, 694 (1995) 151-161.
- 2752 Yashima, E., Yamada, M., Kaida, Y. and Okamoto, Y.: Computational studies on chiral discrimination mechanism of cellulose trisphenylcarbamate. *J. Chromatogr. A*, 694 (1995) 347-354.
- See also 2567, 2608, 2654, 2658, 2664, 2675, 2691, 2765, 2893, 3007, 3032, 3033, 3054, 3087, 3361, 3471, 3475, 3494, 3549, 3558, 3623, 3626, 3633, 3634, 3635, 3636, 3638, 3642, 3661, 3670, 3672, 3673, 3675, 3676, 3686, 3697, 3701, 3703, 3705, 3706, 3715, 3739, 3756, 3770, 3771, 3773, 3837, 3904.

#### 4h. Other special techniques

- 2753 Abbasi, W.A. and Streat, M.: Comments on extraction chromatography. *J. Radioanal. Nucl. Chem.*, 188 (1994) 163-167; *C.A.*, 122 (1995) 95404j.
- 2754 Deckert, P. and Artl, W.: (Simulated countercurrent chromatography). *Chem. - Ing. - Tech.*, 66 (1994) 1334-1340; *C.A.*, 122 (1995) 84473e - a review with 36 refs.
- 2755 Foucault, A.P.: Solvent systems in centrifugal partition chromatography. *Chromatogr. Sci. Ser.*, 68(Centrifugal Partition Chromatography) (1995) 71-97; *C.A.*, 122 (1995) 155313r - a review with 7 refs.



- 2756 Foucault, A.P. (Editor): *Centrifugal Partition Chromatography. Chromatogr. Sci. Ser.*, Vol. 68, Dekker, New York, 1995, 415 pp.; C.A., 122 (1995) 177422c.
- 2757 Wijnhoven, J.E.G.J., Koorn, J.P., Poppe, H. and Kok, W.T.: Hollow-fibre flow field-flow fractionation of polystyrene sulphonates. *J. Chromatogr. A*, 699 (1995) 119-129.
- 2758 Yuan, N., Liu, Z. and Zhu, D.: Chromatograph-type electrophoresis separation method. *Faming Zhuanli Shenqing Gongkai Shuomingshu* CN 1,083,072 (Cl. CO7K3/14), 02 Mar. 1994; Appl. 93,108,629, 16 Jul. 1993; 8 pp.; C.A., 122 (1995) 50769r.
- 2759 Zibas, S.A. and Love, L.J.C.: Solute-micelle interactions in zwitterionic micellar chromatography. *Anal. Chim. Acta*, 299 (1994) 17-27.
- See also 2551, 2565, 2580, 2620, 2712, 2746, 2763, 2859, 2944, 3036, 3110, 3225, 3278, 3505, 3805, 3856, 3869.
5. HYDROCARBONS AND HALOGEN DERIVATIVES
- 5b. *Cyclic hydrocarbons, fullerenes*
- 2760 Belous, E.F., Lanin, S.N. and Nikitin, Yu.S.: Combined application of normal and reversed-phase high-performance liquid chromatography to determining the group composition of aromatic hydrocarbons in petroleum products. *J. Anal. Chem.*, 50 (1995) 93-100; C.A., 122 (1995) 137743b.
- 2761 Cooper, C.L., Whitaker, K.W. and Sepaniak, M.J.: Separation behavior of common fullerenes in cyclodextrin-HPLC based on computationally-derived interaction energies. *J. Liq. Chromatogr.*, 17 (1994) 4311-4325.
- 2762 Davydov, V.Ya., Filatova, G.N., Khrestaleva, N.M. and Roshchina, T.M.: Separation of fullerenes and study of their intermolecular interactions with organic compounds by chromatography. *Proc.-Electrochem. Soc.*, 94-24 (1994) 1588-1594; C.A., 122 (1995) 204204s.
- 2763 Gasper, M.P., Berthod, A., Talabardon, K. and Armstrong, D.W.: An evaluation of the differential partitioning and separation of C<sub>60</sub> and C<sub>70</sub> fullerenes in a biphasic system using centrifugal partition chromatography (CPC). *J. Liq. Chromatogr.*, 18 (1995) 1019-1034.
- 2764 Heaton, D.M., Bartle, K.D., Clifford, A.A., Myers, P. and King, B.W.: Rapid separation of polycyclic aromatic hydrocarbons by packed column supercritical fluid chromatography. *Chromatographia*, 39 (1994) 607-611.
- 2765 Husain, N., Christian, A.Y. and Warner, I.M.: Effect of co-modifiers in cyclodextrin-modified mobile phases on the reversed-phase high-performance liquid chromatographic separation of polyaromatic hydrocarbons. *J. Chromatogr. A*, 699 (1995) 73-83.
- 2766 Lan, W.G., Chee, K.K., Wong, M.K. and Lee, H.K.: Orthogonal array design as a chemometric method for the optimization of analytical procedures. Part 4. Mixed-level design and its application to the high-performance liquid chromatographic determination of polycyclic aromatic hydrocarbon. *Analyst (Cambridge)*, 120 (1995) 281-287.
- 2767 Lintelmann, J., Boos, K.-S. and Kettrup, A.: (Biomonitoring of exposure to PAH using a coupled-column HPLC-method for the determination of 1-hydroxypyrene - part 1). *Bioforum*, 17 (1994) 442-445; C.A., 122 (1995) 232766v.
- 2768 Mäkelä, M. and Pyy, L.: Effect of temperature on retention time reproducibility and on the use of programmable fluorescence detection of fifteen polycyclic aromatic hydrocarbons. *J. Chromatogr. A*, 699 (1995) 49-57.
- 2769 Masterov, V.F., Chudnovski, F.A., Kozyrev, S.V., Zaidenberg, A.S., Rozhkova, N.N., Podbsenova, N.G. and Stefanovich, G.B.: (Microwave absorption in fullerene-containing shungites). *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. C*, 4 (1994) 213-216; C.A., 122 (1995) 121996a.
- 2770 Mueller, W., Broell, R., Weber, E., Daimer, J. and Mueller, R.: Chromatographic separation of fullerenes with a nonpolar arom. solvent as the elutriating agent. *FCT Int. Appl. WO 94 06,715* (Cl. CO1B31/02), 31 Mar. 1994, DE Appl. 4,230,915, 16 Sep. 1992; 14 pp.; C.A., 120 (1994) 248667h.
- 2771 Ohmacht, R., Kele, M. and Matus, Z.: Reversed phase C<sub>18</sub> packings for the separation of polycyclic aromatic hydrocarbons by high performance liquid chromatography. *Chromatographia*, 39 (1994) 668-672.
- 2772 Podosenova, N.G.: (Separation of fullerenes by liquid chromatography). *Zh. Fiz. Chim.*, 68 (1994) 1298-1303; C.A., 122 (1995) 121984v.
- 2773 Reupert, R. and Brausen, G.: (Determination of polycyclic aromatic hydrocarbons in water, sludges, sediments, and soils by high-performance liquid chromatography). *Acta Hydrochim. Hydrobiol.*, 22 (1994) 202-215; C.A., 122 (1995) 122246t.
- 2774 Rozbeh, M. and Hurtubise, R.J.: Reversed-phase chromatographic separation of benzo[a]pyrene metabolites with  $\beta$ -cyclodextrin as a mobile phase modifier. *J. Liq. Chromatogr.*, 18 (1995) 17-37.
- 2775 Takeuchi, T., Hu, R. and Miwa, T.: Retention behavior of aromatic hydrocarbons in reversed-phase liquid chromatography with ion-exchange-induced stationary phases. *Chromatographia*, 39 (1994) 597-601.
- 2776 Thomas, D., Crain, S.M., Sim, P.G. and Benoit, F.M.: Use of immobilized copper ion chromatography and on-line mass spectrometry with atmospheric pressure chemical ionization for the profiling of complex mixtures of polycyclic aromatic compounds. *J. Chromatogr. A*, 695 (1995) 1-9.
- 2777 Whiton, R.S., Witherspoon, C.L. and Buckley, T.J.: Improved high-performance liquid chromatographic method for the determination of polycyclic aromatic hydrocarbon metabolites in human urine. *J. Chromatogr. B*, 665 (1995) 390-394.
- 2778 Wise, S.A., Deissler, A. and Sander, L.C.: Liquid chromatographic detection of polycyclic aromatic hydrocarbon isomers of molecular weight 278 and 302 in environmental standard reference materials. *Polycyclic Aromat. Compd.*, 3 (1993) 169-184; C.A., 122 (1995) 150587y.
- 2779 Wise, S.A., Schantz, M.M., Benner, B.A., Jr., Hays, M.J. and Schiller, S.B.: Certification of polycyclic aromatic hydrocarbons in a marine sediment standard reference material. *Anal. Chem.*, 67 (1995) 1171-1178.
- 2780 Yamamoto, K., Funasaka, H., Takahashi, T. and Akasaka, T.: Isolation of an ESR-active metallofullerene of La@C<sub>82</sub>. *J. Phys. Chem.*, 98 (1994) 2008-2011; C.A., 120 (1994) 207225m.

- 2781 Zobel, H., Nerlich, B. and Panning, H.: Liquid chromatographic procedure and apparatus for separation and qualitative and quantitative determination of polycyclic aromatic hydrocarbon-containing mixtures. *Ger. Offen.* DE 4,312, 925 (Cl. G01N33/22), 10 Oct. 1994, Appl. 15 Apr. 1993; 7 p.; C.A., 122 (1995) 85295k.

See also 2634.

#### 5c. Halogen derivatives

- 2782 Finizio, A., di Guardo, A. and Vighi, M.: Improved RP-HPLC determination of Kow for some chloroaromatic chemicals using molecular connectivity indices. *SAR QSAR Environ. Res.*, 2 (1994) 249-260; C.A., 122 (1995) 98877b.
- 2783 Golba, W., Przybylski, Z., Ceckiewicz, S. and Krasodowski, M.: (Chromatographic methods for determination of polychlorinated biphenyls). *Nafta-Gaz*, 50 (1994) 405-410; C.A., 122 (1995) 204161a a review with 33 refs.
- 2784 Takasuga, T., Ohi, E. and Inoue, T.: Complete isolation and determination of *mono-ortho* and *non-ortho* substituted PCBs and PCDDs/PCDFs by HPLC, on porous graphitic carbon with HRGC/HRMS. *Organohalogen Compd.*, 11 (1993) 101-104; C.A., 122 (1995) 204237e.

See also 3267.

#### 5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

- 2785 Fabinska, M.: (Identification of the group components from extrographic separation of the low-temperature coal tar). *Karbo-Energochem.-Ekol.*, 38 (1993) 55-57; C.A., 122 (1995) 165202t.
- 2786 Whittaker, M. and Pollard, S.J.T.: Characterization of refractory wastes at hydrocarbon-contaminated sites: 1. Rapid column fractionation and thin layer chromatography of reference oils. *J. Planar Chromatogr.*, 7 (1994) 354-361.

See also 3467, 3838.

#### 6. ALCOHOLS

- 2787 Bourdeaux, R., Capolaghi, B., Lacroix, C., Younos, C. and Lehr, P.R.: Assay of urinary free and conjugated 3-methoxy-4-hydroxyphenylethyleneglycol by high-performance liquid chromatography with amperometric detection. *J. Chromatogr. B*, 665 (1995) 45-51.
- 2788 Kok, S.J., Posthumus, R., Bakker, I., Gooijer, C., Brinkman, U.A.T. and Velthorst, N.H.: Identification of stereoisomeric benzo[a]pyrene tetrols by reversed-phase liquid chromatography coupled semi-on-line to fluorescence line-narrowing spectroscopy. *Anal. Chim. Acta*, 303 (1995) 3-10.
- 2789 Tjioe, S.W. and Hurtubise, R.J.: Separation and characterization of tetrol metabolites of benzo[a]pyrene-DNA adducts using HPLC and solid-matrix room temperature luminescence. *Talanta*, 42 (1995) 59-64; C.A., 122 (1995) 125436f.

- 2790 Van Damme, F. and Oomens, A.C.: Determination of residual free epoxide in polyether polyols by derivatization with diethylammonium N,N-diethylthiocarbamate and liquid chromatography. *J. Chromatogr. A*, 696 (1995) 41-47.

- 2791 Wanaka, K. and Murui, T.: (Determination of fluorescence labeled alcohols from waxes in vegetable oils by high performance liquid chromatography). *Yukagaku*, 44 (1995) 43-48; C.A., 122 (1995) 104198z.

For additional information see C.A.:  
122 (1995) 105252z.

See also 3784, 3830.

#### 7. PHENOLS

- 2792 Alymova, A.T., Korenman, Ya.I., Bobrinskaya, E.V., Khizver, D.R. and Li, A.: (Separation and preconcentration of traces of aromatic hydroxy compounds by extraction and chromatography with mixed stationary phases). *Zh. Anal. Khim.*, 49 (1994) 1197-1201; C.A., 122 (1995) 204215w.
- 2793 Amarowicz, R., Wanasundara, P.K.J.P.D. and Shahidi, F.: Chromatographic separation of flaxseed phenolics. *Nahrung*, 38 (1994) 520-526; C.A., 122 (1995) 104164k.
- 2794 Galceran, M.T. and Jáuregui, O.: Determination of phenols in sea water by liquid chromatography with electrochemical detection after enrichment by using solid-phase extraction cartridges and disks. *Anal. Chim. Acta*, 304 (1995) 75-84.
- 2795 Kawahara, A., Yamada, M., Kato, Y., Nakamura, M., Nishijima, M. and Ito, Y.: (Systematic determination of diphenyl, o-phenylphenol, thiabendazole and imazalil in citrus fruits). *Foods Food Ingredients J. Jpn.*, 163 (1995) 104-110; C.A., 122 (1995) 185656x.
- 2796 Krämer, P.M., Li, Q.X. and Hammock, B.D.: Integration of liquid chromatography with immunoassay: an approach combining the strengths of both methods. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1275-1287.
- 2797 Pocerull, E., Sánchez, G., Borrull, F. and Marcé, R.M.: Automated on-line trace enrichment and determination of phenolic compounds in environmental waters by high-performance liquid chromatography. *J. Chromatogr. A*, 696 (1995) 31-39.
- 2798 Rozbeh, M. and Hurtubise, R.J.: Optimum separation and compound class separation of the metabolites of benzo[a]pyrene-DNA adducts with reversed-phase liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3351-3367.
- 2799 Sakuma, S., Kikuchi, C. and Kowaka, M.: Automated analysis of the total polyphenols content of beer and wort using an intelligent autosampler. *J. Am. Soc. Brew. Chem.*, 53 (1995) 29-32; C.A., 122 (1995) 158915z.
- 2800 Variyar, P.S. and Bandyopadhyay, C.: Estimation of phenolic compounds in green pepper berries (*Piper nigrum* L.) by high-performance liquid chromatography. *Chromatographia*, 39 (1994) 743-746.
- 2801 Yang, Y. and He, J.: (Direct determination of microamount of phenol in urine by reversed-phase high performance liquid chromatography). *Sepu*, 13 (1995) 70-71; C.A., 122 (1995) 153472t.

See also 2653, 2774, 2825, 2833, 3000, 3799, 3804, 3823, 3825.

## 8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

### 8a. Flavonoids

- 2802 Bongartz, D. and Hesse, A.: Boronic acid affinity chromatography as a tool for determination of quercetrin in urine. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A17.
- 2803 Cui, J.M., Chung, H.S. and Woo, W.S.: Determination of isoflavonoid glucosides in rhizomes of *Belamcanda chinensis* by high-performance liquid chromatography. *Saengyak Hakhoechi*, 24 (1993) 309-312; C.A., 120 (1994) 280388d.
- 2804 Del Rio, J.A., Castillo, J. and Benavente-Garcia, O.: Elucidation by reverse phase HPLC of some citrus flavanones and their respective dihydrochalcones: structural study. *J. Liq. Chromatogr.*, 17 (1994) 3461-3477.
- 2805 Hammouda, F.M., El-Nasr, M.M.S., Ismail, S.I. and Shahat, A.A.: Quantitative determination of the active constituent in Egyptian cultivated *Cynara scolymus*. *Int. J. Pharmacogn.*, 31 (1993) 299-304; C.A., 120 (1994) 200497a.
- 2806 Kurkin, V.A., Braslavskii, V.B. and Zapesochnaya, G.G.: (Study of extracts of propolis and *Populus balsamifera* buds by high-performance liquid chromatography). *Zh. Fiz. Khim.*, 68 (1994) 1816-1818; C.A., 122 (1995) 75690w.
- 2807 Liu, B., Anderson, D., Ferry, D.R., Seymour, L.W., de Takats, P.G. and Kerr, D.J.: Determination of quercetin in human plasma using reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 149-155.
- 2808 Pietrogrande, M.C. and Kahie, Y.D.: Effect of the mobile and stationary phases on RP-HPLC retention and selectivity of flavonoid compounds. *J. Liq. Chromatogr.*, 17 (1994) 3655-3670.
- 2809 Rehwald, A., Meier, B. and Sticher, O.: Qualitative and quantitative reversed-phase high-performance liquid chromatography of flavonoids in *Passiflora incarnata* L. *Pharm. Acta Helv.*, 69 (1994) 153-158; C.A., 122 (1995) 142702s.
- 2810 Supko, J.G. and Phillips, L.R.: High-performance liquid chromatographic assay for genistein in biological fluids. *J. Chromatogr. B*, 666 (1995) 157-167.

See also 3769, 3803, 3804.

### 8b. Aflatoxins and other mycotoxins

- 2811 Alachaher, L., Abdeddaim, K. and Guermouche, M.H.: (Behavior of aflatoxins in liquid chromatography). *J. Soc. Alger. Chim.*, 4 (1994) 57-96; C.A., 122 (1995) 212352d.
- 2812 Barbieri, G., Bergamini, C., Ori, E. and Resca, P.: Aflatoxin M<sub>1</sub> in Parmesan cheese: HPLC determination. *J. Food Sci.*, 59 (1994) 1313, 1331; C.A., 122 (1995) 131397v.
- 2813 Dragacci, S., Gleizes, E., Fremy, J.M. and Candish, A.A.G.: Use of immunoaffinity chromatography as a purification step for the determination of aflatoxin M<sub>1</sub> in cheeses. *Food Addit. Contam.*, 12 (1995) 59-65; C.A., 122 (1995) 238045f.

- 2814 Guo, Y. and Fu, C.: (HPLC determination of trace trichothecene mycotoxin in corn). *Fenxi Ceshi Xuebao*, 13 (1994) 59-62; C.A., 122 (1995) 79317c.
- 2815 Maksimovic, M.Z. and Vuksanovic, M.: Application of some extraction techniques in analysis of aflatoxins (B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub>) in animal food by HPLC using UV and fluorescence detectors connected in series. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A23.
- 2816 Maragos, C.M. and Richard, J.L.: Quantitation and stability of fumonisins B<sub>1</sub> and B<sub>2</sub> in milk. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1162-1167.
- 2817 Milat, M.-L. and Blein, J.-P.: *Cercospora beticola* toxins. III. Purification, thin-layer and high-performance liquid chromatographic analyses. *J. Chromatogr. A*, 699 (1995) 277-283.
- 2818 Roch, O.G., Blunden, G., Coker, R.D. and Nawaz, S.: The validation of a solid phase clean-up procedure for the analysis of aflatoxins in groundnut cake using HPLC. *Food Chem.*, 52 (1995, Pub. 1994) 93-98; C.A., 122 (1995) 79369w.
- 2819 Trucksess, M.W.: Mycotoxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 135-141 - a review with 120 refs.
- 2820 Wilkes, J.G., Sutherland, J.B., Churchwell, M.I. and Williams, A.J.: Determination of fumonisins B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> by high-performance liquid chromatography with evaporative light-scattering detection. *J. Chromatogr. A*, 695 (1995) 319-323.
- 2821 Zimmerli, B. and Dick, R.: Determination of ochratoxin A at the ppt level in human blood, serum, milk and some foodstuffs by high-performance liquid chromatography with enhanced fluorescence detection and immunoaffinity column cleanup: methodology and Swiss data. *J. Chromatogr. B*, 666 (1995) 85-99.

### 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 2822 Clement, B. and Mladek, C.: Biotransformationsuntersuchungen im Hühnerrei. Teil 1: Metabolische Umwandlung von 7-Ethoxycoumarin. *Pharmazie*, 50 (1995) 207-210.
- 2823 Hiller, D.L. and Cole, R.O.: Short-columns gradient elution high-performance liquid chromatography for the rapid detection of 7-ethoxycoumarin metabolites in liver-slice incubates. *Anal. Biochem.*, 227 (1995) 251-254.
- 2824 Maroni, T. and Lazzari, P.: (Isocratic determination of furosinone using HPLC). *Ind. Aliment. (Pinerolo)*, 33 (1994) 964-966; C.A., 122 (1995) 238011s.
- 2825 Wolbis, M. and Nowak, S.: Coumarins and phenolic acids from the underground part of *Scopolia lurida*. *Acta Pol. Pharm.*, 51 (1994) 171-174; C.A., 122 (1995) 114731g.
- 2826 Xu, Y., Mirocha, C.J. and Xie, W.: Analysis by liquid chromatography of fusarochromanone (TDP-1) added to corn and wheat. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1179-1183.

For additional information see C.A.:

122 (1995) 100764h, 127804h, 142705v, 155422a.

See also 3776, 3805.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 2827 Aboull-Enein, H.Y. and Abou-Basha, L.I.: Simple HPLC method for the determination of thymoquinone in black seed oil (*Nigella sativa* Linn). *J. Liq. Chromatogr.*, 18 (1995) 895-902.

- 2828 Berger, G., Kléo, J., Breton, J., Gilles, N. and Lirsac, P.N.:  $^{18}\text{O}$  and  $^{13}\text{C}$  labeling of photosynthetic and related quinones and their purification by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4531-4539.
- 2829 Coco, F.L., Valentini, C., Novelli, V. and Ceccon, L.: Liquid chromatographic determination of 2-furaldehyde and 5-hydroxymethyl-2-furaldehyde in beer. *Anal. Chim. Acta*, 306 (1995) 57-64.
- 2830 Forgacs, E.: Use of porous graphitized carbon column for determining taxol in *Taxus baccata*. *Chromatographia*, 39 (1994) 740-742.
- 2831 Granados, J.W., Mir, M.V., Lopez Garcia Serrana, H. and Lopez Martinez, M.C.: Comparison of spectrophotometric and chromatographic methods of determination of furanic aldehydes in wine distillates. *Food Chem.*, 52 (1995, Pub. 1994) 203-208; *C.A.*, 122 (1995) 79364r.
- 2832 Liu, G.: Oxidation of acetophenone in a high temperature open-tubular capillary column liquid chromatographic system. *Chromatographia*, 40 (1995) 99-101.
- 2833 Masselter, S., Zemann, A. and Bobleter, O.: Analysis of lignin degradation products by capillary electrophoresis. *Chromatographia*, 40 (1995) 51-57.
- 2834 Sitholé, B.B., Nyarku, S. and Allen, L.H.: Chromatographic methods for the determination of alkylketene dimer in pulp and paper matrices. *Analyst (Cambridge)*, 120 (1995) 1163-1169.
- 2835 Toyo'oka, T. and Liu, Y.-M.: Determination of aldehydes by high-performance liquid chromatography with fluorescence detection after labelling with 4-(2-carbazoylpyrrolidin-1-yl)-7-(N,N-dimethylaminosulfonyl)-2,1,3-benzoxadiazole. *J. Chromatogr. A*, 695 (1995) 11-18.
- 2836 Winkler, R. and Maier, K.: (Testing an HPLC method for determination of malondialdehyde (MDA) in urine, plasma and serum with regard to possible interferences by glucose and streptozotocin (STZ)). *Klin. Labor.*, 40 (1994) 351-355; *C.A.*, 122 (1995) 182302t.
- 2837 Wu, H.-Y. and Jin, J.-K.: Determination of aldehydic lipid peroxidation products with dabsylhydrazine by high-performance liquid chromatography. *Anal. Chem.*, 67 (1995) 1603-1612.
- See also 2643, 2798, 2817, 2896, 3616, 3799, 3830.
10. CARBOHYDRATES
- 10a. Mono and oligosaccharides. Structural studies
- 2838 Buongiorno, A., Morviducci, L., Zorretta, D., Leonetti, F., Tamburrano, G. and Giaccari, A.: Novel method for the determination of hepatic glycogenolysis and gluconeogenesis *in vivo* in rats: preliminary results in hypoglycemia. *Front. Endocrinol.*, 9(Highlights in Molecular and Clinical Endocrinology) (1994) 383-386; *C.A.*, 122 (1995) 234517p.
- 2839 Chusney, G.D., Philippa, M. and Pickup, J.C.: Comparison of micro-enzymatic and high-performance liquid chromatographic methods for the assay of serum 1,5-anhydroglucitol. *Clin. Chim. Acta*, 235 (1995) 91-99.
- 2840 Cooper, G.A. and Rohrer, J.S.: Separation of neutral asparagine-linked oligosaccharides by high-pH anion-exchange chromatography with pulsed amperometric detection. *Anal. Biochem.*, 226 (1995) 182-184.
- 2841 Corradini, C.: Recent advances in carbohydrate analysis by high-performance anion-exchange chromatography coupled with pulsed amperometric detection (HPAEC-PAD). *Ann. Chim. (Rome)*, 84 (1994) 385-396; *C.A.*, 122 (1995) 187876t.
- 2842 Evangelisto, M.F., Adams, R.E., Murray, W.V. and Caldwell, G.W.: Preparative high-performance liquid chromatographic separation of fluorodeoxy sugars. *J. Chromatogr. A*, 695 (1995) 128-131.
- 2843 Fleming, S.C., Gill, M. and Laker, M.F.: A rapid HPLC method for the determination of free fucose in urine. A marker of malignancy in children. *J. Liq. Chromatogr.*, 18 (1995) 173-180.
- 2844 Hino, K., Yamamoto, S., Sano, O., Taniguchi, Y., Kohno, K., Usui, M., Fukuda, S., Hanzawa, H., Haruyama, H. and Kurimoto, M.: Carbohydrate structures of the glycoprotein allergen *Cry j 1* from Japanese cedar (*Cryptomeria japonica*) pollen. *J. Biochem. (Tokyo)*, 117 (1995) 289-295.
- 2845 Hooper, L.V., Beranek, M.C., Manzella, S.M. and Baenziger, J.U.: Differential expression of GalNAc-4-sulfotransferase and GalNAc-transferase results in distinct glycoforms of carbonic anhydrase VI in parotid and submaxillary glands. *J. Biol. Chem.*, 270 (1995) 5985-5993.
- 2846 Karamanos, N.K., Vanky, P., Syrokou, A. and Hjerpe, A.: Identity of dermatan and chondroitin sequences in dermatan sulfate chains determined by using fragmentation with chondroitinases and ion-pair high-performance liquid chromatography. *Anal. Biochem.*, 225 (1995) 220-230.
- 2847 Karlsson, N.G. and Hansson, G.C.: Analysis of monosaccharide composition of mucin oligosaccharide alditols by high-performance anion-exchange chromatography. *Anal. Biochem.*, 224 (1995) 538-541.
- 2848 Kobata, A., Kochibe, N. and Endo, T.: Affinity chromatography of oligosaccharides on *Psathyrella velutina* lectin column. *Methods Enzymol.*, 247(Neoglycoconjugates, Pt. B) (1994) 228-237; *C.A.*, 122 (1995) 182313x.
- 2849 Lopez-Hernandez, J., Gonzalez-Castro, M.J., Vazquez-Blanco, M.E., Vazquez-Oderiz, M.L. and Simal-Lozano, J.: HPLC determination of sugars and starch in green beans. *J. Food Sci.*, 59 (1994) 1048-1049; *C.A.*, 122 (1995) 104173n.
- 2850 Molla, E., Camara, M.M., Diez, C. and Torija, M.E.: (Studies on the determination of sugars in fruits and their products). *Alimentaria (Madrid)*, 254 (1994) 87-93; *C.A.*, 122 (1995) 54322n.
- 2851 Nakagawa, H., Kawamura, Y., Kato, K., Shimada, I., Arata, H. and Takahashi, N.: Identification of neutral and sialyl N-linked oligosaccharide structures from human serum glycoproteins using three kinds of high-performance liquid chromatography. *Anal. Biochem.*, 226 (1995) 130-138.
- 2852 Natunen, J., Niemela, R., Penttila, L., Seppo, A., Ruohtula, T. and Renkonen, O.: Enzymic synthesis of two lacto-N-neo-hexaose-related Lewis x heptasaccharides and their separation by chromatography on immobilized wheat germ agglutinin. *Glycobiology*, 4 (1994) 577-583; *C.A.*, 122 (1995) 133607n.
- 2853 Neville, D.C.A., Field, R.A. and Ferguson, M.A.J.: Hydrophobic glycosides of N-acetylglucosamine can act as primers for polylactosamine synthesis and can affect glycolipid synthesis *in vivo*. *Biochem. J.*, 307 (1995) 791-797.

- 2854 Nojima, K., Isogami, C. and Kobashi, M.: (Application of 4-phenyl-1,2,4-triazoline-3,5-dione for the analysis of sorbic acid in foods). *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 467-471; C.A., 122 (1995) 29975v.
- 2855 Price, N.P.J. and Carlson, R.W.: Rhizobial lipo-oligosaccharide nodulation factors: multidimensional chromatographic analysis of symbiotic signals involved in the development of legume root nodules. *Glycobiology*, 5 (1995) 233-242; C.A., 122 (1995) 238564z.
- 2856 Rodríguez-Aparicio, L.B., Ferrero, M.A. and Reglero, A.: N-Acetyl-D-neuraminic acid synthesis in *Escherichia coli* K1 occurs through condensation of N-acetyl-D-mannosamine and pyruvate. *Biochem. J.*, 308 (1995) 501-505.
- 2857 Romppanen, J. and Mononen, I.: Age-related reference values for urinary excretion of sialic acid and deoxysialic acid: application to diagnosis of storage disorders of free sialic acid. *Clin. Chem. (Washington)*, 41 (1995) 544-547.
- 2858 Saitoh, H., Takagaki, K., Majima, M., Nakamura, T. and Matsuki, A., Kasai, M., Narita, H. and Endo, M.: Enzymic reconstruction of glycosaminoglycan oligosaccharide chains using the transglycosylation reaction of bovine testicular hyaluronidase. *J. Biol. Chem.*, 270 (1995) 3741-3747.
- 2859 Solbrig-Lebuhn, H.: (Two dimensional HPLC of fluorescent labeled oligosaccharides). *BioTec (Wuerzburg)*, 6 (1994) 22; C.A., 122 (1995) 75704d.
- 2860 Takahashi, N., Nakagawa, H., Fujikawa, K., Kawamura, Y. and Tomia, N.: Three-dimensional elution mapping of pyridylaminated N-linked neutral and sialyl oligosaccharides. *Anal. Biochem.*, 226 (1995) 139-146.
- 2861 Takeuchi, T., Kusakabe, I., Yoshida, S.Y.S. and Murata, K.: Separation of oligomannuronic acid based on degree of polymerization. *Jpn. Kokai Tokkyo Koho JP 06,172,375 [94,172,375] (Cl. C07H1/06)*, 21 Jun. 1994, Appl. 92/325,289, 04 Dec. 1992; 4 pp.; C.A., 122 (1995) 106386h.
- 2862 Yutani, S., Toh, G. and Honda, S.: (Analysis of fructooligosaccharides by HPLC and HPCE). *Kuromatogurafi*, 15 (1994) 210-211; C.A., 122 (1995) 127992t.
- For additional information see C.A.:  
122 (1995) 79347n, 101143s, 158794j, 187881r, 187890t.
- See also 2588, 2869, 2872, 2876, 2939, 3206, 3226, 3232, 3405, 3417, 3596, 3812.

#### 10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 2863 Dubin, P.L.: Nonionic polysaccharides as calibration standards for aqueous size exclusion chromatography. *Carbohydr. Polym.*, 25 (1994) 295-303; C.A., 122 (1995) 208918a.
- 2864 Hashimoto-Uoshima, M., Hascall, V.C., MacCallum, D.K. and Yanagishita, M.: Biosynthesis of proteoglycans and hyaluronic acid by rat oral epithelial cells (keratinocytes) *in vitro*. *Arch. Biochem. Biophys.*, 316 (1995) 724-732.
- 2865 Kasinath, B.S.: Effect of insulin on high-glucose medium-induced changes in rat glomerular epithelial cell metabolism of glycoconjugates. *Arch. Biochem. Biophys.*, 318 (1995) 286-294.

- 2866 Lagana, A., Marino, A., Fago, G. and Pardo Martinez, B.: Determination of free N-acetyl neuraminic acid by ultrafiltration and liquid chromatography in the serum of normal and cancer patients. *Anal. Chim. Acta*, 306 (1995) 65-71.
- 2867 Midura, R.J., Salustri, A., Calabro, A., Yanagishita, M. and Hascall, V.C.: High-resolution separation of disaccharide and oligosaccharide alditols from chondroitin sulfate, dermatan sulfate and hyaluronan using CarboPac PA1 chromatography. *Glycobiology*, 4 (1994) 333-342; C.A., 122 (1995) 75676w.
- 2868 Müller, A., Pretus, H.A., McNamee, R.B., Jones, E.L., Browder, I.W. and Williams, D.L.: Comparison of the carbohydrate biological response modifiers Krestin, schizophyllan and glucan phosphate by aqueous size exclusion chromatography with inline argon-ion multi-angle laser light scattering photometry and differential viscometry detectors. *J. Chromatogr. B*, 666 (1995) 283-290.
- 2869 Ng, C.K., Handley, C.J., Preston, B.N., Robinson, H.C., Bolis, S. and Parker, G.: Effect of exogenous hyaluronan and hyaluronan oligosaccharides on hyaluronan and aggrecan synthesis and catabolism in adult articular cartilage explants. *Arch. Biochem. Biophys.*, 316 (1995) 596-606.
- 2870 Shao, M.-C. and Wold, F.: The effect of the protein matrix proximity on glycan reactivity in a glycoprotein model. *Eur. J. Biochem.*, 228 (1995) 79-85.
- 2871 Van Det, N.F., van den Born, J., Tamsma, J.T., Verhagen, N.A.M., van den Heuvel, L.P.W.J., Berden, J.H.M., Bruijn, J.A., Daha, M.R. and van der Woupe, F.J.: Proteoglycan production by human glomerular visceral epithelial cells and mesangial cells *in vitro*. *Biochem. J.*, 307 (1995) 759-768.
- 2872 Volpi, N., Cusmano, M. and Venturelli, T.: Qualitative and quantitative studies of heparin and chondroitin sulfates in normal human plasma. *Biochim. Biophys. Acta*, 1243 (1995) 49-58.
- 2873 Wong, K.S. and Jane, J.: Effects of pushing agents on the separation and detection of debranched amylopectin by high-performance anion-exchange chromatography with pulsed amperometric detection. *J. Liq. Chromatogr.*, 18 (1995) 63-80.
- 2874 Yang, J. and Z.: (Determination of molecular weights and their distributions of polysaccharose in astragalus by using 150-C ALC/GPC gel permeation chromatography). *Taiwan Gongye Daxue Xuebao*, 25 (1994) 77-84; C.A., 122 (1995) 100763g.

For additional information see C.A.:

122 (1995) 185680a.

See also 2629, 2846, 2949, 3139, 3206, 3232, 3475, 3818.

#### 10c. Glycoproteins and their constituents

- 2875 Altman, E., Schäffer, C., Brisson, J.-R. and Messner, P.: Characterization of the glycan structure of a major glycopeptide from the surface layer glycoprotein of *Clostridium thermosaccharolyticum*. *Eur. J. Biochem.*, 229 (1995) 308-315.
- 2876 Ashton, D.S., Beddell, C.R., Cooper, D.J. and Lines, A.C.: Determination of carbohydrate heterogeneity in the humanised antibody CAMPATH 1H by liquid chromatography and matrix-assisted laser desorption ionisation mass spectrometry. *Anal. Chim. Acta*, 306 (1995) 43-48.

- 2877 Cho, M. and Cummings, R.D.: Galectin-1, a  $\beta$ -galactoside-binding lectin in chinese hamster ovary cells. I. Physical and chemical characterization. *J. Biol. Chem.*, 270 (1995) 5198-5206.
- 2878 Coppolino, M., Migliorini, M., Argraves, W.S. and Dedhar, S.: Identification of a novel form of the  $\alpha_3$  integrin subunit: covalent association with transferrin receptor. *Biochem. J.*, 306 (1995) 129-134.
- 2879 Hokke, C.H., Bergwerff, A.A., van Dedem, G.W.K., Kamerling, J.P. and Vliegthart, J.F.G.: Structural analysis of the sialylated N- and O-linked carbohydrate chains of recombinant human erythropoietin expressed in Chinese hamster ovary cells. Sialylation patterns and branch location of dimeric N-acetylglucosamine units. *Eur. J. Biochem.*, 228 (1995) 981-1008.
- 2880 Nakazawa, K., Isomura, T. and Shimoeda, S.: Proteoglycan synthesis by chicken corneal explants. *J. Biochem. (Tokyo)*, 117 (1995) 697-706.
- 2881 Nakazawa, K., Suzuki, S., Wada, K. and Nakazawa, K.: Proteoglycan synthesis by corneal explants from developing embryonic chicken. *J. Biochem. (Tokyo)*, 117 (1995) 707-718.
- 2882 Osman, M.E., Menzies, A.R., Martin, B.A., Williams, P.A., Phillips, G.O. and Baldwin, T.C.: Characterization of gum arabic fractions obtained by anion-exchange chromatography. *Phytochemistry*, 38 (1995) 409-417; *C.A.*, 122 (1995) 209779t.
- 2883 Parekh, R. and Ventom, A.: Glycotechnology: an unambiguous technique for glycan analysis. *Am. Biotechnol. Lab.*, 13 (1995) 9-10; *C.A.*, 122 (1995) 155376p.
- 2884 Paszkiewicz-Gadek, A., Gindzienski, A. and Porowska, H.: The use of preparative polyacrylamide gel electrophoresis and electroelution for purification of mucus glycoproteins. *Anal. Biochem.*, 226 (1995) 263-267.
- 2885 Sareneva, T., Pirhonen, J., Cantell, K. and Julkunen, I.: N-Glycosylation of human interferon- $\gamma$ : glycans at Asn-25 are critical for protease resistance. *Biochem. J.*, 308 (1995) 9-14.
- 2886 Sato, T., Takio, K., Kobata, A., Greenwalt, D.E. and Furukawa, K.: Site-specific glycosylation of bovine butyrophilin. *J. Biochem. (Tokyo)*, 117 (1995) 147-157.
- 2887 Sharom, F.J., Yu, X., Chu, J.W.K. and Doige, C.A.: Characterization of the ATPase activity of P-glycoprotein from multidrug-resistant Chinese hamster ovary cells. *Biochem. J.*, 308 (1995) 381-390.
- 2888 Weber, P.L., Kornfelt, T., Klausen, N.K. and Lunte, S.M.: Characterization of glycopeptides from recombinant coagulation factor VIIa by high-performance liquid chromatography and capillary zone electrophoresis with ultraviolet and pulsed electrochemical detection. *Anal. Biochem.*, 225 (1995) 135-142.
- 2889 Young, N.M., Watson, D.C., Yaguchi, M., Adar, R., Arango, R., Rodriguez-Arango, E., Sharon, N., Blay, P.K.S. and Thibault, P.: C-terminal post-translational proteolysis of plant lectins and their recombinant forms expressed in *Escherichia coli*. Characterization of "ragged ends" by mass spectrometry. *J. Biol. Chem.*, 270 (1995) 2563-2570.
11. ORGANIC ACIDS AND LIPIDS
- 11a. Organic acids and simple esters
- 2890 Bertolini, L., Castagnetti, G.B. and Gambini, G.: (Organic acids in vinegars: HPLC identification). *Ind. Bevande*, 23 (1994) 324-326; *C.A.*, 122 (1995) 54327t.
- 2891 Bieganska, M.L., Petruczynik, A. and Glowinski, K.: Reversed phase retention behavior of  $\beta$ -cyclodextrin complexes of phenolic and cinnamic acids. *J. Planar Chromatogr.*, 8 (1995) 63-68.
- 2892 Buiarelli, F., Cartoni, G., Coccioli, F. and Levetsovitou, Z.: Determination of phenolic acids in wine by high-performance liquid chromatography with a microbore column. *J. Chromatogr. A*, 695 (1995) 229-235.
- 2893 Castellani, L., Flieger, M. and Sinibaldi, M.: Enantiomer separation of 2-arylpropionic acids on an ergot alkaloid-based stationary phase microbore column application. *J. Liq. Chromatogr.*, 17 (1994) 3695-3703.
- 2894 Chavis, C., Chanez, P., Vachier, I., Boosquet, J., Michel, F.B. and Godard, P.: 5-15-Dihete and lipoxins generated by neutrophils from endogenous arachidonic acid as asthma biomarkers. *Biochem. Biophys. Res. Commun.*, 207 (1995) 273-279.
- 2895 Coudray, C., Talla, M., Martin, S., Fatome, M. and Favier, A.: High-performance liquid chromatography-electrochemical determination of salicylate hydroxylation products as an *in vivo* marker of oxidative stress. *Anal. Biochem.*, 227 (1995) 101-111.
- 2896 Demini, P., Reynaud, D. and Pace-Asciak, C.R.: Extractive derivatization of the 12-lipoxygenase products, hepxilins, and related compounds into fluorescent anthryl esters for their complete high-performance liquid chromatography profiling in biological systems. *Anal. Biochem.*, 226 (1995) 252-255.
- 2897 Ding, M.-Y., Koizumi, H. and Suzuki, Y.: Comparison of three chromatographic systems for determination of organic acids in wine. *Anal. Sci.*, 11 (1995) 239-243; *C.A.*, 122 (1995) 238049k.
- 2898 Fernandez-Garcia, E. and McGregor, J.U.: Determination of organic acids during the fermentation and cold storage of yogurt. *J. Dairy Sci.*, 77 (1994) 2934-2939; *C.A.*, 122 (1995) 79319e.
- 2899 Gao, Z. and Fu, C.: (Determination of organic acids and inorganic anions by single-column ion chromatography). *Fenxi Ceshi Xuebao*, 13 (1994) 32-36; *C.A.*, 122 (1995) 122251r.
- 2900 Gensler, M. and Schmidt, H.-L.: Isolation of the main organic acids from fruit juices and nectars for carbon isotope ratio measurements. *Anal. Chim. Acta*, 299 (1994) 231-237.
- 2901 Gronert, K., Virk, S.M. and Herman, C.A.: Endogenous sulfidopeptide leukotriene synthesis and 12-lipoxygenase activity in bullfrog (*Rana catesbeiana*) erythrocytes. *Biochim. Biophys. Acta*, 1255 (1995) 311-319.
- 2902 Hesse, A., Heynck, H. and Berg, W.: Measurement of urinary oxalic acid: a comparison of four methods. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A19.
- 2903 Huang, S.-Y. and Jin, J.-D.: Purification of methyl esters of polyunsaturated fatty acids by displacement chromatography. *Bioseparation*, 4 (1994) 343-351; *C.A.*, 122 (1995) 50412u.
- 2904 Jansen, E.H.J.M., Laan, C.A. and de Fluiter, P.: Determination of phthalate-induced rat liver cytochrome P-450 IVA1 by a fluorimetric enzymatic assay and by chemiluminescence detection on Western blots. *Anal. Chim. Acta*, 303 (1995) 99-102.

See also 2851.

- 2905 Kadkhodaei, M., Endre, Z.H., Towner, R.A. and Cross, M.: Hydroxyl radical generation following ischaemia-reperfusion in cell-free perfused rat kidney. *Biochim. Biophys. Acta*, 1243 (1995) 169-174.
- 2906 Kondo, Y., Hashidoko, Y. and Mizutani, J.: An enzymatic formation of 13-oxo-trideca-9,11-dienoic acid from 13-hydroperoxylinolenic acid by a homolytic hydroperoxide lyase in elicitor-treated soybean cotyledons. *Biochim. Biophys. Acta*, 1255 (1995) 9-15.
- 2907 Kroumova, A.B. and Wagner, G.J.: Methods for separation of free, short, medium, and long chain fatty acids and for their decarboxylation. *Anal. Biochem.*, 225 (1995) 270-276.
- 2908 Lee, H.S.: Liquid chromatographic determination of benzoic acid in orange juice: interlaboratory study. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 80-82.
- 2909 Lin, J.-T., McKeon, T.A. and Stafford, A.E.: Gradient reversed-phase high-performance liquid chromatography of saturated, unsaturated and oxygenated free fatty acids and their methyl esters. *J. Chromatogr. A*, 699 (1995) 85-91.
- 2910 Olawoye, T.L., Minnikin, D.E. and Christie, W.W.: HPLC separation of *tert*-butyl dimethylsilyl (TBDMS) ether homologs of mycolic acids, using a silver impregnated normal phase column. *Biochim. Biophys. Acta*, 122 (1995) 26980g.
- 2911 Park, P.W. and Goins, R.E.: *In situ* preparation of fatty acid methyl esters for analysis of fatty acid composition in foods. *J. Food Sci.*, 59 (1994) 1262-1266; *C.A.*, 122 (1995) 131279h.
- 2912 Perrut, M., Nicoud, R.-M. and Breivik, H.: Process for chromatographic fractionation of polyunsaturated fatty acids and their derivatives. *PCT Int. Appl. WO 94 25,552* (Cl. C11C1/08), 10 Nov. 1994, GB Appl. 93/8,912, 29 Apr. 1993; 70 p.; *C.A.*, 122 (1995) 109324x.
- 2913 Pouliot, J.-F. and Béliveau, R.: Palmitoylation of the glucose transporter in blood-brain barrier capillaries. *Biochim. Biophys. Acta*, 1234 (1995) 191-196.
- 2914 Rao, M.S., Hidajat, K. and Ching, C.B.: Reversed-phase HPLC: the separation method for the characterization and purification of long chain polyunsaturated fatty acids - a review. *J. Chromatogr. Sci.*, 33 (1995) 9-21 - a review with 108 refs.
- 2915 Sacconi, G., Gherardi, S., Trifiro, A., Bordini, C.S., Calza, M., Zoni, C. and Freddi, C.: (Determination of organic acids in fruit juices by ion chromatography). *Ind. Bevande*, 23 (1994) 423-429; *C.A.*, 122 (1995) 212369q.
- 2916 Saito, M., Ushijima, T., Sasamoto, K., Ohkura, Y. and Ueno, K.: 2-(5-Hydrazinocarbonyl-2-furyl)-5,6-dimethoxybenzothiazole as a precolumn fluorescence derivatization reagent for carboxylic acids in high-performance liquid chromatography and its application to the assay of fatty acids in human serum. *Anal. Sci.*, 11 (1995) 103-107; *C.A.*, 122 (1995) 208892n.
- 2917 See, M.M., Thompson, R., Grinberg, N., Perpall, H.J., Bicker, G. and Tway, P.: Chromatographic analysis of residual acetate in bulk drugs. *J. Liq. Chromatogr.*, 18 (1995) 137-154.
- 2918 Skotty, D.R. and Nieman, T.A.: Determination of oxalate in urine and plasma using reversed-phase ion-pair high-performance liquid chromatography with tris(2,2'-bipyridyl)ruthenium(II)-electrogenerated chemiluminescence detection. *J. Chromatogr. B*, 665 (1995) 27-36.
- 2919 Sushchik, N.N., Gladyshev, M.I., Kalachova, G.S. and Guseynova, V.E.: Rapid assay of fatty acid composition using a portable high-performance liquid chromatograph for monitoring aquatic ecosystems. *J. Chromatogr. A*, 695 (1995) 223-228.
- 2920 Vichot, R. and Furton, K.G.: Variables influencing the direct determination of haloacetic acids in water by reversed-phase ion-pair chromatography with indirect UV detection. *J. Liq. Chromatogr.*, 17 (1994) 4405-4429.
- 2921 Walker, T.A.: A liquid chromatographic assay for citric acid in over-the-counter carbamide peroxide products using indirect UV detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 171-176.
- 2922 Yu, W. and Powell, W.S.: Analysis of leukotrienes, lipoxins, and monooxygenated metabolites of arachidonic acid by reversed-phase high-pressure liquid chromatography. *Anal. Biochem.*, 226 (1995) 241-251.

For additional information see C.A.:  
122 (1995) 50452g, 208914w.

See also 2625, 3033, 3268, 3361, 3410, 3565, 3616, 3802, 3810, 3816, 3888.

#### 11b. Prostaglandins

- 2923 Capdevila, J.H., Morrow, J.D., Belosludtsev, Y.Y., Beauchamp, D.R., DuBois, R.N. and Falck, J.R.: The catalytic outcomes of the constitutive and the mitogen inducible isoforms of prostaglandin H<sub>2</sub> synthase are markedly affected by glutathione and glutathione peroxidase(s). *Biochemistry*, 34 (1995) 3325-3337.
- 2924 Gatti, R., Gotti, R., Cavrini, V., Soli, M., Bertaccini, A. and Carparelli, F.: Stability study of prostaglandin E<sub>1</sub> (PGE<sub>1</sub>) in physiological solutions by liquid chromatography (HPLC). *Int. J. Pharm.*, 115 (1995) 113-117; *C.A.*, 122 (1995) 142298q.
- 2925 Yamane, M., Shimizu, S., Abe, A., Sugiura, H., Miyaoka, M. and Saitoh, T.: High-performance liquid chromatography-thermospray mass spectrometry of  $\omega$ -carboxyleukotriene B<sub>4</sub> and  $\omega$ -hydroxyleukotriene B<sub>4</sub> from an incubation mixture of human colonic well-differentiated adenocarcinoma homogenate. *J. Chromatogr. B*, 666 (1995) 197-202.

See also 2922.

#### 11c. Lipids and their constituents

- 2926 Abidi, S.L., Mounts, T.L. and Rennick, K.A.: Separations of major soybean phospholipids on  $\beta$ -cyclodextrin-bonded silica. *J. Liq. Chromatogr.*, 17 (1994) 3705-3725.
- 2927 Adlof, R.O.: Analysis of triacylglycerol positional isomers by silver ion high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 105-107.
- 2928 Adlof, R.O.: The preparative separation of lipids by silver resin chromatography. *Process Technol. Proc.*, 11 (1994) 77-781; *C.A.*, 122 (1995) 189497u.
- 2929 Akasaka, K., Ohata, A., Ohrai, H. and Meguro, H.: Automatic determination of hydroperoxides of phosphatidylcholine and phosphatidylethanolamine in human plasma. *J. Chromatogr. B*, 665 (1995) 37-43.
- 2930 Alter, C.A. and Wolf, B.A.: Identification of phosphatidylinositol 3,4,5-trisphosphate in pancreatic islets and insulin secreting  $\beta$ -cells. *Biochem. Biophys. Res. Commun.*, 208 (1995) 190-197.

- 2931 Bernhard, W., Postle, A.D., Linck, M. and Sewing, K.-F.: Composition of phospholipid classes and phosphatidylcholine molecular species of gastric mucosa and mucus. *Biochim. Biophys. Acta*, 1255 (1995) 99-104.
- 2932 Christie, W.W. and Urwin, R.A.: Separation of lipid classes from plant tissues by high performance liquid chromatography on chemically bonded stationary phases. *J. High Resolut. Chromatogr.*, 18 (1995) 97-100.
- 2933 Eaton, S., Shmueli, E., Al-Mardini, H., Bartlett, K. and Record, C.O.: An HPLC assay for sn-1,2-diacylglycerol. *Clin. Chim. Acta*, 234 (1995) 71-78.
- 2934 Firestone, D.: Fats and oils. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 150-155 - a review with 26 refs.
- 2935 Fukami, M.H. and Holmsen, H.: Diacylglycerol elevations in control platelets are unaccompanied by pleckstrin phosphorylation. Implications for the role of diacylglycerol in platelet activation. *Eur. J. Biochem.*, 228 (1995) 579-586.
- 2936 Grob, K. and Mariani, C.: LC-GC method for the determination of adulterated edible oils and fats. *Spec. Publ. - R. Soc. Chem.*, 160(Developments in the Analysis of Lipids) (1994) 73-81; C.A., 122 (1995) 131220g - a review with 13 refs.
- 2937 Hayashi, K., Ohnishi, T., Tabata, M., Yoshita, K., Nakagawa, H., Kawano, S., Nagase, H., Okazawa, T., Yoshida, M. et al.: (Studies on the analysis of serum alkenal and serum lipid peroxides in migrant workers exposed to dust). *Hokuriku Kosho Eisei Gakkaishi*, 20 (1993) 29-36; C.A., 122 (1995) 48562z.
- 2938 Heron, S., Lesellier, E. and Tchaplal, A.: Analysis of triacylglycerols of borage oil by RPLC identification by coinjection. *J. Liq. Chromatogr.*, 18 (1995) 599-611.
- 2939 Holst, O., Bock, K., Brade, L. and Brade, H.: The structure of oligosaccharide bisphosphates isolated from the lipopolysaccharide of a recombinant *Escherichia coli* strain expressing the gene *gseA* [3-deoxy-D-manno-octulopyranosonic acid (Kdo) transferase I] of *Chlamydia psittaci* 6B. *Eur. J. Biochem.*, 229 (1995) 194-200.
- 2940 Itonori, S., Shiray, T., Kiso, Y., Oha, Y., Shiota, K. and Ogawa, T.: Glycosphingolipid composition of rat placenta: changes associated with stage of pregnancy. *Biochem. J.*, 307 (1995) 399-405.
- 2941 Jalink, K., Hengeveld, T., Mulder, S., Postma, F.R., Simon, M.-F., Chap, H., van der Marel, G.A., van Boom, J.H., van Blitterswijk, W.J. and Moolenaar, W.H.: Lysophosphatidic acid-induced Ca<sup>2+</sup> mobilization in human A431 cells: structure-activity analysis. *Biochem. J.*, 307 (1995) 609-616.
- 2942 Justin, A.-M., Hmyene, A., Kader, J.-C. and Mazliak, P.: Compared selectivities of the phosphatidylinositol-synthase from maize coleoptiles either in microsomal membranes or after solubilization. *Biochim. Biophys. Acta*, 1255 (1995) 161-166.
- 2943 Krainev, A.G., Ferrington, D.A., Williams, T.D., Squier, T.C. and Bigelow, D.J.: Adaptive changes in lipid composition of skeletal sarcoplasmic reticulum membranes associated with aging. *Biochim. Biophys. Acta*, 1235 (1995) 406-418.
- 2944 Leopold, K.: (Hydrophobic interaction chromatography (HIC) of lipid macroamphiphiles). *GIT Fachz. Lab.*, 38 (1994) 455-460; C.A., 122 (1995) 182303u.
- 2945 Luciani, S., Antolini, M., Bova, S., Cargnelli, G., Cusinato, F., Debetto, P., Trevisi, L. and Varotto, R.: Inhibition of cardiac sarcolemmal sodium-calcium exchanger by glycerophosphoinositol 4-phosphate and glycerophosphoinositol 4,5-bisphosphate. *Biochem. Biophys. Res. Commun.*, 206 (1995) 674-680.
- 2946 Ma, Y.-C. and Kim, H.-Y.: Development of the on-line high-performance liquid chromatography/thermospray mass spectrometry method for the analysis of phospholipid molecular species in rat brain. *Anal. Biochem.*, 226 (1995) 293-301.
- 2947 Meyuhas, D. and Lichtenberg, D.: The effect of albumin on the state of aggregation and phase transformations in phosphatidylcholine-sodium cholate mixtures. *Biochim. Biophys. Acta*, 1234 (1995) 203-213.
- 2948 Nikolova-Damyanova, B., Christie, W.W. and Herslöf, B.G.: Retention properties of triacylglycerols on silver ion high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 375-380.
- 2949 Proudfoot, L., Schneider, P., Ferguson, M.A.J. and McConville, M.J.: Biosynthesis of the glycolipid anchor of lipophosphoglycan and the structurally related glycoinositolphospholipids from *Leishmania major*. *Biochem. J.*, 308 (1995) 45-55.
- 2950 Quarfordt, S.H., Landis, B.A. and Oswald, B.S.: The influence of cholesteryl ester content on hepatocyte triolein emulsion uptake. *Biochim. Biophys. Acta*, 1255 (1995) 82-86.
- 2951 Redden, P.R., Huang, Y.-S., Lin, X. and Horrobin, D.F.: Separation and quantification of the triacylglycerols in evening primrose and borage oils by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 381-389.
- 2952 Spanos, G.A., Schwartz, S.J., van Breemen, R.B. and Huang, C.-H.: High-performance liquid chromatography with light-scattering detection and desorption chemical-ionization tandem mass spectrometry of milk fat triacylglycerols. *Lipids*, 30 (1995) 85-90; C.A., 122 (1995) 131316t.
- 2953 Xu, G. and Chang, L.: (Progress in high performance liquid chromatography-precolumn derivatization for gangliosides). *Sepu*, 13 (1995) 26-29; C.A., 122 (1995) 234472v - a review with 5 refs.

For additional information see C.A.:

122 (1995) 27201r, 155416b, 212490x.

See also 2791, 2853, 2855, 2909, 2971, 3144, 3477.

#### 11d. Lipoproteins and their constituents

- 2954 Anchisi, C., Batetta, B., Sanna, F., Fadda, A.M., Maccioni, A.M. and Dessi, S.: HDL subfractions as altered in cancer patients. *J. Pharm. Biomed. Anal.*, 13 (1995) 65-71.
- 2955 Atwal, A.S., Blais, L. and Jenkins, K.J.: Direct determination of apolipoproteins in plasma by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 579-587.
- 2956 Nagayoshi, A., Matsuki, N., Saito, H., Tsukamoto, K., Kaneko, K., Wakashima, M., Kinoshita, M., Yamanaka, M. and Teramoto, T.: Defect in assembly process of very-low-density lipoprotein in suncus liver: an animal model of fatty liver. *J. Biochem. (Tokyo)*, 117 (1995) 787-793.
- 2957 Panteghini, M., Bonora, R. and Pagani, F.: Determination of glycated apolipoprotein B in serum by a combination of affinity chromatography and immunonephelometry. *Ann. Clin. Biochem.*, 31 (1994) 544-549; C.A., 122 (1995) 75915y.
- 2958 Rinninger, F., Brundert, M., Jäckle, S., Kaiser, T. and Greten, H.: Selective uptake of low-density lipoprotein-associated cholesteryl esters by human fibroblasts, human HepG2 hepatoma cells and J774 macrophages in culture. *Biochim. Biophys. Acta*, 1255 (1995) 141-153.



- 2959 Triccerri, A., Garda, H.A. and Brenner, R.R.: A rapid and efficient procedure for the purification of human apolipoprotein A-I using gel filtration HPLC. *Int. J. Bio-Chromatogr.*, 1 (1994) 159-166; *C.A.*, 122 (1995) 75707g.

## 12. ORGANIC PEROXIDES

- 2960 Zhang, J.-R., Cazars, A.R., Lutzke, B.S. and Hall, E.D.: HPLC-chemiluminescence and thermospray LC/MS study of hydroperoxides generated from phosphatidylcholine. *Free Radical Biol. Med.*, 18 (1995) 1-10; *C.A.*, 122 (1995) 75987y.

See also 2641, 2929, 2971.

## 13. STEROIDS

### 13a. General techniques

See 2685.

### 13b. Pregnane and androstane derivatives

- 2961 Bidart, M. and Lesgards, G.: Direct injection analysis of 6 $\beta$ -hydroxycortisol and cortisol in urine by HPLC-UV with on-line ISRP precolumn. *J. Liq. Chromatogr.*, 18 (1995) 725-738.
- 2962 Dwyer, J.L., Chapman, A.E. and Liu, X.: Analysis of steroids by combined chromatography-infrared spectroscopy. *LC-GC*, 13 (1995) 240-250; *C.A.*, 122 (1995) 197098e.
- 2963 Heilmann, P., Vilemeyer, O., Lessing, U. and Schöneshöfer, M.: Fully automated determination of testosterone levels in male urine using the ALCA™ system. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A20.
- 2964 Loftsson, T., Gudmundsdottir, T.K., Fridriksdottir, H., Sigurdardottir, A.M., Thorkelsson, J., Gudmundsson, G. and Hjaltason, B.: Fatty acids from cod-liver oil as skin penetration enhancers. *Pharmazie*, 50 (1995) 188-190.
- 2965 Santos-Montes, A., Gasco-Lopez, A.I. and Izquierdo-Hornillos, R.: Simultaneous determination of dexamethasone and betamethasone in pharmaceuticals by reversed-phase HPLC. *Chromatographia*, 39 (1994) 539-542.
- 2966 Shimada, K., Satoh, Y. and Nishimura, S.: Studies on neurosteroids. I. Retention behavior of derivatized 17-oxosteroids using high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 713-723.
- 2967 Torres Cartas, S., Garcia Alvarez-Coque, M.C. and Villanueva Cernañas, R.M.: Determination of anabolic steroids in pharmaceuticals by liquid chromatography with a microemulsion of sodium dodecyl sulfate and pentanol as mobile phase. *Anal. Chim. Acta*, 302 (1995) 163-172.
- 2968 Wienkers, L.C., Stenwyk, R.C., Miszak, S.A. and Pearson, P.G.: *In vitro* metabolism of tirilazad mesylate in male and female rats. Contribution of cytochrome P4502C11 and 5 $\alpha$ -reductase. *Drug Metab. Disp.*, 23 (1995) 383-392.

For additional information see *C.A.*:  
122 (1995) 115142c.

See also 3648, 3754, 3819.

### 13c. Estrogens

- 2969 Miller, R.B., Delker, G., Chen, C. and Sherwood, C.H.: Development of a HPLC method for the determination of 17 $\beta$ -estradiol-3-phosphate in pharmaceutical preparations. *J. Liq. Chromatogr.*, 18 (1995) 127-136.
- 2970 Sheikh, S.U. and Touchstone, J.C.: Determination of free estriol in amniotic fluid by high performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3813-3820.

See also 2964.

### 13d. Sterols

- 2971 Bartolomeazzi, R., Pizzale, L., Vichi, S. and Lercker, G.: Analysis of isomers of cholesteryl acetate hydroperoxides by HPLC and GC-ITDMS. *Chromatographia*, 39 (1994) 577-580.
- 2972 Dinchuk, J., Hart, J., Gonzalez, G., Karmann, G., Schmidt, D. and Wirak, D.O.: Remodelling of lipoproteins in transgenic mice expressing human cholesteryl ester transfer protein. *Biochim. Biophys. Acta*, 1255 (1995) 301-310.
- 2973 Hamill, T.W. and Soliman, A.M.: Determination of cholesterol by *p*-nitrobenzoate derivatization and liquid chromatography. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1190-1196.
- 2974 Nielsen, J.H., Olsen, C.E., Duedahl, C. and Skibsted, L.H.: Isolation and quantification of cholesterol oxides in dairy products by selected ion monitoring mass spectrometry. *J. Dairy Res.*, 62 (1995) 101-113; *C.A.*, 122 (1995) 131321r.

### 13e. Bile acids and alcohols

- 2975 Iida, T., Yamaguchi, Y., Maruyama, J., Nishio, M., Goto, J. and Nambara, T.: High performance liquid chromatographic separation of nonamidated and glycine- and taurine-amidated bile acid 3-glucosides. *J. Liq. Chromatogr.*, 18 (1995) 701-711.
- 2976 Roda, A., Gioacchini, A.M., Cerrè, C. and Baraldini, M.: High-performance liquid chromatographic-electrospray mass spectrometric analysis of bile acids in biological fluids. *J. Chromatogr. B*, 665 (1995) 281-294.
- 2977 Scalia, S., Tirendi, S., Pazzi, P. and Bousquet, E.: Assay of free bile acids in pharmaceutical preparations by HPLC with electrochemical detection. *Int. J. Pharm.*, 115 (1995) 249-253; *C.A.*, 122 (1995) 170319r.

### 13g. Other steroids

- 2978 Gamoh, K., Yamaguchi, I. and Takatsuto, S.: Rapid and selective sample preparation for the chromatographic determination of brassinosteroids from plant material using solid-phase extraction method. *Anal. Sci.*, 10 (1994) 913-917; *C.A.*, 122 (1995) 50408x.

## 14. STEROID GLYCOSIDES AND SAPONINS

- 2979 Fujii, Y., Ikeda, Y. and Yamazaki, M.: High-performance liquid chromatographic determination of lanatosides in *Digitalis lutea* and *Digitalis ambigua* leaves. *J. Liq. Chromatogr.*, 17 (1994) 4451-4461.
- 2980 Kockar, O.M., Kara, M., Kara, S., Bozan, B. and Baser, K.H.: Quantitative determination of escin. A comparative study of HPLC and TLC densitometry. *Fitoterapia*, 65 (1994) 439-443; *C.A.*, 122 (1995) 170314k.
- 2981 Muhr, P., Michelitsch, A., Likussar, W. and Schubert-Zsilavecz, M.: Ein einfacher Zugang zu reinem Digitonin und Digalotin. *Pharmazie*, 50 (1995) 295-296.

For additional information see *C.A.*:  
122 (1995) 150713m.

## 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

## 15a. Terpenes

- 2982 Cai, J.J., Tao, X. and Lipsky, P.E.: High performance liquid chromatographic determination of triptolide and triptolidide in an ethyl acetate extract of *Tripterygium wilfordii* Hook F. *J. Liq. Chromatogr.*, 17 (1994) 4479-4487.
- 2983 Cheung, H.T.A., Fu, S.-L. and Smal, M.A.: Inhibition of platelet aggregation by diterpene acids from *Pinus massoniana* resin. *Arzneim.-Forsch.*, 44 (1994) 17-25.
- 2984 Green, M.D., Mount, D.L., Todd, G.D. and Capornacchia, A.C.: Chemiluminescent detection of artemisinin. Novel endoperoxide analysis using luminol without hydrogen peroxide. *J. Chromatogr. A*, 695 (1995) 237-242.
- 2985 Mondello, L., Dugo, P., Bartle, K.D., Frere, B. and Dugo, G.: On-line high performance liquid chromatography coupled with high resolution gas chromatography and mass spectrometry (HPLC-HRGC-MS) for the analysis of complex mixtures containing high volatile compounds. *Chromatographia*, 39 (1994) 529-538.
- 2986 Nowak, G.: Isolation and chromatography of 15-deoxyrepin and 25 other sesquiterpene lactones from *Centaurea bella*. *Acta Soc. Bot. Pol.*, 62 (1993) 33-36; *C.A.*, 122 (1995) 89205d.
- 2987 O'Mathuna, D.P. and Doskotch, R.W.: New labdane diterpene glycosides from *Amphiachyris amoena*. *J. Natural Prod.*, 58 (1995) 82-92.
- 2988 Rodriguez, A.D., Cobar, O.M. and Martinez, N.: Isolation and structures of sixteen new asbestinin diterpenes from the caribbean gorgonian *Briareum asbestinum*. *J. Natural Prod.*, 57 (1994) 1638-1655.

## 15b. Essential oils

- 2989 Zekovic, Z., Pekic, B., Ledojevic, Z. and Petrovic, L.: Chromatography in our investigations of camomile (*Metricaria chamomilla* L.). *Chromatographia*, 39 (1994) 587-590.

See also 2985.

## 15c. Bitter substances

- 2990 Buckee, G.K.: Determination of iso-alpha-, alpha-, and beta-acids in hop and isomerized hop extracts. A collaborative trial. *Monatsschr. Brauwiss.*, 47 (1994) 284-286; *C.A.*, 122 (1995) 185640n.

## 16. NITRO AND NITROSO COMPOUNDS

- 2991 Fu, C.-G. and Xu, H.-D.: High-performance liquid chromatography with post-column chemiluminescence detection for simultaneous determination of trace N-nitrosamines and corresponding secondary amines in groundwater. *Analyst (Cambridge)*, 120 (1995) 1147-1151.

See also 2653, 2796.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

## 17a. Amines and polyamines

- 2992 Cheng, F.-C. and Kuo, J.-S.: High-performance liquid chromatographic analysis with electrochemical detection of biogenic amines using microbore columns. *J. Chromatogr. B*, 665 (1995) 1-13 - a review with 141 refs.
- 2993 Dobberpuhl, D.A. and Johnson, D.C.: Pulsed electrochemical detection of alkanolamines separated by multimodal high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 391-398.
- 2994 Gennaro, M.C., Giacosa, D. and Abrigo, C.: The role of pH of the mobile-phase in ion-interaction RP-HPLC. *J. Liq. Chromatogr.*, 17 (1994) 4365-4380.
- 2995 Hayashi, T., Amino, M., Uchida, G. and Sato, M.: High-performance liquid chromatographic determination of primary aromatic amines in urine after derivatization to an azo dye with 2-aminoanthracene. *J. Chromatogr. B*, 665 (1995) 209-212.
- 2996 Hermann, K., Frank, G. and Ring, J.: High performance liquid chromatography for the separation of histamine, its precursor, and metabolites: application to biological samples. *J. Liq. Chromatogr.*, 18 (1995) 189-204.
- 2997 Hiramatsu, K., Sugimoto, M., Kamei, S., Hoshino, M., Kinoshita, K., Iwasaki, K. and Kawakita, M.: Determination of amounts of polyamines excreted in urine: demonstration of N<sup>1</sup>,N<sup>8</sup>-diacetylspermidine and N<sup>1</sup>,N<sup>12</sup>-diacetylspermine as components commonly occurring in normal human urine. *J. Biochem. (Tokyo)*, 117 (1995) 107-112.
- 2998 Huhn, G., Mattusch, J. and Schulz, H.: Determination of polyamines in biological materials by HPLC with 9-fluorenylmethyl chloroformate precolumn derivatization. *Fresenius J. Anal. Chem.*, 351 (1995) 563-566.
- 2999 Iniguez Crespo, M. and Vazquez Lasa, B.: Determination of biogenic amines and other amines in wine by an optimized HPLC method with polarity gradient elution. *Am. J. Enol. Vitic.*, 45 (1994) 460-463; *C.A.*, 122 (1995) 158852b.

- 3000 Lu, C.-S. and Huang, S.-D.: Trace determination of aromatic amines or phenolic compounds in dyestuffs by high-performance liquid chromatography with on-line preconcentration. *J. Chromatogr. A*, 696 (1995) 201-208.
- 3001 Marcé, M., Brown, D.S., Capell, T., Figueras, X. and Tiburcio, A.F.: Rapid high-performance liquid chromatographic method for the quantitation of polyamines as their dansyl derivatives: application to plant and animal tissues. *J. Chromatogr. B*, 666 (1995) 329-335.
- 3002 Qi, X. and Baldwin, R.P.: Liquid chromatography/electrochemical detection of hydroxylamines by oxidation at a cobalt phthalocyanine chemically modified electrode. *Electroanalysis (N.Y.)*, 6 (1994) 353-360; *C.A.*, 122 (1995) 150578w.
- 3003 Skarping, G., Dalene, M. and Brunmark, P.: Liquid chromatography and mass spectrometry determination of aromatic amines in hydrolysed urine from workers exposed to thermal degradation products of polyurethane. *Chromatographia*, 39 (1994) 619-623.
- 3004 Tang, Q., Zhuang, L., Xu, X., Shi, Y. and Li, S.: (Measurement of urinary polyamines by high performance liquid chromatography with electrochemical detection). *Sepu*, 12 (1994) 431-432; *C.A.*, 122 (1995) 50423y.
- 3005 Tomaselli, M., Cozzolino, A., Calemme, I. and Stasio, A.: (Determination of benzidine and other aromatic amines in dyes and dyed leathers). *Cuoio, Pelli, Mater. Concianti*, 69 (1993) 253-263; *C.A.*, 122 (1995) 190912g.
- 3006 Wintersteiger, R., Barary, M.H., El-Yazbi, F.A., Sabry, S.M. and Wahbi, A.-A.M.: Reversed phase-liquid chromatography of primary and secondary aliphatic amines after derivatization combined with electrochemical detection. *Anal. Chim. Acta*, 306 (1995) 273-283.
- 3007 Yamazaki, S., Ozaki, K., Saito, K. and Tanimura, T.: Enantiomeric separation of  $\beta$ -amino alcohols with a primary or tertiary amine moiety by reversed phase liquid chromatography with a chiral mobile phase containing copper(II) and (R)-mandelic acid. *J. High Resolut. Chromatogr.*, 18 (1995) 68-70.
- See also 2566, 2608, 2636, 2653, 3037, 3054, 3616.
- 17b. *Catecholamines and their metabolites*
- 3008 Coward, S., Boa, F.G. and Sherwood, R.A.: Sulfasalazine interference with HPLC assay of 5-hydroxyindole-3-acetic acid. *Clin. Chem. (Washington)*, 41 (1995) 765-766.
- 3009 Dryhurst, G.: Experimental techniques to measure low molecular weight neurotransmitters and their precursors and metabolites. *NATO ASI Ser., Ser. A*, 267(Bioelectrochemistry IV) (1994) 171-193; *C.A.*, 122 (1995) 151453p - a review with 41 refs.
- 3010 Elkihel, L., Grosclaude, J.M., Nembo, B., Jacquot, C., Gayral, P., Goudey-Perriere, F., Letourneux, Y. and Barreteau, H.: Synthesis and determination of N-acetyloctopamine by HPLC with electrochemical detection. Bioassay in *Nippostrongylus brasiliensis* (nematoda). *J. Liq. Chromatogr.*, 17 (1994) 3421-3435.
- 3011 Feier, U. and Goetsch, P.H.: Interlaboratory studies on precision characteristics of analytical methods. Determination of biogenic amines in fish and fish products - HPLC method. *Arch. Lebensmittelhyg.*, 44 (1993) 134-135; *C.A.*, 122 (1995) 131262x.
- 3012 Kirschbaum, J., Luckas, B. and Beinert, W.D.: (Application of automatic precolumn derivatization with 9-fluorenylmethyl chloroformate (FMOC-C1) for the determination of biogenic amines and amino acids in food by HPLC). *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 224-228; *C.A.*, 122 (1995) 29955p.
- 3013 Kirschbaum, J., Luckas, B. and Beinert, W.D.: HPLC analysis of biogenic amines and amino acids in food after automatic precolumn derivatization with 9-fluorenylmethyl chloroformate. *Am. Lab. (Shelton)*, 26 (1994) 28C-28D, 28F, 28H; *C.A.*, 122 (1995) 104168q.
- 3014 Murai, S., Saito, H., Masuda, Y. and Itoh, T.: An improved method for the rapid assay of brain monoaminergic makers using HPLC coupled with coulometric detection. *Biog. Amines*, 10 (1994) 499-508; *C.A.*, 122 (1995) 26991m.
- 3015 Radjaipour, M. and Raster, H.: Separation of urine-catecholamines and their abundant metabolites by uniform HPLC-conditions and fluorimetric detection. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A28.
- 3016 Sanchis Malloles, J.M., Torres Lapasio, J.R., Villanueva Camanas, R.M. and Ramis-Ramos, G.: Determination of catecholamines in urine by micellar liquid chromatography with coulometric detection. *Chromatographia*, 39 (1994) 591-596.
- For additional information see *C.A.*:  
122 (1995) 204407k.
- See also 2992, 2999.
- 17c. *Urea and guanidine derivatives*
- See 2566.
- 17d. *Other amine derivatives and amides (excl. peptides)*
- See 2643, 2685, 2994.
18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 3017 Bartók, T., Szalai, G., Lőrincz, Z., Börscsök, G. and Sági, F.: High-speed RP-HPLC/FL analysis of amino acids after automated two-step derivatization with  $\alpha$ -phthalaldehyde/3-mercaptopropionic acid and 9-fluorenylmethyl chloroformate. *J. Liq. Chromatogr.*, 17 (1994) 4391-4403.
- 3018 Bianchi, V. and Mazza, L.: Rapid reversed-phase high-performance liquid chromatographic method with double derivatization for the assay of urinary hydroxyproline. *J. Chromatogr. B*, 665 (1995) 295-302.
- 3019 Campos, K.L., Giovanelli, J. and Kaufman, S.: Characteristics of the nitric oxide synthase-catalyzed conversion of arginine to N-hydroxyarginine, the first oxygenation step in the enzymic synthesis of nitric oxide. *J. Biol. Chem.*, 270 (1995) 1721-1728.
- 3020 Carta, G. and Dinerman, A.A.: Displacement chromatography of amino acids: effects of selectivity reversal. *AIChE J.*, 40 (1994) 1618-1628; *C.A.*, 122 (1995) 31898r.

- 3021 Castelain, S., Kamel, S., Picard, C., Desmet, G., Sebert, J.L. and Brazier, M.: A simple and automated HPLC method for determination of total hydroxyproline in urine. Comparison with excretion of pyridinolines. *Clin. Chim. Acta*, 235 (1995) 81-90.
- 3022 Chaves das Neves, H.J.: Amino acids. The new methods in chromatography: GLC and SFC. *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, 1992, (1993) 67-79; C.A., 122 (1995) 79290p - a review with 6 refs.
- 3023 Collar, C. and Martínez-Anaya, M.A.: Influence of the microbial starter and the breadmaking step on the free amino acid profiles of wheat sours, doughs, and breads by reversed-phase high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3437-3460.
- 3024 Daniels, D.H., Joe, F.L., Jr. and Diachenko, G.W.: Determination of free glutamic acid in a variety of foods by high-performance liquid chromatography. *Food Addit. Contam.*, 12 (1995) 21-29; C.A., 122 (1995) 238044e.
- 3025 De Witte, P.A., Cuveele, J.F., Merlevede, W.J. and Vandenheede, J.R.: Analysis of phosphorylhydroxyamino acids present in hydrolyzed cell extracts using dansyl derivatization. *Anal. Biochem.*, 226 (1995) 1-9.
- 3026 Egorova, T.A., Eremin, S.V., Mitsner, B.I., Zvonkova, E.N. and Shvets, V.I.: Isolation of individual amino acids from various microbiological sources using reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 53-62.
- 3027 Evrovski, J., Callaghan, M. and Cole, D.E.C.: Determination of homocysteine by HPLC with pulsed integrated amperometry. *Clin. Chem. (Washington)*, 41 (1995) 757-759.
- 3028 Fujimura, S., Yamamoto, A., Kobayashi, K. and Ishibashi, T.: Comparison of performance of amino acid analysis with HPLC equipped with sodium- and lithium-type columns. *Niigata Dai-gaku Nogakubu Kenkyu Hokoku*, 46 (1994) 71-77; C.A., 122 (1995) 234532q.
- 3029 Gasinska, E.: (Affinity chromatography on special polymer beds loaded with ions of transition metals and its usability in the process of separating amino acids in protein hydrolyzates). *Zesz. Nauk. - Politech. Lodz., Technol. Chem. Spozyw.*, 657 (1994) 83-91; C.A., 122 (1995) 158884p.
- 3030 Gerrits, M.I., Thijssen, J.H.H. and van Rijn, H.J.M.: Determination of pyridinoline and deoxypyridinoline in urine, with special attention to retaining their stability. *Clin. Chem. (Washington)*, 41 (1995) 571-574.
- 3031 Inou, H., Moritani, K., Date, Y., Kohashi, K. and Tsuruta, Y.: Determination of free hydroxyproline and proline in human serum by high-performance liquid chromatography using 4-(5,6-dimethoxy-2-phthalimidinyl)phenyl-sulfonyl chloride as pre-column fluorescent labelling reagent. *Analyst (Cambridge)*, 120 (1995) 1141-1145.
- 3032 Kera, Y., Aoyama, H., Matsumura, H., Hasegawa, A., Nagasaki, H. and Yamada, R.-h.: Presence of free D-glutamate and D-aspartate in rat tissues. *Biochim. Biophys. Acta*, 1243 (1995) 282-286.
- 3033 Knox, J.H. and Wan, W.-H.: Chiral chromatography of amino and hydroxy-acids on surface modified porous graphite. *Chromatographia*, 40 (1995) 9-14.
- 3034 Lau, O.-W. and Mok, C.-S.: Indirect conductimetric detection of amino acids after liquid chromatographic separation. *Anal. Chim. Acta*, 300 (1995) 183-191.
- 3035 Lau, O.-W. and Mok, C.-S.: Indirect conductometric detection of amino acids after liquid chromatographic separation. Part II. Determination of monosodium glutamate in foods. *Anal. Chim. Acta*, 302 (1995) 45-52.
- 3036 Ma, Y., Ito, Y., Torok, D.S. and Ziffer, H.: Separation of the 2- and 6-nitro-3-acetamido-4-chlorobenzoic acid precursors of a potent hydroxy-anthranilic acid oxygenase inhibitor by pH-zone-refining-counter-current chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3507-3517.
- 3037 McAllister, T.A., Samuels, S.E., Sedgwick, G.W., Fenton, T.W. and Thompson, J.R.: High-performance liquid chromatographic analysis of  $\beta$ -phenylethylamine for the estimation of *in vivo* protein synthesis. *J. Chromatogr. B*, 666 (1995) 336-341.
- 3038 Michalsky, M., Stepan, J.J., Wilczek, H., Formankova, J. and Moro, L.: Galactosyl hydroxylysine in assessment of Paget's bone disease. *Clin. Chim. Acta*, 234 (1995) 101-108.
- 3039 Miles, D.R.: Amino acid and protein chromatography using immobilized platinum(II) ions. Avail. *Univ. Microfilms Int.*, Order No. DA9424133, 1994, 225 p.; C.A., 122 (1995) 150568t.
- 3040 Misra, I., Charlier, H.A., Jr. and Mizioro, H.M.: Avian cytosolic 3-hydroxy-3-methylglutaryl-CoA synthase: evaluation of the role of cysteines in reaction chemistry. *Biochim. Biophys. Acta*, 1247 (1995) 253-259.
- 3041 Mollé, D., Léonil, J. and Bouhallab, S.: Separation of two-proline-containing peptides by capillary electrophoresis. *Anal. Biochem.*, 225 (1995) 161-162.
- 3042 Mornstad, H., Pfeiffer, H. and Teivens, A.: Estimation of dental age using HPLC-technique to determine the degree of aspartic acid racemization. *J. Forensic Sci.*, 39 (1994) 1425-1431; C.A., 122 (1995) 48621t.
- 3043 Niwa, T., Shiobara, K., Hamada, T., Miyazaki, T., Tsukushi, S., Uema, K. and Tsuzuki, T.: Serum pyridinolines as specific markers of bone resorption in hemodialyzed patients. *Clin. Chim. Acta*, 235 (1995) 33-40.
- 3044 Peaper, D.R. and Jones, H.L.: Determination of S-methylmethionine in malt by alkaline hydrolysis and headspace gas chromatography - a collaborative trial. *J. Inst. Brew.*, 100 (1994) 397-398; C.A., 122 (1995) 158849f.
- 3045 Rawal, N., Rajpurohit, R., Lischwe, M.A., Williams, K.R., Paik, W.K. and Kim, S.: Structural specificity of substrate for S-adenosylmethionine:protein arginine N-methyltransferases. *Biochim. Biophys. Acta*, 1248 (1995) 11-18.
- 3046 Sarvar, G. and Botting, H.G.: Reversed-phase liquid chromatographic determination of hypoglycin A (HG-1) in canned ackee fruit samples. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1175-1179.
- 3047 Sillanpaa, M., Kokkonen, R. and Sihvonen, M.-L.: Determination of EDTA and DTPA as their Fe(III) complexes in pulp and paper mill process and waste waters by liquid chromatography. *Anal. Chim. Acta*, 303 (1995) 187-192.
- 3048 Sottrup-Jensen, L.: A low-pH reverse-phase high-performance liquid chromatography system for analysis of the phenylthiohydantoins of S-carboxymethylcysteine and S-carboxyamidomethylcysteine. *Anal. Biochem.*, 225 (1995) 187-188.
- 3049 Sprongl, L.: Determination of carnitine by high performance liquid chromatography after derivatization with anthryldiazomethan. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A17.

- 3050 Takeuchi, T. and Miwa, T.: (On-column fluorimetric detection of dansylamine acids with bovine serum albumin in liquid chromatography). *Kuromatogurafi*, 15 (1994) 236-237; C.A., 122 (1995) 127788f.
- 3051 Te Poele-Pothoff, M.T.W.B., van den Berg, M., Franken, D.G., Boers, G.H.J., Jakobs, C., de Kroon, I.F.I., Eskes, T.K.A.B., Trijbels, J.M.F. and Blom, H.J.: Three different methods for the determination of total homocysteine in plasma. *Ann. Clin. Biochem.*, 32 (1995) 218-220; C.A., 122 (1995) 234526r.
- 3052 Uchikura, K. and Kirisawa, M.: (Ru(bpy)<sub>3</sub>)<sup>2+</sup> chemiluminescence detection HPLC for D,L-amino acid). *Kuromatogurafi*, 15 (1994) 232-233; C.A., 122 (1995) 155448p.
- 3053 Van de Merbel, N.C., Zuur, P., Frijlink, M., Holthuis, J.J.M., Lingeman, H. and Brinkman, U.A.T.: Automated monitoring of amino acids during fermentation processes using on-line ultrafiltration and column liquid chromatography: application to fermentation medium improvement. *Anal. Chim. Acta*, 303 (1995) 175-185.
- 3054 Yasuhara, F., Yamaguchi, S., Takeda, M., Ochiai, Y. and Miyano, S.: Gas chromatographic separation of enantiomeric amino acids and amines with  $\alpha$ -methoxy- $\alpha$ -trifluoromethylpropionic acid as a chiral derivatizing agent. *J. Chromatogr. A*, 694 (1995) 227-236.
- For additional information see C.A.:  
122 (1995) 104191s, 182748t.
- See also 2590, 2635, 2653, 2658, 2665, 2732, 2745, 2824, 2994, 3007, 3012, 3176, 3409, 3462, 3465, 3476, 3481, 3489, 3586, 3802, 3811.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 3055 Abdel-Kader, H., Kobylkevich, M.M., Wigman, L.S. and Menon, G.K.: Gradient multiplex liquid chromatographic method for determination of ardacin in bulk chemical and premix formulations. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 289-293.
- 3056 Aguilar, M.-I., Richards, K.L., Round, A.J. and Hearn, M.T.W.: Molecular definition of the retention parameters of peptides separated by RP-HPLC. *Pept. Res.*, 7 (1994) 207-217; C.A., 122 (1995) 161287b.
- 3057 Basiuk, V.A. and Gromovoy, T.Yu.: Free energies of adsorption of dipeptides and 2,5-piperazinediones on silica from neutral aqueous solutions as estimated from high-performance liquid-chromatographic retention data. *Collect. Czech. Chem. Commun.*, 59 (1994) 1721-1728; C.A., 122 (1995) 90309x.
- 3058 Cherkasov, A.V., Nedospasov, V.A., Yakhimovich, A.D. and Nedospasov, A.A.: (ANSA-analysis. VI. Activation, inhibition and interaction of proteases in enzyme mixtures). *Biokhimiya (Moscow)*, 59 (1994) 1574-1588.
- 3059 Congote, L.F.: Increased heparin binding by site directed mutagenesis of a recombinant chimera of bombyxin and insulin-like growth factor II. *Biochim. Biophys. Acta*, 1243 (1995) 538-542.
- 3060 Crimmins, D.L., Saylor, M., Rush, J. and Thoma, R.S.: Facile, *in situ* matrix-assisted laser desorption ionization-mass spectrometry analysis and assignment of disulfide pairings in heteropeptide molecules. *Anal. Biochem.*, 226 (1995) 355-361.
- 3061 De Ceuninck, F., Willeput, J. and Corvol, M.: Purification and characterization of insulin-like growth factor II (IGF II) and an IGF II variant from human placenta. *J. Chromatogr. B*, 666 (1995) 203-214.
- 3062 De Collongue, B., Gosselet, N.M., Sébille, N. and Schoot, B.: Comparative study of different predictive methods of peptides retention time on chromatographic reversed-phase columns. *J. Liq. Chromatogr.*, 17 (1994) 4349-4364.
- 3063 Fainzilber, M., van der Schors, R., Lodder, J.C., Li, K.W., Geraerts, W.P.M. and Kits, K.S.: New sodium channel-blocking conotoxins also affect calcium currents in *Lymnaea* neurons. *Biochemistry*, 34 (1995) 5364-5371.
- 3064 Fialaire, A. and Postaire, E.: Hydrolysis of peptide binding by phosphotungstic acid. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1338-1340.
- 3065 Goetze, J.P. and Rehfeld, J.F.: Displaced chromatographic elution of gastrin-17 in normal human serum. *Scand. J. Clin. Lab. Invest.*, 54 (1994) 673-676; C.A., 122 (1995) 151799f.
- 3066 Hänslér, M., Keilhack, H. and Jakubke, H.-D.: Die Verwendung von Ficin zur Knüpfung von Peptidbindungen im gefrorenen wässrigen System. *Pharmazie*, 50 (1995) 184-187.
- 3067 Hoyer, G.L., Nolan, P.E., Jr., LeDoux, J.H. and Moore, L.A.: Selective stability-indicating high-performance liquid chromatographic assay for recombinant human regular insulin. *J. Chromatogr. A*, 699 (1995) 383-388.
- 3068 Johnston, R.N., Shaw, C., Halton, D.W., Verhaert, P. and Baguna, J.: GYIRFamide: a novel FMRFamide-related peptide (FaRP) from the triclud turbellarian, *Dugesia tigrina*. *Biochem. Biophys. Res. Commun.*, 209 (1995) 689-697.
- 3069 Klyushnichenko, V.E., Yakimov, S.A., Arutyunyan, A.M., Ivanov, A.E., Maltsev, K.V. and Wulfson, A.N.: (Recombinant human insulin. IV. Design and optimization of the system of control by reverse-phase high performance liquid chromatography). *Bioorg. Khim.*, 20 (1994) 1080-1088; C.A., 122 (1995) 79157a.
- 3070 Kos, P., Gorzo, G., Suranyi, G. and Borbely, G.: Simple and efficient method for isolation and measurement of cyanobacterial hepatotoxins by plant tests (*Sinapis alba* L.) *Anal. Biochem.*, 225 (1995) 49-53.
- 3071 Li, K.W., El Filali, Z., van Golen, F.A. and Geraerts, W.P.M.: Identification of a novel amide peptide, GLTPNMNSLFF-NH<sub>2</sub>, involved in the control of vas deferens motility in *Lymnaea stagnalis*. *Eur. J. Biochem.*, 229 (1995) 70-72.
- 3072 Manita, H.: Simple immunochromatographic method and apparatus for semiquantitative assay. *PCT Int. Appl. WO 94 28,415 (Cl. G01N33/543)*, 08 Dec. 1994, JP Appl. 93/131,590, 02 Jun. 1993; 54 pp.; C.A., 122 (1995) 101155x.
- 3073 Mitev, V., le Panse, R., Coulomb, B., Miteva, L. and Houdebine, L.-M.: Epidermal growth factor stimulates mitogen-activated protein kinase by PKC-dependent pathway in human keratinocytes. *Biochem. Biophys. Res. Commun.*, 208 (1995) 245-252.
- 3074 Nedospasov, A.A., Rodina, E.V. and Cherkasov, A.V.: (ANSA-analysis. V. Secondary specificity of peptidases for the structure of detectable groups of aminonaphthalenesulfonamide chromogenic substrates). *Biokhimiya (Moscow)*, 59 (1994) 1560-1573.
- 3075 Nguyen, T.S., Hossain, M. and Do, D.D.: Interference phenomena and non-ideal effects in displacement chromatography: experimental study of dipeptides. *Isol. Purif.*, 2 (1994) 27-44; C.A., 122 (1995) 82037s.

- 3076 Nirenberg, U.: Reversed-phase HPLC. Analytical procedure. *Methods Mol. Biol. (Totowa)*, 36(Peptide Analysis Protocols) (1994) 23-35; C.A., 122 (1995) 75701a.
- 3077 Okuno, S., Kitani, T. and Fujisawa, H.: Full activation of brain calmodulin-dependent protein kinase IV requires phosphorylation of the amino-terminal serine-rich region by calmodulin-dependent protein kinase IV kinase. *J. Biochem. (Tokyo)*, 117 (1995) 686-690.
- 3078 Paroni, R., de Vecchi, E., Cighetti, G., Arcelloni, C., Fermo, I., Grossi, A. and Bonini, P.: HPLC with o-phthalaldehyde precolumn derivatization to measure total, oxidized, and protein-bound glutathione in blood, plasma, and tissue. *Clin. Chem. (Washington)*, 41 (1995) 448-454.
- 3079 Ploemen, J.H.T.M., Wormhoudt, L.W., van Ommen, B., Commandeur, J.N.M., Vermeulen, N.P.E. and van Bladeren, P.J.: Polymorphism in the glutathione conjugation activity of human erythrocytes towards ethylene dibromide and 1,2-epoxy-3-(p-nitrophenoxy)-propane. *Biochim. Biophys. Acta*, 1243 (1995) 469-476.
- 3080 Polo, M.C., Gonzalez de Llano, D. and Ramos, M.: Peptides analysis by HPLC. *Acquis. Recent. Chromatogr. Vin Cours Eur. Form. Contin.*, 1992, (1993) 81-85; C.A., 122 (1995) 75622a - a review with 18 refs.
- 3081 Rissler, K.: Sample preparation, high-performance liquid chromatographic separation and determination of substance P-related peptides. *J. Chromatogr. B*, 665 (1995) 233-270 - a review with 186 refs.
- 3082 Rodina, E.V., Cherkasov, A.V. and Nedospasov, A.A.: (ANSA-analysis. IV. Specificity spectra in characterization of proteases and their mixtures). *Biokhimiya (Moscow)*, 59 (1994) 1544-1559.
- 3083 Roturier, J.M., le Bars, D. and Gripon, J.C.: Separation and identification of hydrophilic peptides in dairy products using FMOc derivatization. *J. Chromatogr. A*, 696 (1995) 209-217.
- 3084 Sereda, T.J., Mant, C.T. and Hodges, R.S.: Selectivity due to conformational differences between helical and non-helical peptides in reversed-phase chromatography. *J. Chromatogr. A*, 695 (1995) 205-221.
- 3085 Shon, K., Grilley, M.M., Marsh, M., Yoshikami, D., Hall, A.R., Kurz, B., Gray, W.R., Imperial, J.S., Hillyard, D.R. and Olivera, B.M.: Purification, characterization, synthesis, and cloning of the lockjaw peptide from *Conus purpurascens* venom. *Biochemistry*, 34 (1995) 4913-4918.
- 3086 Szókán, G., Horváth, J., Almás, M., Saftics, G. and Palócz, A.: Liquid chromatographic analysis and separation of polypeptide components from honey bee venoms. *J. Liq. Chromatogr.*, 17 (1994) 3333-3349.
- 3087 Szókán, G., Majer, Z., Kollát, E., Kajtár, M., Hollósi, M. and Peredy-Kajtár, M.: Chromatographic studies on the racemization of thiopeptides. *J. Liq. Chromatogr.*, 18 (1995) 941-964.
- 3088 Thim, L., Wöldike, H.F., Nielsen, P.F., Christensen, M., Lynch-Devaney, K. and Podolsky, D.K.: Characterization of human and rat intestinal trefoil factor produced in yeast. *Biochemistry*, 34 (1995) 4757-4764.
- 3089 Waldmann, T., Nimmesgern, E., Mitsch, M., Peters, J., Pfeifer, G., Müller, S., Kellermann, J., Engel, A., Hartl, F.-U. and Baumeister, W.: The thermosome of *Thermoplasma acidophilum* and its relationship to the eukaryotic chaperonin TRiC. *Eur. J. Biochem.*, 227 (1995) 848-856.
- 3090 Woodward, C. and Geiser, F.O.: Minimization of baseline disturbances in peptide mapping by means of principal-component multivariate visualization. *J. Chromatogr. A*, 699 (1995) 11-20.
- 3091 Zhao, Q., Sannier, F., Ricart, G. and Piot, J.M.: A rapid detection and identification of hemorphins released from bovine hemoglobin enzymatic hydrolysis by use of HPLC coupled with photodiode array detector. *J. Liq. Chromatogr.*, 18 (1995) 93-103.
- For additional information see C.A.:  
122 (1995) 29801k, 178566q.
- See also 2614, 2629, 2639, 2705, 3109, 3120, 3125, 3199, 3243, 3244, 3387, 3795.
- 18c. *Elucidation of structure of proteins and enzymes*
- 3092 Boots, J.-W.P., van Dongen, W.D., Verheij, H.M., de Haas, G.H., Haverkamp, J. and Slotboom, A.J.: Identification of the active site histidine in *Staphylococcus hyicus* lipase using chemical modification and mass spectrometry. *Biochim. Biophys. Acta*, 1248 (1995) 27-34.
- 3093 Carrea, G., D'Arrigo, P., Piergianni, V., Roncaglio, S., Secundo, F. and Servi, S.: Purification and properties of two phospholipases D from *Streptomyces* sp. *Biochim. Biophys. Acta*, 1255 (1995) 273-279.
- 3094 Chang, L.-S., Kuo, K.-W., Lin, J., Lin, S.-R. and Chang, C.-C.: Analysis of a conformation-independent epitope and a conformational epitope in a protein: a study on cobrotoxin from Taiwan cobra venom. *J. Biochem. (Tokyo)*, 117 (1995) 863-868.
- 3095 Chobert, J.-M., Briand, L., Grinberg, V. and Haertlé, T.: Impact of esterification on the folding and the susceptibility to peptic proteolysis of  $\beta$ -lactoglobulin. *Biochim. Biophys. Acta*, 1248 (1995) 170-176.
- 3096 Giorgianni, F., Beranova, S., Wesdemiotis, C. and Viola, R.E.: Elimination of the sensitivity of L-aspartase to active-site-directed inactivation without alteration of catalytic activity. *Biochemistry*, 34 (1995) 3529-3535.
- 3097 Huang, D.-Y. and Ichikawa, Y.: Identification of essential lysyl and cysteinyl residues, and the amino acid sequence at the substrate-binding site of retinal oxidase. *Biochim. Biophys. Acta*, 1243 (1995) 431-436.
- 3098 Jaquet, K., Fukunaga, K., Miyamoto, E. and Meyer, H.E.: A site phosphorylated in bovine cardiac troponin T by cardiac CaM kinase II. *Biochim. Biophys. Acta*, 1248 (1995) 193-195.
- 3099 Mellet, P., Ermolieff, J. and Bieth, J.G.: Mapping the heparin-binding site of mucus proteinase inhibitor. *Biochemistry*, 34 (1995) 2645-2652.
- 3100 Miller, C.G.: Adsorptive biphasic column technology for protein sequence analysis and protein chemical modification. *Methods (San Diego)*, 6 (1994) 315-333; C.A., 122 (1995) 26994q.
- 3101 Murphy, C.M. and Fenselau, C.: Recognition of the carboxy-terminal peptide in cyanogen bromide digests of proteins. *Anal. Chem.*, 67 (1995) 1644-1645.
- 3102 Ojima, T., Tanaka, H. and Nishita, K.: Amino acid sequence of C-terminal 17 kDa CNBr-fragment of Akazara scallop troponin-I. *J. Biochem. (Tokyo)*, 117 (1995) 158-162.

- 3103 Pederson, J., Filimonova, M., Roepstorff, P. and Biedermann, K.: (Isoforms of *Serratia marcescens* nuclease produced by natural and recombinant strains. Comparative characterization by plasma desorption mass spectrometry). *Biokhimiya (Moscow)*, 160 (1995) 450-461.
- 3104 Suzuki, K., Mizuguchi, M., Gomi, T. and Itagaki, E.: Identification of a lysine residue in the NADH-binding site of salicylate hydroxylase from *Pseudomonas putida* S-1. *J. Biochem. (Tokyo)*, 117 (1995) 579-585.
- 3105 Swoboda, I., Jilek, A., Ferreira, F., Engel, E., Hoffmann-Sommergruber, K., Scheiner, O., Kraft, D., Breiteneder, H., Pittenauer, E., Schmid, E. et al.: Isoforms of Bet v 1, the major birch pollen allergen, analyzed by liquid chromatography, mass spectrometry, and cDNA cloning. *J. Biol. Chem.*, 270 (1995) 2607-2613.
- 3106 Vaintraub, I.A. and Lapteva, N.A.: (Kinetics of one-by-one proteolysis. Ovalbumin proteolysis). *Biokhimiya (Moscow)*, 59 (1994) 1406-1413.
- 3107 Yamashita, T., Yomogida, S., Nagaoka, I. and Saito, K.: Effect of a chemical or proteolytic modification on the biological activity of guinea-pig cationic peptide. *Biochim. Biophys. Acta*, 1243 (1995) 295-299.
- 3108 Young, P.M. and Gorenstein, M.V.: Tryptic mapping by reversed-phase HPLC with photodiode-array detection incorporating the spectral-contrast technique. *LC-GC*, 12 (1994) 832-838; *C.A.*, 122 (1995) 208891m.
- See also 3090, 3176, 3194, 3200, 3206, 3208, 3222, 3318.
19. PROTEINS
- 19a. General techniques
- 3109 Bailey, J.M., Tu, O., Issai, G., Ha, A. and Shively, J.E.: Automated carboxy-terminal sequence analysis of polypeptides containing C-terminal proline. *Anal. Biochem.*, 224 (1995) 588-596.
- 3110 Belenkii, B.G. and Malt'sev, V.G.: High-performance membrane chromatography. *BioTechniques*, 18 (1995) 288-291; *C.A.*, 122 (1995) 127808n.
- 3111 Bollag, D.M.: Ion-exchange chromatography. *Methods Mol. Biol. (Totowa)*, 36 (Peptide Analysis Protocols) (1994) 11-22; *C.A.*, 122 (1995) 75700z.
- 3112 Chang, J.-H. and Han, Y.-P.: (Studies on new packings for high-performance ion-exchange chromatography for the separation of proteins. (II). Synthesis and characteristics of new packings for weak cation-exchange chromatography). *Gaodeng, Xuexiao Huaxue Xuebao*, 15 (1994) 1317-1321; *C.A.*, 122 (1995) 100738c.
- 3113 Corradini, D., Capitani, D. and Cellai, L.: Salt concentration effects in high-performance hydrophobic-interaction chromatography in comparison with NMR of proteins in solution. *J. Liq. Chromatogr.*, 17 (1994) 4335-4347.
- 3114 Feng, W., Jiang, X. and Geng, X.: The influence of urea on the retention of protein in hydrophobic interaction chromatography. *J. Liq. Chromatogr.*, 18 (1995) 217-226.
- 3115 Gerstner, J.A., Morris, J., Hunt, T., Hamilton, R. and Afeyan, N.B.: Rapid ion-exchange displacement chromatography of proteins on perfusive chromatographic supports. *J. Chromatogr. A*, 695 (1995) 195-204.
- 3116 Jacob, L.R., Mack, M. and Rathgeber, G.: Large-scale purification of proteins from egg white by tentacle ion-exchangers. *Am. Biotechnol. Lab.*, 12 (1994) 37; *C.A.*, 122 (1995) 75698e.
- 3117 Johnson, E.M., Berk, D.A., Jain, R.K. and Deen, W.M.: Diffusion and partitioning of proteins in charged agarose gels. *Biophys. J.*, 68 (1995) 1561-1568; *C.A.*, 122 (1995) 209067j.
- 3118 Kabzinski, A.K.M., Tagaki, T. and Tanaka, K.: The application of HPLC-LALLS method for determination of changes in thiolproteins and metallothiopropeins during separations by covalent affinity chromatography in example of metallothioneine. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A22-A23.
- 3119 Kasai, K.-i.: (Intelligent chromatography and electrophoresis as tools for understanding of biomolecules). *Kuromatogurafi*, 15 (1994) 206-209; *C.A.*, 122 (1995) 127787e.
- 3120 Kawakatsu, M., Kotaniguchi, H., Freiser, H.H. and Gooding, K.M.: The effects of pore diameter and ligand chain length on fast liquid chromatography of proteins and peptides. *J. Liq. Chromatogr.*, 18 (1995) 633-647.
- 3121 Kulish, D.M., Klyushnichenko, V.E. and Vul'fson, A.N.: (Separation of proteins by liquid chromatography and capillary electrophoresis). *Zh. Fiz. Khim.*, 68 (1994) 1841-1847; *C.A.*, 122 (1995) 50407w.
- 3122 Kurnik, R.T., Yu, A.W., Blank, G.S., Burton, A.R., Smith, D., Athalye, A.M. and van Reis, R.: Buffer exchange using size exclusion chromatography, countercurrent dialysis, and tangential flow filtration: models development, and industrial application. *Biotechnol. Bioeng.*, 45 (1995) 149-157; *C.A.*, 122 (1995) 170139g.
- 3123 Laubscher, J.C.: (Optimization of HPLC [for protein separations]). *Schweiz. Lab.-Z.*, 51 (1994) 162-166; *C.A.*, 122 (1995) 208797k - a review with no refs.
- 3124 Nakazato, K., Mohammad, J. and Hjerten, S.: Studies of some experimental conditions for the polymerization of the continuous beads (cross-linked polyacrylamide gels) in order to improve their chromatographic properties. *Chromatographia*, 39 (1994) 655-662.
- 3125 Narayanan, S.R.: Application of liquid chromatography to the purification of proteins and peptides. *Prog. Pharm. Biomed. Anal.*, 1 (1994) 279-312; *C.A.*, 122 (1995) 234471u.
- 3126 Oscarsson, S.: Influence of salts on protein interactions at interfaces of amphiphilic polymers and adsorbents. *J. Chromatogr. B*, 666 (1995) 21-31.
- 3127 Patrickios, C.S., Gadam, S.D., Cramer, S.M., Hertler, W.R. and Hatton, T.A.: Block methacrylic polyampholytes as protein displacers in ion-exchange chromatography. *Biotechnol. Prog.*, 11 (1995) 33-38; *C.A.*, 122 (1995) 100732w.
- 3128 Tauer, C., Buchacher, A. and Jungbauer, A.: DNA clearance in chromatography of proteins, exemplified by affinity chromatography. *J. Biochem. Biophys. Methods*, 30 (1995) 75-78; *C.A.*, 122 (1995) 182347m.
- 3129 Uversky, V.N.: Gel-permeation chromatography as a unique instrument for quantitative and qualitative analysis of protein denaturation and unfolding. *Int. J. Bio-Chromatogr.*, 1 (1994) 103-114; *C.A.*, 122 (1995) 50340u - a review with 45 refs.

- 3130 Yu, Q. and Wang, F.-Y.: Optimization of ionic strength gradient elution chromatography of proteins. In: *Off. Proc. Comb. Conf., 6th Conf. Asia Pac. Confed. Chem. Eng., 21st Australas. Chem. Eng. Conf., Inst. Eng. Aust., Barton, 1993*, pp. 113/3-118/3; C.A., 122 (1995) 26979p.

For additional information see C.A.:

122 (1995) 26973g, 50428d, 75710c, 182241x, 212099b.

See also 2571, 2629, 2649, 2684, 2758, 3039, 3084, 3218, 3225, 3814.

#### 19b. Proteins of cells, viruses and subcellular particles

- 3131 Cai, S.-J., McAndrew, R.S., Leonard, B.P., Chapman, K.D. and Pidgeon, C.: Rapid purification of cotton seed membrane-bound N-acylphosphatidylethanolamine synthase by immobilized artificial membrane chromatography. *J. Chromatogr. A*, 696 (1995) 49-62.
- 3132 Kandror, K.V., Coderre, L., Pushkin, A.V. and Pilch, P.F.: Comparison of glucose-transporter-containing vesicles from rat fat and muscle tissue: evidence for a unique endosomal compartment. *Biochem. J.*, 307 (1995) 383-390.
- 3133 Krantz, S., Salazar, R., Brandt, R., Kellermann, J. and Lottspeich, F.: Purification and partial amino acid sequencing of a fructosyllysine-specific binding protein from cell membranes of the monocyte-like cell line U937. *Biochim. Biophys. Acta*, 1266 (1995) 109-112.
- 3134 Naylor, D.J., Ryan, M.T., Condron, R., Hoogenraad, N.J. and Høj, P.B.: Affinity-purification and identification of GrpE homologues from mammalian mitochondria. *Biochim. Biophys. Acta*, 1248 (1995) 75-79.
- 3135 Suzuki, T. and Kawakita, M.: Sites of labeling with N-(3-pyrene)maleimide on Ca<sup>2+</sup>-transporting ATPase of the sarcoplasmic reticulum. *J. Biochem. (Tokyo)*, 117 (1995) 881-887.

For additional information see C.A.:

122 (1995) 155383p.

#### 19c. Proteins synthesized by genetic manipulation, monoclonal antibodies

- 3136 Crowe, J., Doebeli, H., Gentz, R., Hochuli, E., Stueber, D. and Henco, K.: 6xHis-Ni-NTA chromatography as a superior technique in recombinant protein expression/purification. *Methods Mol. Biol. (Totowa)*, 31(Protocols for Gene Analysis) (1994) 371-387; C.A., 122 (1995) 97826x.
- 3137 Klyushnichenko, V.E., Wulfson, A.N., Titel, C., Ehwald, R. and Miroshnikov, A.I.: (Separation of recombinant proteins by vesicle chromatography). *Bioorg.Khim.*, 20 (1994) 894-898; C.A., 122 (1995) 100769p.
- 3138 Kwon, K.-S., Lee, S. and Yu, M.-H.: Refolding of  $\alpha$ -antitrypsin expressed as inclusion bodies in *Escherichia coli*: characterization of aggregation. *Biochim. Biophys. Acta*, 1247 (1995) 179-184.

See also 3059, 3067, 3069, 3271.

#### 19d. Microbial and plant proteins

- 3139 Adam, O., Vercellone, A., Paul, F., Monsan, P.F. and Puzo, G.: A nondegradative route for the removal of endotoxin from exopolysaccharides. *Anal. Biochem.*, 225 (1995) 321-327.
- 3140 Ahmed, Md.S., Khan, S., Arakawa, O. and Onoue, Y.: Properties of hemagglutinins newly separated from toxic phytoplankton. *Biochim. Biophys. Acta*, 1243 (1995) 509-512.
- 3141 Feldwisch, J., Zettl, R., Campos, N. and Palme, K.: Identification of a 23 kDa protein from maize photoaffinity-labelled with 5-azido-[7-<sup>3</sup>H]indol-3-ylacetic acid. *Biochem. J.*, 305 (1995) 853-857.
- 3142 Gassenschmidt, U., Jany, K.D., Tauscher, B. and Niebergall, H.: Isolation and characterization of a flocculating protein from *Moringa oleifera* Lam. *Biochim. Biophys. Acta*, 1243 (1995) 477-481.
- 3143 Hamato, N., Koshiba, T., Pham, T.-N., Tatsumi, Y., Nakamura, D., Takano, R., Hayashi, K., Hong, Y.-M. and Hara, S.: Trypsin and elastase inhibitors from bitter melon (*Momordica charantia* Linn.) seeds: purification, amino acid sequences, and inhibitory activities of four new inhibitors. *J. Biochem. (Tokyo)*, 117 (1995) 432-437.
- 3144 Hirayama, C., Sakata, M., Morimoto, S., Tachibe, H., Ihara, H., Ohkuma, K. and Miyachi, Y.: The removal of endotoxins from protein solutions using column packings with aminated poly( $\gamma$ -methyl L-glutamate) spheres. *Chromatographia*, 40 (1995) 33-38.
- 3145 Jekow, P., Fromme, P., Witt, H.T. and Saenger, W.: Photosystem I from *Synechococcus elongatus*: preparation and crystallization of monomers with varying subunit compositions. *Biochim. Biophys. Acta*, 1229 (1995) 115-120.
- 3146 Ko, J.-L., Hsu, C.-I., Lin, R.-H., Kao, C.-L. and Lin, J.Y.: A new fungal immunomodulatory protein, FIP-fve isolated from the edible mushroom, *Flammulina velutipes* and its complete amino acid sequence. *Eur. J. Biochem.*, 228 (1995) 244-249.
- 3147 Nitti, G., Orru, S., Bloch, C., Jr., Morhy, L., Marino, G. and Pucci, P.: Amino acid sequence and disulphide-bridge pattern of three  $\gamma$ -thionins from *Sorghum bicolor*. *Eur. J. Biochem.*, 228 (1995) 250-256.
- 3148 Osman, A.A., Börner, H. and Mothes, T.: Isolation of two gamma-gliadins and determination of their N-terminal amino acid sequence. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A20.
- 3149 Terabe, M., Kojima, S., Taguchi, T., Momose, H. and Miura, K.-i.: A subtilisin inhibitor produced by *Streptomyces bikiniensis* possesses a glutamine residue at reactive site P1. *J. Biochem. (Tokyo)*, 117 (1995) 609-613.
- 3150 Weegels, P.L., Marseille, J.P., Bosveld, P. and Hamer, R.J.: Large-scale separation of gliadins and their bread-making quality. *J. Cereal Sci.*, 20 (1994) 253-264; C.A., 122 (1995) 159163q.
- 3151 Xu, Y.-H., Lu, Z.-Y. and Ihler, G.M.: Purification of deformin, an extracellular protein synthesized by *Bartonella bacilliformis* which causes deformation of erythrocyte membranes. *Biochim. Biophys. Acta*, 1234 (1995) 173-183.
- 3152 Zeidler, W., Egle, C., Ribeiro, S., Wagner, A., Katunin, V., Kreuzer, R., Rodnina, M., Wintermeyer, W. and Sprinzl, M.: Site-directed mutagenesis of *Thermus thermophilus* elongation factor Tu. Replacement of His85, Asp 81 and Arg300. *Eur. J. Biochem.*, 229 (1995) 596-604.



For additional information see C.A.:

122 (1995) 98897h, 207166s.

See also 3070, 3105, 3187, 3239, 3240.

19e. *Proteins of blood, serum and blood cells*

- 3153 Altman, E. and Bundle, D.R.: Polysaccharide affinity columns for purification of lipopolysaccharide-specific murine monoclonal antibodies. *Methods Enzymol.*, 247(Neoglycoconjugates, Pt. B) (1994) 243-253; C.A., 122 (1995) 184908u.
- 3154 Asano, S.: Antibody purification by affinity chromatography. *Jpn. Kokai Tokkyo Koho JP 06,172,399 [94,172,399]* (Cl. CO7K15/14), 21 Jun. 1994, Appl. 92/325,416, 04 Dec. 1992; 4 pp.; C.A., 122 (1995) 130998e.
- 3155 Brumeanu, T.D., Zaghouani, H. and Bona, C.: Purification of antigenized immunoglobulins derivatized with monomethoxy-polyethylene glycol. *J. Chromatogr. A*, 696 (1995) 219-225.
- 3156 Chenou, C. and Fick, M.: (Kinetics and modeling of albumin adsorption in ion-exchange chromatography). *Recents Prog. Genie Procedes*, 7(27, Phenomenes Fondamentaux: Developpements Methodologiques) (1993) 37-42; C.A., 122 (1995) 155399y.
- 3157 Chronos, Z.C., Baynes, J.W. and Thorpe, S.R.: Identification of liver endothelial cells as a primary site of IgM catabolism in the rat. *Arch. Biochem. Biophys.*, 319 (1995) 63-73.
- 3158 Denizli, A. and Piskin, E.: DNA-immobilized polyhydroxyethyl-methacrylate microbeads for affinity sorption of human immunoglobulin G and anti-DNA antibodies. *J. Chromatogr. B*, 666 (1995) 215-222.
- 3159 Ernst, A., Helmhold, M., Brunner, C., Pethö-Schramm, A., Armstrong, V.W. and Müller, H.-J.: Identification of two functionally distinct lysine-binding sites in kringle 37 and in kringles 32-36 of human apolipoprotein(a). *J. Biol. Chem.*, 270 (1995) 6227-6234.
- 3160 Garcia de Frutos, P. and Dahlbäck, B.: cDNA structure of rabbit C4b-binding protein  $\alpha$ -chain. Preserved sequence motive in complement regulatory protein modules which bind C4b. *Biochim. Biophys. Acta*, 1261 (1995) 285-289.
- 3161 Higgins, D.A., Cromie, R.L., Liu, S.S., Magor, K.E. and Warr, G.W.: Purification of duck immunoglobulins: an evaluation of protein A and protein G affinity chromatography. *Vet. Immunol. Immunopathol.*, 44 (1995) 169-180; C.A., 122 (1995) 211516y.
- 3162 Ingendoh, A., Nakayama, H., Isobe, T., Ishioka, N., Kanai, M., Seta, K. and Okuyama, T.: (Analysis of human plasma proteins utilizing HPLC-ESI/MS and MALDI/TOFMS). *Kuromatogurafi*, 15 (1994) 260-261; C.A., 122 (1995) 127793d.
- 3163 Jacob, L., Schmitt, E. and Bruemmer, W.: Purification of antibodies by new chromatography techniques. *Am. Biotechnol. Lab.*, 12 (1994) 44-45; C.A., 122 (1995) 50390k.
- 3164 Kent, U.M.: Purification of antibodies using affinity chromatography. *Methods Mol. Biol. (Totowa)*, 34(Immunocytochemical Methods and Protocols) (1994) 29-35; C.A., 122 (1995) 103414e.
- 3165 Kent, U.M.: Purification of antibodies using ion-exchange chromatography. *Methods Mol. Biol. (Totowa)*, 34(Immunocytochemical Methods and Protocols) (1994) 23-27; C.A., 122 (1995) 103413d.
- 3166 Kuroma, T., Kurata, S. and Natori, S.: Molecular characterization of an insect transferrin and its selective incorporation into eggs during oogenesis. *Eur. J. Biochem.*, 228 (1995) 229-235.
- 3167 Litvinovich, S.V., Henschen, A.H., Kriegstein, K.G., Ingham, K.C. and Medved, L.V.: Structural and functional characterization of proteolytic fragments derived from the C-terminal regions of bovine fibrinogen. *Eur. J. Biochem.*, 229 (1995) 605-614.
- 3168 Loun, B. and Hage, D.S.: Characterization of thyroxine-albumin binding using high-performance affinity chromatography. II. Comparison of the binding of thyroxine, triiodothyronines and related compounds at the warfarin and indole sites of human serum albumin. *J. Chromatogr. B*, 665 (1995) 303-314.
- 3169 Martin, S.E.V., Thomas, O.R.T., Owens, R., Hill, C.R. and Dunnill, P.: Engineering and expression of a single-chain antibody for purification by immobilized metal affinity chromatography. In: *ICHEM Res. Event, Two-Day Symp.*, 1, Inst. Chem. Eng., Rugby, 1994, pp. 85-87; C.A., 122 (1995) 29504r.
- 3170 Miyata, T., Funatsu, A. and Kato, H.: Chemical cross-linking of activated coagulation factor VII with soluble tissue factor: calcium ions are not essential for full amidolytic activity of the factor VIIA-tissue factor complex after complex formation. *J. Biochem. (Tokyo)*, 117 (1995) 836-844.
- 3171 Motojima, K. and Goto, S.: Characterization of serum proteins down-regulated by peroxisome proliferators: transient repression of apoE gene expression in the rat liver. *J. Biochem. (Tokyo)*, 117 (1995) 597-602.
- 3172 Nassander, U.K., Steerenberg, P.A., de Jong, W.H., van Overveld, W.O.W.M., te Boekhorst, C.M.E., Poels, L.G., Jap, P.H.K. and Storm, G.: Design of immunoliposomes directed against human ovarian carcinoma. *Biochim. Biophys. Acta*, 1235 (1995) 126-139.
- 3173 Oono, Y.: (Preparation of active blood coagulation factor XII). *Jpn. Kokai Tokkyo Koho JP 06,174,727 [94,174,727]* (Cl. G01N33/86), 24 Jun. 1994, Appl. 92/329,180, 09 Dec. 1992; 5 p.; C.A., 122 (1995) 142481u.
- 3174 Pirie-Shepherd, S.R., Jett, E.A., Andon, N.L. and Pizzo, S.V.: Sialic acid content of plasminogen 2 glycoforms as a regulator of fibrinolytic activity. Isolation, carbohydrate analysis, and kinetic characterization of six glycoforms of plasminogen 2. *J. Biol. Chem.*, 270 (1995) 5877-5881.
- 3175 Tharakan, J.P. and Ayers, V.: Immobilized ligand penetration and capacity in resin-based immunoaffinity purification of factor IX. *Int. J. Bio-Chromatogr.*, 1 (1994) 93-102; C.A., 122 (1995) 50421w.
- 3176 Wang, X., DeVries, A.L. and Cheng, C.-H.: Antifreeze peptide heterogeneity in an antarctic eel pout includes an unusually large major variant comprised of two 7 kDa type III AFPs linked in tandem. *Biochim. Biophys. Acta*, 1247 (1995) 163-172.
- 3177 Yasuda, F., Hayashi, T., Tanitame, K., Nishioka, J. and Suzuki, K.: Molecular cloning and functional characterization of rat plasma protein S. *J. Biochem. (Tokyo)*, 117 (1995) 374-383.

For additional information see C.A.:

122 (1995) 29509w, 50736c, 184909v.

See also 2713, 3138, 3186, 3189, 3192, 3197, 3198, 3229, 3351.

19f. *Structural and muscle proteins*

- 3178 Ahmad, F. and Goldstein, B.J.: Purification, identification and subcellular distribution of three predominant protein-tyrosine phosphatase enzymes in skeletal muscle tissue. *Biochim. Biophys. Acta*, 1248 (1995) 57-69.
- 3179 Bailey, A.J., Sims, T.J., Avery, N.C. and Halligan, E.P.: Non-enzymic glycation of fibrous collagen: reaction products of glucose and ribose. *Biochem. J.*, 305 (1995) 385-390.
- 3180 Di Lisa, F., De Tullio, R., Salamino, F., Barbato, R., Melloni, E., Siliprandi, N., Schiaffino, S. and Pontremoli, S.: Specific degradation of troponin T and I by  $\mu$ -calpain and its modulation by substrate phosphorylation. *Biochem. J.*, 308 (1995) 57-61.
- 3181 Postnikova, G.B.: (Chemical modification of myoglobin by isothiocyanate reagents. Effect of the N-terminal amino group modification on protein conformation). *Biokhimiya (Moscow)*, 59 (1994) 1445-1457.
- 3182 Potter, J.D., Sheng, Z., Pan, B. and Zhao, J.: A direct regulatory role for troponin T and a dual role for troponin C in the  $\text{Ca}^{2+}$  regulation of muscle contraction. *J. Biol. Chem.*, 270 (1995) 2557-2562.
- 3183 Skolysheva, L.K., Smirnova, E.A., Medvedeva, M.V. and Gusev, N.B.: (Isolation and some properties of bovine aorta caldesmon). *Biokhimiya (Moscow)*, 60 (1995) 339-348.
- 3184 Tillet, E., Mann, K., Nischt, R., Pan, T.-c., Chu, M.-L. and Timpl, R.: Recombinant analysis of human  $\alpha 1(\text{XVI})$  collagen. Evidence for processing of the N-terminal globular domain. *Eur. J. Biochem.*, 228 (1995) 160-168.

For additional information see C.A.:  
122 (1995) 159742c.

See also 2719, 3098, 3132, 3185, 3387.

19h. *Chromoproteins and metalloproteins*

- 3185 Baksh, S., Spamer, C., Oikawa, K., McCubbin, W.D., Heilmann, C., Kay, C.M. and Michalak, M.:  $\text{Zn}^{2+}$  binding to cardiac calsequestrin. *Biochem. Biophys. Res. Commun.*, 209 (1995) 310-315.
- 3186 Bissé, E., Huaman-Guillen, P. and Wieland, H.: Chromatographic evaluation of minor hemoglobins: clinical significance of hemoglobin A<sub>1d</sub>, comparison with hemoglobin A<sub>1c</sub>, and possible interferences. *Clin. Chem. (Washington)*, 41 (1995) 658-663.
- 3187 Chan, C., Willis, A.C., Robinson, C.V., Aplin, R.T., Radford, S.E. and Ferguson, S.J.: The complete amino acid sequence confirms the presence of pseudoazurin in *Thiosphaera pantotropa*. *Biochem. J.*, 308 (1995) 585-590.
- 3188 De Pina, K., Navarro, C., McWalter, L., Boxer, D.H., Price, N.C., Kelly, S.M., Mandrand-Berthelot, M.-A. and Wu, L.-F.: Purification and characterization of the periplasmic nickel-binding protein NikA of *Escherichia coli* K12. *Eur. J. Biochem.*, 227 (1995) 857-865.
- 3189 Huaman-Guillen, P., Bissé, E. and Wieland, H.: Determination of carbamylated hemoglobin (CarHb) by HPLC. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A19-A20.
- 3190 Hunziker, P.E., Kaur, P., Wan, M. and Känzig, A.: Primary structures of seven metallothioneins from rabbit tissue. *Biochem. J.*, 306 (1995) 265-270.

- 3191 Lee, W.-Y., Brune, D.C., Lobrutto, R. and Blankenship, R.E.: Isolation, characterization and primary structure of rubredoxin from the photosynthetic bacterium, *Helicobacillus mobilis*. *Arch. Biochem. Biophys.*, 318 (1995) 80-88.
- 3192 Li, M., Lin, Z. and Johnson, M.E.: Photoaffinity labelling of cyanomethaemoglobin with derivatives of tryptophan and 5-bromotryptophan. *Biochem. J.*, 308 (1995) 251-260.
- 3193 Mateescu, M.A., Chakine, R., Roger, S., Atanasiu, R., Yamaguchi, N., Lamumiere, G. and Nadeau, R.: Protection of myocardial tissue against deleterious effects of oxygen free radicals by ceruloplasmin. *Arzneim. Forsch.*, 45 (1995) 476-480.
- 3194 Qian, Z.-Y., Jollès, P., Migliore-Samour, D. and Fiat, A.-M.: Isolation and characterization of sheep lactoferrin, an inhibitor of platelet aggregation and comparison with human lactoferrin. *Biochim. Biophys. Acta*, 1243 (1995) 25-32.
- 3195 Robinson, K., Jones, D., Howell, S., Soneji, Y., Martin, S. and Aitken, A.: Expression and characterization of maize ZBP14, a member of a new family of zinc-binding proteins. *Biochem. J.*, 307 (1995) 267-272.
- 3196 Tashiro, H. and Sone, N.: Preparation and characterization of the hydrophilic Cu<sub>A</sub>-cytochrome c domain of subunit II of cytochrome c oxidase from thermophilic *Bacillus* P53. *J. Biochem. (Tokyo)*, 117 (1995) 521-526.
- 3197 Turpeinen, U., Karjalainen, U. and Stenman, U.-H.: Three assays for glycohemoglobin compared. *Clin. Chem. (Washington)*, 41 (1995) 191-195.
- 3198 Turpeinen, U., Sipilä, I., Anttila, P., Karjalainen, U., Kuronen, B., Kalkkinen, N., Ahola, T. and Stenman, U.-H.: Two  $\alpha$ -chain hemoglobin variants, Hb Broussais and Hb Cemenelum, characterized by cation-exchange HPLC, isoelectric focusing, and peptide sequencing. *Clin. Chem. (Washington)*, 41 (1995) 532-536.

For additional information see C.A.:  
122 (1995) 26987q, 50419b.

See also 3118, 3166, 3313.

19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*

- 3199 Cervenansky, C., Engström, A. and Karlsson, E.: Role of arginine residues for the activity of fasciculins. *Eur. J. Biochem.*, 229 (1995) 270-275.
- 3200 Chu, C.-C., Li, S.-H. and Chen, Y.-H.: Resolution of isotoxins in the  $\beta$ -bungarotoxin family. *J. Chromatogr. A*, 694 (1995) 492-497.
- 3201 Datta, G., Dong, A., Witt, J. and Tu, A.T.: Biochemical characterization of basilase, a fibrinolytic enzyme from *Crotalus basiliscus basiliscus*. *Arch. Biochem. Biophys.*, 317 (1995) 365-373.
- 3202 Escriva, H., Pierce, A., Coddeville, B., Gonzalez, F., Benaissa, A., Leger, D., Wieruszkeski, J.-M., Spik, G. and Pamblanco, M.: Rat mammary-gland transferrin: nucleotide sequence, phylogenetic analysis and glycan structure. *Biochem. J.*, 307 (1995) 47-55.
- 3203 Fainzilber, M., Lodder, J.C., Kits, K.S., Kofman, O., Vinnitsky, I., van Rietschoten, J., Zlotkin, E. and Gordon, D.: A new conotoxin affecting sodium current inactivation interacts with the  $\delta$ -conotoxin receptor site. *J. Biol. Chem.*, 270 (1995) 1123-1129.

- 3204 Fiorucci, L., Erba, F., Falasca, L., Dini, L. and Ascoli, F.: Localization and interaction of bovine pancreatic trypsin inhibitor and trypsin in the granules of bovine mast cells. *Biochim. Biophys. Acta*, 1243 (1995) 407-413.
- 3205 Hay, D.I., Ahern, J.M., Schluckebier, S.K. and Schlesinger, D.H.: Human salivary acidic proline-rich protein polymorphisms and biosynthesis studied by high-performance liquid chromatography. *J. Dent. Res.*, 73 (1994) 1717-1726; *C.A.*, 122 (1995) 182305w.
- 3206 Lochnit, G. and Geyer, R.: Carbohydrate structure analysis of batroxobin, a thrombin-like serine protease from *Bothrops moojeni* venom. *Eur. J. Biochem.*, 228 (1995) 805-816.
- 3207 Machold, J., Weise, C., Utkin, Y.N., Franke, P., Tsetlin, V.I. and Hucho, F.: A new class of photoactivable and cleavable derivatives of neurotoxin II from *Naja naja oxiana*. Synthesis, characterizations, and application for affinity labelling of the nicotinic acetylcholine receptor from *Torpedo californica*. *Eur. J. Biochem.*, 228 (1995) 947-954.
- 3208 Marangoni, S., Toyama, M.H., Arantes, E.C., Giglio, J.R., da Silva, C.A., Carneiro, E.M., Goncalves, A.A. and Oliveira, B.: Amino acid sequence of TsTX-V, an  $\alpha$ -toxin from *Tityus serrulatus* scorpion venom, and its effect on  $K^+$  permeability of  $\beta$ -cells from isolated rat islets of Langerhans. *Biochim. Biophys. Acta*, 1243 (1995) 309-314.
- 3209 Nikai, T., Ohara, A., Komori, Y., Fox, J.W. and Sugihara, H.: Primary structure of a coagulant enzyme, bilineobin, from *Agkistrodon bilineatus* venom. *Arch. Biochem. Biophys.*, 318 (1995) 89-96.
- 3210 Ota, Y., Ido, K., Kimura, K., Fukui, E. and Ikemoto, S.: Electrophoretic analysis of bile proteins from patients with and without gallstones. *Electrophoresis (Weinheim)*, 16 (1995) 402-406.
- 3211 Palumbo, A. and Jackson, I.J.: Peroxidase activity in the ink gland of *Sepia officinalis* and partial nucleotide sequence of a candidate cDNA encoding the enzyme. *Biochim. Biophys. Acta*, 1247 (1995) 173-178.
- 3212 Schulze-Forster, K., Eckert, K. and Maurer, H.R.: The thymus extract Thymex-L potentiates the retinoic acid-induced differentiation of the human myeloid leukemia cell line HL-60. *Biochim. Biophys. Acta*, 1265 (1995) 110-116.
- 3213 Urbanke, W. and Luf, W.: (Identification of  $\alpha$ -lactalbumin and  $\beta$ -lactoglobulin in sheep, goat, and mare milk by reversed-phase HPLC and derivative spectroscopy). *Lebensmittelind. Milchwirtsch.*, 114 (1993) 403-407; *C.A.*, 122 (1995) 131253v.
- 3214 Yu, H., Diamandis, E.P., Monne, M. and Croce, C.M.: Oral contraceptive-induced expression of prostate-specific antigen in the female breast. *J. Biol. Chem.*, 270 (1995) 6615-6618.
- 3215 Zietlow, M.F. and Etzel, M.R.: Evaluation of sulfopropyl ion-exchange membrane cartridges for isolation of proteins from bovine whey. *J. Liq. Chromatogr.*, 18 (1995) 1001-1018.
- See also 3094, 3095.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 3216 Borghini, I., Barja, F., Pometta, D. and James, R.W.: Characterization of subpopulations of lipoprotein particles isolated from human cerebrospinal fluid. *Biochim. Biophys. Acta*, 1255 (1995) 192-200.
- 3217 Chao, D., Severson, D.L., Zwiers, H. and Hollenberg, M.D.: Radiolabelling of bovine myristoylated alanine-rich protein kinase C substrate (MARCKS) in an ADP-ribosylation reaction. *Biochem. Cell Biol.*, 72 (1994) 391-396.
- 3218 Isobe, T., Nakayama, H., Ishioka, N., Kanai, M., Ingendoh, A., Seta, K. and Okuyama, T.: (Analysis of brain proteins utilizing HPLC-ES/MS and MALDI/TOFMS). *Kuromatogurafi*, 15 (1994) 262-263; *C.A.*, 122 (1995) 127794e.
- 3219 Lambrechts, A., van Damme, J., Goethals, M., Vandekerckhove, J. and Ampe, C.: Purification and characterization of bovine profilin II. Actin, poly(L-proline) and inositolphospholipid binding. *Eur. J. Biochem.*, 230 (1995) 281-286.
- 3220 Melloni, E., Sparatore, B., Patrone, M., Pessino, A., Passalacqua, M. and Pontremoli, S.: Identity in molecular structure between "differentiation enhancing factor" of murine erythroleukemia cells and the 30 kD heparin-binding protein of developing rat brain. *Biochem. Biophys. Res. Commun.*, 210 (1995) 82-89.
- 3221 Smith, J.B., Yang, Z., Lin, P., Zaidi, Z., Abbasi, A. and Russell, P.: The complete sequence of human lens  $\gamma$ -S crystallin. *Biochem. J.*, 307 (1995) 407-410.
- 3222 Strub, J.-M., Garcia-Sablone, P., Lonning, K., Taupenot, L., Hubert, P., van Dorsselaer, A., Aunis, D. and Metz-Boutigue M.-H.: Processing of chromogranin B in bovine adrenal medulla. Identification of secretolytin, the endogenous C-terminal fragment of residues 614-626 with antibacterial activity. *Eur. J. Biochem.*, 229 (1995) 356-368.
- See also 3157, 3315.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 3223 Agrez, M.V., Meldrum, C.J., Sim, A.T.R., Aebersold, R.H., Clark, I.M., Cawston, T.E. and Burns, G.F.: A fibroblast elongation factor purified from colon carcinoma cells shares sequence identity with TIMP-1. *Biochem. Biophys. Res. Commun.*, 206 (1995) 590-600.
- 19l. *Specific binding and receptor proteins*
- 3224 Berggren Söderlund, M., Johannesson, G. and Fex, G.: Expression of humal *all-trans*-retinoic acid receptor and its ligand-binding domain in *Escherichia coli*. *Biochem. J.*, 308 (1995) 353-359.
- 3225 Buus, S., Stryhn, A., Winther, K., Kirkby, N. and Pedersen, L.O.: Receptor-ligand interactions measured by an improved spun column chromatography technique. A high efficiency and high throughput size separation method. *Biochim. Biophys. Acta*, 1243 (1995) 453-460.
- 3226 Dostalova, Z., Calvete, J.J., Sanz, L. and Töpfer-Petersen, E.: Boar spermadhesin AWN-1. Oligosaccharide and zona pellucida binding characteristics. *Eur. J. Biochem.*, 230 (1995) 329-336.
- 3227 Dotzlaw, J.E., Ghag, S.K. and Yeh, W.-K.: Process for purification of water-soluble penicillin-binding protein 2a. *Eur. Pat. Appl. EP 627,445 (Cl. C07K15/00)*, 07 Dec. 1994, US Appl. 61,894, 14 May 1993; 14 pp.; *C.A.*, 122 (1995) 128084k.
- 3228 Feldwisch, O., Lamertz, M., Hartmann, E., Feldwisch, J., Palme, K., Jastorff, B. and Jaenicke, L.: Purification and characterization of a cAMP-binding of *Volvox carteri* f. *nagariensis* lyengar. *Eur. J. Biochem.*, 228 (1995) 480-489.

- 3229 Greengard, J.S., Fernandez, J.A., Radtke, K.-P. and Griffin, J.H.: Identification of candidate residues for interaction of protein S with C4b binding protein and activated protein C. *Biochem. J.*, 305 (1995) 397-403.
- 3230 Gregoriou, M., Jones, P.F., Timms, J.F., Yang, J.J., Radford, S.E. and Rees, A.R.: Physicochemical characterization of the cytoplasmic domain of the epidermal growth factor receptor and evidence for conformational changes associated with its activation by ammonium sulphate. *Biochem. J.*, 306 (1995) 667-678.
- 3231 Joseph, S.K., Pierson, S. and Samanta, S.: Trypsin digestion of the inositol trisphosphate receptor: implications for the conformation and domain organization of the proteins. *Biochem. J.*, 307 (1995) 859-865.
- 3232 Karamanos, N.K., Hjerpe, A., Aletras, A., Tseggenidis, T., Anastasiou, E.D. and Antonopoulos, C.A.: Antibodies to three chondroitin sulfate-containing proteoglycans in squid skin recognize hexa- or longer chondroitin oligosaccharides as major antigenic determinants. *Arch. Biochem. Biophys.*, 316 (1995) 100-109.
- 3233 Kuo, K.-W., Chang, L.-S. and Chang, C.-C.: The structural loop II of cobrotoxin is the main binding region for nAChR and epitope in the region is conformation-dependent. *J. Biochem. (Tokyo)*, 117 (1995) 438-442.
- 3234 Langland, J.O., Pettiford, S.M. Jacobs, B.L.: Nucleic acid affinity chromatography: preparation and characterization of double-stranded RNA agarose. *Protein Expression Purif.*, 6 (1995) 25-32; C.A., 122 (1995) 127807m
- 3235 Lesca, P., Peryt, B., Larrieu, G., Alvinerie, M., Galtier, P., Daujat, M., Maurel, P. and Hoogenboom, L.: Evidence for the ligand-independent activation of the AH receptor. *Biochem. Biophys. Res. Commun.*, 209 (1995) 474-482.
- 3236 Ren, L., Chen, H. and Sternberg, E.A.: Tethered bandshift assay and affinity purification of a new DNA-binding protein. *BioTechniques*, 16 (1994) 852-855; C.A., 122 (1995) 75675v.
- 3237 Sasaki, T., Shimazawa, R., Sawada, T., Iijima, T., Fukasawa, H., Shudo, K., Hashimoto, Y. and Iwasaki, S.: Determination of the photoaffinity-labeled site on the ligand-binding domain of retinoic acid receptor  $\alpha$ . *Biochem. Biophys. Res. Commun.*, 207 (1995) 444-451.
- 3238 Tani, A., Lee, S., Oishi, O., Aoyagi, H. and Ohno, M.: Interaction of the fragments characteristic of batenecin 7 with phospholipid bilayers and their antimicrobial activity. *J. Biochem. (Tokyo)*, 117 (1995) 560-565.
- 3239 Tomioka, H. and Sasabe, H.: Isolation of photochemically active archaeobacterial photoreceptor, pharaonis phoborhodopsin from *Natronobacterium pharaonis*. *Biochim. Biophys. Acta*, 1234 (1995) 261-267.
- 3240 Yang, Z. and Yang, Y.: (Cytokinin binding proteins from *Phaseolus vulgaris* hypocotyls). *Zhiwu Xuebao*, 36 (1994) 878-883; C.A., 122 (1995) 235275v.
- 3241 Yu, H. and Diamandis, E.P.: Prostate-specific antigen immunoreactivity in amniotic fluid. *Clin. Chem. (Washington)*, 41 (1995) 204-210.
- 19m. *Urinary proteins*
- 3242 Jäckle-Meyer, I., Szukics, B., Neubauer, K., Metzke, V., Petzoldt, R. and Stolte, H.: Extracellular matrix protein as early markers in diabetic nephropathy. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 211-219.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 3243 Heath, R.L., Barton, P.A., Simpson, R.J., Reid, B.E., Lim, G. and Anderson, M.A.: Characterization of the protease processing sites in a multidomain proteinase inhibitor precursor from *Nicotiana glauca*. *Eur. J. Biochem.*, 230 (1995) 250-257.
- 3244 Jean, F., Basak, A., DiMaio, J., Seidah, N.G. and Lazure, C.: An internally quenched fluorogenic substrate of prohormone convertase 1 and furin leads to a potent prohormone convertase inhibitor. *Biochem. J.*, 307 (1995) 689-695.
- 3245 Levison, P.R., Badger, S.E., Toome, D.W. and Jones, R.M.H.: Strategies in the scale-up of the anion-exchange chromatography of hen egg-white proteins. *Spec. Publ. - R. Soc. Chem.*, 158(Separations for Biotechnology 3) (1994) 267-272; C.A., 122 (1995) 131366j.
- 3246 Pinto, R.A., Hawgood, S., Clements, J.A., Benson, B.J., Naidu, A., Hamilton, R.L. and Wright, J.R.: Association of surfactant protein C with isolated alveolar type II cells. *Biochim. Biophys. Acta*, 1255 (1995) 16-22.
- 3247 Ribeiro, J.M.C., Schneider, M. and Guimaraes, J.A.: Purification and characterization of prolixin 5 (nitroprolin 2), the salivary anticoagulant of the blood-sucking bug *Rhodnius prolixus*. *Biochem. J.*, 308 (1995) 243-249.
- 3248 Rodgers, K.J., Melrose, J. and Ghosh, P.: Biotin-labeled potato chymotrypsin inhibitor-1: a useful probe to the detection and quantitation of chymotrypsin-like serine proteinases on western blots and its application in the detection of a serine proteinase synthesized by articular chondrocytes. *Anal. Biochem.*, 227 (1995) 129-134.
- 3249 Subramanian, N. and Adiga, P.R.: Simultaneous purification of biotin-binding proteins-I and -II from chicken egg yolk and their characterization. *Biochem. J.*, 308 (1995) 573-577.
- See also 3099, 3106, 3143, 3238.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 3250 Rudometova, N.V., Glazova, N.V. and Dmitrenko, L.V.: (Use of sorption-chromatographic methods for purification and analysis of enzyme preparations). *Zh. Fiz. Khim.*, 68 (1994) 1848-1850; C.A., 122 (1995) 54047b.
- 20a. *Oxidoreductases*
- 3251 Almira Correia, M., Yao, K., AllenToff, A.J., Wrighton, S.A. and Thompson, J.A.: Interactions of peroxyquinols with cytochromes P450 2B1, 3A1, and 3A5: influence of the apoprotein on heterolytic versus homolytic O-O bond cleavage. *Arch. Biochem. Biophys.*, 317 (1995) 471-478.
- For additional information see C.A.:  
122 (1995) 101133p.
- See also 2878, 3168, 3249, 3302, 3315, 3828.

- 3252 Avezoux, A., Goodwin, M.G. and Anthony, C.: The role of the novel disulphide ring in the active site of the quinoprotein methanol dehydrogenase from *Methylobacterium extorquens*. *Biochem. J.*, 307 (1995) 735-741.
- 3253 Bagley, F.J., Hirschberger, L.L. and Stipanuk, M.H.: Evaluation and modification of an assay procedure for cysteine dioxygenase activity: high-performance liquid chromatography method for measurement of cysteine sulfinate and demonstration of physiological relevance of cysteine dioxygenase activity in cysteine catabolism. *Anal. Biochem.*, 227 (1995) 40-48.
- 3254 Bartholomew, B.A., Smith, M.J., Long, M.T., Darcy, P.J., Trudgill, P.W. and Hopper, D.J.: Tropine dehydrogenase: purification, some properties and an evaluation of its role in the bacterial metabolism of tropine. *Biochem. J.*, 307 (1995) 603-608.
- 3255 Camadro, J.-M., Matringe, M., Thome, F., Brouillet, N., Mornet, R. and Labbe, P.: Photoaffinity labeling of protoporphyrinogen oxidase, the molecular target of diphenylether-type herbicides. *Eur. J. Biochem.*, 229 (1995) 669-674.
- 3256 Camardella, L., Damonte, G., Carratore, V., Benatti, U., Tonetti, M. and Moneti, G.: Glucose 6-phosphate dehydrogenase from human erythrocytes: identification of N-acetyl-alanine at the N-terminus of the nature protein. *Biochem. Biophys. Res. Commun.*, 207 (1995) 331-338.
- 3257 Chaga, G.: A general method for immobilization of glycoproteins on regenerable immobilized metal-ion carriers: application to glucose oxidase from *Penicillium chrysogenum* and horseradish peroxidase. *Biotechnol. Appl. Biochem.*, 20 (1994) 43-53; *C.A.*, 122 (1995) 154367z.
- 3258 De Stefano, G., Pizzuto, P. and Sciancalepore, V.: (Recovery of lactoperoxidase from acid whey by immobilized copper affinity chromatography). *Ind. Aliment. (Pinerolo)*, 33 (1994) 387-389; *C.A.*, 122 (1995) 49731r.
- 3259 Emanuele, J.J., Heasley, C.J. and Fitzpatrick, P.F.: Purification and characterization of the flavoprotein tryptophan 2-monooxygenase expressed at high levels in *Escherichia coli*. *Arch. Biochem. Biophys.*, 316 (1995) 241-248.
- 3260 Gillam, E.M.J., Guo, Z., Ueng, Y.-F., Yamazaki, H., Cock, I., Reilly, P.E.B., Hooper, W.D. and Guengerich, F.P.: Expression of cytochrome P450 3A5 in *Escherichia coli*: effects of 5'-modification, purification, spectral characterization, reconstitution conditions, and catalytic activities. *Arch. Biochem. Biophys.*, 317 (1995) 374-384.
- 3261 Hallahan, D.L., West, J.M., Wallsgrove, R.M., Smiley, D.W.M., Dawson, G.W., Pickett, J.A. and Hamilton, J.G.C.: Purification and characterization of an acyclic monoterpene primary alcohol:NADP oxidoreductase from catmint (*Nepeta racemosa*). *Arch. Biochem. Biophys.*, 318 (1995) 105-112.
- 3262 Hiromasa, Y., Aso, Y., Yamashita, S. and Aikawa, Y.: Homogeneity of the pyruvate dehydrogenase multienzyme complex from *Bacillus stearothermophilus*. *J. Biochem. (Tokyo)*, 117 (1995) 467-470.
- 3263 Huang, Y.-C., Haselbeck, R.J., McAlister-Henn, L. and Colman, R.F.: N-Ethylmaleimide profiling of yeast NADP-dependent isocitrate dehydrogenase. *Arch. Biochem. Biophys.*, 316 (1995) 485-492.
- 3264 Il'chenko, A.P., Morgunov, I.G., Honeck, H., Mauersberger, S., Vasilkova, N.N. and Müller, H.-G.: (Purification and some properties of alcohol oxidase from the yeast *Yarrowia lipolytica* H-222). *Biokhimiya (Moscow)*, 59 (1994) 1312-1319.
- 3265 Jansson, I., Mole, J.E. and Schenkman, J.B.: The isolation and comparison of multiple forms of CYP-2B from untreated and phenobarbital-treated rabbit liver microsomes. *Arch. Biochem. Biophys.*, 316 (1995) 275-284.
- 3266 Kabzinski, A.K.N. and Paryjzak, T.: The preparation of enzymes electrodes with application of the fast immobilization of enzymes of affinity chromatography supports. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A22.
- 3267 Koga, N., Kikuichi-Nishimura, N., Hara, T., Harada, N., Ishii, Y., Yamada, H., Oguri, K. and Yoshimura, H.: Purification and characterization of a newly identified isoform of cytochrome P450 responsible for 3-hydroxylation of 2,5,2',5'-tetrachlorobiphenyl in hamster liver. *Arch. Biochem. Biophys.*, 317 (1995) 464-470.
- 3268 Lomnitski, L., Sklan, D. and Grossman, S.: Lipoxygenase activity in rat dermis and epidermis: partial purification and characterization. *Biochim. Biophys. Acta*, 1255 (1995) 351-359.
- 3269 Maier, T., Lottspeich, F. and Böck, A.: GTP hydrolysis by HypB is essential for nickel insertion into hydrogenases of *Escherichia coli*. *Eur. J. Biochem.*, 230 (1995) 133-138.
- 3270 Martinez, A., Knappskog, P.M., Olafsdottir, S., Doskeland, A.P., Eiken, H.G., Svebak, R.M., Bozzini, M.L., Apold, J. and Flatmark, T.: Expression of recombinant human phenylalanine hydroxylase as fusion protein in *Escherichia coli* circumvents proteolytic degradation by host cell proteases. Isolation and characterization of the wild-type enzyme. *Biochem. J.*, 306 (1995) 589-597.
- 3271 Notheis, C., Drewke, C. and Leistner, E.: Purification and characterization of the pyridoxol-5'-phosphate:oxygen oxidoreductase (deaminating) from *Escherichia coli*. *Biochim. Biophys. Acta*, 1247 (1995) 265-271.
- 3272 Ōba, K., Ishikawa, S., Nishikawa, M., Mizuno, H. and Yamamoto, T.: Purification and properties of L-galactono- $\gamma$ -lactone dehydrogenase, a key enzyme for ascorbic acid biosynthesis, from sweet potato roots. *J. Biochem. (Tokyo)*, 117 (1995) 120-124.
- 3273 Obata, T., Shibusawa, Y. and Ito, Y.: Purification of soluble monoamine oxidase in rat liver cytosol by countercurrent chromatography. *Chromatographia*, 39 (1994) 624-626.
- 3274 Padiglia, A., Medda, R., Pazzaglia, G., Rescigno, A., Cruciani, E. and Floris, G.: Fractionation and characterization of two forms of peroxidase from *Oryza sativa*. *Prepar. Biochem.*, 25 (1995) 11-19.
- 3275 Peters, D., Frank, R. and Hengstenberg, W.: Lactose-specific enzyme II of the phosphoenolpyruvate-dependent phosphotransferase system of *Staphylococcus aureus*. Purification of the histidine-tagged transmembrane component IICB<sup>46</sup> and its hydrophilic IIB domain by metal-affinity chromatography, and functional characterization. *Eur. J. Biochem.*, 228 (1995) 798-804.
- 3276 Rescigno, A., Sollai, F., Masala, S., Porcu, M.C., Sanjust, E., Rinaldi, A.C., Curreli, N., Grifi, D. and Rinaldi, A.: Purification and characterization of an NAD(P)H:quinone oxidoreductase from *Glycine max* seedlings. *Prepar. Biochem.*, 25 (1995) 57-67.
- 3277 Riveros-Moreno, V., Heffernan, B., Torres, B., Chubb, A., Charles, I. and Moncada, S.: Purification to homogeneity and characterisation of rat brain recombinant nitric oxide synthase. *Eur. J. Biochem.*, 230 (1995) 52-57.

- 3278 Russell, R. and Scopes, R.K.: Use of hydrophobic chromatography for purification of the membrane-located choline dehydrogenase from a *Pseudomonas* strain. *Bioseparation*, 4 (1994) 279-284; *C.A.*, 122 (1995) 127113g.
- 3279 Schilling, B. and Lerch, K.: Amine oxidases from *Aspergillus niger*: identification of a novel flavin-dependent enzyme. *Biochim. Biophys. Acta*, 1243 (1995) 529-537.
- 3280 Schoch, S., Helfrich, M., Wiktorsson, B., Sundqvist, C., Rüdiger, W. and Ryberg, M.: Photoreduction of zinc protoporphorbide with NADPH-protochlorophyllide oxidoreductase from etiolated wheat (*Triticum aestivum* L.). *Eur. J. Biochem.*, 229 (1995) 291-298.
- 3281 Sharma, M.C. and Shapiro, B.H.: Purification and characterization of constituent testosterone 2 $\alpha$ -hydroxylase (cytochrome P450 $_{20}$ ) from mouse liver. *Arch. Biochem. Biophys.*, 316 (1995) 478-484.
- 3282 Simons, B.H., Barnett, P., Vollenbroek, E.G.M., Dekker, H.L., Muijsers, A.O., Messerschmidt, A. and Wever, R.: Primary structure and characterization of the vanadium chloroperoxidase from the fungus *Curvularia inaequalis*. *Biochem. J.*, 229 (1995) 566-574.
- 3283 Sinhababu, A.K., Boehlert, C.C., Gan, L.-S., Yanni, S.B. and Thakker, D.R.: High-performance liquid chromatographic purification, optimization of the assay, and properties of ribonucleoside diphosphate reductase from rabbit bone marrow. *Arch. Biochem. Biophys.*, 317 (1995) 285-291.
- 3284 Soukri, A., Valverde, F., Hafid, N., Elkebbaj, M.S. and Serrano, A.: Characterization of muscle glyceraldehyde-3-phosphate dehydrogenase isoforms from euthermic and induced hibernating *Jaculus orientalis*. *Biochim. Biophys. Acta*, 1243 (1995) 161-168.
- 3285 Swarts, J.C., Aquino, M.A.S., Han, J.-Y., Lam, K.-Y. and Sykes, A.G.: Kinetic studies on the reduction of the tyrosyl radical of the R2 subunit of *E. coli* ribonucleotide reductase. *Biochim. Biophys. Acta*, 1247 (1995) 215-224.
- 3286 Tachibana, M., Onoe, H., Okubo, A. and Yamazaki, S.: Application of micellar electrokinetic chromatography as a novel assay method for dimethyl sulfoxide reductase. *Biosci., Biotechnol., Biochem.*, 59 (1995) 282-284; *C.A.*, 122 (1995) 208205r.
- 3287 Takeuchi, A., Miyamoto, T., Yamaji, K., Masuho, Y., Hayashi, M., Hayashi, H. and Onozaki, K.: A human erythrocyte-derived growth-promoting factor with a wide target cell spectrum: identification as catalase. *Cancer Res.*, 55 (1995) 1586-1589.
- 3288 Wan, L., Gijzen, M. and van Huystee, R.S.: Heterogeneous glycosylation of cationic peanut peroxidase. *Biochem. Cell Biol.*, 72 (1994) 411-417.
- 3289 Wang, H.-t., Liu, W.-y. and Ulbrich, N.: Isolation and characterization of a tyrosinase from the skin of the white silky fowl (*Gallina lanigera*) employing copper saturated diethylaminoethyl-cellulose. *Biochim. Biophys. Acta*, 1243 (1995) 251-255.
- For additional information see *C.A.*:  
122 (1995) 100181r, 127125n.
- See also 3104, 3193.
- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 3290 Bico, P., Erhardt, J., Kaplan, W. and Dirr, H.: Porcine class  $\pi$  glutathione S-transferase: anionic ligand binding and conformational analysis. *Biochim. Biophys. Acta*, 1247 (1995) 225-230.
- 3291 Deigner, H.P., Mato, J.M. and Pajares, M.A.: Study of the rat liver S-adenosylmethionine synthetase active site with 8-azido ATP. *Biochem. J.*, 308 (1995) 565-571.
- 3292 Egaas, E., Falls, J.G., Svendsen, N.O., Ramstad, H., Skaare, J.U. and Dauterman, W.C.: Strain- and sex-specific differences in the glutathione S-transferase class pi in the mouse examined by gradient elution of the glutathione-affinity matrix and reverse-phase high performance liquid chromatography. *Biochim. Biophys. Acta*, 1243 (1995) 256-264.
- 3293 Habuchi, H., Habuchi, O. and Kimata, K.: Purification and characterization of heparan sulfate 6-sulfotransferase from the culture medium of chinese hamster ovary cells. *J. Biol. Chem.*, 270 (1995) 4172-4179.
- 3294 Herold, M., Grimm, R., Heiger, D.N., Bayerbach, S., Fischer, S. and Miller, C.E.: Purification and characterization of aspartate aminotransferase from *Escherichia coli* by chromatographic methods. *Int. J. Bio-Chromatogr.*, 1 (1994) 29-45; *C.A.*, 122 (1995) 154858k.
- 3295 Liu, Y., van Heeswijck, R., Hoj, P. and Hoogenraad, N.: Purification and characterization of ornithine acetyltransferase from *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 228 (1995) 291-296.
- 3296 Palmerini, C.A., Datti, A., Alunni, S., Vabder Elst, I.E. and Orlandino, A.: A fluorescent assay for the determination of UDPGalNAc: Gal $\beta$ 1,3GalNAc-R(GlcNAc to GalNAc) $\beta$ 1,6-N-acetylglucosaminyltransferase activity. *Anal. Biochem.*, 225 (1995) 315-320.
- 3297 Slany, R.K. and Müller, S.O.: tRNA-guanine transglycosylase from bovine liver. Purification of the enzyme to homogeneity and biochemical characterization. *Eur. J. Biochem.*, 230 (1995) 221-228.
- 3298 Stadie, T.R.E., Chai, W., Lawson, A.M., Byfield, P.G.H. and Hanisch, F.-G.: Studies on the order and site specificity of GalNAc transfer to MUC1 tandem repeats by UDP-GalNAc: polypeptide N-acetylgalactosaminyltransferase from milk or mammary carcinoma cells. *Eur. J. Biochem.*, 229 (1995) 140-147.
- 3299 Taniguchi, M., Kobe, A., Kato, M. and Sugiyama, T.: Aspartate aminotransferase isozymes in *Panicum miliaceum* L., an NAD-malic enzyme-type C $_4$  plant: comparisons of enzymatic properties, primary structures, and expression patterns. *Arch. Biochem. Biophys.*, 318 (1995) 295-306.
- 3300 Tsvetnitsky, V., Auch, L., Nicolaou, A. and Gibbons, W.A.: Characterization of phospholipid methylation in rat brain myelin. *Biochem. J.*, 307 (1995) 239-244.
- 3301 Varone, C.L., Canépa, E.T., Llambías, E.B.C. and Grinstein, M.: cAMP regulation of phenobarbital-mediated induction of  $\delta$ -aminolevulinic synthase mRNA in hepatocytes from normal and experimental diabetic rats. *Biochem. Cell Biol.*, 72 (1994) 381-390.
- 3302 Weifold, P.A. and Feldman, D.A.: Fatty acids promote the formation of complexes between choline-phosphate cytidylyltransferase and cytidylyltransferase binding protein. *Arch. Biochem. Biophys.*, 318 (1995) 147-156.

- 3303 Yamamoto, T., Moriwaki, Y., Takahashi, S., Nasako, Y., Yamakita, J.-i., Hiroishi, K. and Higashino, K.: Determination of plasma purine nucleosides phosphorylase activity by high-performance liquid chromatography. *Anal. Biochem.*, 227 (1995) 135-139.
- 3304 Yeh, H.-I., Hsieh, C.-H., Wang, L.-Y., Tsai, S.-P., Hsu, H.-Y. and Tam, M.F.: Mass spectrometric analysis of rat liver cytosolic glutathione S-transferases: modifications are limited to N-terminal processing. *Biochem. J.*, 308 (1995) 69-75.
- See also 3420.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 3305 Asano, Y., Mizumoto, K., Maruyama, T. and Ishihama, A.: Photoaffinity labeling of influenza virus RNA polymerase PB1 subunit with 8-azido GTP. *J. Biochem. (Tokyo)*, 117 (1995) 677-682.
- 3306 Bergwerff, A.A., van Oostrum, J., Kamerling, J.P. and Vliegthart, J.F.G.: The major N-linked carbohydrate chains from human urokinase. The occurrence of 4-O-sulfated, ( $\alpha$ 2-6)-sialylated or ( $\alpha$ 1-3)-fucosylated N-acetylgalactosamine( $\beta$ 1-4)-N-acetylglucosamine elements. *Eur. J. Biochem.*, 228 (1995) 1009-1019.
- 3307 Brunati, A.M., Ruzzene, M., James, P., Guerra, B. and Pinna, L.A.: Hierarchical phosphorylation of a 50-kDa protein by protein tyrosine kinases TPK-IIb and C-Fgr, and its identification as hematopoietic-lineage cell-specific protein. *Eur. J. Biochem.*, 229 (1995) 164-170.
- 3308 Chernov, A.V., Matvienko, N.N., Zheleznyaya, L.A. and Matvienko, N.I.: (A new site-specific endonuclease-methylase from the thermophilic strain of *Bacillus* species LU11). *Biokhimiya (Moscow)*, 59 (1994) 1714-1729.
- 3309 Garg, N., Galey, I.Yu. and Mattiasson, B.: Effect of poly(vinyl alcohol) treatment of the dye-matrix on the chromatography of pyruvate kinase. *Biotechnol. Tech.*, 8 (1994) 645-650; *C.A.*, 122 (1995) 233947s.
- 3310 Hsuan, J.J., Totty, N.F., Truong, O. and Waterfield, M.D.: Purification of phosphatidylinositol 3-kinase by SDA-agarose gel electrophoresis and HPEC. *Protein Pept. Lett.*, 1 (1994) 25-32; *C.A.*, 122 (1995) 154733r.
- 3311 Ipata, P.L., Gualerzi, C., Scolozzi, C., Tozzi, M.G., Trinei, M. and Barsacchi, D.: Occurrence of inosine kinase as a distinct enzyme in *Spirulina platensis*. *Biochem. Biophys. Res. Commun.*, 209 (1995) 547-553.
- 3312 Joshi, B., Morley, S.J., Rgoads, R.E. and Pain, V.M.: Inhibition of protein synthesis by the heme-controlled eIF-2 $\alpha$  kinase leads to the appearance of mRNA-containing 48S complexes that contain eIF-4E but lack methionyl-tRNA<sub>i</sub>. *Eur. J. Biochem.*, 228 (1995) 31-38.
- 3313 Lecocq, R., Lamy, F., Erneux, C. and Dumont, J.E.: Rapid purification and identification of calyphosine, a Ca<sup>2+</sup>-binding protein phosphorylated by protein kinase A. *Biochem. J.*, 306 (1995) 147-151.
- 3314 Price, D.J., Kawakami, Y., Kawakami, T. and Rivnay, B.: Purification of a major tyrosine kinase from RBL-2H3 cells phosphorylating Fc $\epsilon$ RI  $\gamma$ -cytoplasmic domain and identification as the Btk tyrosine kinase. *Biochim. Biophys. Acta*, 1265 (1995) 135-142.
- 3315 Seki, K., Chen, H.-C. and Huang, K.-P.: Dephosphorylation of protein kinase C substrates, neurogranin, neuromodulin, and MARCKS, by calcineurin and protein phosphatases 1 and 2A. *Arch. Biochem. Biophys.*, 316 (1995) 673-679.
- 3316 Shapovalova, N.I., Zheleznyaya, L.A. and Matvienko, N.I.: (A new site-specific endonuclease and methylase from the thermophilic strain of *Bacillus* species KT6). *Biokhimiya (Moscow)*, 59 (1994) 1730-1738.
- 3317 Stathopoulos, C., Kalpaxis, D.L. and Drinas, D.: Partial purification and characterization of RNase P from *Dictyostelium discoideum*. *Eur. J. Biochem.*, 228 (1995) 976-980.
- 3318 Zhang, J., Fukui, T. and Ichikawa, A.: A third type of nucleoside diphosphate kinase from spinach leaves: purification, characterization and amino-acid sequence. *Biochim. Biophys. Acta*, 1248 (1995) 19-26.
- See also 3180.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 3319 Basu, P.S., Batabyal, S.K., Bhattacharya, A. and Datta, T.K.: Cholinesterase activities in cerebrospinal fluid of patients with idiopathic convulsive disorders. *Clin. Chim. Acta*, 235 (1995) 107-112.
- 3320 Beil, S., Kehrl, H., James, P., Staudenmann, W., Cook, A.M., Leisinger, T. and Kertesz, M.A.: Purification and characterization of the arylsulfatase synthesized by *Pseudomonas aeruginosa* PAO during growth in sulfate-free medium and cloning of the arylsulfatase gene (atsA). *Eur. J. Biochem.*, 229 (1995) 385-394.
- 3321 Cal, S., Aparicio, J.F., de Los Reyes-Gavilan, C.G., Nicieza, R.G. and Sanchez, J.: A novel exocyttoplasmic endonuclease from *Streptomyces antibioticus*. *Biochem. J.*, 306 (1995) 93-100.
- 3322 Cernia, E., Ortaggi, G., Soro, S. and Castagnola, M.: Esters of crosslinked poly(vinyl alcohol) with fatty acids: new stationary phases for the separation of lipolytic enzymes. *Tetrahedron Lett.*, 35 (1994) 9051-9054; *C.A.*, 122 (1995) 181711p.
- 3323 Craxton, A., Ali, N. and Shears, S.B.: Comparison of the activities of a multiple inositol polyphosphate phosphatase obtained from several sources: a search for heterogeneity in this enzyme. *Biochem. J.*, 305 (1995) 491-498.
- 3324 Gijon, M.A., Perez, C., Mendez, E. and Sanchez Crespo, M.: Phospholipase A<sub>2</sub> from plasma of patients with septic shock is associated with high-density lipoproteins and C3 anaphylatoxin: some implications for its functional role. *Biochem. J.*, 306 (1995) 167-175.
- 3325 Johansson, F., Sommarin, M. and Larsson, C.: Rapid purification of the plasma membrane H<sup>+</sup>-ATPase in its non-activated form using FPLC. *Physiol. Plant.*, 92 (1994) 389-396; *C.A.*, 122 (1995) 49754a.
- 3326 Koblan, K.S., Schaber, M.D., Edwards, G., Gibbs, J.B. and Pompiano, D.L.: src-Homology 2 (SH2) domain ligation as an allosteric regulator: modulation of phosphoinositide-specific phospholipase C $\gamma$ 1 structure and activity. *Biochem. J.*, 305 (1995) 745-751.

- 3327 Larnkjaer, A., Nykjaer, A., Olivecrona, G., Thogersen, H. and Ostergaard, P.B.: Structure of heparin fragments with high affinity for lipoprotein lipase and inhibition of lipoprotein lipase binding to  $\alpha_2$ -macroglobulin-receptor/low density lipoprotein-receptor-related protein by heparin fragments. *Biochem. J.*, 307 (1995) 205-214.
- 3328 Lindberg, A. and Olivecrona, G.: Lipase evolution: trout, *Xenopus* and chicken have lipoprotein lipase and apolipoprotein C-II-like activity but lack hepatic lipase-like activity. *Biochim. Biophys. Acta*, 1255 (1995) 205-211.
- 3329 McIntosh, J.M., Ghomashchi, F., Gelb, M.H., Dooley, D.J., Stoehr, S.J., Giordani, A.B., Naisbitt, S.R. and Olivera, B.M.: Conodipine-M, a novel phospholipase A<sub>2</sub> isolated from the venom of the marine snail *Conus magus*. *J. Biol. Chem.*, 270 (1995) 3518-3526.
- 3330 Mehrani, H. and Storey, K.B.: Effects of anoxia on protein phosphatase in turtle organs: purification and properties of protein phosphatase type-1 from turtle liver. *Arch. Biochem. Biophys.*, 316 (1995) 836-843.
- 3331 Mingarro, I., González-Navarro, H. and Braco, L.: Direct HPLC monitoring of lipase activity in reverse micellar media. *J. Liq. Chromatogr.*, 18 (1995) 235-244.
- 3332 Moorhead, G., MacKintosh, R.W., Morrice, N., Gallagher, T. and MacKintosh, C.: Purification of type 1 protein (serine/threonine) phosphatases by microcystin-Sepharose affinity chromatography. *FEBS Lett.*, 356 (1994) 46-50; *C.A.*, 122 (1995) 181618p.
- 3333 Pedersen, J., Andersen, J., Roepstorff, P., Filimonova, M. and Biedermann, K.: (Characterization of *Serrata marcescens* nuclease isoforms by electrospray mass spectrometry). *Biokhimiya (Moscow)*, 160 (1995) 462-469.
- 3334 Pozidis, C., Vlatakis, G. and Bouriotis, V.: Sequence-specific DNA affinity chromatography: application of a group specific adsorbent for the isolation of restriction endonucleases. *Prog. Biotechnol.*, 9(CEB6: Proceedings of the 6th European Congress on Biotechnology, 1993, Pt. 1) (1994) 543-546; *C.A.*, 122 (1995) 127098f.
- 3335 Recktenwald, A., Vernet, T., Storer, A.C. and Ziomek, E.: Reduction of strong lipase-polyclonal antibodies binding by limited proteolysis. *Anal. Biochem.*, 226 (1995) 31-34.
- 3336 Rugani, N., Carrière, F., Thim, L., Borgstrom, B. and Sarda, L.: Lipid binding and activating properties of porcine pancreatic colipase split at the Ile<sup>79</sup>-Thr<sup>80</sup> bond. *Biochim. Biophys. Acta*, 1247 (1995) 185-194.
- 3337 Sunaga, H., Sugimoto, H., Nagamachi, Y. and Yamashita, S.: Purification and properties of lysophospholipase isoenzymes from pig gastric mucosa. *Biochem. J.*, 308 (1995) 551-557.
- 3338 Talesa, V., Grauso, M., Giovannini, E., Rosi, G. and Toutant, J.-P.: Solubilization, molecular forms, purification and substrate specificity of two acetylcholinesterases in the medicinal leech (*Hirudo medicinalis*). *Biochem. J.*, 306 (1995) 687-692.
- 3339 Teissère, M., Borel, M., Caillol, B., Nari, J., Gardies, A.M. and Noat, G.: Purification and characterization of a fatty acyl-ester hydrolase from post-germinated sunflower seeds. *Biochim. Biophys. Acta*, 1255 (1995) 105-112.
- 3340 Thomson, F.J. and Clark, M.A.: Purification of a phosphatidic-acid-hydrolysing phospholipase A<sub>2</sub> from rat brain. *Biochem. J.*, 306 (1995) 305-309.
- 3341 Witmer, M.R., Micanovic, R., Tredup, J., Lin, E., Hail, M. and Villafranca, J.J.: Active recombinant human cytosolic phospholipase A<sub>2</sub> is expressed in *Escherichia coli*. *Arch. Biochem. Biophys.*, 318 (1995) 430-438.
- 3342 Yamazaki, Y., Ogawa, Y., Afify, A.S., Kageyama, Y.-i., Okada, T., Okuno, H., Yoshii, Y. and Nose, T.: Difference between cancer cells and the corresponding normal tissue in view of stereoselective hydrolysis of synthetic esters. *Biochim. Biophys. Acta*, 1243 (1995) 300-308.

See also 3092, 3093, 3103, 3275.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.)*

- 3343 Ara, K., Saeki, K., Igarashi, K., Takaiwa, M., Uemura, T., Hagi-hara, H., Kawai, S. and Ito, S.: Purification and characterization of an alkaline amylopullulanase with both  $\alpha$ -1,4 and  $\alpha$ -1,6 hydrolytic activity from alkalophilic *Bacillus* sp. KSM-1378. *Biochim. Biophys. Acta*, 1243 (1995) 315-324.
- 3344 Blaak, H. and Schrempf, H.: Binding and substrate specificities of a *Streptomyces olivaceoviridis* chitinase in comparison with its proteolytically processed form. *Eur. J. Biochem.*, 229 (1995) 132-139.
- 3345 Braithwaite, K.L., Black, G.W., Hazlewood, G.P., Ali, B.R.S. and Gilbert, H.J.: A non-modular endo- $\beta$ -1,4-mannanase from *Pseudomonas fluorescens* subspecies *cellulosa*. *Biochem. J.*, 305 (1995) 1005-1010.
- 3346 Christakopoulos, P., Kekos, D., Macris, B.J., Claeysens, M. and Bhat, M.K.: Purification and characterization of a less randomly acting endo-1,4- $\beta$ -D-glucanase from the culture filtrates of *Fusarium oxysporum*. *Arch. Biochem. Biophys.*, 316 (1995) 428-433.
- 3347 Gottvaldova, M., Hrabova, H. and Kucera, J.: (Amylase isolation by affinity chromatography on immobilized lectin of pea (*Pisum sativum* L.) saturated with beta-cyclodextrin). *Potravin. Vedy*, 12 (1994) 137-142; *C.A.*, 122 (1995) 100171n.
- 3348 Katoh, S. and Terashima, M.: Purification of secreted  $\alpha$ -amylases by immunoaffinity chromatography with cross-reactive antibody. *Appl. Microbiol. Biotechnol.*, 42 (1994) 36-39; *C.A.*, 122 (1995) 181674d.
- 3349 Kucera, J.: (Tandem column bioaffinity separation of amylases, agglutinin and chymotrypsin inhibitor of rye). *Potravin. Vedy*, 12 (1994) 339-350; *C.A.*, 122 (1995) 104189x.
- 3350 Martino, S., Emiliani, C., Orlacchio, A., Hosseini, R. and Stirling, J.L.:  $\beta$ -N-Acetylhexosaminidases A and S have similar sub-cellular distributions in HL-60 cells. *Biochim. Biophys. Acta*, 1243 (1995) 489-495.
- 3351 Moriyama, T., Tozawa, T., Nobuoka, M. and Ikeda, H.: Sialyl salivary-type amylasemia associated with immunoglobulin D-type multiple myeloma. *Clin. Chim. Acta*, 233 (1995) 127-134.
- 3352 Ribeiro, M.G., Pinto, R., Sa Miranda, M.C. and Suzuki, K.: Tay-Sachs disease: intron 7 splice junction mutation in two Portuguese patients. *Biochim. Biophys. Acta*, 1270 (1995) 44-51.
- 3353 Sova, V.V., Elyakova, L.A., Ivanova, E.P., Fedosov, Yu.V. and Mikhailov, V.V.: (Induced  $\beta$ -1,3-glucanase from the marine bacterium *Alteromonas* sp.) *Biokhimiya (Moscow)*, 59 (1994) 1369-1377.
- 3354 Tomizawa, H., Yamada, H., Tanigawa, K. and Imoto, T.: Effects of additives on irreversible inactivation of lysozyme at neutral pH and 100°C. *J. Biochem. (Tokyo)*, 117 (1995) 369-373.



- 3355 Vandersall, A.S., Cameron, R.G., Nairn C.J., III, Yelenosky, G. and Wodzinski, R.J.: Identification, characterization, and partial purification of glucoamylase from *Aspergillus niger* (syn. *A. ficuum*) NRRL 3135. *Prepar. Biochem.*, 25 (1995) 29-55.
- 3356 Wood, P.J., Erfle, J.D., Teather, R.M., Weisz, J. and Miller, S.S.: Comparison of (1→3)(1→4)- $\beta$ -D-glucan-4-glucanohydrolases (E.C. 3.2.1.73) from *Fibrobacter succinogenes* and from *Bacillus subtilis* and use of high-performance anion exchange chromatography in product characterization. *J. Cereal Sci.*, 19 (1994) 65-75; *C.A.*, 122 (1995) 208354p.
- See also 2569, 3371.
- 20f. *Other hydrolases*
- 3357 Akasaki, K., Yoshimoto, H., Nakamura, A., Shiomi, H. and Tsuji, H.: Purification and characterization of a major kyotorphin-hydrolyzing peptidase of rat brain. *J. Biochem. (Tokyo)*, 117 (1995) 897-902.
- 3358 Ardo, Y. and Jonsson, L.: Chromatographic profiling of aminopeptidolytic activities in lactobacilli as a tool for strain identification. *J. Dairy Res.*, 61 (1994) 573-579; *C.A.*, 122 (1995) 29988b.
- 3359 Becker, C., Shutov, A.D., Nong, V.H., Senyuk, V.I., Jung, R., Horstmann, C., Eischer, J., Nielsen, N.C. and Müntz, K.: Purification, cDNA cloning and characterization of proteinase B, an asparagine-specific enolpeptidase from germinating vetch (*Vicia sativa* L.) seeds. *Eur. J. Biochem.*, 228 (1995) 456-462.
- 3360 Belinga, H.F., Steghens, J.P. and Collombel, C.: Firefly luciferase purification using polyethylene glycol and Dymatex Orange A. *J. Chromatogr. A*, 695 (1995) 33-40.
- 3361 Blée, E. and Schuber, F.: Stereocontrolled hydrolysis of the linoleic acid monoepoxide regioisomers catalyzed by soybean epoxide hydrolase. *Eur. J. Biochem.*, 230 (1995) 229-234.
- 3362 Buttle, D.J.: Affinity chromatography of cysteine peptidases. *Methods Enzymol.*, 244 (Proteolytic Enzymes: Serine and Cysteine Peptidases) (1994) 639-648; *C.A.*, 122 (1995) 181495w.
- 3363 Chiba, Y., Midorikawa, T. and Ichishima, E.: Cloning and expression of the carboxypeptidase gene from *Aspergillus saitoi* and determination of the catalytic residues by site-directed mutagenesis. *Biochem. J.*, 308 (1995) 405-409.
- 3364 Clark, I.M., Mitchell, R.E., Posell, L.K., Bigg, H.F., Cawston, T.E. and O'Hare, M.C.: Recombinant porcine collagenase: purification and autolysis. *Arch. Biochem. Biophys.*, 316 (1995) 123-127.
- 3365 De Saizieu, A., Vankan, P. and van Loon, A.P.G.M.: Enzymic characterization of *Bacillus subtilis* GTP cyclohydrolase I. Evidence for a chemical dephosphorylation of dihydroneopterin triphosphate. *Biochem. J.*, 306 (1995) 371-377.
- 3366 Depertthes, D., Gauthier, E.R., Chapdelaine, P., Lazure, C., Tremblay, R.R. and Dubé, J.Y.: Identification of glandular kallikrein in dog pancreas and determination of its tissue distribution. *Biochim. Biophys. Acta*, 1243 (1995) 291-294.
- 3367 Diefenbach-Jagger, H., Brenner, C., Kemp, B.E., Baron, W., McLean, J., Martin, T.J. and Moseley, J.M.: Arg21 is the preferred kexin-cleavage site in parathyroid-hormone-related protein. *Eur. J. Biochem.*, 229 (1995) 91-98.
- 3368 Ding, Z. and Xu, Y.: A rapid sensitive purification of bovine thrombin. *Prepar. Biochem.*, 25 (1995) 21-28.
- 3369 Fujimura, S., Rikimaru, T., Baba, S., Mori, J., Hao, X.-Q., Terada, S. and Kimoto, E.: Purification and characterization of a non-hemorrhagic metalloprotease from *Agkistrodon halys brevicaudus* venom. *Biochim. Biophys. Acta*, 1243 (1995) 94-100.
- 3370 Gary, J.D. and Clarke, S.: Purification and characterization of an isoaspartyl dipeptidase from *Escherichia coli*. *J. Biol. Chem.*, 270 (1995) 4076-4087.
- 3371 Gey, M.H. and Unger, K.K.: Calculation of the molecular masses of two newly synthesized thermostable enzymes isolated from thermophilic microorganisms. *J. Chromatogr. B*, 666 (1995) 188-193.
- 3372 Grigoleit, V., Möder, M. and Reissmann, S.: Studies on pseudopeptidic inhibitors of renin as a potential antihypertensives by HPLC and mass spectrometry. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A25.
- 3373 Hashimoto, T., Yamamoto, Y., Yoshida, Y. and Tagawa, K.: Cleavage of bovine mitochondrial ATPase inhibitor with endopeptidases, and binding of the resulting peptides to the interface between the  $\alpha$ - and  $\beta$ -subunits of  $F_1$ ATPase. *J. Biochem. (Tokyo)*, 117 (1995) 641-647.
- 3374 Hirose, T., Yamasaki, K. and Yamamoto, T.: Irradiation with ultraviolet light in the presence of vanadate increases  $Ca^{2+}$  permeability of the sarcoplasmic reticulum membrane via  $Ca^{2+}$ -ATPase. *J. Biochem. (Tokyo)*, 117 (1995) 324-330.
- 3375 Inaba, K. and Morisawa, M.: Identification of a 10S trypsin-like protease that cross-reacts with anti-proteasome antibody in sea urchin egg jelly. *Arch. Biochem. Biophys.*, 319 (1995) 177-184.
- 3376 Itoh, C. and Nagamatsu, A.: An aminopeptidase activity from porcine kidney that hydrolyzes oxytocin and vasopressin: purification and partial characterization. *Biochim. Biophys. Acta*, 1243 (1995) 203-208.
- 3377 Itoh, Y., Binner, S. and Nagase, H.: Steps involved in activation of the complex of pro-matrix metalloproteinase 2 (progelatinase A) and tissue inhibitor of metalloproteinases (TIMP)-2 and 4-aminophenylmercuric acetate. *Biochem. J.*, 308 (1995) 645-651.
- 3378 Katsumi, Y., Kihara, H., Ochiai, M. and Ashida, M.: A serine protease zymogen in insect plasma. Purification and activation by microbial cell wall components. *Eur. J. Biochem.*, 228 (1995) 870-877.
- 3379 Kim, Y.-T., Muramatsu, T. and Takahashi, K.: Leader peptidase from *Escherichia coli*: overexpression, characterization, and inactivation by modification of tryptophan residues 300 and 310 with N-bromosuccinimide. *J. Biochem. (Tokyo)*, 117 (1995) 535-544.
- 3380 Mäkinen, K.K., Mäkinen, P.-L., Loesche, W.J. and Syed, S.A.: Purification and general properties of an oligopeptidase from *Treponema denticola* ATCC 35405. A human oral spirochete. *Arch. Biochem. Biophys.*, 316 (1995) 689-698.
- 3381 Mars, I. and Monnet, V.: An aminopeptidase P from *Lactococcus lactis* with original specificity. *Biochim. Biophys. Acta*, 1243 (1995) 209-215.
- 3382 Meyer, W., Moll, R., Kath, T. and Schäfer, G.: Purification, cloning, and sequencing of archaeobacterial pyrophosphatase from the extreme thermoacidophile *Sulfolobus acidocaldarius*. *Arch. Biochem. Biophys.*, 319 (1995) 149-156.
- 3383 Moser, T.L., Enghild, J.J., Pizzo, S.V. and Stack, M.S.: Specific binding of urinary-type plasminogen activator (u-PA) to vitronectin and its role in mediating u-PA-dependent adhesion of U937 cells. *Biochem. J.*, 307 (1995) 867-873.

- 3384 Proudfoot, A.E.I., Magnetat, E., Haley, T.M., Maione, T.E. and Wells, T.N.C.: The complete primary structure of glycosylated porcine platelet factor 4. *Eur. J. Biochem.*, 228 (1995) 658-664.
- 3385 Stewart, A.J., Piggott, N.H., May, F.E.B. and Westley, B.R.: Mitogenic activity of procathepsin D purified from conditioned medium of breast-cancer cells by affinity chromatography on pepstatinyl agarose. *Int. J. Cancer*, 57 (1994) 715-718; *C.A.*, 122 (1995) 77980q.
- 3386 Thomas, M.P., Verma, C., Boyd, S.M. and Brocklehurst, K.: The structural origins of the unusual specificities observed in the isolation of chymopapain M and actinidin by covalent chromatography and the lack of inhibition of chymopapain M by cystatin. *Biochem. J.*, 306 (1995) 39-46.
- 3387 Tyagi, S.C. and Simon, S.R.: Elastin-derived peptide binding to a hydrophobic domain on neutrophil elastase. *Biochem. Cell Biol.*, 72 (1994) 419-427.
- 3388 Vergas Romero, C., Neudorfer, I., Mann, K. and Schäfer, W.: Purification and amino acid sequence of aminopeptidase P from pig kidney. *Eur. J. Biochem.*, 229 (1995) 262-269.
- 3389 Yamaguchi, N., Koyama, K., Yamamura, Y., Okui, A., Konishi, E., Katayama, T., Katsuragi, N., Kurihara, T., Nakazato, H. and Imanishi, J.: A novel proteinase, glucagon-degrading enzyme, secreted by a human pancreatic cancer cell line, HPC-YO. *J. Biochem. (Tokyo)*, 117 (1995) 7-10.
- 3390 Zamolodchikova, T.S., Vorotyntseva, T.I. and Antonov, C.K.: Duodenase, a new serine protease of unusual specificity from bovine duodenal mucosa. Purification and properties. *Eur. J. Biochem.*, 227 (1995) 866-872.
- See also 3058, 3074, 3082, 3206.
- 20g. *Lyases*
- 3391 Broux, M., Ravel, S. and Douce, R.: Methionine biosynthesis in higher plants. II. Purification and characterization of cystathionine  $\beta$ -lyase from spinach chloroplasts. *Arch. Biochem. Biophys.*, 316 (1995) 585-595.
- 3392 De Waal, A., Meijer, A.H. and Verpoorte, R.: Strictosidine synthase from *Catharanthus roseus*: purification and characterization of multiple forms. *Biochem. J.*, 306 (1995) 571-580.
- 3393 Euerink, G.J.W., Wolters, D.J. and Dijkhuizen, L.: Prephenate dehydratase of the actinomycete *Amycolatopsis methanolica*: purification and characterization of wild-type and deregulated mutant proteins. *Biochem. J.*, 308 (1995) 313-320.
- 3394 He, Z. and Wiegel, J.: Purification and characterization of an oxygen-sensitive reversible 4-hydroxybenzoate decarboxylase from *Clostridium hydroxybenzoicum*. *Eur. J. Biochem.*, 229 (1995) 77-82.
- 3395 Law, R.D. and Plaxton, W.C.: Purification and characterization of a novel phosphoenolpyruvate carboxylase from banana fruit. *Biochem. J.*, 307 (1995) 807-816.
- 3396 Ravel, S., Droux, M. and Douce, R.: Methionine biosynthesis in higher plants. I. Purification and characterization of cystathionine  $\gamma$ -synthase from spinach chloroplasts. *Arch. Biochem. Biophys.*, 316 (1995) 572-584.
- 3397 Shelton, M.C. and Toone, E.J.: Differential dye-ligand chromatography as a general purification protocol for 2-keto-3-deoxy-6-phosphogluconate aldolases. *Tetrahedron: Asymmetry*, 6 (1995) 207-211; *C.A.*, 122 (1995) 154855g.
- 3398 Zhang, Z., Schofield, C.J., Baldwin, J.E., Thomas, P. and John, P.: Expression, purification and characterization of 1-aminocyclopropane-1-carboxylate oxidase from tomato in *Escherichia coli*. *Biochem. J.*, 307 (1995) 77-85.
- See also 2856, 3096.
- 20h. *Isomerases*
- 3399 Ashiuchi, M., Yoshimura, T., Kitamura, T., Kawata, Y., Nagai, J., Gorlatov, S., Esaki, N. and Soda, K.: *In vivo* effect of GroESL on the folding of glutamate racemase of *Escherichia coli*. *J. Biochem. (Tokyo)*, 117 (1995) 495-498.
- 3400 Huang, Y. and Dennis, D.T.: Histidine residues 139, 363 and 500 are essential for catalytic activity of cofactor-independent phosphoglyceromutase from developing endosperm of the castor plant. *Eur. J. Biochem.*, 229 (1995) 395-402.
- 3401 Saito, M., Hori, K., Kurotsu, T., Kanda, M. and Saito, Y.: Three conserved glycine residues in valine activation of gramicidin S synthetase 2 from *Bacillus brevis*. *J. Biochem. (Tokyo)*, 117 (1995) 276-282.
- 3402 Stein, T., Kluge, B., Vater, J., Franke, P., Otto, A. and Wittmann-Liebold, B.: Gramicidin S synthetase 1 (phenylalanine racemase), a prototype of amino acid racemases containing the cofactor 4'-phosphopantetheine. *Biochemistry*, 34 (1995) 4633-4642.
- 20i. *Ligases*
- See 3131.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 3403 Alexiou, M. and Leese, H.J.: Enzymes of purine salvage and catabolism in the mouse preimplantation embryo measured by high performance liquid chromatography. *J. Reprod. Fertil.*, 101 (1994) 151-158; *C.A.*, 122 (1995) 100153h.
- 3404 Gao, X.-D., Katsumoto, T. and Onodera, K.: Purification and characterization of chitin deacetylase from *Absidia coerulea*. *J. Biochem. (Tokyo)*, 117 (1995) 257-263.
21. **PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS**
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 3405 Bedford, C.T., Hickman, A.D. and Logan, C.J.: Mixed ion-pair high-performance liquid chromatography of uridine 5'-diphospho- $\alpha$ -D-glucuronic acid and its hydrolysis products. *J. Chromatogr. A*, 695 (1995) 123-127.
- 3406 Boulieu, R. and Lenoir, A.: Determination of thiopurine nucleotides in human lung tissue by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 213-216.
- 3407 Chimi, E.N. and Dousa, T.P.: Enzymatic synthesis and degradation of nicotinate adenine dinucleotide phosphate (NAADP), a  $Ca^{2+}$ -releasing agonist, in rat tissues. *Biochem. Biophys. Res. Commun.*, 209 (1995) 167-174.
- 3408 Dolan, J.W.: Questions about ion pairing. *LC-GC*, 12 (1994) 820-824; *C.A.*, 122 (1995) 100761e.

- 3409 Edwards, R.: Determination of S-adenosyl-L-methionine and S-adenosyl-L-homocysteine in plants. *Phytochem. Anal.*, 6 (1995) 25-30; C.A., 122 (1995) 127811h.
- 3410 Fernandez, B., Centeno, M.L., Sanchez-Tames, R. and Rodriguez, A.: Simultaneous analysis of cytokinins, auxins and abscisic acid by combined immunoaffinity chromatography, high-performance liquid chromatography and immunoassay. *Phytochem. Anal.*, 6 (1995) 49-54; C.A., 122 (1995) 127812j.
- 3411 Fischer, J.H.: Specific detection of nucleotides, creatine phosphate, and their derivatives from tissue samples in a simple, isocratic, recycling, low-volume system. *LC-GC Int.*, 8, No. 5 (1995) 254-264.
- 3412 Gajewska, M., Pawinski, T., Dziergowska-Szmidt, A., Grieb, P. and Ryba, M.: Determination of 2-bromo and 2-chloro derivatives of 2'-deoxyadenosine in human blood serum by high performance liquid chromatography. *Acta Pol. Pharm.*, 51 (1994) 109-114; C.A., 122 (1995) 100746d.
- 3413 Irisawa, M. and Komiyama, M.: Hydrolysis of DNA through cooperation of two metal ions: a novel mimic of phosphoesterases. *J. Biochem. (Tokyo)*, 117 (1995) 465-466.
- 3414 Iwai, S., Maeda, M., Shirai, M., Shimada, Y., Osafune, T., Murata, T. and Ohtsuka, E.: Reaction mechanism of T4 endonuclease V determined by analysis using modified oligonucleotide duplexes. *Biochemistry*, 34 (1995) 4601-4609.
- 3415 Krohn, M. and Wagner, R.: A procedure for the rapid preparation of guanosine tetraphosphate (ppGpp) from *Escherichia coli* ribosomes. *Anal. Biochem.*, 225 (1995) 188-190.
- 3416 Kyosseff, Z.N., Drake, R.R., Kyosseva, S.V. and Elbein, A.D.: Synthesis of a new photoaffinity probe, 5-azido-[<sup>32</sup>P]UDP xylose, by UDPglucuronate carboxylase from wheat germ. *Eur. J. Biochem.*, 228 (1995) 109-112.
- 3417 Liljebjelke, K., Adolphson, R., Baker, K., Doong, R.L. and Mohnen, D.: Enzymatic synthesis and purification of uridine diphosphate [<sup>14</sup>C]galacturonic acid: a substrate for pectin biosynthesis. *Anal. Biochem.*, 225 (1995) 296-304.
- 3418 Lou, S.N. and Montag, A.: (Determination of the status of nucleoside compounds in mushrooms). *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 212-217; C.A., 122 (1995) 29956q.
- 3419 Meredith, M., Rabaglia, M. and Metz, S.: Cytosolic biosynthesis of GTP and ATP in normal rat pancreatic islets. *Biochim. Biophys. Acta*, 1266 (1995) 16-22.
- 3420 Meynial, Y., Paquet, V. and Combes, D.: Simultaneous separation of nucleotides and nucleotide sugars using an ion-pair reversed-phase HPLC: application for assaying glycosyltransferase activity. *Anal. Chem.*, 67 (1995) 1627-1631.
- 3421 Rush, J.S. and Waechter, C.J.: Method for the determination of cellular levels of guanosine-5'-diphosphate-mannose based on a weak interaction with concanavalin A at low pH. *Anal. Biochem.*, 224 (1995) 494-501.
- 3422 Schubert, B.: Enzymatic synthesis of ATP analogs and their purification by reverse-phase high-performance liquid chromatography. *Anal. Biochem.*, 226 (1995) 288-292.
- 3423 Shah, G.M., Poirier, D., Duchaine, C., Brochu, G., Desnoyers, S., Laguex, J., Verreault, A., Hofflack, J.-C., Kirkland, J.B. and Poirier, G.G.: Methods for biochemical study of poly(ADP-ribose) metabolism *in vitro* and *in vivo*. *Anal. Biochem.*, 227 (1995) 1-13, - review with some papers cited.
- 3424 Stolk, J.N., de Abreu, R.A., Boerbooms, A.M.T., de Koning, D.G.M., de Graaf, R., Kerstens, P.J.S.M. and van de Putte, L.B.A.: Purine enzyme activities in peripheral blood mononuclear cells: comparison of a new non-radiochemical high-performance liquid chromatography procedure and a radiochemical thin-layer chromatography procedure. *J. Chromatogr. B*, 666 (1995) 33-43.
- 3425 Streeper, R.T., Cotter, R.J., Colvin, M.E., Hilton, J. and Colvin, O.M.: Molecular pharmacology of hepsulfam, NSC 3296801: identification of alkylated nucleosides, alkylation site, and site of DNA cross-linking. *Cancer Res.*, 55 (1995) 1491-1498.
- 3426 Suzuki, J., Knoue, Y. and Suzuki, S.: Changes in the urinary excretion level of 8-hydroxyguanine by exposure to reactive oxygen-generating substances. *Free Radical Biol. Med.*, 18 (1995) 431-436; C.A., 122 (1995) 127810g.
- 3427 Warren, D.J., Andersen, A. and Slordal, L.: Quantitation of 8-thioguanine residues in peripheral blood leukocyte DNA obtained from patients receiving 6-mercaptopurine-based maintenance therapy. *Cancer Res.*, 55 (1995) 1670-1674.
- 3428 Yhang, Y., Lu, H. and Wang, J.: (Determination of nucleotides in food by HPLC). *Shipin Kexue (Beijing)*, (1994) 59-62; C.A., 122 (1995) 79324c.

For additional information see C.A.:

122 (1995) 50384m, 101064s, 150721h, 208913v.

See also 2994, 3466, 3483, 3721, 3727, 3742, 3748.

#### 21b. Nucleic acids, RNA

- 3429 Dewanjee, M.K., Ghafouripour, A.K., Kapadvanjwala, M. and Samy, A.T.: Kinetics of hybridization of mRNA of c-myc oncogene with <sup>111</sup>In-labeled antisense oligodeoxynucleotide probes by high-pressure liquid chromatography. *BioTechniques*, 16 (1994) 844-850; C.A., 122 (1995) 73083b.
- 3430 Feuser, P., Hermann, R., Schorr, J., Colpan, M. and Bastian, H.: Chromatographic purification and separation of nucleic acid mixtures. *Ger. Offen. DE 4,321,904 (Cl. CO7H21/04)*, 12 Jan. 1995, Appl. 01 Jul. 1993; 9 pp.; C.A., 122 (1995) 101132n.
- 3431 Kanduc, D., Bracalello, A. and Mosavi, A.: HPLC fraction of rat liver tRNA. Isolation of tRNA<sup>Glu</sup>, tRNA<sup>Gly</sup>, tRNA<sup>Phe</sup>, and tRNA<sup>Thr</sup>. *Biochem. Arch.*, 10 (1994) 243-248; C.A., 122 (1995) 50388r.
- 3432 Kanduc, D., Bracalello, A., Troccoli, G., de Benedetto, C., Mosavi, A. and Azzarone, A.: Separation of rat liver cytoplasmic tRNAs by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 505-513.
- 3433 Oefner, P.J. and Bonn, G.K.: High-resolution liquid chromatography of nucleic acids. *Am. Lab. (Shelton)*, 26 (1994) 28C, 28D, 28F-28J; C.A., 122 (1995) 155374m.
- 3434 Yamakawa, Y. and Yamada, Y.: (Separation of *E. coli* tRNAs by two dimensional chromatography on hydroxyapatite and hydrophobic interaction chromatography). *Kuromatogurafi*, 15 (1994) 246-249; C.A., 122 (1995) 127790a.

See also 3119, 3413, 3418.

## 21c. Nucleic acids, DNA

- 3435 Douki, T., Voituriez, L. and Cadet, J.: Measurement of pyrimidine (6-4) photoproducts in DNA by a mild acidic hydrolysis-HPLC fluorescence detection assay. *Chem. Res. Toxicol.*, 8 (1995) 244-253; C.A., 122 (1995) 127596s.
- 3436 Milliman, A., Huang, C.-R., Kim, Y. and LeBreton, P.R.: HPLC preparation of highly purified single-stranded M13 DNA. *BioTechniques*, 18 (1995) 308-315; C.A., 122 (1995) 127809p.
- 3437 Oefner, P.J., Huber, C.G., Puchhammer-Stoeckl, E., Umlauf, F., Gruenwald, K., Bonn, G.K. and Kunz, C.: High-performance liquid chromatography for routine analysis of hepatitis C virus cDNA/PCR products. *BioTechniques*, 16 (1994) 898-908; C.A., 122 (1995) 47659z.
- 3438 Sundaram, K.M.S. and Sloane, L.: Liquid chromatographic assay for the separation of single- and double-stranded DNA by using UV and UV diode-array detectors and hydroxylapatite column. *J. Liq. Chromatogr.*, 18 (1995) 925-939.

See also 3119, 3128, 3413, 3418, 3430, 3433.

## 21e. Structural studies on DNA and DNA mapping

- 3439 Ferrero, E.A., Fitch, M.M.M., Moore, P.H. and Wu, K.-K.: Resin-purified digoxigenin-labeled DNA probes: an efficient protocol for mapping and fingerprinting sugarcane. *BioTechniques*, 18 (1995) 266-273; C.A., 122 (1995) 206312z.

See also 2629.

## 22. ALKALOIDS

- 3440 Asano, N., Kato, A., Oseki, K., Kizo, H. and Matsui, K.: Calistegins of *Physalis alkekengi* var. *francheti* (Solanaceae). Structure determination and their glycosidase inhibitory activities. *Eur. J. Biochem.*, 229 (1995) 269-376.
- 3441 Chen, C.-M. and Chang, H.-C.: Determination of berberine in plasma, urine and bile by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 117-123.
- 3442 Draeger, B.: Identification and quantification of calystegines, polyhydroxyl nortropane alkaloids. *Phytochem. Anal.*, 6 (1995) 31-37; C.A., 122 (1995) 155671f.
- 3443 Fernandez, P., Bermejo, A.M., Lopez-Rivadulla, M. and Landeira, A.C.: Determination of colchicine in biological fluids by reverse-phase HPLC: variation of colchicine levels in rats. In: Oliver, J.S. (Editor), *Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol., 26th 1992*, Scott. Acad. Press, Edinburgh, 1992, pp. 240-243; C.A., 122 (1995) 232820h.
- 3444 Gontier, E., Sangwan, B.S. and Barbotin, J.N.: Simultaneous determination of some tropane alkaloids by reversed-phase HPLC of *Datura innoxia* Mill. plant material. *Agro-Food-Ind. Hi-Tech*, 5 (1994) 26-28; C.A., 122 (1995) 182311v.
- 3445 Imamura, T., Nagata, T., Tokunaga, H., Kudo, K., Jitsufuchi, N. and Nakasono, I.: (Forensic toxicological analysis of colchicine in human samples using liquid chromatography-mass spectrometry). *Nippon Iyo Masu Supekutoru Gakkai Koenshu*, 19 (1994) 155-158; C.A., 122 (1995) 74091q.

- 3446 Kuczynski, J.: (Retention of some alkaloids in aqueous chromatographic systems with reversed phases. I. The acetate buffer-methanol system (RP-19). *Acta Pol. Pharm.*, 50 (1993) 131-136; C.A., 122 (1995) 150536f.
- 3447 Kuczynski, J.: (Retention of some alkaloids in aqueous chromatographic systems with reversed phases. II. The phosphate buffer-methanol system with a counterion (RP-18). *Acta Pol. Pharm.*, 50 (1993) 137-141; C.A., 122 (1995) 150537g.
- 3448 Lame, M.W., Jones, A.D., Morin, D., Segall, H.J. and Wilson, D.W.: Biliary excretion of pyrrolic metabolites of [<sup>14</sup>C]monocrotaline in the rat. *Drug Metab. Disp.*, 23 (1995) 422-429.
- 3449 Mwalupindi, A.G. and Warner, I.M.: Determination of theophylline by liquid chromatography with sensitized lanthanide luminescence detection. *Anal. Chim. Acta*, 306 (1995) 49-56.
- 3450 Perez-Martinez, I., Sagrado, S., Medina-Hernández, M.J.: A rapid procedure for the determination of caffeine, theophylline and theobromine in urine by micellar liquid chromatography and direct sample injection. *Anal. Chim. Acta*, 304 (1995) 195-201.
- 3451 Rehwald, A., Sticher, O. and Meier, B.: Trace analysis of Harman alkaloids in *Passiflora incarnata* by reversed-phase high-performance liquid chromatography. *Phytochem. Anal.*, 6 (1995) 96-100; C.A., 122 (1995) 197108h.
- 3452 Reynolds, J. and Albazi, S.J.: Simultaneous determination of nicotine and cotinine in untreated human urine by micellar liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 537-552.
- 3453 Seregina, Ye.V. and Laipanov, A.Kh.: (Use of HPLC in analysis tropane alkaloids). *Farmatsiya (Moscow)*, 43 (1994) 44-47; C.A., 122 (1995) 170327s.
- 3454 Weingarten, B., Wang, H.-Y. and Roberts, D.M.: Determination of codeine in human plasma by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 696 (1995) 83-92.
- 3455 Yevstratova, K.I. and Kudryashov, M.N.: (Analysis of scopolamine from *Datura indiana* seeds). *Farmatsiya (Moscow)*, 43 (1994) 37-44; C.A., 122 (1995) 170326r.

For additional information see C.A.:

122 (1995) 75715h, 75717k, 98906k, 115093n, 115095q, 177547x, 197102b.

See also 3679, 3681, 3688, 3700, 3709, 3782, 3785, 3786, 3789, 3810.

## 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

## 23a. Porphyrins and other pyrroles

- 3456 Beukeveld, G.J.J. and Wolthers, B.G.: Cholestyramine orally administered to patients with erythropoietic protoporphyria results in urinary excretion of protoporphyrin: to be explained by the Herbst-Volkheimer effect? *Clin. Chim. Acta*, 233 (1995) 119-126.
- 3457 Hori, H., Ishitani, O. and Ibusuki, T.: High performance liquid chromatographic isolation and characterization of (pheophytin)mercury(II). *J. Liq. Chromatogr.*, 18 (1995) 779-789.

3458 Hua, Z., Gibson, S.L., Foster, T.H. and Hilf, R.: Effectiveness of  $\delta$ -aminolevulinic acid-induced protoporphyrin as a photosensitizer for photodynamic therapy *in vivo*. *Cancer Res.*, 55 (1995) 1723-1731.

23c. *Indole derivatives and plant hormones (gibberelins)*

3459 Hansen-Moeller, J. and Andersen, J.R.: Boar taint. Analytical alternatives. *Fleischwirtschaft*, 74 (1994) 963-966, 1005-1009; *C.A.*, 122 (1995) 212340y.

3460 Huang, T. and Kissinger, P.T.: Determination of serotonin in microdialyzates using Unijet fused silica microbore columns. *Curr. Sep.*, 13 (1995) 114-118; *C.A.*, 122 (1995) 230953e.

3461 Julliard, J., Maldiney, R., Sotta, B., Miginiac, E., Kerhoas, L. and Einhorn, J.: HPLC-ELISA and MS as complementary techniques to study plant hormone metabolites. *Analysis*, 22 (1994) 483-489; *C.A.*, 122 (1995) 182656m.

3462 Kagedal, B., Konradsson, P., Shibata, T. and Mishima, Y.: High-performance liquid-chromatographic analysis of dopachrome and dihydroxyphenylalanine. *Anal. Biochem.*, 225 (1995) 264-269.

3463 Oppen, C., Wesemann, W., Barka, G., Kerkdijk, H. and Haak, G.: Analysis of serotonin in whole-blood samples - a novel fully automated method. *LC-GC Int.*, 8, No. 2 (1995) 104-106.

For additional information see *C.A.*:  
122 (1995) 237916d.

See also 2992, 3008, 3410.

23e. *Other N-heterocyclic compounds*

3464 Bellstedt, K. and Seidel, J.: Determination of orotic acid in urine by means of HPLC. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A21-A22.

3465 Busse-Grawitz, A., Bissé, E. and Wieland, H.: HPLC method for a rapid and accurate determination of 3-methylhistidine (3-MH). *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A19.

3466 Kock, R., Seitz, S., Delvoux, B. and Greiling, H.: A method for the simultaneous determination of creatinine and uric acid in serum by high-performance liquid chromatography evaluated versus reference method. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 23-29.

3467 Mao, J., Pacheco, C.R., Traficante, D.D. and Rosen, W.: Analysis of benzoquinolines and acridines in a Brazilian diesel oil by particle beam LC/MS and HPLC/UV. *J. Liq. Chromatogr.*, 18 (1995) 903-916.

3468 Mawatari, K.-i., Oshida, K., Iinuma, F. and Watanabe, M.: Determination of quinolinic acid in human urine by liquid chromatography with fluorimetric detection. *Anal. Chim. Acta*, 302 (1995) 179-183.

3469 Scholl, J.P.: An evaluation of column-to-column retention variability for compounds related to panadiplon (U-78,875) using three different reverse-phase HPLC separation modes. *J. Liq. Chromatogr.*, 17 (1994) 3369-3382.

3470 Thienpont, L.M., van Landuyt, K.G., Stöckl, D., Saeyens, W., de Keukeleire, D. and de Leenheer, A.P.: Evaluation of 2-iminoimidazolidin-4-one and thymine as respective internal standards for normal-phase and reversed-phase high-performance liquid chromatographic determination of creatinine in human serum. *J. Chromatogr. B*, 665 (1995) 63-69.

See also 2639, 2795, 3057, 3715, 3798.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

3471 Gasparrini, F., Misiti, D. and Villani, C.: Behaviour of allyl aryl sulfoxides in high-performance liquid chromatography on a chiral stationary phase. *J. Chromatogr. A*, 694 (1995) 163-167.

3472 Haj-Yehia, A.I. and Benet, L.Z.: 2-(4-N-Maleimidophenyl)-6-methoxybenzofuran: a superior derivatizing agent for fluorimetric determination of aliphatic thiols by high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 45-53.

3473 Howard, A.G. and Russell, D.W.: HPLC-FPD instrumentation for the measurement of the atmospheric dimethyl sulfide precursor  $\beta$ -(dimethylsulfonio)propionate. *Anal. Chem.*, 67 (1995) 1293-1295.

3474 Majima, E., Goto, S., Hori, H., Shinohara, Y., Hong, Y.-M. and Terada, H.: Stabilities of the fluorescent SH-reagent eosin-5-maleimide and its adducts with sulfhydryl compounds. *Biochim. Biophys. Acta*, 1243 (1995) 336-342.

3475 Okafo, G.N., Rana, K.K. and Camilleri, P.: Improved separation of diastereoisomers in capillary electrophoresis using a mixture of  $\beta$ -cyclodextrin and sodium taurodeoxycholate. *Chromatographia*, 39 (1994) 627-630.

3476 Winters, R.A., Zukowski, J., Ercal, N., Matthews, R.H. and Spitz, D.R.: Analysis of glutathione, glutathione disulfide, cysteine, homocysteine, and other biological thiols by high-performance liquid chromatography following derivatization by N-(1-pyrenyl)maleimide. *Anal. Biochem.*, 227 (1995) 14-21.

For additional information see *C.A.*:  
122 (1995) 27151z.

See also 2795, 3715, 3836.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

3477 Abidi, S.L. and Mounts, T.L.: Separations of molecular species of phosphatidic acid by high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 365-373.

3478 Altrichter, J., Guse, A.H., Resch, K., Brock, J., Daeron, M. and Hüffel, C.: Phosphatidylinositol hydrolysis and an increase in  $Ca^{2+}$  concentration in the signal-transduction process triggered by murine Fc $\gamma$ RIII are not required for protein kinase C translocation. *Eur. J. Biochem.*, 228 (1995) 587-595.

3479 Barker, C.J., French, P.J., Moore, A.J., Milsson, T., Berggren, P.-O., Bunce, C.M., Kirk, C.J. and Michell, R.H.: Inositol 1,2,3-trisphosphate and inositol 1,2- and/or 2,3-bisphosphate are normal constituents of mammalian cells. *Biochem. J.*, 306 (1995) 557-564.

- 3480 Burbano, C., Muzquiz, M., Osagie, A., Ayet, G. and Cuadrado, C.: Determination of phytate and lower inositol phosphates in Spanish legumes by HPLC methodology. *Food Chem.*, 52 (1995) 321-325; C.A., 122 (1995) 185705n.
- 3481 Chen, S. and Claeys, M.: Approach to the large-scale preparation of highly pure phosphatidylserine from bovine brain. *J. Chromatogr. B*, 666 (1995) 178-182.
- 3482 Edwards, Y.S. and Murray, A.W.: Accumulation of phosphatidylalcohol in cultured cells: use of subcellular fractionation to investigate phospholipase D activity during signal transduction. *Biochem. J.*, 308 (1995) 473-480.
- 3483 Guranowski, A., Starzyńska, E., Sillero, M.A.G. and Sillero, A.: Conversion of adenosine(5') oligophospho(5') adenosines into inosine(5') oligophospho(5') inosines by non-specific adenylate deaminase from the snail *Helix pomatia*. *Biochim. Biophys. Acta*, 1243 (1995) 78-84.
- 3484 Tul'skaya, E.M., Shashkov, A.S., Evtushenko, L.I., Taran, V.V. and Naumova, I.B.: (Cell wall teichoic acids of *Nocardiosis albus* ssp. *albus* DSM 43120). *Biokhimiya (Moscow)*, 60 (1995) 179-186.
- 3485 Van Dijken, P., Lammers, A.A. and van Haastert, P.J.M.: In *Dicystoselium discoideum* inositol 1,3,4,5-tetrakisphosphate is dephosphorylated by a 3-phosphatase and a 1-phosphatase. *Biochem. J.*, 308 (1995) 127-130.
- See also 2926, 2929, 2930, 2931, 2939, 2942, 2945, 2949, 3025, 3077, 3405, 3417, 3419.
26. ORGANOMETALLIC AND RELATED COMPOUNDS
- 26a. *Organometallic compounds*
- 3486 Compañó, R., Granados, M., Leal, C. and Prat, M.D.: Determination of triphenyltin in sea water samples by liquid chromatography with fluorimetric detection. *Anal. Chim. Acta*, 302 (1995) 185-191.
- 3487 Gamoh, K., Ketuly, K.A., Cole, W.J., Brooks, C.J.W. and Anderson, R.A.: Chromatographic mass-spectrometric studies of cyclic 2-(N,N-dimethylaminomethyl)ferroceneboronates and related esters. *Anal. Sci.*, 10 (1994) 705-711; C.A., 122 (1995) 204200n.
- 3488 Hempel, M., Chau, Y.K., Dutka, R.J., McInnis, R., Kwan, K.K. and Liu, D.: Toxicity of organomercury compounds: Bioassay results as a basis for risk assessment. *Analyst (Cambridge)*, 120 (1995) 721-724.
- 3489 Matni, G., Azani, R., van Calsteren, M.R., Bissonnette, M.C. and Blais, J.S.: Determination of free selenomethionine in nutritional supplements by high-performance liquid chromatography coupled with thermochemical hydride generation atomic absorption spectrometry. *Analyst (Cambridge)*, 120 (1995) 395-401.
- 3490 Padró, A., Rubio, R. and Rauret, G.: Germanium speciation by LC-HG-ICP/OES. *Fresenius J. Anal. Chem.*, 351 (1995) 449-453.
- 3491 Spurnar-Lobinska, J., Witte, C., Lobinski, R. and Adams, F.C.: Separation techniques in speciation analysis for organometallic species. *Fresenius J. Anal. Chem.*, 351 (1995) 351-377.
- 3492 Suzuki, K.T., Itoh, M. and Ohmichi, M.: Detection of selenium-containing biological constituents by high-performance liquid chromatography-plasma source mass spectrometry. *J. Chromatogr. B*, 666 (1995) 13-19.
- See also 3854.
- 26c. *Coordination compounds*
- 3493 Kumar, K., Sukumaran, K., Taylor, S., Chang, C.A., Nunn, A.D. and Tweedle, M.F.: Partition coefficients (log P) and HPLC capacity factors (k') of some Gd(III) complexes of linear and macrocyclic polyamino carboxylates. *J. Liq. Chromatogr.*, 17 (1994) 3735-3746.
- 3494 Nakata, R., Nagayama, M., Naitoh, T., Yasuma, S. and Oki, H.: Liquid chromatography of diastereoisomers of neutral tris- and bis(L- $\alpha$ -amino acidato)chromium(III) complexes. *Anal. Sci.*, 10 (1994) 663-665; C.A., 122 (1995) 150547k.
- 3495 Tallarek, U., Baumeister, E., Albert, K., Bayer, E. and Guiochon, G.: NMR imaging of the chromatographic process. Migration and separation of bands of gadolinium chelates. *J. Chromatogr. A*, 696 (1995) 1-18.
- 3496 Tan, K.-C., Hor, T.S.A. and Lee, H.K.: High-performance liquid chromatography of group 6 dimetal decacarbonyl complexes singly-bridged by diphosphine ligands. *J. Liq. Chromatogr.*, 17 (1994) 3671-3680.
- See also 3862.
27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 3497 Acilu, M. and Font, J.: (Identification of tocopherol isomers or vitamin E in vegetable oil samples by HPLC with photodiode detector). *Tec. Lab.*, 16 (1994) 866-869; C.A., 122 (1995) 185660u.
- 3498 Adamson, P.C., Murphy, R.F., Godwin, K.A., Ulm, E.H. and Balis, F.M.: Pharmacokinetics of 9-cis-retinoic acid in the rhesus monkey. *Cancer Res.*, 55 (1995) 482-485.
- 3499 Ambrosini, L. and Sala, B.: Method for the industrial preparation of (6S) folic acid derivatives by chromatographic separation. *Eur. Pat. Appl. EP 627,435 (Cl. C07D475/04)*, 07 Dec. 1994, Appl. 93/108,752, 01 Jun. 1993; 6 p.; C.A., 122 (1995) 109342b.
- 3500 Bempong, D.K., Honigberg, I.L. and Meltzer, N.M.: Normal phase LC-MS determination of retinoic acid degradation products. *J. Pharm. Biomed. Anal.*, 13 (1995) 285-291.
- 3501 Bergaentzle, M., Arella, F., Bourguignon, J.B. and Hasselmann, C.: Determination of vitamin B6 in foods by HPLC - a collaborative study. *Food Chem.*, 52 (1995, Pub. 1994) 81-86; C.A., 122 (1995) 79368v.
- 3502 Blanco, D., Pajares, M., Escotet, V.J. and Gutiérrez, M.D.: Determination of fat-soluble vitamins by liquid chromatography in pediatric parenteral nutrition. *J. Liq. Chromatogr.*, 17 (1994) 4513-4530.
- 3503 Cahnmann, H.J.: A fast photoisomerisation method for the preparation of tritium-labeled 9-cis-retinoic acid of high specific activity. *Anal. Biochem.*, 227 (1995) 49-53.

- 3504 Delgado-Zamarreño, M.M., Sanchez-Perez, A., Gomez-Perez, M.C. and Hernandez-Mendez, J.: Directly coupled sample treatment-high-performance liquid chromatography for on-line automatic determination of liposoluble vitamins in milk. *J. Chromatogr. A*, 694 (1995) 399-406.
- 3505 Du, Q., Zhang, T.-Y. and Ito, Y.: Purification of 25-hydroxy-cholecalciferol from irradiation of cholesta-5,7-diol by high-speed countercurrent chromatography. *J. Liq. Chromatogr.*, 18 (1995) 181-188.
- 3506 Esteve, M.J., Farre, R., Frigola, A., Lopez, J.C., Romera, J.M., Ramirez, M. and Gil, A.: Comparison of voltammetric and high performance liquid chromatographic methods for ascorbic acid determination in infant formulas. *Food Chem.*, 52 (1995, Pub. 1994) 99-102; C.A., 122 (1995) 79370q.
- 3507 Fujiwara, K., Asao, T., Ikeda, M., Kisara, K. and Ono, S.: (Determination of vitamin K in plasma by HPLC with electrochemical detection). *Bitamin*, 68 (1994) 711-719; C.A., 122 (1995) 100739d.
- 3508 Gehring, T.A., Cooper, W.M., Holder, C.L. and Thompson, H.C., Jr.: Liquid chromatographic determination of thiamine in rodent feed by postcolumn derivatization and fluorescence detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 307-309.
- 3509 Hentz, N.G. and Bachas, L.G.: Class-selective detection system for liquid chromatography based on the streptavidin-biotin interaction. *Anal. Chem.*, 67 (1995) 1014-1018.
- 3510 Hu, O.Y.-P., Wu, C.-Y., Chan, W.-K. and Wu, F.Y.-H.: Determination of anticancer drug vitamin D<sub>3</sub> in plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 299-305.
- 3511 Huber, J.F.K. and Lamprecht, G.: Assay of neopterin in serum by means of two-dimensional high-performance liquid chromatography with automated column switching using three retention mechanisms. *J. Chromatogr. B*, 666 (1995) 223-232.
- 3512 Kraft, J.C., Kimelman, D. and Juchau, M.R.: *Xenopus laevis*: a model system for the study of embryonic retinoid metabolism. I. Embryonic metabolism of 9-*cis* and all-*trans*-retinals and retinols to their corresponding acid form. *Drug Metab. Disp.*, 23 (1995) 72-82.
- 3513 Kraft, J.C., Kimelman, D. and Juchau, M.R.: *Xenopus laevis*: a model system for the study of embryonic retinoid metabolism. II. Embryonic metabolism of all-*trans*-3,4-didehydroretinol to all-*trans*-3,4-didehydroretinoic acid. *Drug Metab. Disap.*, 23 (1995) 83-89.
- 3514 Lefebvre, P., Agadir, A., Cornic, M., Gourmel, B., Hue, B., Dreux, C., Degos, L. and Chomienne, C.: Simultaneous determination of all-*trans* and 13-*cis* retinoic acids and their 4-oxo metabolites by adsorption liquid chromatography after solid-phase extraction. *J. Chromatogr. B*, 666 (1995) 55-61.
- 3515 Li, K. and Zhao, F.-L.: Determination of  $\alpha$ -tocopherol in human and rabbit plasma or serum by high-performance liquid chromatography. *J. Chin. Pharm. Sci.*, 3 (1994) 164-170; C.A., 122 (1995) 155381m.
- 3516 Liaaen-Jensen, S. and Hertzberg, S.: Example 11: carotenoid sulfates. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 283-286; C.A., 122 (1995) 234521k.
- 3517 Manan, F. and Ryley, J.: Development of an improved HPLC method for vitamin A assay in foods. *Discovery Innovation*, 6 (1994) 287-290; C.A., 122 (1995) 104172m.
- 3518 Pfander, H.: Chromatography: Part I. General aspects. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 109-116; C.A., 122 (1995) 234411z - a review with 5 refs.
- 3519 Pfander, H. and Niggli, U.: Chromatography: Part V. Supercritical-fluid chromatography. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 191-198; C.A., 122 (1995) 234415d - a review with 14 refs.
- 3520 Pfander, H. and Riesen, R.: Chromatography: Part IV. High-performance liquid chromatography. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 145-190; C.A., 122 (1995) 234414c - a review with 108 refs.
- 3521 Poon, R., Chu, L., Lecavalier, P., Bergman, A. and Villeneuve, D.C.: Urinary ascorbic acid - HPLC determination and application as a noninvasive biomarker of hepatic response. *J. Biochem. Toxicol.*, 9 (1994) 297-304; C.A., 122 (1995) 74070g.
- 3522 Randolph, R.K. and Simon, M.: Metabolic regulation of active retinoid concentrations in cultured human epidermal keratinocytes by exogenous fatty acids. *Arch. Biochem. Biophys.*, 318 (1995) 6-14.
- 3523 Scalia, S., Renda, A., Ruberto, G., Bonina, F. and Menegatti, E.: Assay of vitamin A palmitate and vitamin E acetate in cosmetic creams and lotions by supercritical fluid extraction and HPLC. *J. Pharm. Biomed. Anal.*, 13 (1995) 273-277.
- 3524 Schuep, W., Hess, D. and Schierle, J.: Example 7: human plasma. Simultaneous determination of carotenoids, retinol and tocopherols. In: Britton, G., Liaaen-Jensen, S. and Pfander, H. (Editors), *Carotenoids*, Birkhaeuser, Basel, 1995, pp. 261-264; C.A., 122 (1995) 234520j.
- 3525 Siddiqui, F.Q., Malik, F. and Fazli, F.R.Y.: Determination of serum retinol by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 342-346.
- 3526 Skurikhin, V.N.: (Determination of vitamins D<sub>2</sub> and D<sub>3</sub> in feed preparations by microcolumn high-performance liquid chromatography with ultraviolet detection). *Bull. Vseross. Nauchno-Issled. Inst. Fiziol. Biokhim. Pitan. S-kh. Zhivotn.*, (1992) 70-76; C.A., 122 (1995) 158878q.
- 3527 Skurikhin, V.N., Dvinskaya, L.M. and Riabikh, T.E.: (Carotenoids determination by high-performance liquid chromatography in plasma and tissues of farm animals). *Bull. Vseross. Nauchno-Issled. Inst. Fizio', Biokhim. Pitan. S-kh. Zhivotn.*, (1992) 76-82; C.A., 122 (1995) 100770g.
- 3528 Stein, J., Hahn, A. and Rehner, G.: High-performance liquid chromatographic determination of nicotinic acid and nicotinamide in biological samples applying post-column derivatization resulting in bathochrome absorption shifts. *J. Chromatogr. B*, 665 (1995) 71-78.
- 3529 Sugimoto, T., Yoshida, A., Teradaira, R., Fujita, K. and Nagatsu, T.: Isolation of deoxybiopterin from urine of cancer patients. *Biog. Amines*, 11 (1995) 1-6; C.A., 122 (1995) 234528t.
- 3530 Thompson, C.O. and Trenerry, V.C.: A rapid method for the determination of total L-ascorbic acid in fruits and vegetables by micellar electrokinetic capillary chromatography. *Food Chem.*, 53 (1995) 43-50; C.A., 122 (1995) 238024y.

- 3531 Wakabayashi, H., Yamato, S., Nakajima, M. and Shimada, K.: Simultaneous determination of oxidized and reduced coenzyme Q and  $\alpha$ -tocopherol in biological samples by high performance liquid chromatography with platinum catalyst reduction and electrochemical detection. *Biol. Pharm. Bull.*, 17 (1994) 997-1002; *C.A.*, 122 (1995) 154758c.
- 3532 Yakushina, L.M. and Taranova, A.G.: (Rapid method for determination of fat-soluble vitamins in human blood serum). *Zh. Fiz. Khim.*, 68 (1994) 1826-1828; *C.A.*, 122 (1995) 50406v.
- 3533 Zhang, Y. and Wang, J.: (Determination of vitamin D in food by high performance liquid chromatography). *Sepu*, 12 (1994) 411-413; *C.A.*, 122 (1995) 54275z - a review with 15 refs.
- 3534 Zhao, H., Li, K., Cao, Y. and Zheng, J.: (Simultaneous determination of vitamins A and D<sub>2</sub> in fortified milk by high performance liquid chromatography). *Sepu*, 13 (1995) 61-62; *C.A.*, 122 (1995) 158912w.
- For additional information see *C.A.*:  
122 (1995) 115138f, 155417c, 158871g, 185697m, 197103c.
- See also 2566, 2828, 3464, 3598.
28. ANTIBIOTICS
- 3535 Anderson, K.L., Moats, W.A., Rushing, J.E., Wesen, D.P. and Papich, M.G.: Potential for oxytetracycline administration by three routes to cause milk residues in lactating cows, as detected by radioimmunoassay (Charm II) and high-performance liquid chromatography test methods. *Am. J. Vet. Res.*, 56 (1995) 70-77; *C.A.*, 122 (1995) 122373g.
- 3536 Baertschi, S.W., Dorman, D.E., Spangle, L.A., Collins, M.W. and Lorenz, L.J.: Formation of fluorescent pyrazine derivatives via a novel degradation pathway of the carbacephalosporin loracarbef. *J. Pharm. Biomed. Anal.*, 13 (1995) 323-328.
- 3537 Becker, M., Junker-Buchheit, A. and Wotschokowsky, M.: (Solid-phase extraction and HPLC analysis of cyclosporin A in blood plasma). *Labor-Med.*, 17 (1994) 169-170; *C.A.*, 122 (1995) 71214w.
- 3538 Blanchflower, W.J. and Kennedy, D.G.: Determination of lasalocid in eggs using liquid-chromatography - electrospray mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 1129-1132.
- 3539 Cobin, J.A. and Johnson, N.A.: Liquid chromatographic method for rapid determination of total avermectin B<sub>1</sub> and 8,9-z-avermectin B<sub>1</sub> residues in apples. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 419-423.
- 3540 Coleman, M.R., Macy, T.D., Moran, J.W. and Rodewald, J.M.: Ruggedness of the monensin and narasin liquid chromatographic assays. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1065-1072.
- 3541 Croubels, S., Baeyens, W. and van Peteghem, C.: Post-column zirconium chelation and fluorescence detection for the liquid chromatographic determination of tetracyclines. *Anal. Chim. Acta*, 303 (1995) 11-16.
- 3542 Emara, S., Morita, I., Tamura, K., Razee, S., Masujima, T., Mohamed, H.A., El-Gizawy, S.M. and El-Rabbat, N.A.: Simultaneous determination of free and bound adriamycin, adriamycinol, adriamycinone and daunorubicin in plasma using column-switching technique and protein-coated pre-columns. *Talanta*, 41 (1994) 1973-1980; *C.A.*, 122 (1995) 71228d.
- 3543 Fagan, S.V.B., Gombatz, C., Abdel-Kader, H. and Menon, G.: Determination of ardacin in various silage food diets by a rapid liquid chromatographic assay. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 16-21.
- 3544 Gräfe, U., Ihn, W., Ritzau, M., Schade, W., Stengel, C., Fleck, W.F., Künkel, W., Härtl, A. and Gutsche, W.: Helioferins, a novel antifungal lipopeptides from *Mycogone rosea*: screening, isolation, structures and biological properties. *J. Antibiot.*, 48 (1995) 126-133.
- 3545 Gräfe, U., Schlegel, R., Ritzau, M., Ihn, W., Dornberger, K., Stengel, C., Fleck, W.F., Gutsche, W., Härtl, A. and Paulus, E.F.R.: Aurantiamicins, new depsipeptide antibiotics from *Streptomyces aurantiacus* IMET 43917. Production, isolation, structure elucidation, and biological activity. *J. Antibiot.*, 48 (1995) 119-125.
- 3546 Guyonnet, J., Richard, M. and Hellings, P.: Determination of fumagillin in muscle tissue of rainbow trout using automated ion-pairing liquid chromatography. *J. Chromatogr. B*, 666 (1995) 354-359.
- 3547 Hormazabal, V. and Yndestad, M.: Rapid assay for monitoring residues of enrofloxacin in milk and meat tissues by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3775-3782.
- 3548 Ikai, Y., Oka, H., Hayakawa, J., Matsumoto, M., Saito, M., Harada, K.-i., Mayumi, T. and Suzuki, M.: Total structures and antimicrobial activity of bacitracin minor components. *J. Antibiot.*, 48 (1995) 233-242.
- 3549 Itoh, T. and Yamada, H.: Diastereomeric  $\beta$ -lactam antibiotics. Analytical methods, isomerization and stereoselective pharmacokinetics. *J. Chromatogr. A*, 694 (1995) 195-208 - a review with 37 refs.
- 3550 Kobayashi, H., Sunaga, R., Furihota, K., Morisaki, N. and Iwasaki, S.: Isolation and structures of an antifungal antibiotic, fusarieline A, and related compounds produced by a *Fusarium* sp. *J. Antibiot.*, 48 (1995) 42-52.
- 3551 Kominato, K., Watanabe, Y., Hirano, S.-i., Kioka, T., Terasawa, T., Yoshioka, T., Okamura, K. and Tone, H.: Mer-A2026A and B, novel piericidins with vasodilating effect I. Producing organism, fermentation, isolation and biological properties. *J. Antibiot.*, 48 (1995) 106-112.
- 3552 La Follette, G., Jayewardene, A.L., Seneviratne, A.K., Lin, E.T. and Gambertoglio, J.G.: Determination of ticarcillin in human plasma and in urine by reversed-phase LC. *J. Pharm. Biomed. Anal.*, 13 (1995) 159-164.
- 3553 Lau, R.C.M. and Rinehart, K.L.: Berninamicins B, C and D, minor metabolites from *Streptomyces bernensis*. *J. Antibiot.*, 47 (1994) 1466-1472.
- 3554 Lee, C.-H., Kim, S., Hyun, B., Suh, J.-W., Yon, C., Kim, C., Lim, C. and Kim, C.: Cepacidina A, a novel antifungal antibiotic produced by *Pseudomonas cepacia*. I. Taxonomy, production, isolation and biological activity. *J. Antibiot.*, 47 (1994) 1402-1405.
- 3555 Mathew, M. and Das Gupta, V.: Stability of vancomycin hydrochloride solutions at various pH values as determined by high-performance liquid chromatography. *Drug. Dev. Ind. Pharm.*, 21 (1995) 257-264; *C.A.*, 122 (1995) 114783a.



- 3556 Matsumoto, K., Tanka, K., Matsutani, S., Sakazaki, R., Hino, H., Uotani, N., Tanimoto, T., Kawamura, Y., Nakamoto, S. and Yoshida, T.: Isolation and biological activity of thielocins: novel phospholipase A<sub>2</sub> inhibitors produced by *Thielavia terricola* RF-143. *J. Antibiot.*, 48 (1995) 106-112.
- 3557 McMullin, C.M., White, L.O., MacGowan, A.P., Holt, H.A., Lovering, A.M. and Reeves, D.S.: Assay of serum teicoplanin concentrations in clinical specimens: a comparison of isocratic high performance liquid chromatography with polarization fluoroimmunoassay and bioassay. *J. Antimicrob. Chemother.*, 34 (1994) 425-429; *C.A.*, 122 (1995) 45512x.
- 3558 Mikisch, E.: (Significance of the 4-epimers in the determination of oxytetracycline, tetracycline, and chlorotetracycline residues in animal-derived food, by HPLC). *Dtsch. Lebensm.-Rundsch.*, 90 (1994) 39-41; *C.A.*, 122 (1995) 54280x.
- 3559 Moats, W.A. and Harik-Khan, R.: Liquid chromatographic determination of  $\beta$ -lactam antibiotics in milk: a multiresidue approach. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 49-54.
- 3560 Nishikawa, T., Suzuki, S. and Ohtani, H.: Reversible isomerization of rapamycin demonstrated by liquid chromatography. *J. Antibiot.*, 47 (1994) 1554-1556.
- 3561 Nozawa, O., Okazaki, T., Sakai, N., Komurasaki, T., Hanada, K., Morimoto, S., Chen, Z.-X., He, P.-M. and Mizoue, K.: A novel bioactive  $\delta$  lactone FD-211. Taxonomy, isolation and characterization. *J. Antibiot.*, 48 (1995) 113-118.
- 3562 Paesen, J., Claeys, P., Cypers, W., Roets, E. and Hoogmartens, J.: Liquid chromatography of tylosin A and related substances on poly(styrene-divinylbenzene). *J. Chromatogr. A*, 699 (1995) 93-97.
- 3563 Paesen, J., Cypers, W., Busson, R., Roets, E. and Hoogmartens, J.: Isolation of decomposition products of tylosin using liquid chromatography. *J. Chromatogr. A*, 699 (1995) 99-106.
- 3564 Petersen, F., Fredenhagen, A., Mett, H., Lydon, N.B., Delmendo, R., Jenny, H.-B. and Peter, H.H.: Paecilquinones A, B, C, D, E and F: new potent inhibitors of protein tyrosine kinases produced by *Paecilomyces carneus*. I. Taxonomy, fermentation, isolation and biological activity. *J. Antibiot.*, 48 (1995) 191-198.
- 3565 Proksa, B. and Fuska, J.: Role of aliphatic acids in the biosynthesis of vermiculin in *Penicillium vermiculatum*. *Pharmazie*, 50 (1995) 215-216.
- 3566 Reeves, V.S.: Liquid chromatographic procedure for the determination of novobiocin residues in bovine milk: interlaboratory study. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 55-58.
- 3567 Saitoh, K., Furumai, T., Oki, T., Nishida, F., Harada, K.-i. and Suzuki, M.: Pradimicin S, a new pradimicin analog. III. Application of the Frit-FAB LC/MS technique to the elucidation of the pradimicin S biosynthetic pathway. *J. Antibiot.*, 48 (1995) 162-168.
- 3568 Sancho-Chust, V., Fabra-Campos, S., Gómez-Meseguer, V., Benigochea, M. and Martín-Villodre, A.: Experimental studies on the influence of surfactants on intestinal absorption of drugs. Cefadroxil as model drug and sodium lauryl sulfate as model surfactant: studies in rat. *Arzneim.-Forsch.*, 45 (1995) 595-601.
- 3569 Seki-Asano, M., Okazaki, T., Yamagishi, M., Sakai, N., Takayama, Y., Hanada, K., Morimoto, S., Takatsuki, A. and Mizoue, K.: Isolation and characterization of a new 12-membered macrolide FD-895. *J. Antibiot.*, 47 (1994) 1395-1401.
- 3570 Tyczkowska, K.L., Voyksner, R.D., Straub, R.F. and Aronson, A.L.: Simultaneous multiresidue analysis of  $\beta$ -lactam antibiotics in bovine milk by liquid chromatography with ultraviolet detection and confirmation by electrospray mass spectrometry. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1122-1131.
- 3571 Vickers, A.E.M., Fisher, R.L., Brendel, K., Guertler, J., Dannecker, R., Keller, B. and Fisher, V.: Sites of biotransformation for the cyclosporin derivative SD2 IMM 125 using human liver and kidney slices and intestine. Comparison with rat liver slices and cyclosporin A metabolism. *Drug Metab. Disp.*, 23 (1995) 327-333.
- 3572 Vickers, A.E.M., Meyer, E., Dannecker, R., Keller, B., Tynes, R.E. and Maurer, G.: Human liver cytochrome P4503A biotransformation of the cyclosporin derivative SD2 IMM 125. *Drug Metab. Disp.*, 23 (1995) 321-326.
- 3573 Yang, C.F., Stassinopoulos, A. and Goldberg, I.H.: Specific binding of the biradical analog of neocarzinostatin chromophore to bulged DNA: implications for thiol-independent cleavage. *Biochemistry*, 34 (1995) 2267-2275.

For additional information see *C.A.*:

122 (1995) 115150d, 204532x, 209170n.

See also 3741, 3766.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

### 29a. General techniques

- 3574 Beneventi, G., Barbieri, C., del Carlo, G., Dondi, C. and Forti, S.: (Pesticides in wine: simultaneous determination of ethylenethiourea (ETU), propylenethiourea (PTU), fungicides and insecticides by SPE extraction with a C18 cartridge). *Ind. Bevande*, 23 (1994) 317-323; *C.A.*, 122 (1995) 54326s.
- 3575 Holstege, D.M., Scharberg, D.L., Tor, E.R., Hart, L.C. and Galey, F.D.: A rapid multiresidue screen for organophosphorus, organochlorine, and N-methyl carbamate insecticides in plant and animal tissues. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1263-1274.
- 3576 Marcé, R.M., Prosen, H., Crespo, C., Calull, M., Borrull, F. and Brinkman, U.A.T.: On-line trace enrichment of polar pesticides in environmental waters by reversed-phase liquid chromatography-diode array detection-particle beam mass spectrometry. *J. Chromatogr. A*, 696 (1995) 63-74.
- 3577 Newsome, W.H.: An evaluation of pesticide residue methodology. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 4-8.
- 3578 Nouri, B., Fouillet, B., Toussaint, G., Chambon, P. and Chambon, R.: High-performance liquid chromatography with diode-array detection for the determination of pesticides in water using automated solid-phase extraction. *Analyst (Cambridge)*, 120 (1995) 1133-1136.
- 3579 Pflab, G. and York, H.: Application of AMD (automated multiple development) for the determination of pesticides in drinking water. Part 3. Solid phase extraction and influencing factors. *Acta Hydrochim. Hydrobiol.*, 22 (1994) 216-223; *C.A.*, 122 (1995) 88874j.

- 3580 Sannino, A.: Investigation into contamination of processed fruit products by carbendazim, methyl thiophanate and thiabendazole. *Food Chem.*, 52 (1995, Pub. 1994) 57-61; C.A., 122 (1995) 79366t.
- 3581 Sennert, S., Volmer, D., Levsen, K. and Wünsch, G.: Multi-residue analysis of polar pesticides in surface and drinking water by on-line enrichment and thermospray LC-MS. *Fresenius J. Anal. Chem.*, 351 (1995) 642-649.
- 3582 Volmer, D., Levsen, K. and Engewald, W.: Analysis of polar pesticides in aqueous samples by combined online trace enrichment and thermospray liquid chromatography mass spectrometry. *Vom Wasser*, 82 (1994) 335-384; C.A., 122 (1995) 114385x.
- 3583 Wilson, S.B. and Rontree, J.: (Analysis of pesticides by electro-spray LC-MS: a sensitive, specific, routine and robust analysis technique). *Foods Food Ingredients J. Jpn.*, 162 (1994) 81-86; C.A., 122 (1995) 158873j.
- 3584 Yamamoto, A., Matsunaga, A., Ohto, M., Mizukami, E., Hayakawa, K. and Miyazaki, M.: Real-time analysis of multicomponent chromatograms: application to high-performance liquid chromatography. *Analyst (Cambridge)*, 120 (1995) 377-380.
- For additional information see C.A.:  
122 (1995) 48963f, 100762f, 212345d.
- See also 3824, 3826.
- 29c. *Phosphorus insecticides*
- For additional information see C.A.:  
122 (1995) 99232z.
- 29d. *Carbamates*
- 3585 Sundaram, K.M.S. and Curry, J.: Fluorometric determination of aminocarb and mexacarbamate and some of their metabolites by liquid chromatography: influence of structural factors on fluorescence intensity. *J. Liq. Chromatogr.*, 17 (1994) 3759-3774.
- For additional information see C.A.:  
122 (1995) 29977x.
- 29e. *Herbicides*
- 3586 Brandsteterova, E., Kralikova, D., Lehotay, J. and Tekel, J.: HPLC analysis of some substituted anilines and phenylurea herbicides in milk. *Nahrung*, 38 (1994) 439-441; C.A., 122 (1995) 54311h.
- 3587 Chiron, S., Papilloud, S., Haerdi, W. and Barceló, D.: Automated on-line liquid-solid extraction followed by liquid chromatography - high-flow pneumatically assisted electrospray mass spectrometry for the determination of acidic herbicides in environmental waters. *Anal. Chem.*, 67 (1995) 1637-1643.
- 3588 Reichert, J., Gernikeites, T. and Winkler, M.: (Application of HPLC with fluorescence detection to environmental analysis exemplified with the herbicide dalapon). *Vom Wasser*, 82 (1994) 37-48; C.A., 122 (1995) 114387z.
- 29f. *Fungicides*
- 3589 Bushway, R.J., Young, B.E.S., Paradis, L.R. and Perkins, L.B.: Determination of thiabendazole in fruits and vegetables by competitive-inhibition enzyme immunoassay. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1243-1248.
- 3590 Bushway, R.J., Young, B.E.S., Paradis, L.R., Perkins, L.B., Martin, S.K. and Brown, M.P.: Determination of methyl 2-benzimidazolecarbamate in bulk fruit juice concentrates by competitive-inhibition enzyme immunoassay. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1237-1243.
- 29g. *Other types of pesticides and various agrochemicals*
- 3591 Blanco Gomis, D., Castano Fernandez, A., Megido Bernardo, V. and Gutierrez Alvarez, M.D.: High performance liquid chromatographic determination of cymiazole in honey with UV and electrochemical detection. *Chromatographia*, 39 (1994) 602-606.
- 3592 Sundaram, K.M.S., Sloane, L. and Curry, J.: Kinetics of azadirachtin hydrolysis in model aquatic systems by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 363-376.
- 3593 Tamiya, M., Kashiwabara, F., Watanabe, Y., Ando, T., Tsutsumi, T., Norizuki, H., Minamisawa, M. and Hisai, S.: (Development of an analytical method for methoprene in rice by HPLC). *Shokuhin Eiseigaku Zasshi*, 35 (1994) 593-598; C.A., 122 (1995) 131286h.
- See also 3808.
30. SYNTHETIC AND NATURAL DYES
- 30a. *Synthetic dyes*
- 3594 Castle, M. and Neuteboom, E.: High-performance liquid chromatography of the fluorescent dyes Fura-2 and Mag-Fura. Stability in organic solvents. *J. Chromatogr. A*, 696 (1995) 93-99.
- 3595 Roybal, J.E., Pfenning, A.P., Munns, R.K., Holland, D.C., Hurlbut, J.A. and Long, A.R.: Determination of malachite green and its metabolite, leucomalachite green, in catfish (*Ictalurus punctatus*) tissue by liquid chromatography with visible detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 453-457.
- See also 2628, 3000, 3005.
- 30b. *Chloroplast and other natural pigments*
- 3596 Bento, L.S.M.: Separation of sugar colorants using chromatographic columns containing sucrose crystals. *Publ. Tech. Pap. Proc. Annu. Meet. Sugar Ind. Technol.*, 53 (1994) 17-39; C.A., 122 (1995) 109191b.
- 3597 Gao, L. and Mazza, G.: Characterization of acetylated anthocyanins in lowbush blueberries. *J. Liq. Chromatogr.*, 18 (1995) 245-259.
- 3598 Klein, R.: Identification, stereoconfiguration, chromatographic and fluorescence properties of natural pterins. *Adv. Exp. Med. Biol.*, 338 (Chemistry and Biology of Pteridines and Foliates) (1993) 43-46; C.A., 122 (1995) 155614q.

- 3599 Mínguez-Mosquera, M.I., Hornero-Méndez, D. and Garrido-Fernández, J.: Detection of bixin, lycopene, cantaxanthin, and  $\beta$ -apo-8'-carotenol in products derived from red pepper. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 491-498.
- 3600 Moskalenko, A.A., Toropygina, O.A. and Kuznetsova, N.Yu.: (Pigment-protein complexes of *Chromatium minutissimum* with extracted carotenoids). *Biokhimiya (Moscow)*, 59 (1994) 1414-1420.
- 3601 Szabo, G., Prosser, S.L. and Bulman, R.A.: Extraction and high-performance size-exclusion chromatography of humic substances from soils. In: Kalasz, H., Ettore, L.S. and Pick, J. (Editors), *New Approaches Chromatogr. '93, [Pap. Budapest Chromatogr. Conf.]*, 1990, Fekete Sas Diado, Budapest, 1993, pp. 171-186; C.A., 122 (1995) 212895h.
- 3602 Van Heukelem, L., Lewitus, A.J., Kana, T.M. and Craft, N.E.: Improved separations of phytoplankton pigments using temperature-controlled high performance liquid chromatography. *Mar. Ecol.: Prog. Ser.*, 114 (1994) 303-313; C.A., 122 (1995) 155379s.
- For additional information see C.A.:  
122 (1995) 158879r.
- See also 3514, 3518, 3519, 3520, 3527, 3803.
31. PLASTICS AND THEIR INTERMEDIATES
- 3603 Allan, D.S., Birchmeier, M., Pribish, J., Priddy, D.B. and Smith, P.: Application of GPC-UV/VIS analysis to investigate the mechanism of styrene-co-acrylonitrile thermal discoloration. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 34 (1993) 258-259; C.A., 121 (1994) 10841f.
- 3604 Bullock, J.: High-temperature reversed-phase high-performance liquid chromatographic analysis of a synthetic copolymer on a non-porous support. *J. Chromatogr. A*, 694 (1995) 415-423.
- 3605 Cutie, S.S. and Martin, S.J.: Size-exclusion chromatography of crosslinked superabsorbent polymers. *J. Appl. Polym. Sci.*, 55 (1995) 605-609; C.A., 122 (1995) 161960r.
- 3606 Glöckner, G., Wolf, D. and Engelhardt, H.: Control of adsorption and solubility in gradient high-performance liquid chromatography. Part V. Separation of styrene/methyl methacrylate copolymers by sudden-transition gradients in normal-phase and reversed-phase mode. *Chromatographia*, 39 (1994) 557-563.
- 3607 Henry, R.A., Noguchi, K. and Stevenson, R.: Solvent scouting in steric exclusion chromatography. Characterization of poly(vinyl acetate) and poly(vinyl alcohol) on the same organic gel column. *Am. Lab. (Shelton)*, 25 (1993) 36Z, 368B-36DD; C.A., 120 (1994) 299734k.
- 3608 Janco, M., Proudskova, T. and Berek, D.: Liquid chromatography of polymer mixtures applying combination of exclusion and full adsorption mechanisms. I. Analysis of polystyrene in its mixture with poly(methyl methacrylate) single column/single eluent approach. *J. Appl. Polym. Sci.*, 55 (1995) 393-397; C.A., 122 (1995) 161958w.
- 3609 Kuzaev, A.I., Ol'khova, O.M., Solomko, S.I. and Baturin, S.M.: (Use of liquid chromatography in the critical region for separation of oligo(epichlorohydrins) with respect to functionality types). *Zh. Fiz. Khim.*, 68 (1994) 1794-1798; C.A., 122 (1995) 106976a.
- 3610 Lehrle, R.S. and Sarson, D.S.: Polymer molecular weight distribution: results from matrix-assisted laser desorption ionization compared with those from gel-permeation chromatography. *Rapid Commun. Mass Spectrom.*, 9 (1995) 91-92; C.A., 122 (1995) 162029f.
- 3611 Martin, L. and Balke, S.T.: Cyclic oligomer considerations in the size exclusion chromatography of poly(ethylene terephthalate). *J. Liq. Chromatogr.*, 18 (1995) 1-15.
- 3612 Mendichi, R., Audisio, G. and Giacometti Schieroni, A.: Molecular mass characterization of poly(4-vinylpyridine). *J. Liq. Chromatogr.*, 17 (1994) 3323-3331.
- 3613 Nagy, D.J.: Characterization of poly(vinyl alcohol) using SEC multiangle laser light scattering. *Int. Lab.*, 25, No. 2 (1995) P6-P11.
- 3614 Okada, T.: Multifunctional separation with polyamine-bonded resin. *Anal. Chim. Acta*, 303 (1995) 193-197.
- 3615 Pasch, H. and Rode, K.: Use of matrix-assisted laser desorption/ionization mass spectrometry for molar mass-sensitive detection in liquid chromatography of polymers. *J. Chromatogr. A*, 699 (1995) 21-29.
- 3616 Shintani, H.: HPLC analysis of toxic additives and residual monomer from dental plate. *J. Liq. Chromatogr.*, 18 (1995) 613-626.
- 3617 Staal, W.J., Cools, P., van Herk, A.M. and German, A.L.: Online extraction of polymers, oligomers, additives and monomers by multiple solvents on packed HPLC columns. *Process Technol. Proc.*, 11 (1994) 857-859; C.A., 122 (1995) 162162u.
- 3618 Sturman, H.W., Geissler, M., Weimar, A., Paul, A. and Nickel, B.: (High performance liquid chromatography (HPLC) and gas chromatography (GC) as methods in plastic material analysis). *Kautsch. Gummi Kunstst.*, 47 (1994) 812-815; C.A., 122 (1995) 107514k.
- 3619 Terneuil, G., Hotier, G. and Lonchamp, D.: Chromatographic separation of mixtures of polyglycerols. *Ger. Offen. DE 4,404,291 (Cl. C07C43/10)*, 18 Aug. 1994, FR Appl. 93/1,755, 15 Feb. 1993; 9 p.; C.A., 122 (1995) 82342n.
- 3620 Van Every, K.W. and Long, W.J.: High-temperature GPC/IR characterization of polyolefines. *Proc. SPIE-Int. Soc. Opt. Eng.*, 2089 (9th International Conference on Fourier Transform Spectroscopy, 1993) (1993) 186-187; C.A., 122 (1995) 82482h.
- For additional information see C.A.:  
122 (1995) 82466f, 107275h, 204273p.
- See also 2671, 2680, 2757, 3003, 3841.

## 32. DRUG ANALYSIS

## 32a. Drug analysis, general techniques

- 3621 Balogh, G., Csizer, E., Ferenczy, G.G., Halmos, Z., Herenyi, B., Horvath, P., Lauko, A. and Gorog, S.: Estimation of impurity profiles of drugs and related materials. 12. Isolation and identification of an isomeric impurity in danazol. *Pharm. Res.*, 12 (1995) 295-298; C.A., 122 (1995) 197141p.
- 3622 Bieganowska, M.L. and Petruczynek, A.: Ion pair chromatography and its application in analysis of pharmaceuticals. *Chem. Anal. (Warsaw)*, 39 (1994) 525-541; C.A., 122 (1995) 26925t.
- 3623 Caldwell, J.: Stereochemical determinants of the nature and consequences of drug metabolism. *J. Chromatogr. A*, 694 (1995) 39-48 - a review with 47 refs.
- 3624 Glazova, N.V., Efimova, T.B. and Dmitrenko, L.V.: (Use of gel chromatography for quantitative analysis of pyrogenic impurities in drugs). *Zh. Fiz. Khim.*, 68 (1994) 1803-1805; C.A., 122 (1995) 89575z.
- 3625 Gluck, S.J., Martin, E. and Benko, M.H.: Centrifugal partition chromatography for the determination of octanol-water partition coefficients. *Chromatogr. Sci. Ser.*, 68(Centrifugal Partition Chromatography) (1995) 199-218; C.A., 122 (1995) 155377q.
- 3626 Hermansson, J. and Grahn, A.: Optimization of the separation of enantiomers of basic drugs. Retention mechanisms and dynamic modification of the chiral bonding properties on an  $\alpha_1$ -acid glycoprotein column. *J. Chromatogr. A*, 694 (1995) 57-69.
- 3627 Herraiz-Hernández, R., van de Merbel, N.C. and Brinkman, U.A.T.: Determination of the total concentration of highly protein-bound drugs in plasma by on-line dialysis and column liquid chromatography: application to non-steroidal anti-inflammatory drugs. *J. Chromatogr. B*, 666 (1995) 127-137.
- 3628 Itoh, H., Kinoshita, T. and Nimura, N.: (Direct serum injection analysis on ODS silica gel column). *Kuromatogurafi*, 15 (1994) 256-257; C.A., 122 (1995) 127791b.
- 3629 Leahy, D.E., Lynch, J., Finney, R.E. and Taylor, D.C.: Estimation of sieving coefficients of convective absorption of drugs in perfused rat jejunum. *J. Pharmacokin. Biopharm.*, 22 (1994) 411-429.
- 3630 Pidgeon, C., Ong, S., Liu, H., Qiu, X., Pidgeon, M., Dantzig, A.H., Munroe, J., Hornback, W.J., Kasher, J.S. et al.: IAM chromatography: an *in vitro* screen for predicting drug membrane permeability. *J. Med. Chem.*, 38 (1995) 590-594; C.A., 122 (1995) 95739x.
- 3631 Qin, Z.: Advances in biopharmaceutical analysis in the People's Republic of China: 1991-1993. *J. Pharm. Biomed. Anal.*, 13 (1995) 1-7 - a review with 58 refs.
- 3632 Steele, K.M. and Lunte, C.E.: Microdialysis sampling coupled to on-line microbore liquid chromatography for pharmacokinetic studies. *J. Pharm. Biomed. Anal.*, 13 (1995) 149-154.
- 3633 Wainer, I.W., Ducharme, J., Granvil, C.P., Parenteau, H. and Abdullah, S.: Using chirality as a unique probe of pharmacological properties. *J. Chromatogr. A*, 694 (1995) 169-179.
- 3634 Zeng, S.: (Separations of drug enantiomers by high-performance liquid chromatography with chiral mobile phase additives). *Sepu*, 13 (1995) 21-25; C.A., 122 (1995) 197088b - a review with 47 refs.
- 3635 Zeng, S., Shen, X. and Liu, Z.: (Pharmacology of chiral pharmaceuticals and HPLC determination of their enantiomers *in vivo*). *Zhejiang Yike Daxue Xuebao*, 23 (1994) 234-238; C.A., 122 (1995) 95711g.

For additional information see C.A.:

122 (1995) 122271x, 229996q, 230008a.

See also 2575, 2613, 2722, 2727, 2741, 2748, 3780, 3781, 3796, 3831.

## 32b. Antirheumatics and antiinflammatory drugs

- 3636 Azzolina, O. and Collina, S.: Chiral recognition of alkyl 2-aryloxypropionates by HPLC. *J. Liq. Chromatogr.*, 18 (1995) 81-92.
- 3637 Blondino, F.E. and Byron, P.R.: The quantitative determination of aspirin and its degradation products in a model solution aerosol. *J. Pharm. Biomed. Anal.*, 13 (1995) 111-119.
- 3638 Caccamese, S., Principato, G., Ottanà, R., Previtera, T. and Zapalà, C.: Liquid chromatographic separation of the enantiomers of antihistaminic 3,3'-di(1,3-thiazolidin-4-one) derivatives with two and four stereogenic centres. *J. Chromatogr. A*, 694 (1995) 355-363.
- 3639 Daffonchio, L., Bestetti, A., Clavenna, G., Fedele, G., Ferrari, M.P. and Omini, C.: Effect of a new foam formulation of keto-profen lysine salt in experimental models of inflammation and hyperalgesia. *Arzneim.-Forsch.*, 45 (1995) 590-594.
- 3640 Deigner, H.P., Freyberg, C.E. and Laufer, S.: Distribution and excretion of [<sup>14</sup>C]-labelled [2,2-dimethyl-6-(4-chlorophenyl)-7-phenyl-2,3-dihydro-1H-pyrrolizine-5-yl]-[2'-<sup>14</sup>C]-acetic acid in rats. *Arzneim.-Forsch.*, 45 (1995) 272-276.
- 3641 Ding, A., Zia-Amirhosseini, P., McDonagh, A.F., Burlingame, A.I. and Benet, L.Z.: Reactivity of tolmetin glucuronide with human serum albumin. Identification of binding sites and mechanisms of reaction by tandem mass spectrometry. *Drug Metab. Disp.*, 23 (1995) 369-376.
- 3642 Hamdoune, M., Mounie, J., Magdalou, J., Masmoudi, T., Goudonnet, H. and Escousse, A.: Characterization of the *in vitro* glucuronidation of flurbiprofen enantiomers. *Drug Metab. Disp.*, 23 (1995) 343-348.
- 3643 Hartmann, D., Stief, G., Lingenfelder, M., Güzelhan, C. and Horsch, A.K.: Study on the possible interaction between tenoxicam and atenolol in hypertensive patients. *Arzneim.-Forsch.*, 45 (1995) 494-498.
- 3644 Mason, J.L. and Hobbs, G.J.: Simple method for the analysis of tenoxicam in human plasma using high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 410-415.
- 3645 Maya, M.T., Pais, J.P. and Morais, J.A.: A rapid method for the determination of piroxicam in plasma by high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 319-322.

- 3646 Teffera, Y. and Abramson, F.: Application of high-performance liquid chromatography/chemical reaction interface mass spectrometry for the analysis of conjugated metabolites: a demonstration using deuterated acetaminophen. *Biol. Mass Spectrom.*, 23 (1994) 776-783; *C.A.*, 122 (1995) 45531c.
- 3647 Velic, I., Metzler, M., Hege, H.G. and Weymann, J.: Separation and identification of phase I and phase II [<sup>14</sup>C]antipyrine metabolites in rat and dog urine. *J. Chromatogr. B*, 666 (1995) 139-147.
- 3648 Volin, P.: Simple and specific reversed-phase liquid chromatographic method with diode-array detection for simultaneous determination of serum hydroxychloroquine, chloroquine and some corticosteroids. *J. Chromatogr. B*, 666 (1995) 347-353.
- For additional information see *C.A.*:  
122 (1995) 89603g, 170366d, 170370a, 170372c, 170373d.
- See also 2741, 2893, 2925, 3008, 3627, 3711, 3755.
- 32c. *Autonomic and cardiovascular drugs*
- 3649 Berndt, A., Oertel, R., Richter, K., Terhaag, B. and Gramatté, T.: Relative Bioverfügbarkeit des Antiarrhythmikums Tiracizin und seiner drei Metabolite. *Arzneim.-Forsch.*, 45 (1995) 250-253.
- 3650 Cavazzutti, G., Gagliardi, L., de Orsi, D. and Tonelli, D.: Simultaneous determination of buzepide, phenylpropanolamine, and clocinizine in pharmaceutical preparations by ion-pair reversed-phase HPLC. *J. Liq. Chromatogr.*, 18 (1995) 227-234.
- 3651 Chatterjee, D.J., Li, W.Y., Hurst, A.K. and Koda, R.T.: High-performance liquid chromatographic method for determination of atenolol from human plasma and urine: simultaneous fluorescence and ultraviolet detection. *J. Liq. Chromatogr.*, 18 (1995) 791-806.
- 3652 Chen, C.L., Sangiah, S., Chen, H., Berlin, K.D., Garrison, G.L., Scherlag, B.J. and Lazzara, R.: Simultaneous determination of a novel antiarrhythmic agent, 7-benzyl-3-thia-7-azabicyclo[3.3.1]nonane and its sulfoxidation metabolite in plasma and urine by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3681-3694.
- 3653 Chilton, A.S., Godward, R.E. and Carey, P.F.: The determination in human plasma of 1-hydroxy-2-naphthoic acid following administration of salmeterol xinafoate. *J. Pharm. Biomed. Anal.*, 13 (1995) 165-169.
- 3654 Cieri, U.R.: Determination of reserpine, hydralazine HCl, and hydrochlorothiazide in tablets by liquid chromatography on a short, normal-phase column. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1104-1108.
- 3655 El Wailly, A.F.M., El-Yazbi, F.A., Belal, S.F. and Abdel Razak, O.: High-performance liquid chromatographic and derivative ultraviolet spectrophotometric determination of pindolol and clopamide mixture in tablets. *Anal. Lett.*, 28 (1995) 893-907.
- 3656 Elm, T. and Hansen, E.L.: Simultaneous determination of lofepramine and desipramine by a high-performance liquid chromatographic method used for therapeutic drug monitoring. *J. Chromatogr. B*, 665 (1995) 355-361.
- 3657 Frey, H.O., Kinkeldei, J. and Quast, M.: (Simple and rapid determination for determining the anticonvulsant lamotrigine). *Labor-Med.*, 17 (1994) 166-168; *C.A.*, 122 (1995) 71213v.
- 3658 Herrmann, R.: Automated HPLC assay for sotalol in human plasma. *J. Pharm. Biomed. Anal.*, 13 (1995) 329-333.
- 3659 Hutt, V., Pabst, G., Salama, Z., Kappler, J. and Jaeger, H.: Untersuchungen zur Pharmakokinetik und Bioverfügbarkeit einer neuen Mexiletin-Zubereitung und gesunden Freiwilligen. *Arzneim.-Forsch.*, 45 (1995) 254-257.
- 3660 Jahn, P., Eckrich, C., Schneidrowski, B., Volz-Zang, C., Schulte, B., Mutschler, E. and Palm, D.:  $\beta_1$ -Adrenoceptor subtype selective antagonism of esmolol and its major metabolite *in vitro* and in man. Investigations using tricresylphosphate as red blood cell carboxylesterase inhibitor. *Arzneim.-Forsch.*, 45 (1995) 536-541.
- 3661 Koytchev, R., Alken, R.-G., Mayer, O., Böhm, R., Ellrich, A. and Waldner-Kölblin, R.G.: Untersuchungen zur Bioäquivalenz von zwei oralen Propafenon-Zubereitungen. *Arzneim.-Forsch.*, 45 (1995) 542-545.
- 3662 Kugler, A.R., Olson, S.C. and Smith, D.E.: Determination of quinapril and quinaprilat by high-performance liquid chromatography with radiochemical detection, coupled to liquid scintillation counting spectrometry. *J. Chromatogr. B*, 666 (1995) 360-367.
- 3663 Leube, J. and Fischer, G.: Determination of the renin inhibitor Ro 42-5892 in human plasma by automated pre-column derivatization, reversed-phase high-performance liquid chromatographic separation and electrochemical detection after post-column irradiation. *J. Chromatogr. B*, 665 (1995) 373-381.
- 3664 Lozano, R., Warren, F.V., Jr., Perlman, S. and Joseph, J.M.: Quantitative analysis of fosinopril sodium by capillary zone electrophoresis and liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 139-148.
- 3665 Misztal, G. and Przyborowski, L.: Determination of aprindine in human plasma using reversed phase HPLC. *Pharmazie*, 50 (1995) 187-188.
- 3666 Miyazaki, M., Hosoki, K., Okunishi, H., Ishii, K., Ikeno, A. and Okazaki, Y.: Anti-atherosclerotic and plasma lipid lowering effects of the novel calcium blocker with  $\alpha_1$ -adrenoreceptor antagonistic activity, monatepil, in high cholesterol diet-fed Japanese *Macaca fuscata* monkeys. *Arzneim.-Forsch.*, 44 (1994) 288-297.
- 3667 Mustafa, A.A., Al-Rashood, K.A., Hagga, M.E.M., Gad-Kariem, E.A. and Al-Awady, M.E.: High-performance liquid chromatographic and polarographic methods for the quantitation of nifedipine hydrochloride in capsules. *Pak. J. Sci. Ind. Res.*, 37 (1994) 1-4; *C.A.*, 122 (1995) 142704u.
- 3668 Mutoh, M., Nakada, N., Matsukuma, S., Ohshima, S., Yoshinari, K., Watanabe, J. and Arisawa, M.: Panclitins, novel pancreatic lipase inhibitors. I. Taxonomy, fermentation, isolation and biological activity. *J. Antibiot.*, 47 (1994) 1369-1375.
- 3669 Niaz, E.M., El-Sayed, Y.M. and Khidr, S.H.: Analysis of prazosin in plasma by high-performance liquid chromatography using fluorescence detection. *J. Liq. Chromatogr.*, 18 (1995) 977-989.
- 3670 Pham-Huy, C., Radenen, B., Sahu-Gnassi, A. and Claude, J.-R.: High-performance liquid chromatographic determination of (S)- and (R)-propranolol in human plasma and urine with a chiral  $\beta$ -cyclodextrin bonded phase. *J. Chromatogr. B*, 665 (1995) 125-132.
- 3671 Ring, J.A., Ghabrial, H., Ching, M.S., Shulkes, A., Smallwood, R.A. and Morgan, D.J.: Fetal hepatic propranolol metabolism. Studies in the isolated fetal sheep liver. *Drug Metab. Disp.*, 23 (1995) 190-196.

- 3672 Sharma, S.C., Evans, M.B. and Evans, S.J.: The enantiomeric separation of metipranolol and desacetylmepipranolol on a cellulose tris-3,5-dimethylphenyl-carbamate chiral stationary phase. *J. Pharm. Biomed. Anal.*, 13 (1995) 129-137.
- 3673 Siret, L., Macaudiere, P., Bargmann-Leyder, N., Tambute, A., Caude, M. and Goudeon, E.: Separation of the optical isomers of a new 1,4-dihydropyridine calcium channel blocker (LF 2.0254) by liquid and supercritical fluid chromatography. *Chirality*, 6 (1994) 440-445; *C.A.*, 122 (1995) 150543f.
- 3674 Stearns, R.A., Chakravarty, P.K., Chen, R. and Chiu, S.-H.L.: Biotransformation of lasartan to its active carboxylic acid metabolites in human liver microsomes. Role of cytochrome P4502C and 3A subfamily members. *Drug Metab. Disp.*, 23 (1995) 207-215.
- 3675 Tokuma, Y. and Noguchi, H.: Stereoselective pharmacokinetics of dihydropyridine calcium antagonists. *J. Chromatogr. A*, 694 (1995) 181-193 - a review with 96 refs.
- 3676 Volz, M., Mitrovic, V., Thiemer, J. and Schlepfer, M.: Steady-state plasma kinetics of slow-release propafenone, its two isomers and its main metabolites. *Arzneim.-Forsch.*, 45 (1995) 246-249.
- For additional information see *C.A.*:  
122 (1995) 45519e, 45530b, 71230y, 197104d.
- See also 2617, 2741, 3372, 3551, 3643, 3793.
- 32d. Central nervous system drugs**
- 3677 Al-Khamis, K.I., El-Sayed, Y.M., Al-Rashood, K.A. and Bawazir, S.A.: High performance liquid chromatographic determination of ranitidine in human plasma. *J. Liq. Chromatogr.*, 18 (1995) 277-286.
- 3678 Amin, R.D., Cheng, H. and Rogers, J.D.: Determination of MK-0476 in human plasma by liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 155-158.
- 3679 Bacracheva, N., Drenska, A., Gorantcheva, J., Tyutyulkova, N. and Vlahov, V.: Comparative bioavailability of two oral metamizole formulations. Influence of the acetylation phenotype. *Arzneim.-Forsch.*, 45 (1995) 282-285.
- 3680 Bansal, R. and Aranda, J.V.: Simultaneous microassay of alfentanil, fentanyl, and sufentanil by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 339-348.
- 3681 Barrett, D.A. and Shaw, P.N.: A stability indicating HPLC assay for diamorphine in aqueous solution. *J. Liq. Chromatogr.*, 17 (1994) 3727-3733.
- 3682 Berliz, G., Wintersteiger, R. and Pacha, W.: Determination of a novel xanthine derivative, MKS 213-492, in plasma by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 666 (1995) 368-373.
- 3683 Boughton, B.J., Sheehan, T.M.T., Wood, J., O'Brien, D., Butler, M., Simpson, A. and Hale, K.A.: High-performance liquid chromatographic assay of plasma thalidomide: stabilization of specimens and determination of a tentative therapeutic range for chronic graft-versus-host disease. *Ann. Clin. Biochem.*, 32 (1995) 79-83; *C.A.*, 122 (1995) 204389f.
- 3684 Bowyer, J.F., Clausing, P. and Newport, G.D.: Determination of d-amphetamine in biological samples using high-performance liquid chromatography after precolumn derivatization with o-phthalaldehyde and 3-mercaptopropionic acid. *J. Chromatogr. B*, 666 (1995) 241-250.
- 3685 Browne, T.R., Szabo, G.K., Davoudi, H. and Josephs, E.G.: Determination of pharmacokinetic drug interactions with carbamazepine and phenytoin using stable isotope labeling and simple high-performance liquid chromatography/ultraviolet detection technique. *Neurology*, 44 (1994) 2410-2411; *C.A.*, 122 (1995) 122371e.
- 3686 Cairns, A.M., Chiou, R.H.-Y., Rogers, C.J.D. and Demetriades, J.L.: Enantioselective high-performance liquid chromatographic determination of omeprazole in human plasma. *J. Chromatogr. B*, 666 (1995) 323-328.
- 3687 Chauert, N., Li, C., Ducharme, Y., Trimble, L.A., Ramachandran, C. and Nicoll-Griffith, D.A.: *In vitro* and *in vivo* biotransformations of the naphthalenic lignan lactone 5-lipoxygenase inhibitor, L-702,539. *Drug Metab. Disp.*, 23 (1995) 65-71.
- 3688 Chen, X., Lu, J. and Li, D.: (Simultaneous determination of theophylline, phenobarbital, phenytoin, and carbamazepine in human serum by high-performance liquid chromatography). *Zhongguo Yiyuan Yaoxue Zazhi*, 14 (1994) 389-391; *C.A.*, 122 (1995) 122396s.
- 3689 Delon, A., Favreliere, S., Couet, W., Courtois, P. and Bouquet, S.: Rapid and sensitive determination of thalidomide in human plasma by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 297-309.
- 3690 Dobrev, D., Berndt, A., Oertel, R., Richter, K., Terhaag, B. and Gramatté, T.: Untersuchung zur Bioäquivalenz von Metoclopramid in zwei Tablettenformulierungen. *Arzneim.-Forsch.*, 45 (1995) 304-306.
- 3691 El-Sayed, Y.M., Niazy, E.M. and Khidir, S.H.: High-performance liquid chromatographic method for the quantitative determination of pheniramine in plasma. *J. Liq. Chromatogr.*, 18 (1995) 763-777.
- 3692 Fadiran, E.O., Leslie, J., Fossler, M. and Young, D.: Determination of clozapine and its major metabolites in human serum and rat plasma by liquid chromatography using solid-phase extraction and ultraviolet detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 185-190.
- 3693 Feng, M.R., Siersma, P.R., Strenkoski, C.A. and Wright, D.S.: Sensitive high-performance liquid chromatographic method for a dopamine receptor agonist, CI-1007, and its metabolite PD 147693 in monkey plasma. *J. Chromatogr. B*, 665 (1995) 193-199.
- 3694 Garner, D.M., Pieniaszek, H.J., Jr., Connell, J.M. and Fiske, W.D.: Determination of linopirdine and its mono-N-oxide metabolite in human plasma and urine by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4541-4555.
- 3695 Jonczyk, A. and Wilcynska-Wojtulewicz, I.: Determination of diphenhydramine and naphazoline in pharmaceutical mixtures by high-performance liquid chromatography. *Acta Pol. Pharm.*, 51 (1994) 115-118; *C.A.*, 122 (1995) 88606k.
- 3696 Kapil, R.P., Lee, W., Jr., Padovani, P.K., Wong, N.Y., Holm, K.A. and Lam, G.N.: Determination of a novel cognitive enhancer, X9121, and its mono N-oxide metabolite, XG696, in dog plasma by reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 665 (1995) 226-231.

- 3697 Kondo, T., Hamasaki, K., Yamaguchi, M., Aoki, I., Yoshida, K., Yoshimura, Y. and Tanayama, S.: Enantioselective determination of pazinaclone, a new isoindoline anxiolytic, and its active metabolite in rat plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 291-297.
- 3698 Lacroix, P.M., Yat, P.N. and Lovering, E.G.: Liquid chromatographic methods for fluoxetine hydrochloride, its meta isomer, and related compounds in raw materials. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 334-339.
- 3699 Lee, D.-S., Kwon, K.-H. and Shin, H.-S.: Determination of cimetidine in human urine by column switching HPLC and UV-detection. *Bull. Korean Chem. Soc.*, 16 (1995) 77-79; *C.A.*, 122 (1995) 177582e.
- 3700 Lehr, G.J., Yuen, S.M. and Lawrence, G.D.: Liquid chromatographic determination of atropine in nerve gas antidotes and other dosage forms. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 339-343.
- 3701 Matsui, K., Oda, Y., Ohe, H., Tanaka, S. and Asakawa, N.: Direct determination of E2020 enantiomers in plasma by liquid chromatography-mass spectrometry and column-switching techniques. *J. Chromatogr. A*, 694 (1995) 209-218.
- 3702 Miratsuka, A., Chu, T.Y.Y., Distefano, E.W., Lin, L.Y., Schmitz, D.A. and Cho, A.K.: Inactivation of constitutive hepatic cytochromes P450 by phencyclidine in the rat. *Drug Metab. Disp.*, 23 (1995) 201-206.
- 3703 Naidong, W., Lee, J.W. and Hulise, J.D.: Development and validation of a chiral HPLC method for the quantitation of methocarbamol enantiomers in human plasma. *J. Liq. Chromatogr.*, 17 (1994) 3747-3758.
- 3704 Nishikawa, M., Tatsuno, M., Katagi, M. and Tsuchihashi, H.: (Determination of triazolam and its metabolites in human urine by EMIT, gas chromatography-mass spectrometry, and thermospray liquid chromatography-mass spectrometry) *Jpn. Toxicol. Environ. Health*, 41 (1995) 59-66; *C.A.*, 122 (1995) 232809m.
- 3705 Sueyasu, M., Ikeda, T., Otsubo, K., Taniyama, T., Aoyama, T. and Oishi, R.: Enantioselective determination of thiamylal in human serum by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 133-137.
- 3706 Terhechte, A. and Blaschke, G.: Investigation of the stereoselective metabolism of the chiral H<sub>1</sub>-antihistaminic drug terfenadine by high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 219-225.
- 3707 Torchio, L., Lombardi, F., Visconti, M. and Doyle, E.: Determination of the polar drug dimiracetam in human plasma and serum by column-switching high-performance liquid chromatography. *J. Chromatogr. B*, 666 (1995) 169-177.
- 3708 Tosunoglu, S. and Ersoy, L.: Determination of baclofen in human plasma and urine by high-performance liquid chromatography with fluorescence detection. *Analyst (Cambridge)*, 120 (1995) 373-375.
- 3709 Venkateshwaran, T.G. and Stewart, J.T.: HPLC determination of morphine-hydromorphone-bupivacaine and morphine-hydromorphone-tetracaine mixtures in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 18 (1995) 565-578.
- 3710 Venkateshwaran, T.G., King, D.T. and Stewart, J.T.: HPLC determination of a metoclopramide and ondansetron mixture in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 18 (1995) 117-126.
- 3711 Walily, A.E., Belal, S.F., Yakout, M.H. and Zamel, S.: Determination of mixture of acetaminofen and ibuprofen in tablets using derivative spectrophotometry and high-performance liquid chromatography. *Bull. Fac. Pharm. (Cairo Univ.)*, 31 (1993) 331-337; *C.A.*, 122 (1995) 197127p.
- 3712 Wang, M.Z., Klee, M.S. and Yang, S.K.: Achiral and chiral analysis of camazepam and metabolites by packed-column supercritical fluid chromatography. *J. Chromatogr. B*, 665 (1995) 139-146.
- 3713 Wu, W.-N., Masucci, J.A., McKown, L.A. and Rodgers, J.D.: Identification of four plasma metabolites of etoperidone in dogs. *J. Pharm. Biomed. Anal.*, 13 (1995) 177-184.

For additional information see C.A.:

122 (1995) 23084g, 115098t, 115128c, 177562y, 204420j.

See also 2583, 2701, 2741, 3621, 3632, 3650, 3659, 3788.

### 32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 3714 Cannavan, A., Blanchflower, W.J. and Kennedy, D.G.: Determination of levamisole in animal tissues using liquid chromatography - thermospray mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 331-333.
- 3715 Ficarra, R., Calabro, M.L., Tommasini, S., Grasso, S., Carulli, M., Monforte, A.M., Constantino, D. and Ficarra, P.: Enantiomeric resolution of anti-HIV agents on a cellulose derivative chiral stationary phase. Note II. *Chromatographia*, 40 (1995) 39-41.
- 3716 Flores-Murrieta, F.J., Granados-Soto, V. and Hong, E.: A simple and rapid method for determination of fluconazole in human plasma samples by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3803-3811.
- 3717 Garcia-Alvarez-Coque, M.C., Simo-Alfonso, E.F., Ramis-Ramos, G. and Esteve-Romero, J.S.: High-performance micellar liquid chromatography determination of sulphonamides in pharmaceuticals after azodye precolumn derivatization. *J. Pharm. Biomed. Anal.*, 13 (1995) 237-245.
- 3718 Holland, D.C., Munns, R.K., Roybal, J.E., Hurlbut, J.A. and Long, A.R.: Liquid chromatographic determination of the anticoccidial drug halofuginone hydrobromide in eggs. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 37-40.
- 3719 Horie, M., Saito, K., Nose, N. and Nakazawa, H.: (Simultaneous determination of sulfonamides and their N<sup>4</sup>-acetylated metabolites in meat by semi-micro high-performance liquid chromatography). *Kuromatogurafi*, 15 (1994) 254-255; *C.A.*, 122 (1995) 158856f.
- 3720 Hormazabal, V. and Yndestad, M.: High-performance liquid chromatographic determination of praziquantel in plasma and tissues of cultured fish for residue and pharmacokinetic studies. *J. Liq. Chromatogr.*, 18 (1995) 589-597.
- 3721 Huelsewede, J.W.: Comparison of high-performance liquid chromatography and bioassay for the determination of 5-fluorocytosine in serum. *Zentralbl. Bakteriol.*, 28 (1994) 513-518; *C.A.*, 122 (1995) 177548y.
- 3722 Humberstone, A.J., Currie, G.J., Porter, C.J.H., Scanlon, M.J. and Charman, W.N.: A simplified liquid chromatography assay for the quantitation of halofantrine and desbutylhalofantrine in plasma and identification of a degradation product of desbutylhalofantrine formed under alkaline conditions. *J. Pharm. Biomed. Anal.*, 13 (1995) 265-272.

- 3723 Idowu, O.R., Peggins, J.O., Brewer, T.G. and Kelley, C.: Metabolism of a candidate 8-aminoquinoline antimalarial agent, WR 238605, by rat liver microsomes. *Drug Metab. Disp.*, 23 (1995) 1-17.
- 3724 Kaka, J.S.: Rapid simplified analysis of pyrazinamide in rat plasma by HPLC. *J. Liq. Chromatogr.*, 17 (1994) 3793-3801.
- 3725 Kotecka, B.M. and Rieckmann, K.H.: *In vivo-in vitro* model to assess chloroquine activity in monkeys. *Chemotherapy (Basel)*, 41 (1995) 134-140; C.A., 122 (1995) 204419r.
- 3726 Munns, R.K., Turnipseed, S.B., Pfenning, A.P., Roybal, J.E., Holland, D.C. and Long, A.R.: Determination of residues of flumequine and nalidixic, oxolinic, and piromidic acids in catfish by liquid chromatography with fluorescence and UV detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 343-352.
- 3727 Nicolas, F., de Sousa, G., Thomas, P., Placidi, M., Lorenzon, G. and Rahmani, R.: Comparative metabolism of 3'-azido-3'-deoxythymidine in cultured hepatocytes from rats, dogs, monkeys, and humans. *Drug Metab. Disp.*, 23 (1995) 308-313.
- 3728 Nishimura, K., Hirama, Y., Kawano, J., Hiratsuka, A. and Nakano, M.: (Simultaneous determination of synthetic antibacterials in meat and fish by HPLC. II. Studies of cleanup methods with liquid-liquid extraction). *Hokkaidoritsu Eisei Kenkyushoho*, 44 (1994) 66-68; C.A., 122 (1995) 79331c.
- 3729 Poirier, J.-M., Lebot, M., Descamps, P., Levy, M. and Cheymol, G.: Determination of itraconazole and its active metabolite in plasma by column liquid chromatography. *Ther. Drug Monit.*, 16 (1994) 596-601; C.A., 122 (1995) 71208x.
- 3730 Rifai, N., Sakamoto, M., Law, T., Platt, O., Mikati, M., Armsby, C.C. and Brugnara, C.: HPLC measurement, blood distribution, and pharmacokinetics of oral clotrimazole, potentially useful anticickling agent. *Clin. Chem. (Washington)*, 41 (1995) 387-391.
- 3731 Roth, J.S. and Kelley, J.A.: Determination of the anti-HIV drug 2'- $\beta$ -fluoro-2',3'-dideoxyadenosine in biological fluids by reversed-phase HPLC. *J. Liq. Chromatogr.*, 18 (1995) 441-462.
- 3732 Schenck, F.J.: Isolation and quantification of ivermectin in bovine milk by matrix solid phase dispersion (MSPD) extraction and liquid chromatographic determination. *J. Liq. Chromatogr.*, 18 (1995) 349-362.
- 3733 Smedley, M.D.: Liquid chromatographic determination of multiple sulfonamide residues in bovine milk. Collaborative study. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1112-1122.
- 3734 Tachibana, M., Aoyama, M., Taniguchi, R., Anabuki, K. and Kumagai, K.: (Simultaneous determination of residual synthetic antibacterials in chicken by HPLC). *Tokyo-to Suginami-ku Eisei Shikensho Nenpo*, 12 (1994) 86-89; C.A., 122 (1995) 103961f.
- 3735 Takahashi, Y., Sekiya, T., Nishikawa, M. and Endoh, Y.S.: Simultaneous high-performance liquid chromatographic determination of amprolium, ethopabate, sulfaquinoxaline and N4-acetylsulfaquinoxaline in chicken tissues. *J. Liq. Chromatogr.*, 17 (1994) 4489-4512.
- 3736 Tendolkar, N.M., Desai, B.S. and Shinde, V.M.: Simultaneous determination of tinidazole and clotrimazole from tablets by RP-HPLC. *Indian Drugs*, 31 (1994) 551-553; C.A., 122 (1995) 89592c.
- 3737 Tsai, C.-E. and Kondo, F.: A sensitive high-performance liquid chromatographic method for detecting sulfonamide residues in swine serum and tissues after fluorescamine derivatization. *J. Liq. Chromatogr.*, 18 (1995) 965-976.
- 3738 Vergin, H., Kikuta, C., Macher, H. and Metz, R.: Pharmacokinetics and bioavailability of different formulations of aciclovir. *Arzneim.-Forsch.*, 45 (1995) 508-515.
- 3739 Wei, Y., Nygard, G.A. and Khalil, S.K.W.: A HPLC method for the separation and quantification of the enantiomers of hydroxychloroquine and its three major metabolites. *J. Liq. Chromatogr.*, 17 (1994) 3479-3490.
- For additional information see C.A.:  
122 (1995) 115116x, 115143d, 197142q, 197143r, 197151s.
- See also 3540, 3591, 3794.
- 32f. *Cytostatics*
- 3740 Albertioni, F., Pettersson, B., Beck, O., Rask, C., Seideman, P. and Peterson, C.: Simultaneous quantitation of methotrexate and its two main metabolites in biological fluids by a novel solid-phase extraction procedure using high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 163-170.
- 3741 Burk, M., Volmer, M., Fartash, K. and Schneider, W.: Ion-pair liquid chromatography of cytarabine and uracil-arabinoside in human plasma. *Arzneim.-Forsch.*, 45 (1995) 616-619.
- 3742 Freeman, K.B., Anliker, S., Hamilton, M., Osborne, D., Dhahir, P.H., Nelson, R. and Allerheiligen, S.R.B.: Validated assays for the determination of gemcitabine in human plasma and urine using high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 665 (1995) 171-181.
- 3743 Himberg, J.-P., Koskinen, M., Riekkola, M.-L. and Elo, H.: High-performance liquid chromatographic separation and determination of salicylaldoxime and  $\beta$ -resorcyldoxime in aqueous solutions and in human urine. *J. Liq. Chromatogr.*, 17 (1994) 3519-3529.
- 3744 Lu, G. and Jun, H.W.: Determination of trace methotrexate and 7-OH-methotrexate in plasma by high-performance liquid chromatography with fluorimetric detection. *J. Liq. Chromatogr.*, 18 (1995) 155-171.
- 3745 Mank, A.J.G., Molenaar, E.J., Gooijer, C., Lingeman, H., Velthorst, N.H. and Brinkman, U.A.T.: Determination of the anticancer drug metabolite WR1065 using pre-column derivatization and diode laser induced fluorescence detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 255-263.
- 3746 Poon, G.K., Walter, B., Lonning, P.E., Horton, M.N. and McCague, R.: Identification of tamoxifen metabolites in human Hep G2 cell line, human liver homogenate, and patients on long-term therapy for breast cancer. *Drug Metab. Disp.*, 23 (1995) 377-382.
- 3747 Qian, M., O'Dwyer, P.J. and Gallo, J.M.: High-performance liquid chromatographic method for the determination of AG-331, a novel anti-cancer agent, in human serum and urine using solid-phase extraction and photodiode-array detection. *J. Chromatogr. B*, 666 (1995) 307-314.
- 3748 Ramsauer, B., Braess, J., Unterhalt, M., Kaufmann, C.C., Hidde-mann, W. and Schleyer, E.: Highly sensitive high-performance liquid chromatographic assay for 1- $\beta$ -D-arabinofuranosylcytosine-5'-stearyl phosphate (cytarabine-ocfosfate). *J. Chromatogr. B*, 665 (1995) 183-192.
- 3749 Saini, P., Abu-Izza, K. and Lu, D.R.: High-performance liquid chromatographic assay for sodium mercaptoundecahydrododecaborate in rat tissues. *J. Chromatogr. B*, 665 (1995) 155-161.



- 3750 Walle, T., Walle, U.K., Kumar, G.N. and Bhalla, K.N.: Taxol metabolism and disposition in cancer patients. *Drug Metab. Disp.*, 23 (1995) 506-512.
- 3751 Webster, L.K., Ellis, A.G., Kestell, P. and Rewcastle, G.W.: Metabolism and elimination of 5,6-dimethylxanthenone-4-acetic acid in the isolated perfused rat liver. *Drug Metab. Disp.*, 23 (1995) 363-368.
- 3752 Yatsевич, O.V., Arzamashev, A.P., Zimakova, N.I., Abrazakov, M.M. and Rodionova, G.M.: Quantitative determination of sarcosyl and its hydrolysis products by high-performance liquid chromatography. *Khim.-Farm. Zh.*, 28 (1994) 56-59; C.A., 122 (1995) 89589g.
- For additional information see C.A.:  
122 (1995) 71221w, 204406j, 230000s, 230001t.
- See also 3510, 3542.
- 32g. *Other drug categories*
- 3753 Bo, L.D., Verga, F., Marzo, A., Ambrosoli, L. and Poli, A.: Determination of ticlopidine in human plasma by high-performance liquid chromatography and ultraviolet absorbance detection. *J. Chromatogr. B*, 665 (1995) 404-409.
- 3754 Brunn, H. and Georgii, S.: (Qualitative and quantitative detection of dexamethasone and related xenobiotic corticosteroids in bovine urine by ELISA and HPLC/ELISA). *Arch. Lebensmittelhyg.*, 45 (1994) 66-71; C.A., 122 (1995) 178543e.
- 3755 Caraballo, I., Fernandez-Arevalo, M., Holgado, M.A., Alvarez-Fuentes, J. and Rabasco, A.M.: Simultaneous HPLC determination of some drugs commonly used in cold medications: dextromethorphan, diphenhydramine, phenylephrine, phenylpropranolamine and pseudoephedrine. *Drug Dev. Ind. Pharm.*, 21 (1995) 605-613; C.A., 122 (1995) 142731a.
- 3756 Chiarotto, J.A. and Wainer, I.W.: Determination of metyrapone and the enantiomers of its chiral metabolite metyrapol in human plasma and urine using coupled achiral-chiral liquid chromatography. *J. Chromatogr. B*, 665 (1995) 147-154.
- 3757 Cohen, H., Armstrong, F. and Campbell, H.: Sensitive fluorescence detection of robenidine by derivatization with dansyl chloride and high-performance liquid chromatography. *J. Chromatogr. A*, 694 (1995) 407-413.
- 3758 Constanzer, M., Chavez, C. and Matuszewski, B.: Development and comparison of high-performance liquid chromatographic methods with tandem mass spectrometric and ultraviolet absorbance detection for the determination of cyclobenzaprine in human plasma and urine. *J. Chromatogr. B*, 666 (1995) 117-126.
- 3759 Crimella, T., Orlandi, R., Bocchiola, G., Anders, U. and Stradi, R.: Analytical and chemical profile of pidotimod. *Arzneim.-Forsch.*, 44 (1994) 1405-1410; C.A., 122 (1995) 197136r.
- 3760 Davis, M.R. and Quigley, M.N.: Liquid chromatographic determination of UV absorbers in sunscreens. *J. Chem. Educ.*, 72 (1995) 279-281; C.A., 122 (1995) 213167j.
- 3761 Dias, V.C., Lucien, J., LeGat, D.F. and Yatscoff, R.W.: Measurement of the active leflunomide metabolite (A77 1726) by reverse-phase high-performance liquid chromatography. *Ther. Drug Monit.*, 17 (1995) 84-88; C.A., 122 (1995) 177553w.
- 3762 El-Ealily, A.F.M., El-Anwar, F.M. and El-Swify, M.: Ion-pair HPLC analysis of multi-ingredient sedative-cough syrup. *Bull. Fac. Pharm. (Cairo Univ.)*, 31 (1993) 297-301; C.A., 122 (1995) 197123j.
- 3763 Farag, S.A.: Liquid chromatographic assay of diatrizoic acid and its diiodo degradation products in radio-opaque solutions. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 328-333.
- 3764 Gliniski, J.A. and Caviness, G.O.: Centrifugal partition chromatography in assay guided isolation of natural products. A case study of immunosuppressive components of *Tripterygium wilfordii*. *Chromatogr. Sci. Ser.*, 68(Centrifugal Partition Chromatography) (1995) 133-165; C.A., 122 (1995) 170309n - a review with 42 refs.
- 3765 Gordon, B., Marchant, N., Richards, R. and Walther, B.: Methods for the analysis of benfluorex metabolites. *Methodol. Surv. Bioanal. Drugs*, 23(Biofluid and Tissue Analysis for Drugs, Including Hypolipidaemics) (1994) 103-112; C.A., 122 (1995) 229986m.
- 3766 Guillaume, Y., Peyrin, E. and Guinchard, C.: Rapid determination of sulbactam and tazobactam in human serum by high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 363-371.
- 3767 Gumieniczek, A. and Przyborowski, L.: Determination of benzobromarone and benzarone in pharmaceuticals by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3411-3419.
- 3768 Herzog, R. and Leuschner, J.: Experimental studies on the pharmacokinetics and toxicity of 5-aminosalicylic acid-O-sulfate following local and systemic application. *Arzneim.-Forsch.*, 45 (1995) 300-303.
- 3769 Ho, Y., Lee, Y.-L. and Hsu, K.-Y.: Determination of (+)-catechin in plasma by high-performance liquid chromatography using fluorescence detection. *J. Chromatogr. B*, 665 (1995) 383-389.
- 3770 Noroski, J.E. and Mayo, D.J.: Chiral LC of a cholesterol-lowering drug using serum albumin mobile phases. *J. Pharm. Biomed. Anal.*, 13 (1995) 83-87.
- 3771 Nowotnik, D.P., Nanjappan, P., Zeng, W. and Ramalingam, K.: Separation of the stereoisomers of hexamethyl-propyleneamine oxime (HM-PAO) by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 673-687.
- 3772 Pesonen, T., Holmalahti, J. and Pohjola, J.: Determination of chlorhexidine in saliva using high-performance liquid chromatography. *J. Chromatogr. B*, 665 (1995) 222-225.
- 3773 Rajewski, R.A., Kosednar, D.G., Matches, T.A., Wong, O.S., Burchett, K. and Thakker, K.: Stereo-specific analysis of a novel protein kinase C inhibitor. *J. Pharm. Biomed. Anal.*, 13 (1995) 247-253.
- 3774 Rastogi, S.C., Schouten, A., de Kruijff, N. and Weijland, J.W.: Contents of methyl-, ethyl-, propyl-, butyl- and benzylparaben in cosmetic products. *Contact Dermatitis*, 32 (1995) 28-30; C.A., 122 (1995) 142007u.
- 3775 Sallustio, B.C. and Fairchild, B.A.: Biosynthesis, characterisation and direct high-performance liquid chromatographic analysis of gemfibrozil 1-O- $\beta$ -acylglucuronide. *J. Chromatogr. B*, 665 (1995) 345-353.

- 3776 Vielhauer, S., Rudolphi, A., Boos, K.-S. and Seidel, D.: Evaluation and routine application of the novel restricted-access precolumn packing material Alkyl-Diol Silica: coupled-column high-performance liquid chromatographic analysis of the photoreactive drug 8-methoxypsoralen in plasma. *J. Chromatogr. B*, 666 (1995) 315-322.
- 3777 Wallner, P.: HPLC analysis of UVA and -B filters in cosmetics. *Dtsch. Lebensm.-Rundsch.*, 89 (1993) 375-379; *C.A.*, 122 (1995) 142002p.
- 3778 Wang, C.P., Chan, K.W., Schiksnis, R.A., Scatina, J. and Sisenwine, S.F.: High performance liquid chromatographic isolation, spectroscopic characterization, and immunosuppressive activities of two rapamycin degradation products. *J. Liq. Chromatogr.*, 17 (1994) 3383-3392.
- For additional information see *C.A.*:  
122 (1995) 45518d, 89602f, 229990h, 229991j, 230002u.
- See also 2725, 2822, 2868, 3406, 3464, 3537, 3560, 3571, 3572, 3650, 3655, 3699, 3729, 3746, 3819, 3835.
- 32h. *Toxicological and forensic applications*
- 3779 Ardrey, R.E., Kimber, K.A. and Kilpatrick, G.: LC-MS - a routine analytical tool in forensic toxicology. In: Oliver, J.S. (Editor), *Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol., 26th 1992*, Scot. Acad. Press, Edinburgh, 1992, pp. 103-119; *C.A.*, 122 (1995) 232815k.
- 3780 Bogusz, M. and Erkens, M.: Influence of biological matrix on chromatographic behavior and detection of selected acidic, neutral, and basic drugs examined by means of a standardized HPLC-DAD system. *J. Anal. Toxicol.*, 19 (1995) 49-55.
- 3781 Carretero, L., Maldonado, M. and Laserna, J.J.: (Laser-induced fluorescence for detection of drugs banned in sports by micellar liquid chromatography). *Quim. Anal. (Barcelona)*, 11 (1992) 107-119; *C.A.*, 122 (1995) 48615u.
- 3782 Casale, J.F. and Moore, J.M.: Detection and determination of pseudococaine in coca leaves and illicit cocaine samples. *J. Forensic Sci.*, 39 (1994) 1537-1543; *C.A.*, 122 (1995) 48624w.
- 3783 Chen, H., Ron, N., Ma, Q., Han, N. and Fan, D.: (Reverse phase liquid chromatography for analyzing Saikuzuo residue in field atmosphere, brown rice, hull and straw). *Huanjing Wuran Yu Fangzhi*, 16 (1994) 35-36; *C.A.*, 122 (1995) 74269d.
- 3784 Chen, H.-M. and Peterson, C.M.: Quantifying ethanol by high-performance liquid chromatography with precolumn enzymic conversion and derivatization with fluorometric detection. *Alcohol (N.Y.)*, 11 (1994) 577-582; *C.A.*, 122 (1995) 48567e.
- 3785 Debruyne, D., Albessard, F., Bigot, M.C. and Moulin, M.: Comparison of three advanced chromatographic techniques for cannabis identification. *Bull. Narc.*, 46 (1994) 109-121; *C.A.*, 122 (1995) 74099y.
- 3786 Furness, J.F., Weekes, A., Clench, M., Wolff, K., Barnes, I., Hay, A.W.M. and Cooke, M.: The determination of methadone and metabolites in human urine by HPLC with ultraviolet, and particle beam mass spectrometric detection. *J. Liq. Chromatogr.*, 17 (1994) 4431-4444.
- 3787 Hallbach, J. and Guder, W.G.: Experiences with HPLC and/or GC-MS as a first step for drugs of abuse testing and clinical toxicology. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A29.
- 3788 Lambert, W.E., Meyer, E., Xue-Ping, Y. and de Leenheer, A.P.: Screening, identification, and quantitation of benzodiazepines in postmortem samples by HPLC with photodiode array detection. *J. Anal. Toxicol.*, 19 (1995) 35-40.
- 3789 Lehman, T. and Brenneisen, R.: High performance liquid chromatographic profiling of cannabis products. *J. Liq. Chromatogr.*, 18 (1995) 689-700.
- 3790 Logan, B.K. and Stafford, D.T.: In-line solid phase sample preparation for analysis of drugs in postmortem fluids by high performance liquid chromatography. In: Oliver, J.S. (Editor), *Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol., 26th 1992*, Scot. Acad. Press, Edinburgh, 1992, pp. 288-294; *C.A.*, 122 (1995) 207218k.
- 3791 Marr, J.C., McDowell, L.M. and Quilliam, M.A.: Investigation of derivatization reagents for the analysis of diarrhetic shellfish poisoning toxins by liquid chromatography with fluorescence detection. *Nat. Toxins*, 2 (1994) 302-311; *C.A.*, 122 (1995) 25421p.
- 3792 Nishikawa, M., Tatsuno, M., Tsuchihashi, H., Igarashi, K., Kasuya, F. and Fukui, M.: (The analysis of quaternary ammonium compounds in human urine by thermospray liquid chromatography-mass spectrometry). *Jpn. J. Toxicol. Environ. Health*, 40 (1994) 534-541; *C.A.*, 122 (1995) 125463x.
- 3793 Poletini, A., Montagna, M., Hogendoorn, E.A., Dijkman, E., van Zoonen, P. and van Ginkel, L.A.: Applicability of coupled-column liquid chromatography to the analysis of  $\beta$ -agonists in urine by direct sample injection. I. Development of a single-residue reversed-phase liquid chromatography-UV method for clenbuterol and selection of chromatographic conditions suitable for multi-residue analysis. *J. Chromatogr. A*, 695 (1995) 191-31.
- 3794 Roenn, A.M., Lemnge, M.M., Angelo, H.R. and Bygbjerg, I.C.: High-performance liquid chromatography determination of dapson, monoacetyldapson, and pyrimethamine in filter paper blood spots. *Ther. Drug Monit.*, 17 (1995) 79-83; *C.A.*, 122 (1995) 177552v.
- 3795 Rositano, J. and Nicholson, B.C.: A method for the detection of cyanobacterial peptide toxins by HPLC. *Spec. Publ. - R. Soc. Chem.*, 149 (Detection Methods for Cyanobacterial Toxins) (1994) 155-157; *C.A.*, 122 (1995) 74064h.
- 3796 Sabater Montesinos, J., Garcia Alvarez-Coque, M.C., Ramis Ramos, G. and Laserna Vazquez, J.J.: On the use of micellar liquid chromatography for detection of drug misuse in sport. *Quim. Anal. (Barcelona)*, 11 (1992) 163-171; *C.A.*, 122 (1995) 48616v.
- 3797 Stockemer, J. and Guerke, M.: (Optimization of a HPLC method for the determination of DSP toxins). *Arch. Lebensmittelhyg.*, 44 (1993) 146-147; *C.A.*, 122 (1995) 131263y.
- 3798 Walker, J.A., Kkrueger, S.T., Lurie, I.S., Marche, H.L. and Newby, N.: Analysis of heroin drug seizures by micellar electrokinetic capillary chromatography (MECC). *J. Forensic Sci.*, 40 (1995) 6-9; *C.A.*, 122 (1995) 207200y.
- For additional information see *C.A.*:  
122 (1995) 95762z.
- See also 2777, 2789, 2801, 2937, 3042, 3443, 3445, 3450, 3521, 3704.

## 32i. Plant extracts

- 3799 Basha, L.I.A., Rashed, M.S. and Aboul-Enein, H.Y.: TLC assay of thymoquinone in black seed oil (*Nigella sativa* L.) and identification of dithymoquinone and thymol. *J. Liq. Chromatogr.*, 18 (1995) 105-115.
- 3800 Collins, M.D., Wasmund, L.M. and Bosland, P.W.: Improved method for quantifying capsaicinoids in *Capsicum* using HPLC. *HortScience*, 30 (1995) 137-139; *C.A.*, 122 (1995) 182312w.
- 3801 Graenicher, F., Christen, P. and Vuagnat, P.: Rapid high performance liquid chromatographic quantification of valepotriates in hair root cultures of *Valeriana officinalis* L. var. *sambucifolia* Mikan. *Phytochem. Anal.*, 5 (1994) 297-301; *C.A.*, 122 (1995) 89574y.
- 3802 Singh, B.K., Teclé, B. and Shaner, D.L.: Determination of 2-keto acids and amino acids in plant extracts. *J. Liq. Chromatogr.*, 17 (1994) 4469-4477.
- 3803 Tarantilis, P.A., Tsoupras, G. and Polissiou, M.: Determination of saffron (*Crocus sativus* L.) components in crude plant extract using high-performance liquid chromatography-UV-visible photodiode-array detection-mass spectrometry. *J. Chromatogr. A*, 699 (1995) 107-118.
- 3804 Tzakou, O., Verekokidou, E. and Harvala, C.: Comparative HPLC analysis of polyphenolic compounds in four species of *Galium* L. *J. Liq. Chromatogr.*, 17 (1994) 4463-4467.
- 3805 Yang, F., Ou, Q. and Yu, W.: Semi-preparative separation of taraxeryl-acetate and coumarins from *Artemisia dalailamae* Kraschen by high-speed countercurrent chromatography. *J. Liq. Chromatogr.*, 18 (1995) 395-403.

For additional information see *C.A.*:

122 (1995) 89595f, 170330n, 170332q, 196659b, 197105e.

See also 2805, 2827, 2882, 2979, 2982, 3764, 3785, 3789.

## 33. CLINICO-CHEMICAL APPLICATIONS

## 33a. General papers and reviews

- 3806 Riley, C.M., Lough, W.J. and Wainer, I.W. (Editors): *Pharmaceutical and Biomedical Applications of Liquid Chromatography Prog. Pharm. Biomed. Anal.*, Vol. 1, Pergamon, Oxford, 1994, 379 pp.; *C.A.*, 122 (1995) 234817m.

## 33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 3807 Idei, M., Pajor, A. and Menyhart, J.: Bioanalysis of human amniotic fluid: chromatographical and biological characterization of a component inhibiting contraction of the myometrium. In: Kalasz, H., Ettre, L.S. and Pick, J. (Editors), *New Approaches Chromatogr. '93, [Pap. Budapest Chromatogr. Conf.], 1990*, Fekete Sas Kiado, Budapest, 1993, pp. 103-123; *C.A.*, 122 (1995) 210391s.

See also 2787, 2836, 2839, 2843, 2857, 2902, 2918, 2933, 2954, 2955, 2961, 2963, 2970, 2996, 2997, 3008, 3021, 3027, 3030, 3038, 3043, 3051, 3078, 3186, 3189, 3197, 3198, 3241, 3242, 3319, 3351, 3426, 3456, 3464, 3468, 3470, 3515, 3746, 3750, 3865, 3899.

## 34. FOOD ANALYSIS

- 3808 Yueming, Q. and Wuji, Z.: (Chemical reaction and chromatographic method for the determination of dimehypo residues in vegetables). *Fenzi Huaxue*, 22 (1994) 899-901; *C.A.*, 122 (1995) 79348p.

See also 3022.

## 34a. General papers and reviews

- 3809 Doneche, B. (Editor): *Modern Advances in the Chromatography of Wine: Applications to the Sensory Analysis of Wines, Porto, Port., Mar. 31-33 April, 1992*. Tec & Doc - Lavoisier, Paris, 1993, 253 pp.; *C.A.*, 122 (1995) 104523b.

## 34b. Complex mixtures (single compounds by cross-reference only)

- 3810 De Maria, C.A.B., Trugo, L.C., Moreira, R.F.A. and Petracco, M.: Simultaneous determination of total chlorogenic acid, trigonelline and caffeine in green coffee samples by high performance gel filtration chromatography. *Food Chem.*, 52 (1995) 447-449; *C.A.*, 122 (1995) 185709s.
- 3811 Golc-Wondra, A., Skocir, E. and Prosek, M.: Determination of monosodium glutamate in food products. *J. Planar Chromatogr.*, 8 (1995) 117-121.
- 3812 Juelich, E. and Lind, G.: (Determination of sugar, GdL, and sodium ascorbate in seasonings for sausages). *Fleischwirtschaft*, 74 (1994) 637-638; *C.A.*, 122 (1995) 185635q.
- 3813 Kermasha, S., Goetghebeur, M. and Dumont, J.: Separation and characterization of pepper-contaminated cinnamon using HPLC analyses. *Food Sci. Technol. (London)*, 27 (1994) 578-582; *C.A.*, 122 (1995) 54386m.
- 3814 Krueger, E., Radau, B. and Roenn, D.: (Investigation of beers and beer heads by methods employed in protein chemistry). *Monatsschr. Brauwiss.*, 46 (1993) 312-319; *C.A.*, 122 (1995) 131260v.
- 3815 Lancaster, F.E. and Lawrence, J.F.: Determination of annatto in high-fat dairy products, margarine and hard candy by solvent extraction followed by high-performance liquid chromatography. *Food Addit. Contam.*, 12 (1995) 9-19; *C.A.*, 122 (1995) 238043d.
- 3816 Madigan, D., McMurrrough, I. and Smyth, M.R.: Rapid determination of 4-vinyl guaiacol and ferulic acid in beers and worts by high-performance liquid chromatography. *J. Am. Soc. Brew. Chem.*, 52 (1994) 152-155; *C.A.*, 122 (1995) 54374f.
- 3817 Rimedio, N.T. and Chen, L.F.: Chromatographic separation of organic nonsugars, colloidal material and inorganic-organic complexes from juice, liquors, syrups and/or molasses. *U.S. US 5,382,294* (Cl. 127-42; C1311/06), 17 Jan. 1995, US Appl. 749,667, 26 Aug. 1991; 25 p.; *C.A.*, 122 (1995) 190875x.

- 3818 Van Loo, J., Booten, K. and Smits, G.: Chromatographic separation of polydispersed saccharide compositions to give products useful in foods. *PCT Int. Appl. WO 94 12,541* (Cl. C08837/18), 09 Jun. 1994, BE Appl. 92/1,021, 24 Nov. 1992; 47 p.; C.A., 122 (1995) 109228u.
- 3819 Zhu, Z.: (Rapid determination of dexamethasone acetate residue in pork by high-performance liquid chromatography). *Fenxi Huaxue*, 22 (1994) 1163-1166; C.A., 122 (1995) 185693g.
- For additional information see C.A.:  
122 (1995) 104175q, 158880j, 163938p, 212360e.
- See also 2793, 2795, 2799, 2812, 2813, 2814, 2815, 2816, 2818, 2824, 2826, 2829, 2831, 2849, 2850, 2854, 2890, 2892, 2897, 2898, 2900, 2908, 2911, 2915, 2926, 2936, 2952, 2974, 2990, 2999, 3011, 3013, 3023, 3024, 3035, 3044, 3083, 3245, 3358, 3428, 3459, 3480, 3501, 3504, 3506, 3517, 3526, 3530, 3533, 3534, 3535, 3546, 3547, 3558, 3559, 3566, 3574, 3580, 3586, 3589, 3590, 3591, 3593, 3719, 3728, 3732, 3733, 3734, 3735, 3737, 3797, 3844, 3850, 3861, 3889, 3891, 3901.
- 34c. *Organoleptically important compounds (flavors, odors, volatiles)*
- See 3459.
35. ENVIRONMENTAL ANALYSIS
- 35a. *General papers and reviews*
- 3820 Conrad, E.E., Kelly, K.P. and Schwartz, N.L.: GPC and oily samples. *Environ. Test. Anal.*, 3 (1994) 34-39; C.A., 122 (1995) 204159f - a review with 3 refs.
- 3821 Fort, D.J., Delphon, J., Powers, C.R., Helems, R., Gonzalez, R. and Stover, E.L.: Development of automated methods of identifying toxicants in the environment. *Bull. Environ. Contam. Toxicol.*, 54 (1995) 104-111; C.A., 122 (1995) 232799h.
- 3822 Hayakawa, K.: (Development of new photometric detection in liquid chromatography for the determination of environmental compounds). *Yakugaku Zasshi*, 114 (1994) 669-680; C.A., 122 (1995) 182254d - a review with 39 refs.
- See also 2624, 2709, 2767, 2773, 2778, 2781, 3867.
- 35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*
- See 2697, 3473, 3900.
- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*
- 3823 Elvira-Cozar, C., Cano-Faura, P., Perez-Arribas, L.V., Leon-Gonzalez, M.E. and Polo-Diez, L.M.: Trace priority pollutant phenols enrichment from water by ion chromatography. *Chromatographia*, 40 (1995) 91-95.
- 3824 Minnaard, W.A., Slobodnik, J., Vreuls, J.J., Hupe, K.-P. and Brinkman, U.A.T.: Rapid liquid chromatographic screening of organic micropollutants in aqueous samples using a single short column for trace enrichment and separation. *J. Chromatogr. A*, 696 (1995) 333-340.
- 3825 Pocurull, E., Marce, R.M. and Borrull, F.: Liquid chromatography of phenolic compounds in natural water using on-line trace enrichment. *Chromatographia*, 40 (1995) 85-90.
- 3826 Zapf, A., Heyer, R. and Stan, H.-J.: Rapid micro liquid-liquid extraction method for trace analysis of organic contaminants in drinking water. *J. Chromatogr. A*, 694 (1995) 453-461.
- See also 2764, 2794, 2797, 2919, 2920, 3047, 3576, 3578, 3579, 3582, 3585, 3587, 3878, 3882, 3885, 3898.
- 35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*
- 3827 Osipov, G.A., Nazina, T.N. and Ivanova, A.E.: (Analysis of microbial community from water-flooded oil field by chromatography-mass spectrometry). *Mikrobiologiya*, 63 (1994) 876-882; C.A., 122 (1995) 75689c.
- See also 2764, 2786, 3857.
36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES
- 36a. *Surfactants*
- 3828 Brenner, C., Jan, G., Chevalier, Y. and Wroblewski, H.: Evaluation of the efficacy of zwitterionic dodecyl carboxybetaine surfactants for the extraction and the separation of mycoplasma membrane protein antigens. *Anal. Biochem.*, 224 (1995) 515-523.
- 3829 Forgács, E. and Cserháti, T.: Retention behaviour of nonyl-phenyl ethyleneoxide oligomers on alumina columns. *Fresenius J. Anal. Chem.*, 351 (1995) 688-689.
- 3830 Marcomini, A. and Zanette, M.: Derivatization procedures and HPLC separations for environmental analysis of aliphatic alcohols polyethylenglycol ethers (AE). *Riv. Ital. Sostanze Grasse*, 71 (1994) 203-208; C.A., 122 (1995) 136796j.
- 3831 Matsuzaki, M., Ishii, K., Yoshimura, H. and Hashimoto, S.: HPLC determination of cationic and amphoteric surfactants in cosmetics. *J. SCCJ*, 27 (1993) 494-497; C.A., 120 (1994) 307031w.
- 3832 McPherson, B.P. and Rasmussen, H.T.: Chromatography of cationic surfactants: HPLC, TLC and GLC. *Surfactant Sci. Ser.*, 53 (1994) 289-326; C.A., 122 (1995) 136787g - a review with 89 refs.
- See also 2934, 3843.
- 36b. *Antioxidants and preservatives*
- 3833 Parhizkari, G., Miller, R.B. and Chen, C.: A stability-indicating HPLC method for the determination of benzalkonium chloride in phenylephrine HCl 10% ophthalmic solution. *J. Liq. Chromatogr.*, 18 (1995) 553-563.

- 3834 Rose, R.C. and Bode, A.M.: Analysis of water-soluble antioxidants by high-pressure liquid chromatography. *Biochem. J.*, 306 (1995) 101-105.
- 3835 Weyland, J.W., Stern, A. and Rooselaar, J.: Determination of bonopol, bronidox, and methyl dibromo glutaronitrile in cosmetics by liquid chromatography with electrochemical detection. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1132-1136.
- 36c. *Complex mixtures, technical products and unidentified compounds*
- 3836 Heikkilä, H., Hyöky, G. and Kuisma, J.: Fractionating sulphite cooking liquors by simulated moving bed chromatography. *PCT Int. Appl. WO 94 26,380* (Cl. B01D15/08) 24 Nov. 1994, FI Appl. 93/2,108, 10 May 1993; 38 p.; C.A., 122 (1995) 109957f.
- 3837 Lawrence, J.F., Ménard, C. and Cleroux, C.: Evaluation of pre-chromatographic oxidation for liquid chromatographic determination of paralytic shellfish poisons in shellfish. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 514-520.
- 3838 Lehotay, J. and Oktavec, D.: Application of HPLC with diode array detection in tribology. *J. Liq. Chromatogr.*, 17 (1994) 3783-3792.
- 3839 Oshima, Y.: Postcolumn derivatization liquid chromatographic method for paralytic shellfish toxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 528-532.
- 3840 Quilliam, M.A.: Analysis of diarrhetic shellfish poisoning toxins in shellfish tissue by liquid chromatography with fluorometric and mass spectrometric detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 555-570.
- 3841 Sassiati, P., Machtalere, G., Hui, F., Kolodziejczyk, H. and Rosset, R.: Liquid chromatographic determination of base oil composition and content in lubricating oils containing dispersants of the polybutenylsuccinimide type. *Anal. Chim. Acta.*, 306 (1995) 73-79.
- 3842 Swerin, A. and Waagberg, L.: Size-exclusion chromatography for characterization of cationic polyelectrolytes used in papermaking. *Nord. Pulp. Pap. Res. J.*, 9 (1994) 18-25; C.A., 122 (1995) 84033m.
- 3843 Taylor, C.D., Anderson, H.A. and Brydon, D.L.: Characterization of textile lubricants by HPLC. Part III: investigation of emulsions. *J. Text. Inst.*, 85 (1994) 49-56; C.A., 122 (1995) 204239g.
- 3844 Yasumoto, T., Fukui, M., Sasaki, K. and Sugiyama, K.: Determination of marine toxins in foods. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 574-582.
- For additional information see C.A.:  
122 (1995) 75712e, 85174v, 165048x.
- See also 2588, 2615.
37. **CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES**
- 3845 Chun, J.-H. and Yang, S.-M.: Electrokinetic effect on separation of colloidal particles in capillary hydrodynamic chromatography (CHDC). In: *Off. Proc. Comb. Conf., 6th Conf. Asia Pac. Confed. Chem. Eng., 21st Australas Chem. Eng. Conf., Inst. Eng., Aust., Barton, 1993*, pp. 283/2-287/2; C.A., 122 (1995) 84499t.
- 3846 Mandrusov, E., Houng, A., Klein, E. and Leonard, E.F.: Membrane-based cell affinity chromatography to retrieve viable cells. *Biotechnol. Prog.*, 11 (1995) 208-213; C.A., 122 (1995) 208889s.
- 3847 Nandan, D., Daubenberger, C., Mpimbaza, G. and Pearson, T.W.: A rapid single-step purification method for immunogenic members of the hsp 70 family: validation and application. *J. Immunol. Methods*, 176 (1994) 255-263; C.A., 122 (1995) 26992n.
- 3848 Qin, A. and Cui, Z.: (Purification of Marek's disease virus pp38 expressed in insect cells infected with recombinant baculovirus through affinity column chromatography). *Shengwu Gongcheng Xuebao*, 10 (1994) 239-243; C.A., 122 (1995) 75705e.
- 3849 Quilliam, M.A.: Seafood toxins. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 144-148 - a review with 56 refs.
- 3850 Quilliam, M.A., Xie, M. and Hardstaff, W.R.: Rapid extraction and cleanup for liquid chromatographic determination of domoic acid in unsalted seafood. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 543-554.
- 3851 Shibusawa, Y., Suzuki, K., Kinoshita, H. and Matsumoto, U.: Selective separation of human peripheral platelets, granulocytes and lymphocytes by surface affinity chromatography. *J. Chromatogr. B.* 666 (1995) 233-239.
- For additional information see C.A.:  
122 (1995) 50734a.
38. **INORGANIC COMPOUNDS**
- 38a. **Cations**
- 3852 Alberti, J., Rubio, R. and Rauret, G.: Arsenic speciation in marine biological materials by LC-UV-HG-ICP/OES. *Fresenius J. Anal. Chem.*, 351 (1995) 415-419.
- 3853 Arakawa, K.: Analysis of cations and anions by ion-exchange chromatography. *Jpn. Kokai Tokkyo Koho JP 06,273,401* [94,273,401] (Cl. G01N30/02), 30 Sep. 1994, Appl. 93/57,372, 17 Mar. 1993; 6 p.; C.A., 122 (1995) 122162n.
- 3854 Cobo-Fernández, M.G., Palacios, M.A., Chakraborti, D., Quevauviller, P. and Cámara, C.: On line speciation of Se(VI), Se(IV) and trimethylselenium by HPLC - microwave oven - hydride generation - atomic absorption spectrometry. *Fresenius J. Anal. Chem.*, 351 (1995) 438-442.
- 3855 Ding, H., Wang, J., Dorsey, J.G. and Caruso, J.A.: Arsenic speciation by micellar liquid chromatography with inductively coupled plasma mass spectrometric detection. *J. Chromatogr. A*, 694 (1995) 425-431.
- 3856 Fedotov, P.S., Maryutina, T.A., Pukhovskaya, V.M. and Spivakov, B.Y.: Influence of kinetic properties of extraction systems on the separation and preconcentration of some elements by countercurrent chromatography. *J. Liq. Chromatogr.*, 17 (1994) 3491-3506.
- 3857 Fodor, P. and Fischer, L.: Problems of the speciation of chromium soil samples. *Fresenius J. Anal. Chem.*, 351 (1995) 454-455.

- 3858 High, K.A., Blais, J.-S., Methven, B.A.J. and McLaren, J.W.: Probing the characteristics of metal-binding proteins using high-performance liquid chromatography - atomic absorption spectroscopy and inductively coupled plasma mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 629-634.
- 3859 High, K.A., Methven, B.A., McLaren, J.W., Siu, K.W.M., Wang, J., Klaverkamp, J.F. and Blais, J.S.: Physico-chemical of metal binding proteins using HPLC-ICP-MS, HPLC-MA-AAS and electrospray-MS. *Fresenius J. Anal. Chem.*, 351 (1995) 393-402.
- 3860 Hu, W., Tao, H., Tominaga, M., Miyazaki, A. and Haraguchi, H.: A new approach for the simultaneous determination of inorganic cations and anions using ion chromatography. *Anal. Chim. Acta*, 299 (1994) 249-256.
- 3861 Jeter, H.W. and Grob, B.: Determination of radiostrotrium in milk using an extraction chromatography column. *Radioact. Radiochem.*, 5 (1994) 8-17; *C.A.*, 122 (1995) 79342g.
- 3862 Khuhawar, M.Y. and Lanjwani, S.N.: High-performance liquid chromatographic separation and determination of cobalt(II), cobalt(III) and iron(II) using bis(salicylaldehyde)tetramethylethylenediimine. *J. Chromatogr. A*, 695 (1995) 132-135.
- 3863 Lintschinger, J., Kalcher, K., Gössler, W., Kölbl, G. and Novic, M.: Simultaneous determination of chromium(III) and chromium(VI) by reversed-phase ion-pair HPLC with chromium-specific detection. *Fresenius J. Anal. Chem.*, 351 (1995) 604-609.
- 3864 Martin, I., López-González, M.A., Gómez, M., Cámara, C. and Palacios, M.A.: Evaluation of high-performance liquid chromatography for the separation and determination of arsenic species by on-line high-performance liquid chromatographic-hydride generation-atomic absorption spectrometry. *J. Chromatogr. B*, 666 (1995) 101-109.
- 3865 Mathieu, C. and Mathieu, H.: Aluminium in serum determined by HPLC. *Clin. Chem. (Washington)*, 41 (1995) 625.
- 3866 Mimura, H., Kobayashi, T. and Akiba, K.: Chromatographic separation of strontium and cesium with mixed zeolite column. *J. Nucl. Sci. Technol.*, 32 (1995) 60-67; *C.A.*, 122 (1995) 145124r.
- 3867 Mohite, B.S., Patil, J.M. and Zambare, D.N.: Column chromatographic separation of lead from alloy and environmental samples using poly(dibenzo-18-crown-6). *J. Indian Chem. Soc.*, 71 (1994) 289-291; *C.A.*, 122 (1995) 121985w.
- 3868 Myasoedov, B.F., Chmutova, M.K., Smirnov, I.V. and Shadrin, A.Yu.: Isolation of actinides and separation from rare earth elements and selected fission products by bidentate neutral organophosphorus compounds. *Proc. Int. Conf. Technol. Expo. Future Nucl. Syst.: Emerging Fuel Cycles Waste Disposal Options*, 1 (1993) 581-587; *C.A.*, 120 (1994) 333498y.
- 3869 Nakamura, S., Watanabe, M. and Akiba, K.: Separation of yttrium from lanthanoid elements by centrifugal partition chromatography. *J. Liq. Chromatogr.*, 18 (1995) 377-394.
- 3870 Neely, F.L.: Determination of Fe(III) in fermentation broth by ion-interaction chromatography. *J. Liq. Chromatogr.*, 17 (1994) 4381-4390.
- 3871 Owen, L.M.W., Crews, H.M., Massey, R.C. and Bishop, N.J.: Determination of copper, zinc and aluminium from dietary sources in femur, brain and kidney and guinea pigs and study of some elements in *in vivo* intestinal digests by size-exclusion chromatography inductively coupled plasma mass spectrometry. *Analyst (Cambridge)*, 120 (1995) 705-712.
- 3872 Peterson, J.A. and Risley, D.S.: Validation of an HPLC method for the determination of sodium in LY293111 sodium, a novel LTB<sub>4</sub> receptor antagonist, using evaporative light scattering detection. *J. Liq. Chromatogr.*, 18 (1995) 331-338.
- 3873 Prasad, B.B., Kumar, A. and Sundd, S.: Preparation, characterization and performance of silica gel-immobilized poly(N-chloranil-N,N,N',N'-tetramethylethylenediammonium disulphosalicylate) for chromatographic separation of heavy metals. *React. Polym.*, 23 (1994) 229-238; *C.A.*, 122 (1995) 177128m.
- 3874 Renko, M., Osojnik, A. and Hudnik, V.: ICP emission spectrometric analysis of rare earth elements in permanent magnet alloys. *Fresenius J. Anal. Chem.*, 351 (1995) 610-613.
- 3875 Schilling, T., Schramel, P., Michalke, B. and Knapp, G.: Separation of noble metal ions on silica with chemically bonded ligands. *Microchim. Acta*, 116 (1994) 83-90; *C.A.*, 122 (1995) 150303c.
- 3876 Subramanian, S. and Woititz, J.R.W.: Determination of cobalt in biological samples by radiochemical neutron activation analysis employing reversed-phase chromatography. *Biol. Trace Elem. Res.*, 43-45 (1994) 117-124; *C.A.*, 122 (1995) 50697r.
- 3877 Szczepaniak, W. and Szymanski, A.: Chromatographic separation of platinum metals on silica modified with macrocyclic compounds. *Acta Chromatogr.*, 2 (1993) 7-20; *C.A.*, 122 (1995) 150301a.
- 3878 Vasconcelos, M.T. and Gornes, C.A.R.: Limitations of ion chromatography with post-column reaction for determination of heavy metals in waters containing strong chelating agents. *J. Chromatogr. A*, 696 (1995) 227-234.
- 3879 Voloshchik, I.N.: Method of determination of ammonium and alkali metal cations. *U.S.S.R. SU 1,810,822 (Cl. G01N30/06)*, 23 Apr. 1993, Appl. 4,757,658, 10 Nov. 1989; *C.A.*, 122 (1995) 122186y.
- 3880 Wang, H., Zhang, H.S. and Cheng, J.K.: Determination of V(V), Nb(V) and Ta(V) as their 2-(5-nitro-2-pyridylazo)-5-diethylaminophenol complexes by reversed-phase high-performance liquid chromatography. *Fresenius J. Anal. Chem.*, 351 (1995) 683-685.
- 3881 Wang, J. and Marshall, W.D.: Supercritical fluid extraction with carbon dioxide for the characterisation of cadmium, copper and zinc bound to metallothioneins. *Analyst (Cambridge)*, 120 (1995) 623-628.
- 3882 Watanabe, K.: (Determination of trace level chromium(VI) and cadmium ions by ion chromatography). *Kogyo Yosui*, 433 (1994) 55-62; *C.A.*, 122 (1995) 88657r - a review with 15 refs.
- 3883 Wróbel, K., González, E.B., Wróbel, K. and Sanz-Medel, A.: Aluminium and silicon speciation in human serum by ion-exchange high-performance liquid chromatography - electrothermal atomic absorption spectrometry and gel electrophoresis. *Analyst (Cambridge)*, 120 (1995) 809-815.
- 3884 Yang, X.-J.: Extractive chromatographic separation and inductively coupled plasma atomic emission spectrometric determination of trace impurities in high purity europium oxide. *Talanta*, 41 (1994) 1807-1813; *C.A.*, 122 (1995) 150412n.
- 3885 Zernichow, L. and Lund, W.: Size exclusion chromatography of aluminium species in natural waters. *Anal. Chim. Acta*, 300 (1995) 167-171.

- 3886 Zhang, X.S., Lin, X.S., Quiao, G.J. and Liu, L.Z.: Determination of palladium and gold as 4,4'-bis(dimethylamino)thiobenzophenone complex by ion-pair reversed phase liquid chromatography. *Chromatographia*, 39 (1994) 693-696.
- 3887 Zuo, X.J., Chen, L.R. and Wang, Q.: System peaks in single column ion chromatography (SCIC) with indirect spectrophotometric detection. *Chromatographia*, 39 (1994) 713-718.

For additional information see C.A.:  
122 (1995) 95400e, 122169v.

See also 2627, 2685, 2723, 3906.

### 38b. Anions

- 3888 Albazi, S.J., Gharib, T. and Cortes, J.C.: Chromate as a mobile phase for the determination of anions using cetyltrimethylammonium-coated columns and indirect photometric detection. *J. Liq. Chromatogr.*, 18 (1995) 489-503.
- 3889 Anonymous: Anions in beer by ion chromatography. *J. Am. Soc. Brew. Chem.*, 52 (1994) 178-180; C.A., 122 (1995) 54378k.
- 3890 DeVargas, M.C., DeMedina, H.L., Villalobos, E., Gutierrez, E. and Mazurek, R.: Determination of total nitrogen in sediments by using high-pressure bombs and ion chromatography. *Analyst (Cambridge)*, 120 (1995) 761-763.
- 3891 Dolenc, J. and Gorenc, D.: Ion chromatographic determination of inorganic anions in vinegar samples. *Nahrung*, 38 (1994) 434-438; C.A., 122 (1995) 54310g.
- 3892 Edgell, K.W., Longbottom, J.E. and Pfaff, J.D.: Determination of inorganic anions in water by ion-chromatography: collaborative study. *J. Assoc. Off. Anal. Chem.*, 77 (1994) 1253-1263.
- 3893 Fung, Y.S. and Dao, K.L.: Ion chromatographic determination of traces of some oxoanions with direct spectrophotometric detection. *Anal. Chim. Acta*, 300 (1995) 207-214.
- 3894 Karpuyuk, A.D., Vakhrusheva, M.V. and Sazhina, N.K.: (Determination of chloride ions by ion chromatography in the coolant of the primary circuit of a nuclear power plant). *Zavod. Lab.*, 60 (1994) 8-9, 62; C.A., 122 (1995) 177260y.
- 3895 Li, W. and Rossi, T.M.: Determination of sulfamate and sulfate as degradation products in an antiepileptic drug using ion chromatography and indirect UV detection. *J. Liq. Chromatogr.*, 18 (1995) 917-923.
- 3896 Liu, A., Xu, L., Li, T., Dong, S. and Wang, E.: Electrocatalytic oxidation and ion chromatographic detection of Br<sup>-</sup>, I<sup>-</sup>, SO<sub>3</sub><sup>2-</sup>, S<sub>2</sub>O<sub>3</sub><sup>2-</sup> and SCN<sup>-</sup> at a platinum particle-based glassy carbon modified electrode. *J. Chromatogr. A*, 699 (1995) 39-47.
- 3897 Mehra, M.C. and Landry, E.: Novel UV absorbing eluents for speciation and ion chromatographic determinations of anions. *Fresenius J. Anal. Chem.*, 351 (1995) 584-586.
- 3898 Murayama, T., Oohara, H. and Aoyama, Y.: Determination of free cyanide in water. *Jpn. Kokai Tokkyo Koho JP 06,300,742 [94,300,742] (Cl. G01N30/14)*, 28 Oct. 1994, JP Appl. 93/26,403, 16 Feb. 1993; 7 p.; C.A., 122 (1995) 114538z.
- 3899 Nagashima, K., Fukushima, K. and Kamaya, M.: Determination of trace amounts of sulfide in human serum by high-performance liquid chromatography with fluorometric detection after derivatization with 2-amino-5-N,N-diethylaminotoluene and iron(III). *J. Liq. Chromatogr.*, 18 (1995) 515-526.

- 3900 Poovey, H.G. and Rando, R.J.: Determination of chlorine and chlorine dioxide by non-suppressed ion chromatography and application to exposure assessment in the paper industry. *J. Liq. Chromatogr.*, 18 (1995) 261-275.
- 3901 Ruiz, E., Santillana, M.I., Nieto, M.T. and Sastre, I.: Determination of water-soluble inorganic phosphates in fresh vegetables by ion chromatography. *J. Liq. Chromatogr.*, 18 (1995) 989-1000.
- 3902 Sakai, H., Fujiwara, T. and Kumamaru, T.: Sensitivity enhancement in the determination of silicate by ion-exclusion chromatography using luminol chemiluminescence detection. *Anal. Chim. Acta*, 302 (1995) 173-177.

For additional information see C.A.:

122 (1995) 95458e, 95459f, 122163p, 122164q, 177336c.

See also 2566, 2685, 2899, 3836, 3853, 3860, 3879, 3887.

### 38d. Volatile inorganic compounds

- 3903 Charbonneau, S., Gilbert, R. and Lépine, L.: Determination of carbonic acid in steam-condensate cycle samples using nonsuppressed ion chromatography. *Anal. Chem.*, 67 (1995) 1204-1209.
- 3904 Huszthy, P., Bradshaw, J.S., Bordunov, A.V. and Izatt, R.M.: Enantiomeric separation of chiral [α-(1-naphthyl)ethyl]ammonium perchlorate by silica gel-bound chiral-pyridino-18-crown-6 ligands. *ACH - Models Chem.*, 131 (1994) 445-454; C.A., 122 (1995) 159871u.

For additional information see C.A.:

122 (1995) 122165r.

See also 2641.

## 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 3905 Dadachova, E., Mirzadeh, S., Lambrecht, R.M., Hetherington, E.L. and Knapp, F.F., Jr.: Separation of carrier-free <sup>166</sup>Ho from Dy<sub>2</sub>O<sub>3</sub> targets by partition chromatography and electrophoresis. *J. Radioanal. Nucl. Chem.*, 199 (1995) 155-123; C.A., 122 (1995) 144930p.
- 3906 Mathus, J.N., Murali, M.S., Iyer, R.H., Ramanujam, A., Dhani, P.S., Gopalakrishnan, V., Badheka, L.P. and Banerji, A.: Extraction chromatographic separation of minor actinides from PUREX high-level wastes using CMPO. *Nucl. Technol.*, 109 (1995) 216-225; C.A., 122 (1995) 117198z.

For additional information see C.A.:

122 (1995) 117178t.

See also 2828, 3861, 3866.