

Electrophoresis

1. REVIEWS AND BOOKS

- 2671 Cai, J. and Henion, J.: Capillary electrophoresis-mass spectrometry. *J. Chromatogr. A*, 703 (1995) 667-692 - a review with 145 refs.
- 2672 Terabe, S., Chen, N. and Otsuka, K.: Micellar electrokinetic chromatography. *Adv. Electrophor.*, 7 (1994) 87-153; C.A., 122 (1995) 229671e - a review with 228 refs.

See also 2696, 2698, 2713, 2720, 2722, 2723, 2726, 2730, 2732, 2736, 2741, 2754, 2790, 2793, 2794, 2797, 2813, 2830, 2857, 2885, 2887, 2894, 2898, 2936, 3038, 3156, 3187, 3198, 3237, 3239, 3241, 3272, 3280, 3309.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 2673 Finsy, R., Xu, R. and Deriemacker, L.: Effect of laser beam dimensions on electrophoretic mobility measurements. *Part. Part Syst. Charact.*, 11 (1994) 375-378; C.A., 123 (1995) 10514t.
- 2674 Gombinsky, M. and Lev, Adi Se-lavan: Electrophoresis means and method for separating biological macromolecules. *U.S. US 5,395,498* (Cl. 204-182.8; C25B7/00), 07 Mar. 1995, IL Appl. 99,974, 06 Nov. 1991; 12 pp.; C.A., 122 (1995) 286048f.
- 2675 Hengen, P.N.: Agarose gel electrophoresis in your kitchen. *Trends Biochem. Sci.*, 20 (1995) 202-203; C.A., 122 (1995) 309965u.
- 2676 Heukeshoven, J. and Dernick, R.: Effective blotting of ultrathin polyacrylamide gels anchored to a solid matrix. *Electrophoresis (Weinheim)*, 16 (1995) 748-756.
- 2677 Imawaka, F.: Electrophoresis method. *Jpn. Kokai Tokkyo Koho JP 07,110,297* [95,110,297] (Cl. G01N21/640), 25 Apr. 1995, Appl. 92/200,706, 03 Jul. 1992, 4 p.; C.A., 123 (1995) 47084d.
- 2678 Shiue, M.-P. and Pearlstein, A.J.: Free-solution electrophoresis with amplification of small mobility differences by helical flow. *J. Chromatogr. A*, 707 (1995) 87-103.
- 2679 Voelkel, A.R. and Noolandi, J.: On the mobility of stiff polyelectrolytes. *J. Chem. Phys.*, 102 (1995) 5506-5511; C.A., 122 (1995) 310102s.
- 2680 Zhou, H., Zhang, Y. and Lu, P.: Effect of molecular structure on the solute-micelle and solute-stationary phase binding constants in micellar liquid chromatography. *Anal. Chim. Acta*, 310 (1995) 461-471.

See also 2706, 2763.

2b. Thermodynamics and theoretical relationships

- 2681 Egorova, E.M.: The Dukhin-Deryaguin equation for the electrophoretic mobility in monovalent electrolytes with arbitrary ion mobilities. *Electrophoresis (Weinheim)*, 16 (1995) 905-910.
- 2682 Gas, B., Stedry, M., Rizzi, A. and Kenndler, E.: Dynamics of peak dispersion in capillary zone electrophoresis including wall adsorption. I. Theoretical model and results of simulation. *Electrophoresis (Weinheim)*, 16 (1995) 958-967.
- 2683 Russell, A.S., Scales, P.J., Mangelsdorf, C.S. and Underwood, S.M.: An electrophoretic investigation of the relaxation term in electrokinetic theory. *Langmuir*, 11 (1995) 1112-1115; C.A., 123 (1995) 18716f.
- 2684 Zharkikh, N.I., Lyklema, J. and Dukhin, S.S.: (Electrophoresis and electrical conductivity of suspensions under the conditions of quasi-equilibrium polarization of the double layer as a whole). *Kolloidn. Zh.*, 56 (1994) 761-766; C.A., 122 (1995) 249144p.

See also 2679, 2717, 2720, 2721, 2728, 2739, 2749, 2758, 2760, 2847, 3203, 3209, 3234.

2d. Measurement of physico-chemical and related values

See 2683, 2735, 2869, 2947, 3260.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 2685 Chang, H.-T. and Chrambach, A.: Horizontal gel electrophoresis with sample volumes up to 1.5 ml, using a discontinuous buffer system and automated apparatus. *Electrophoresis (Weinheim)*, 16 (1995) 952-957.
- 2686 Gjerde, D.T. and Yengoyan, L.: Beaded capillaries for capillary electrophoresis and chromatographic separations. *PCT Int. Appl. WO 95 10,344* (Cl. B01D15/08), 20 Apr. 1995, US Appl. 135,971, 14 Oct. 1993, 35 p.; C.A., 123 (1995) 12529a.
- 2687 Jacobson, S.C., Moore, A.W. and Ramsey, J.M.: Fused quartz substrates for microchip electrophoresis. *Anal. Chem.*, 67 (1995) 2059-2063.
- 2688 Nishikawa, T., Takahashi, S. and Kanbara, H.: Capillary electrophoresis apparatus. *Jpn. Kokai Tokkyo Koho JP 07 20,090* [95 20,090] (Cl. G01N27/447), 24 Jan. 1995, Appl. 93/164,414, 02 Jul. 1993; 8 p.; C.A., 122 (1995) 255089r.
- 2689 Sarrine, R.J.: Platform for conducting electrophoresis, and electrophoresis plate for use therewith. *U.S. US 5,399,255* (Cl. 204-299R; G01N27/26), 21 Mar. 1995, US Appl. 79,229, 21 Jun. 1993; 10 pp.; C.A., 122 (1995) 286049g.
- 2690 Tanaka, M. and Nega, N.: Fabrication of electrophoretic display panel. *Jpn. Kokai Tokkyo Koho JP 07 72,807* [95 72,807] (Cl. G09F9/30), 17 Mar. 1995, Appl. 93/221,289, 06 Sep. 1993, 6 p.; C.A., 123 (1995) 44507h.

- 2691 Watson, A.R. and Smith, A.R.: Apparatus for simultaneously supplying a plurality of samples to an electrophoretic gel. *PCT Int. Appl.* WO 95 11,450 (Cl. G01N27/447), 27 Apr. 1995, GB Appl. 93/21,650, 20 Oct. 1993; 18 pp.; C.A., 123 (1995) 29022c.
- 2692 Yasuda, K., Tsuruta, H., Takada, Y. and Oonuma, S.: Device comprising chromatographic and electrophoretic separation for analysis and purification. *Jpn. Kokai Tokkyo Koho* JP 07,103,958 [95,103,958] (Cl. G01N30/84), 21 Apr. 1995, Appl. 93/249,334, 05 Oct. 1993; 10 pp.; C.A., 123 (1995) 51707b.
- See also 2694, 2724, 2740, 2753, 2754, 2758, 2779, 3045, 3299.
- 3b. Detectors and detection procedures**
- 2693 Banks, P.R. and Paquette, D.M.: Comparison of three common amine reactive fluorescent probes used for conjugation to biomolecules by capillary zone electrophoresis. *Bioconjugate Chem.*, 6 (1995) 447-458; C.A., 123 (1995) 51522n.
- 2694 Griess, G.A., Louie, D. and Serwer, P.: A desktop, low-cost video fluorometer for quantitation of macromolecules after gel electrophoresis. *Appl. Theor. Electrophor.*, 4 (1995) 175-177; C.A., 122 (1995) 309963s.
- 2695 Hanning, A.: A new method of detecting substances in a liquid. *PCT Int. Appl.* WO 95 09,355 (Cl. G01N21/41), 06 Apr. 1995, SE Appl. 93/3,149, 28 Sep. 1993; 25 p.; C.A., 122 (1995) 329580t.
- 2696 Kaneda, T. and Imasaka, T.: (A detection of semiconductor laser excited fluorescence. Capillary electrophoresis). *Bunko Kenkyu*, 44 (1995) 21-22; C.A., 122 (1995) 301475p - a review with 9 refs.
- 2697 Ruberto, M.A.: Acridinium chemiluminescence detection with capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9428016, 1994, 81 p.; C.A., 122 (1995) 255059f.
- 2698 Schwartz, H.E., Ulfelder, K.J., Chen, F.-T.A. and Pentoney, S.L., Jr.: The utility of laser-induced fluorescence detection in applications of capillary electrophoresis. *J. Capillary Electrophor.*, 1 (1994) 36-54; C.A., 122 (1995) 285706g - a review with 122 refs.
- 2699 Seto, E.K., Damaskinos, S., Dixon, A.E., Diehl-Jones, W.L. and Mandato, C.A.: Imaging electrophoretic gels with a scanning beam laser microscope. *Electrophoresis (Weinheim)*, 16 (1995) 934-940.
- 2700 Swain, M. and Ross, N.W.: A silver stain protocol for proteins yielding high resolution and transparent background in sodium dodecyl sulfate-polyacrylamide gels. *Electrophoresis (Weinheim)*, 16 (1995) 948-951.
- 2701 Xue, Y.: Novel absorption detection techniques for capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9503611, 1994, 164 p.; C.A., 122 (1995) 305504v.
- 2702 Zhou, W., Wu, M. and Wang, E.: (Amperometric detection for high-performance capillary electrophoresis). *Fenxi Huaxue*, 23 (1995) 343-348; C.A., 122 (1995) 254779d.
- See also 2677, 2708, 2712, 2758, 2800, 2803, 2816, 2840, 2847, 2848, 2850, 2889, 3045, 3140, 3190, 3307.
- 3c. Stabilization media for electrophoresis**
- 2703 Chen, Y., Höltje, J.-V. and Schwarz, U.: Improving the UV detection sensitivity of condensed polyacrylamide gel-filled capillaries using non-flow buffer-filled capillaries as a detection cell. *J. Chromatogr. A*, 700 (1995) 35-42.
- 2704 Gu, J., Liu, Yu. and Fu, R.: (Application of gelatin/ β -cyclodextrin polymer as a chiral selector in capillary zone electrophoresis). *Fenxi Huaxue*, 22 (1994) 1098-1101; C.A., 122 (1995) 281036r.
- 2705 Liu, X.-H. and Yan, T.-T.: (Studies on grafting ethyl acrylate onto gelatin by sodium dodecyl sulphate-polyacrylamide gel electrophoresis). *Gaodeng Xuexiao Huaxue Xuebao*, 16 (1995) 156-158; C.A., 122 (1995) 241129s.
- 2706 Starr, C.M.: Utilization of additives and defined storage systems to increase the stability and performance of electrophoresis media. *PCT Int. Appl.* WO 95 07,458 (Cl. G01N27/447), 16 Mar. 1995, US Appl. 118,957, 09 Sep. 1993; 43 pp.; C.A., 122 (1995) 260550p.
- 2707 Vanek, P.G., Fabian, S.J., Fisher, C.L., Chirikjian, J.G. and Collier, G.B.: Alternative to polyacrylamide gels improves the electrophoretic mobility shift assay. *BioTechniques*, 18 (1995) 704-706; C.A., 122 (1995) 260341w.
- See also 2739, 2753, 2757, 2847, 3166, 3182.
- 3d. Quantitative analysis**
- See 2721, 2760, 3288.
- 3e. Preparative scale electrophoresis**
- See 2820, 2886, 2889, 2931.
- 4. SPECIAL TECHNIQUES**
- 4a. Automation**
- 2708 Rappaport, B.N.: Automated reading of electrophoresis gels and southern blots. Avail. *Univ. Microfilms Int.*, Order No. DA9510416, 1994, 114 pp.; C.A., 123 (1995) 26491u.
- See also 2685.
- 4b. Computerization and modelling**
- 2709 Allison, S.A. and Tran, V.T.: Modeling the electrophoresis of rigid polyions: application to lysozyme. *Biophys. J.*, 68 (1995) 2261-2270; C.A., 123 (1995) 51530p.
- 2710 Jensen, K., Søndergaard, I., Skovgaard, I.M. and Nielsen, H.B.: From image processing to classification: I. Modeling disturbances of isoelectric focusing patterns. *Electrophoresis (Weinheim)*, 16 (1995) 921-926.
- 2711 Kesmir, C., Søndergaard, I. and Jensen, K.: From image processing to classification: II. Classification of electrophoretic patterns using self-organizing feature maps and feed-forward neural networks. *Electrophoresis (Weinheim)*, 16 (1995) 927-933.

- 2712 Van de Lest, C.H.A., Veerkamp, J.H. and Kuppevelt, T.H.: ImageCalc: a Microsoft Windows application for quantitative image analysis and comparison. *BioTechniques*, 18 (1995) 1050-1055; C.A., 123 (1995) 29004y.
- See also 2678, 2695, 2884, 2890, 2894, 2902, 2935, 2964, 2980, 3000, 3184.
- 4c. *Combination with other physicochemical techniques, (MS, IR etc.)*
- 2713 Albert, K.: On-line use of NMR detection in separation chemistry. *J. Chromatogr. A*, 703 (1995) 123-147 - a review with 27 refs.
- 2714 Liu, Y., Lopez-Avila, V., Zhu, J.J., Wiederin, D.R. and Beckert, W.F.: Capillary electrophoresis coupled on-line with inductively coupled plasma mass spectrometry for elemental speciation. *Anal. Chem.*, 67 (1995) 2020-2025.
- 2715 Lord, G.A., Gordon, D.B., Tetler, L.W. and Carr, C.M.: Electrochromatography-electrospray mass spectrometry of textile dyes. *J. Chromatogr. A*, 700 (1995) 27-33.
- 2716 Tomlinson, A.J. and Naylor, S.: Enhanced performance membrane preconcentration-capillary electrophoresis-mass spectrometry (mPC-CE-MS) in conjunction with transient isotachopheresis for analysis of peptide mixtures. *J. High Resolut. Chromatogr.*, 18 (1995) 384-386.
- See also 2671, 2748, 2755, 2759, 2785, 2786, 2841, 2855, 2893, 2895, 3236, 3242, 3244.
- 4d. *Affinity electrophoresis*
- See 2812, 2813, 2855, 3090.
- 4e. *Capillary zone electrophoresis and electrokinetic chromatography*
- 2717 Altria, K.D. and Fabre, H.: Approaches to optimisation of precision in capillary electrophoresis. *Chromatographia*, 40 (1995) 313-320.
- 2718 Armstrong, D.W., Rundlett, K.L. and Chen, J.-R.: Evaluation of the macrocyclic antibiotic vancomycin as a chiral selector for capillary electrophoresis. *Chirality*, 6 (1994) 496-509; C.A., 123 (1995) 24944v.
- 2719 Bächmann, K., Bazzanella, A., Haag, I., Han, K.-Y., Arnecke, R., Böhmer, V. and Vogt, W.: Resorcarenes as pseudostationary phases with selectivity for electrokinetic chromatography. *Anal. Chem.*, 67 (1995) 1722-1726.
- 2720 Baechmann, K., Boden, J., Ehmann, T., Goettlicher, B., Groh, T., Han, K.Y., Hauman, I., Hensel, W., Mainka, A. et al.: (Limits of and opportunities with capillary electrophoresis). *CLB Chem. Labor. Biotech.*, 46 (1995) 77-79; C.A., 122 (1995) 305490n - a continued review with refs.
- 2721 Baumann, K. and Wätzig, H.: Appropriate calibration functions for capillary electrophoresis. II. Heteroscedasticity and its consequences. *J. Chromatogr. A*, 700 (1995) 9-20.
- 2722 Beachmann, K., Boden, J., Ehmann, T., Goettlicher, B., Groh, T., Han, K.Y., Haumann, I., Hensel, W., Mainka, A. et al.: (Capillary electrophoresis. Limits and possibilities). *CLB, Chem. Labor Biotech.*, 46 (1995) 121-123; C.A., 122 (1995) 329219g - a review with refs.
- 2723 Belen'kii, B.G.: (Capillary electric separation methods and their use in analytical chemistry). *Ross Khim. Zh.*, 38 (1994) 25-32; C.A., 123 (1995) 24661a - a review with 29 refs.
- 2724 Boughtflower, R.J., Underwood, T. and Paterson, C.J.: Capillary electrochromatography - some important consideration in the preparation of packed capillaries and the choice of mobile phase buffers. *Chromatographia*, 40 (1995) 329-335.
- 2725 Brondz, I., Dahle, U.R., Greibrokk, T. and Olsen, I.: Capillary zone electrophoresis as a new tool in the chemotaxonomy of oral treponemes. *J. Chromatogr. B*, 667 (1995) 161-165.
- 2726 Bruin, G.J.M. and Paulus, A.: Biopolymer separations with capillary electrophoresis. *Anal. Methods Instrum.*, 2 (1995) 3-26; C.A., 123 (1995) 28661y - a review with 175 refs.
- 2727 Chang, H.-T.: Development of novel separation techniques for biological samples in capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9503536, 1994, 273 pp.; C.A., 122 (1995) 285763y.
- 2728 Cross, R.F. and Ricci, M.C.: pH and pKa limitations in the CZE analysis of sulphonamides. *LC-GC*, 13 (1995) 132-142; C.A., 122 (1995) 305528f.
- 2729 Effenhauser, C.S., Manz, A. and Widmer, H.M.: Manipulation of sample fractions on a capillary electrophoresis chip. *Anal. Chem.*, 67 (1995) 2284-2287.
- 2730 Fanali, S., Cristalli, M., Vespalet, R. and Bocek, P.: Chiral separations in capillary electrophoresis. *Adv. Electrophor.*, 7 (1994) 1-86; C.A., 122 (1995) 254991s - a review with 156 refs.
- 2731 Fujimoto, C. and Sawada, H.: (Capillary electrophoresis of small molecules in high-viscosity polymer solutions). *Bunseki Kagaku*, 44 (1995) 75-78; C.A., 122 (1995) 329495u.
- 2732 Gilman, S.D. and Ewing, A.G.: Recent advances in the application of capillary electrophoresis to neuroscience. *J. Capillary Electrophor.*, 2 (1995) 1-13; C.A., 122 (1995) 309901v - a review with 74 refs.
- 2733 Gu, J., Fu, R., Huang, Z. and Xu, W.: (Study of water-soluble crown ether used as chiral selector in capillary zone electrophoresis (CZE)-study of chiral selector in the view of supramolecular chemistry). *Sepu*, 12 (1994) 391-394; C.A., 122 (1995) 329501t.
- 2734 Guttman, A. and Schwartz, H.E.: Artifacts related to sample introduction in capillary gel electrophoresis affecting separation performance and quantitation. *Anal. Chem.*, 67 (1995) 2279-2283.
- 2735 Holloway, R.R.: Capillary electrophoresis using zwitterion-coated capillary tubes. *Eur. Pat. Appl. EP 639,768 (Cl. G01N27/447)*, 22 Feb. 1995, US Appl. 107,393, 16 Aug. 1993; 10 p.; C.A., 122 (1995) 229796z.
- 2736 Issaq, H.J., Chan, K.C., Muschik, G.M. and Janini, G.M.: Applications of capillary zone electrophoresis and micellar electrokinetic chromatography in cancer research. *J. Liq. Chromatogr.*, 18 (1995) 1273-1288 - a review with 33 refs.
- 2737 Iwata, T. and Kuroshu, Y.: Enhancement of the sensitivity and resolution by flow-counterbalanced capillary electrophoresis. *Anal. Sci.*, 11 (1995) 131-133; C.A., 122 (1995) 281000a.
- 2738 Janini, G.M., Muschik, G.M. and Issaq, H.J.: Sample matrix effects in capillary zone electrophoresis. Effect of chloride ion on nitrate and nitrite. *J. Capillary Electrophor.*, 1 (1994) 116-120; C.A., 122 (1995) 310101r.
- 2739 Jansson, M. and Roeraade, J.: N-Methylformamide as a separation medium in capillary electrophoresis. *Chromatographia*, 40 (1995) 163-169.

- 2740 Kaljurand, M., Ebber, A. and Somer, T.: An automatic sampling device for capillary zone electrophoresis. *J. High Resolut. Chromatogr.*, 18 (1995) 263-265.
- 2741 Karger, B.L., Chu, Y.-H. and Foret, F.: Capillary electrophoresis of proteins and nucleic acids. *Annu. Rev. Biophys. Biomol. Struct.*, 24 (1995) 579-610; *C.A.*, 123 (1995) 28670a - a review with 99 refs.
- 2742 Katsuta, S., Tsumura, T., Saitoh, K. and Teramae, N.: Control of selectivity in micellar electrokinetic chromatography by modification of sodium dodecyl sulfate micelles with organic hydroxy compounds. *J. Chromatogr. A*, 705 (1995) 319-324.
- 2743 Kenndler, E. and Poppe, H.: Entangled, noncross-linked polymer solutions used as physical network in capillary zone electrophoresis. *J. Capillary Electrophor.*, 1 (1994) 144-157; *C.A.*, 122 (1995) 309903x.
- 2744 Lue, J. and Fu, X.: (Effect of sample solution matrix on separation in capillary electrophoresis). *Fenxi Huaxue*, 22 (1994) 1231-1233; *C.A.*, 123 (1995) 24954y.
- 2745 Macka, M., Haddad, P.R. and Buchberger, W.: Separation of some metallochromic ligands by capillary zone electrophoresis and micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 706 (1995) 493-501.
- 2746 Nilsson, S., Johansson, J., Mecklenburg, M., Birnbaum, S., Svanberg, S., Wahlund, K.-G., Mosbach, K., Miyabayashi, A. and Larsson, P.-O.: Real-time fluorescence imaging of capillary electrophoresis. *J. Capillary Electrophor.*, 2 (1995) 46-52; *C.A.*, 122 (1995) 310097u.
- 2747 Otsuka, K., Kigashimori, M., Okada, Y. and Terabe, S.: (Effect of organic modifiers in micellar electrokinetic chromatography). *Kuromatogurafi*, 15 (1995) 120-121; *C.A.*, 122 (1995) 281014g.
- 2748 Ozaki, H. and Terabe, S.: (Micellar electrokinetic chromatography using high-molecular surfactants: Applicability of coupling MEKC with mass spectrometry). *Kuromatogurafi*, 15 (1994) 116-118; *C.A.*, 122 (1995) 281013f.
- 2749 Pyell, U. and Bütchorn, U.: Optimization strategies in micellar electrokinetic capillary chromatography. One-parameter optimizations of the concentration of sodium dodecyl sulfate and the concentration of modifiers (urea and glucose). *Chromatographia*, 40 (1995) 175-184.
- 2750 Saari-Nordhaus, R. and Anderson, J.M., Jr.: Membrane-based solid-phase extraction as a sample clean-up technique for anion analysis by capillary electrophoresis. *J. Chromatogr. A*, 706 (1995) 563-569.
- 2751 Sepaniak, M.J., Copper, C.L., Whitaker, K.W. and Anigbogu, V.C.: Evaluation of a dual-cyclodextrin phase variant of capillary electrokinetic chromatography for separations of nonionizable solutes. *Anal. Chem.*, 67 (1995) 2037-2041.
- 2752 Shihabi, Z.K. and Hinsdale, M.E.: Sample matrix effects in micellar electrokinetic capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 75-83.
- 2753 Smith, J.T.: Development of micellar phases and capillary columns for high-performance capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9508648, 1994, 286 p.; *C.A.*, 123 (1995) 18699c.
- 2754 Stevenson, R.: A critical review of the development of HPCE instrumentation. *J. Capillary Electrophor.*, 1 (1994) 169-174; *C.A.*, 122 (1995) 317499k - a review with 24 refs.
- 2755 Tetler, L.W., Cooper, P.A. and Powell, B.: Influence of capillary dimensions on the performance of a coaxial capillary electrophoresis-electrospray mass spectrometry interface. *J. Chromatogr. A*, 700 (1995) 21-26.
- 2756 Tomlinson, A.J., Benson, L.M., Braddock, W.D., Oda, R.P. and Naylor, S.: Improved on-line membrane preconcentration-capillary electrophoresis (mPC-CE). *J. High Resolut. Chromatogr.*, 18 (1995) 381-383.
- 2757 Tsuda, T., Kitagawa, S. and Nakasima, M.: (Chromatographic behavior and column supports in electrochromatography (electrochromatography XVI)). *Kuromatogurafi*, 15 (1994) 112-113; *C.A.*, 122 (1995) 280802a.
- 2758 Van den Moolen, J.N., Louwse, D.J., Poppe, H. and Smit, H.C.: Correlation capillary zone electrophoresis, a novel technique to decrease detection limits. *Chromatographia*, 40 (1995) 368-374.
- 2759 Wahl, J.H. and Smith, R.D.: Comparison of buffer systems and interface designs for capillary electrophoresis-mass spectrometry. *J. Capillary Electrophor.*, 1 (1994) 62-71; *C.A.*, 122 (1995) 310200x.
- 2760 Wätzig, H.: Appropriate calibration functions for capillary electrophoresis. I. Precision and sensitivity using peak areas and heights. *J. Chromatogr. A*, 700 (1995) 1-7.
- 2761 Xian, J.: Application of capillary electrophoresis for the separation of biomolecules (DNA). Avail. *Univ. Microfilms Int.*, Order No. DA9505102, 1994, 175 p.; *C.A.*, 122 (1995) 329296e.
- 2762 Zhang, C.-X., von Heeren, F. and Thormann, W.: Separation of hydrophobic, positively chargeable substances by capillary electrophoresis. *Anal. Chem.*, 67 (1995) 2070-2077.
- 2763 Zhu, M.-D. and Siebert, C.J.: Suppression of electroendosmosis during electrophoresis in gel-free polymer media by use of charged polymers. *PCT Int. Appl.* WO 95 10,771 (Cl. G01N27/447), 20 Apr. 1995, US Appl. 136,689, 14 Oct. 1993; 23 pp.; *C.A.*, 123 (1995) 51703x.
- See also 2671, 2672, 2680, 2682, 2686, 2687, 2688, 2693, 2696, 2697, 2698, 2701, 2702, 2703, 2704, 2714, 2716, 2766, 2767, 2768, 2769, 2770, 2771, 2775, 2777, 2780, 2781, 2782, 2783, 2784, 2785, 2786, 2787, 2788, 2789, 2790, 2791, 2792, 2793, 2795, 2796, 2797, 2798, 2799, 2800, 2801, 2802, 2803, 2813, 2825, 2826, 2828, 2829, 2838, 2840, 2841, 2842, 2843, 2844, 2847, 2848, 2849, 2850, 2851, 2852, 2855, 2857, 2858, 2859, 2861, 2863, 2864, 2867, 2868, 2878, 2879, 2880, 2882, 2886, 2887, 2888, 2895, 2896, 2897, 2929, 2937, 2947, 2952, 2961, 2962, 2966, 2977, 2991, 2996, 2997, 2998, 3007, 3010, 3011, 3027, 3039, 3040, 3065, 3070, 3081, 3090, 3093, 3098, 3099, 3100, 3102, 3103, 3106, 3112, 3125, 3128, 3129, 3131, 3135, 3145, 3160, 3173, 3174, 3177, 3180, 3182, 3183, 3190, 3195, 3198, 3203, 3211, 3212, 3213, 3215, 3216, 3217, 3218, 3219, 3220, 3221, 3222, 3223, 3226, 3227, 3228, 3230, 3231, 3232, 3233, 3235, 3236, 3237, 3238, 3239, 3240, 3241, 3242, 3243, 3244, 3245, 3246, 3247, 3248, 3249, 3250, 3251, 3253, 3254, 3255, 3256, 3257, 3258, 3259, 3261, 3262, 3263, 3264, 3265, 3266, 3267, 3268, 3269, 3270,

3271, 3272, 3273, 3274, 3275, 3278, 3279, 3280, 3281, 3282, 3284, 3286, 3287, 3288, 3289, 3290, 3291, 3292, 3295, 3296, 3297, 3298, 3299, 3300, 3301, 3302, 3303, 3304, 3305, 3306, 3307, 3308, 3309.

4f. Isotachopheresis

- 2764 Deshmukh, R.R.: Experimental and theoretical studies in isotachopheresis. Avail. *Univ. Microfilms Int.*, Order No. DA9426339, 1994, 325 p.; C.A., 123 (1995) 18685v.
- 2765 Sauve, D.M., Ho, D.T. and Roberge, M.: Concentration of dilute protein for gel electrophoresis. *Anal. Biochem.*, 226 (1995) 382-383.

See also 2716, 2723, 2839, 2993, 3252, 3260, 3293, 3294.

4g. Enantiomers, separation

- 2766 Agyei, N.M., Gahm, K.H. and Stalcup, A.M.: Chiral separations using heparin and dextran sulfate in capillary zone electrophoresis. *Anal. Chim. Acta*, 307 (1995) 185-191.
- 2767 Baumy, P., Morin, P., Dreux, M., Viaud, M.C., Boye, S. and Guillaumet, G.: Determination of β -cyclodextrin inclusion complex constants for 3,4-dihydro-2-H-1-benzopyran enantiomers by capillary electrophoresis. *J. Chromatogr. A*, 707 (1995) 311-326.
- 2768 Ingelse, B.A., Everaerts, F.M., Sevcik, J., Stransky, Z. and Fanali, S.: A further study on the chiral separation power of a soluble neutral β -cyclodextrin polymer. *J. High Resolut. Chromatogr.*, 18 (1995) 348-352.
- 2769 Krasensky, S., Fanali, S., Krivankova, L. and Bocek, P.: Highly sensitive chiral analysis in on-line combined chiral and achiral media by capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 968-973.
- 2770 Rundlett, K.L. and Armstrong, D.W.: Effect of micelles and mixed micelles on efficiency and selectivity of antibiotic-based capillary electrophoretic enantioseparations. *Anal. Chem.*, 67 (1995) 2088-2095.
- 2771 Wu, W. and Stalcup, A.M.: Capillary electrophoretic chiral separations using a sulfated β -cyclodextrin-containing electrolyte. *J. Liq. Chromatogr.*, 18 (1995) 1289-1315.

See also 2704, 2718, 2730, 2733, 2780, 2782, 2798, 2841, 2842, 2844, 2850, 2851, 2852, 3213, 3216, 3217, 3218, 3250, 3251, 3254, 3261, 3265, 3266.

4h. Two dimensional electrophoresis

- 2772 Burggraf, D. and Lottspeich, F.: The further construction of the two-dimensional database of common human proteins. *Electrophoresis (Weinheim)*, 16 (1995) 831-840.
- 2773 Castro, A. and Shera, E.B.: Single-molecule electrophoresis: applications to biomolecular detection. *Proc. SPIE-Int. Soc. Opt. Eng.*, 2386 (1995) 79-85; C.A., 123 (1995) 28852m.
- 2774 Klose, J. and Kobalz, U.: Two-dimensional electrophoresis of proteins: an updated protocol and implications for a functional analysis of the genome. *Electrophoresis (Weinheim)*, 16 (1995) 1034-1059.

- 2775 Lada, M.W., Schaller, G., Carriger, M.H., Vickroy, T.W. and Kennedy, R.T.: On-line interface between microdialysis and capillary zone electrophoresis. *Anal. Chim. Acta*, 307 (1995) 217-225.

- 2776 Schägger, H.: Quantification of oxidative phosphorylation enzymes after blue native electrophoresis and two-dimensional resolution: normal complex I protein amounts in Parkinson's disease conflict with reduced catalytic activities. *Electrophoresis (Weinheim)*, 16 (1995) 763-770.

See also 2809, 2832, 2837, 2872, 2877, 2884, 2893, 2894, 2901, 2902, 2904, 2908, 2915, 2916, 2917, 2918, 2919, 2931, 2932, 2934, 2935, 2938, 2940, 2942, 2943, 2944, 2945, 2948, 2954, 2963, 2964, 2965, 2967, 2968, 2970, 2974, 2979, 2980, 2983, 2987, 2989, 3000, 3001, 3002, 3003, 3008, 3014, 3017, 3018, 3019, 3041, 3042, 3047, 3068, 3091, 3150.

4i. Other special techniques

- 2777 Fujimoto, C.: Charged polyacrylamide gels for capillary electrochromatographic separations of uncharged low molecular weight compounds. *Anal. Chem.*, 67 (1995) 2050-2053.
- 2778 Hartig, R., Hausmann, M. and Cremer, C.: Continuous focusing of biological particles by continuous immuno magnetic sorter: technique and applications. *Electrophoresis (Weinheim)*, 16 (1995) 789-792.
- 2779 Wu, J. and Pawliszyn, J.: A capillary cartridge with an on-line desalting device that allows fast sampling for capillary isoelectric focusing. *Anal. Chem.*, 67 (1995) 2010-2014.

See also 2687, 2781, 2875.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. Cyclic hydrocarbons, fullerenes

- 2780 Desiderio, C., Fanali, S., Sinibaldi, M. and Polcaro, C.: Cyclodextrin modified micellar electrokinetic chromatography for the chiral direct resolution of (+), (-)-trans-1,2-dihydrodiol metabolite of chrysene *in vitro* activated by rat liver microsome S9 fraction. *Electrophoresis (Weinheim)*, 16 (1995) 784-788.
- 2781 Yan, C., Dadoo, R., Zhao, H., Zare, R.N. and Rakestraw, D.J.: Capillary electrochromatography: analysis of polycyclic aromatic hydrocarbons. *Anal. Chem.*, 67 (1995) 2026-2029.

See also 2719.

6. ALCOHOLS

See 2780.

7. PHENOLS

- 2782 Chankvetadze, B., Endresz, G. and Blaschke, G.: Enantiomeric resolution of anionic R/S-1,1'-binaphthyl-2,2'-diyl hydrogen phosphate by capillary electrophoresis using anionic cyclodextrin derivatives as chiral selectors. *J. Chromatogr. A*, 704 (1995) 234-237.
- 2783 Li, G. and Locke, D.C.: Separation of the eleven priority pollutant phenols by capillary zone electrophoresis. *J. Chromatogr. B*, 669 (1995) 93-102.
- 2784 Lin, C.-e., Lin, W.-c. and Chiou, W.-c.: Migration behaviour and optimization of selectivity of dichlorophenols in capillary zone electrophoresis. *J. Chromatogr. A*, 705 (1995) 325-333.

See also 2826.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

- 2785 Aramendia, M.A., Garcia, I., Lafont, F. and Marinas, J.M.: Determination of isoflavones using capillary electrophoresis in combination with electrospray mass spectrometry. *J. Chromatogr. A*, 707 (1995) 327-333.

8b. Aflatoxins and other mycotoxins

- 2786 Hines, H.B., Brueggemann, E.E., Holcomb, M. and Holder, C.L.: Fumonisin B₁ analysis with capillary electrophoresis-electrospray ionization mass spectrometry. *Rapid Commun. Mass Spectrom.*, 9 (1995) 519-524; C.A., 123 (1995) 8146f.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 2787 Deasy, B., Bogan, D.P., Smyth, M.R., O'Kennedy, R. and Fuhr, U.: Study of coumarin metabolism by human liver microsomes using capillary electrophoresis. *Anal. Chim. Acta*, 310 (1995) 101-107.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 2788 Sheu, S.-J. and Chen, H.-R.: Determination of five major anthraquinoids in Chinese herbal preparations by micellar electrokinetic capillary electrophoresis. *Anal. Chim. Acta*, 309 (1995) 361-367.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 2789 Chang, L., Yao, R., Guo, Q. and Yan, F.: (Analysis of oligosaccharides by capillary electrophoresis). *Fenxi Huaxue*, 22 (1994) 975-979; C.A., 122 (1995) 305526d.

- 2790 El Rassi, Z. and Nashabeh, W.: High performance capillary electrophoresis of carbohydrates and glycoconjugates. *J. Chromatogr. Libr.*, Vol. 58, 1995, pp. 267-360; C.A., 122 (1995) 309920a - a review with 219 refs.

- 2791 Evangelista, R.A., Liu, M.-S. and Chen, F.-T.A.: Characterization of 9-aminopyrene-1,4,6-trisulfonate-derivatized sugars by capillary electrophoresis with laser-induced fluorescence detection. *Anal. Chem.*, 67 (1995) 2239-2245.

- 2792 Guttman, A. and Starr, C.: Capillary and slab gel electrophoresis profiling of oligosaccharides. *Electrophoresis (Weinheim)*, 16 (1995) 993-997.

- 2793 Honda, S.: (Analysis of sugar chains by capillary electrophoresis). *Rinsho Kensa*, 39 (1995) 86-88; C.A., 123 (1995) 4776p - a review with 11 refs.

- 2794 Hu, G.-F.: Fluorophore-assisted carbohydrate electrophoresis. Technology and applications. *J. Chromatogr. A*, 705 (1995) 89-103 - a review with 89 refs.

- 2795 Huang, X. and Kok, W.T.: Determination of sugars by capillary electrophoresis with electrochemical detection using cuprous oxide modified electrodes. *J. Chromatogr. A*, 707 (1995) 335-342.

- 2796 Lin, Q., Zhang, R. and Liu, G.: (Trace analysis of amino sugars and neutral saccharides by high performance capillary electrophoresis and high performance liquid chromatography). *Sepu*, 13 (1995) 12-15; C.A., 123 (1995) 47025k.

- 2797 Mort, A.J. and Pierce, M.L.: Preparation of carbohydrates for analysis by HPLC and HPCE. *J. Chromatogr. Libr.*, Vol. 58, 1995, pp. 3-37; C.A., 122 (1995) 309913a - a review with 91 refs.

- 2798 Noe, C.R. and Freissmuth, J.: Capillary zone electrophoresis of aldose enantiomers: separation after derivatization with S-(-)-1-phenylethylamine. *J. Chromatogr. A*, 704 (1995) 503-512.

- 2799 Oefner, P., Chiesa, C., Bonn, G. and Horvath, C.: Developments in capillary electrophoresis of carbohydrates. *J. Capillary Electrophor.*, 1 (1994) 5-26; C.A., 122 (1995) 285704e.

- 2800 Roberts, R.E.: Development of fast pulsed electrochemical detection of carbohydrates separated by liquid chromatography and capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9503589, 1994, 146 p.; C.A., 122 (1995) 30519d.

- 2801 Suzuki, S. and Honda, S.: Two-dimensional mapping of N-glycosidically linked oligosaccharides in glycoproteins by high-performance capillary electrophoresis. *TrAC*, 14 (1995) 279-288.

- 2802 Zhang, Y., Le, X., Dovichi, N.J., Compston, C.A., Palcic, M.M., Diedrich, P. and Hindsgaul, O.: Monitoring biosynthetic transformations of N-acetyllactosamine using fluorescently labeled oligosaccharides and capillary electrophoretic separation. *Anal. Biochem.*, 227 (1995) 368-376.

- 2803 Zhaolou, Y., Liwen, C., Xizhang, H. and Tiansheng, S.: (Indirect ultraviolet detection of monosaccharides by high-performance capillary electrophoresis using β -indoleacetic acid-phosphate as background electrolyte). *Fenxi Huaxue*, 22 (1994) 755-758, C.A., 122 (1995) 255055b.

See also 2808.

10b. Polysaccharides, mucopolysaccharides, lipopolysaccharides

- 2804 Fernandez-Busquets, X. and Burger, M.M.: Use of Rhodamine B isocyanate to detect proteoglycan core proteins in polyacrylamide gels. *Anal. Biochem.*, 227 (1995) 394-396.

- 2805 Lebowhl, M., Longas, M.O., Konstadt, J., Koreen, R., Grabowski, G.A., Phelps, R., Lemlich, G. and Schwartz, E.: Hyaluronic acid and dermatan sulfate in non-lesional pseudoxanthoma elasticum skin. *Clin. Chim. Acta*, 238 (1995) 101-107.
- 2806 Moller, H.J., Moller-Pedersen, T., Damsgaard, T.E. and Poulsen, J.H.: Demonstration of immunogenic keratan sulphate in commercial chondroitin 6-sulphate from shark cartilage. Implications for ELISA assays. *Clin. Chim. Acta*, 236 (1995) 195-204.
- 2807 Volpi, N., Fregni, R. and Venturelli, T.: Activity of chondroitin ABC lyase on dermatan sulfate partially degraded by cupric-ion-mediated free-radical treatment. *J. Chromatogr. B*, 669 (1995) 197-205.
- See also 2794, 3037.
- 10c. *Glycoproteins and their constituents*
- 2808 Friedman, Y. and Higgins, E.A.: A method for monitoring the glycosylation of recombinant glycoproteins from conditioned medium, using fluorophore-assisted carbohydrate electrophoresis. *Anal. Biochem.*, 228 (1995) 221-225.
- 2809 Golaz, O., Gravel, P., Walzer, C., Turler, H., Balant, L. and Hochstrasser, D.F.: Rapid detection of the main human plasma glycoproteins by two-dimensional polyacrylamide gel electrophoresis lectin affinity blotting. *Electrophoresis (Weinheim)*, 16 (1995) 1187-1189.
- 2810 Gu, J. and Wada, Y.: Aberrant expressions of decorin and biglycan genes in the carbohydrate-deficient glycoprotein syndrome. *J. Biochem. (Tokyo)*, 117 (1995) 1276-1279.
- 2811 Hagihara, Y., Goto, Y., Kato, H. and Yoshimura, T.: Role of the N- and C-terminal domains of bovine β_2 -glycoprotein I in its interaction with cardiolipin. *J. Biochem. (Tokyo)*, 118 (1995) 129-136.
- 2812 Hänsler, M., Kötz, K. and Hantzschel, H.: Detection of immunoglobulin G glycosylation changes in patients with rheumatoid arthritis by means of isoelectric focusing and lectin-affinoblotting. *Electrophoresis (Weinheim)*, 16 (1995) 811-812.
- 2813 Honda, S.: Basic studies on carbohydrate-protein interaction by high performance affinity chromatography and high performance capillary affinity electrophoresis using lectins as protein models. *J. Chromatogr. Libr.*, Vol. 58, 1995, pp. 211-231; C.A., 122 (1995) 309918f - a review with 10 refs.
- 2814 Kamoshita, K., Shiota, M., Sasaki, M., Koga, Y., Okumura, Y. and Kido, H.: Calcium requirement and inhibitor spectrum for intracellular HIV type 1 gp160 processing in cultured HeLa cells and CD4 lymphocytes: similarity to those of viral envelope glycoprotein maturase. *J. Biochem. (Tokyo)*, 117 (1995) 1244-1253.
- 2815 Lee, T.K., Drohan, W.N. and Lubon, H.: Proteolytic processing of human protein C in swine mammary gland. *J. Biochem. (Tokyo)*, 118 (1995) 81-87.
- 2816 Moller, H.J. and Poulsen, J.H.: Improved method for silver staining of glycoproteins in thin sodium dodecyl sulfate polyacrylamide gels. *Anal. Biochem.*, 226 (1995) 371-374.
- 2817 Nakashima, Y.: (Analysis of urinary glycoproteins for diabetic nephropathy. Western blotting analysis using peroxidase labeled lectins). *Nara Igaku Zasshi*, 45 (1994) 672-683; C.A., 122 (1995) 262872u.
- 2818 Odo, S., Kamino, K., Kanai, S., Maruyama, T. and Harayama, S.: Biochemical characterization of a Ca^{2+} -dependent lectin from the hemolymph of a photosymbiotic marine bivalve, *Tridacna derasa* (Röding). *J. Biochem. (Tokyo)*, 117 (1995) 965-973.
- 2819 Ohannesian, D.W., Lotan, D., Thomas, P., Jessup, J.M., Fukuda, M., Gabius, H.-J. and Lotan, R.: Carcinoembryonic antigen and other glycoconjugates act as ligands for galectin-3 in human colon carcinoma cells. *Cancer Res.*, 55 (1995) 2191-2199.
- 2820 Paszkiewicz-Gadek, A., Gindzienski, A. and Porowska, H.: The use of preparative polyacrylamide gel electrophoresis and electroelution for purification of mucus glycoproteins. *Anal. Biochem.*, 226 (1995) 263-267.
- 2821 Pilatte, Y., Rostoker, G., Violleau, K., Basta, M. and Hammer, C.H.: Further identification of human plasma glycoproteins interacting with the galactose-specific lectin Jacalin. *J. Chromatogr. B*, 668 (1995) 1-11.
- 2822 Rall, C.J.N. and Rustgi, A.K.: CD44 Isoform expression in primary and metastatic pancreatic adenocarcinoma. *Cancer Res.*, 55 (1995) 1831-1835.
- 2823 Scott, P.G., Nakano, T. and Dodd, C.M.: Small proteoglycans from different regions of the fibrocartilaginous temporomandibular joint disc. *Biochim. Biophys. Acta*, 1244 (1995) 121-128.
- 2824 To, W.Y., Leung, J.C.K. and Lai, K.N.: Identification and characterization of human serum alpha2-HS glycoprotein as a jacalin-bound protein. *Biochim. Biophys. Acta*, 1249 (1995) 58-64.
- See also 2790, 2921, 2924, 2953, 3107.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 2825 Gutnikov, G., Beck, W. and Engelhardt, H.: Separation of homologous fatty acids by capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 565-570; C.A., 122 (1995) 229725a.
- 2826 Morin, P. and Dreux, M.: Separation of phenolic acids by micellar electrokinetic capillary chromatography. *Bull. Liaison - Groupe Polyphenols*, 16(Pt. 1) (1992) 201-206; C.A., 123 (1995) 4937s.
- 2827 Petersen, M., Haeusler, E., Meinhard, J., Karwatzki, B. and Gertlowski, C.: The biosynthesis of rosmarinic acid in suspension cultures of *Coleus blumei*. *Plant Cell, Tissue Organ Cult.*, 38 (1994) 171-179; C.A., 122 (1995) 286707v.
- 2828 Petucci, C.J., Kantes, H.L., Strein, T.G. and Veening, H.: Capillary electrophoresis as a clinical tool. Determination of organic anions in normal and uremic serum using photodiode-array detection. *J. Chromatogr. B*, 668 (1995) 241-251.
- 2829 Roldan-Assad, R. and Gareil, P.: Capillary zone electrophoretic determination of C_2 - C_{18} linear saturated free fatty acids with indirect absorbance detection. *J. Chromatogr. A*, 708 (1995) 339-350.
- See also 2739, 2769, 3302.

11d. Lipoproteins and their constituents

- 2830 Brouillette, C.G. and Anantharamiah, G.M.: Structural models of human apolipoprotein A-I. *Biochim. Biophys. Acta*, 1256 (1995) 103-129 - a review with 219 refs.
- 2831 Cavalcanti, S., Cavalcanti, E., Angotti, E., Liparoti, A. and Colonna, A.: Identification of apolipoproteins A-I and B using high resolution electrophoresis on supported cellulose acetate. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 275-279.
- 2832 Golaz, O., Sanchez, J.-C., James, R.W. and Hochstrasser, D.F.: Phenotyping of apolipoprotein E using immobilized pH gradient gels for one-dimensional and two-dimensional separations. *Electrophoresis (Weinheim)*, 16 (1995) 1184-1186.
- 2833 Huby, T., Schroöder, W., Doucet, C., Chapman, J. and Thillet, J.: Characterization of the N-terminal and C-terminal domains of human apolipoprotein(a): relevance to fibrin binding. *Biochemistry*, 34 (1995) 7385-7393.
- 2834 Ji, Y. and Jonas, A.: Properties of an N-terminal proteolytic fragment of apolipoprotein AI in solution and in reconstituted high density lipoproteins. *J. Biol. Chem.*, 270 (1995) 11290-11297.
- 2835 LaDu, M.J., Pederson, T.M., Frail, D.E., Reardon, C.A., Getz, G.S. and Falduto, M.T.: Purification of apolipoprotein E attenuates isoform-specific binding to β -amyloid. *J. Biol. Chem.*, 270 (1995) 9039-9042.
- 2836 Meng, Q., Bergeron, J., Sparks, D.L. and Marcel, Y.L.: Role of apolipoprotein A-I in cholesterol transfer between lipoproteins. Evidence for involvement of specific apoA-I domains. *J. Biol. Chem.*, 270 (1995) 8588-8596.
- 2837 Sviridov, D. and Fidge, N.: Pathway of cholesterol efflux from human hepatoma cells. *Biochim. Biophys. Acta*, 1256 (1995) 210-220.

See also 2920, 2962.

13. STEROIDS

13c. Estrogens

- 2838 Ji, A.J., Nunez, M.F., Machacek, D., Ferguson, J.E., Iossi, M.F., Kao, P.C. and Landers, J.P.: Separation of urinary estrogens by micellar electrokinetic chromatography. *J. Chromatogr. B*, 669 (1995) 15-26.

16. NITRO AND NITROSO COMPOUNDS

- 2839 Spicka, J., Lukavska, A. and Helclova, Z.: (Comparison of photometric and isotachophoretic determinations of nitrates in waters). *Vodni Hospod. Ochr. Ovzduši*, 43 (1993) 36-38; *C.A.*, 122 (1995) 222250h.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

See 2693.

17b. Catecholamines and their metabolites

- 2840 Robert, F., Bert, L., Denoroy, L. and Renaud, B.: Capillary zone electrophoresis with laser-induced fluorescence detection for the determination of nanomolar concentrations of noradrenaline and dopamine: application to brain microdialysate analysis. *Anal. Chem.*, 67 (1995) 1838-1844.
- 2841 Sheppard, R.L., Tong, X., Cai, J. and Henion, J.D.: Chiral separation and detection of terbutaline and ephedrine by capillary electrophoresis coupled with ion spray mass spectrometry. *Anal. Chem.*, 67 (1995) 2054-2058.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

18a. Amino acids and their derivatives

- 2842 Bonfichi, R., Dallanocce, C., Lociuo, S., Spada, A.: Free-solution capillary electrophoretic resolution of chiral amino acids via derivatization with homochiral isothiocyanates. *J. Chromatogr. A*, 707 (1995) 355-365.
- 2843 Dawson, L.A., Stow, J.M., Dourish, C.T. and Routledge, C.: Analysis of glutamate in striatal microdialysates using capillary electrophoresis and laser-induced fluorescence detection. *J. Chromatogr. A*, 700 (1995) 81-87.
- 2844 De Silva, K.H.: Capillary electrophoretic separation and mechanism study of chiral amino acids and peptides with cyclodextrins. Avail. *Univ. Microfilms Int.*, Order No. DA9504007, 1994, 284 p.; *C.A.*, 122 (1995) 329498x.
- 2845 De Witte, P.A., Cuveele, J.F., Merlevede, W.J. and Vandenneede, J.R.: Analysis of phosphorylhydroxyamino acids present in hydrolyzed cell extracts using dansyl derivatization. *Anal. Biochem.*, 226 (1995) 1-9.
- 2846 Kitaoka, Y., Kobayashi, M., Kawamoto, K. and Ichimoto, I.: Studied of the complexation of the hydroxyboryl compounds by zone electrophoresis: electrophoretic behavior of *p*-boronophenylalanine in various carboxylic acid supporting solutions. *Annu. Rep. Res. React. Inst., Kyoto Univ.*, 27 (1994) 224-228; *C.A.*, 123 (1995) 56517z.
- 2847 Lalljie, S.P.D. and Sandra, P.: MEKC analysis of FITC and DTAF amino acid derivatives with LIF detection. *Chromatographia*, 40 (1995) 513-518.
- 2848 Lalljie, S.P.D. and Sandra, P.: MEKC analysis of FITC and DTAF amino acid derivatives with LIF detection: practical and quantitative aspects in the analysis of FITC and DTAF amino acid derivatives by capillary electrophoresis and LIF detection. *Chromatographia*, 40 (1995) 519-526.
- 2849 Malone, M.A., Zuo, H., Lunte, S.M. and Smyth, M.R.: Determination of tryptophan and kynurenine in brain microdialysis samples by capillary electrophoresis with electrochemical detection. *J. Chromatogr. A*, 700 (1995) 73-80.

- 2850 Tivesten, A. and Folestad, S.: Separation of precolumn-labelled D- and L-amino acids by micellar electrokinetic chromatography with UV and fluorescence detection. *J. Chromatogr. A*, 708 (1995) 323-337.
- 2851 Vogt, C., Georgi, A. and Werner, G.: Enantiomeric separation of D/L-carnitine using HPLC and CZE after derivatization. *Chromatographia*, 40 (1995) 287-295.
- 2852 Wan, H., Andersson, P.E., Engström, A. and Blomberg, L.G.: Direct and indirect chiral separation of amino acids by capillary electrophoresis. *J. Chromatogr. A*, 704 (1995) 179-193.
- See also 2770, 2857, 2864, 2916, 2918.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 2853 Asamoah, K.A., Atkinson, P.G.P., Carter, W.G. and Sale, G.J.: Studies on an insulin-stimulated insulin receptor serine kinase activity: separation of the kinase activity from the insulin receptor and its reconstitution back to the insulin receptor. *Biochem. J.*, 308 (1995) 915-922.
- 2854 Borromeo, V., Berrini, A., Secchi, C., Brambilla, G.F. and Cantafora, A.: Matrix-assisted laser desorption mass spectrometry for the detection of recombinant bovine growth hormone in sustained-release form. *J. Chromatogr. B*, 669 (1995) 366-371.
- 2855 Chu, Y.-H., Kirby, D.P. and Karger, B.L.: Free solution identification of candidate peptides from combinatorial libraries by affinity capillary electrophoresis/mass spectrometry. *J. Am. Chem. Soc.*, 117 (1995) 5419-5420; *C.A.*, 123 (1995) 33635x.
- 2856 Churcher, Y. and Moss, S.E.: Inhibition of interleukin-2-mediated DNA synthesis in activated human T-lymphoblasts by okadaic acid is accompanied by hyperphosphorylation of Ick. *Biochem. J.*, 308 (1995) 777-783.
- 2857 Cui, H., Leon, J., Reusaet, E. and Bult, A.: Selective determination of peptides containing specific amino acid residues by high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr. A*, 704 (1995) 27-36 - a review with 51 refs.
- 2858 Dette, C. and Wätzig, H.: Separation of r-hirudin from similar substances by capillary electrophoresis. *J. Chromatogr. A*, 700 (1995) 89-94.
- 2859 Dupin, P., Galinou, F. and Bayol, A.: Analysis of recombinant human growth hormone and its related impurities by capillary electrophoresis. *J. Chromatogr. A*, 707 (1995) 396-400.
- 2860 Fleming, L.M. and Johnson, G.V.W.: Modulation of the phosphorylation state of tau *in situ*: the roles of calcium and cyclic AMP. *Biochem. J.*, 309 (1995) 41-47.
- 2861 Huang, M.X.: Reliable peptide separations using a polar bonded-phase capillary electrophoresis column. *Am. Biotechnol. Lab.*, 13 (1995) 80; *C.A.*, 122 (1995) 309989e.
- 2862 Khalkhali-Ellis, Z.: An improved SDS-polyacrylamide gel electrophoresis for resolution of peptides in the range of 3.5-200 kDa. *Prepar. Biochem.*, 25 (1995) 1-9.
- 2863 Lowther, W.T. and Dunn, B.M.: Kinetics of enzyme-catalyzed oligopeptide cleavage monitored by capillary zone electrophoresis: comparison to spectrophotometric and HPLC methods. *Lett. Pept. Sci.*, 1 (1994) 89-94; *C.A.*, 122 (1995) 309339t.
- 2864 Mank, A.J.G. and Yeung, E.S.: Diode laser-induced fluorescence detection in capillary electrophoresis after pre-column derivatization of amino acids and small peptides. *J. Chromatogr. A*, 708 (1995) 309-321.
- 2865 Nishi, N., Inui, M., Miyataka, H., Oya, H. and Wada, F.: Western blot analysis of epidermal growth factor using gelatin-coated polyvinylidene difluoride membrane. *Anal. Biochem.*, 227 (1995) 401-402.
- 2866 Saito, T., Kawabata, S.-i., Shigenaga, T., Takayenoki, Y., Cho, J., Nakajima, H., Hirata, M. and Iwanaga, S.: A novel big defensin identified in horseshoe crab hemocytes: isolation, amino acid sequence, and antibacterial activity. *J. Biochem. (Tokyo)*, 117 (1995) 1131-1137.
- 2867 Shippy, S.A., Jankowski, J.A. and Sweedler, J.V.: Analysis of trace level peptides using capillary electrophoresis with UV laser-induced fluorescence. *Anal. Chim. Acta*, 307 (1995) 163-171.
- 2868 Suzuki, K. and Sato, T.: Method and apparatus for detection of cytokines after capillary electrophoresis. *Jpn. Kokai Tokkyo Koho JP 07 98,319 [95 98,319] (Cl. G01N33/68)*, 11 Apr. 1995, Appl. 93/263,084, 28 Sep. 1993; 4 pp.; *C.A.*, 123 (1995) 5108j.
- 2869 Van Orman, H.B., Liversidge, G.G. and McIntire, G.L.: The electrophoretic mobility of tripeptides as a function of pH and ionic strength: comparison with iontophoretic flux data. *Pharm. Res.*, 12 (1995) 751-755; *C.A.*, 122 (1995) 299243r.
- See also 2716, 2756, 2844, 2874, 2882.
- 18c. *Elucidation of structure of proteins and enzymes*
- 2870 Aggeler, R., Haughton, M.A. and Capaldi, R.A.: Disulfide bond formation between the COOH-terminal domain of the β subunits and the γ and ϵ subunits of the *Escherichia coli* F₁-ATPase. Structural implications and functional consequences. *J. Biol. Chem.*, 270 (1995) 9185-9191.
- 2871 Arystarhova, E., Gibbons, D.L. and Sweadner, K.J.: Topology of the Na,K-ATPase. Evidence for externalization of a labile transmembrane structure during heating. *J. Biol. Chem.*, 270 (1995) 8785-8796.
- 2872 Gorg, A., Boguth, G., Obermaier, C., Posch, A. and Weiss, W.: Two-dimensional polyacrylamide gel electrophoresis with immobilized pH gradients in the first dimension (IPG-Dalt): the state of the art and the controversy of vertical versus horizontal systems. *Electrophoresis (Weinheim)*, 16 (1995) 1079-1086.
- 2873 Zhou, S. and Admon, A.: Enzymic digestion of proteins in zinc chloride and Ponceau S stained gels. In: Crabb, J.W. (Editor), 2Tech. Protein Chem. VI, [Pap. Symp. Protein Soc.] 8th 1994, Academic, San Diego, 1995, pp. 161-167; *C.A.*, 123 (1995) 28754f.
- See also 2884, 2894, 2913, 2919, 2965, 2966, 3019, 3074.
19. PROTEINS
- 19a. *General techniques*
- 2874 Basak, S.K. and Ladisch, M.R.: Correlation of electrophoretic mobilities of proteins and peptides with their physicochemical properties. *Anal. Biochem.*, 226 (1995) 51-58.
- 2875 Basak, S.K., Velayudhan, A., Kohlmann, K. and Ladisch, M.R.: Electrochromatographic separation of proteins. *J. Chromatogr. A*, 707 (1995) 69-76.

- 2876 Blackledge, J.A. and Alexander, A.J.: Investigation of polyethylene membranes as potential sample support for linking SDS-PAGE with MALDI-TOF MS for mass measurement of proteins. In: Crabb, J.W. (Editor), 2Tech. Protein Chem. VI, [Pap. Symp. Protein Soc.], 8th 1994, Academic, San Diego, 1995, pp. 13-19; C.A., 123 (1995) 51432h.
- 2877 Burggraf, D., Weber, G. and Lottspeich, F.: Free flow-isoelectric focusing of human cellular lysates as sample preparation for protein analysis. *Electrophoresis (Weinheim)*, 16 (1995) 1010-1015.
- 2878 Chiu, R.W., Jimenez, J.C. and Monnig, C.A.: High molecular weight polyarginine as a capillary coating for separation of cationic proteins by capillary electrophoresis. *Anal. Chim. Acta*, 307 (1995) 193-201.
- 2879 Cohen, N. and Grushka, E.: Capillary electrophoresis of proteins using polydimethyldiallylammonium chloride as a buffer additive. *J. Capillary Electrophor.*, 1 (1994) 112-115; C.A., 122 (1995) 285884p.
- 2880 Cole, K.D., Todd, P., Srinivasan, K. and Dutta, B.K.: Free-solution electrophoresis of proteins in an improved density gradient column and by capillary electrophoresis. *J. Chromatogr. A*, 707 (1995) 77-85.
- 2881 Douglas, N.G., Humffray, A.A., Pratt, H.R.C. and Stevens, G.W.: Electrophoretic mobilities of proteins and protein mixtures. *Chem. Eng. Sci.*, 50 (1995) 743-754; C.A., 122 (1995) 260335x.
- 2882 Erim, F.B., Cifuentes, A., Poppe, H. and Kraak, J.C.: Performance of a physically adsorbed high-molecular-mass polyethylenimine layer as coating for the separation of basic proteins and peptides by capillary electrophoresis. *J. Chromatogr. A*, 708 (1995) 356-361.
- 2883 Fernandez-Patron, C., Madrazo, J., Hardy, E., Mendez, E., Frank, R. and Castellanos-Serra, L.: Single-step electrotransfer of reverse-stained proteins from sodium dodecyl sulfate-polyacrylamide gel onto reversed-phase minicardridge and subsequent desalting and elution with a conventional high-performance liquid chromatography gradient system for analysis. *Electrophoresis (Weinheim)*, 16 (1995) 911-920.
- 2884 Galat, A., Bouet, F. and Rivière, S.: Amino acid compositions of proteins and their identities. *Electrophoresis (Weinheim)*, 16 (1995) 1095-1103.
- 2885 Gianazza, E.: Isoelectric focusing as a tool for the investigation of post-translational processing and chemical modifications of proteins. *J. Chromatogr. A*, 705 (1995) 67-87 - a review with 261 refs.
- 2886 Grimm, R. and Herold, M.: Micropreparative single run fraction collection of peptides separated by CZE for protein sequencing. *J. Capillary Electrophor.*, 1 (1994) 79-82; C.A., 122 (1995) 310098v.
- 2887 Hiraoka, A.: (Analysis of proteins in clinical samples by capillary electrophoresis). *Kensa to Gijutsu*, 22 (1994) 869-872; C.A., 122 (1995) 260045c - a review with 17 refs.
- 2888 Hjerten, S., Valtcheva, L. and Li, Y.-M.: A simple method for desalting and concentration of microliter volumes of protein solutions with special reference to capillary electrophoresis. *J. Capillary Electrophor.*, 1 (1994) 83-89; C.A., 122 (1995) 310201y.
- 2889 Jansen, S., Kriegesmann, B., Jones, R. and Brenig, B.: Ultraviolet shadowing of proteins in preparative sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Anal. Biochem.*, 228 (1995) 177-178.
- 2890 Lemkin, P.F., Orr, G.A., Goldstein, M.P., Creed, J., Whitley, E., Myrick, J.E. and Merrill, C.R.: The protein disease database. *Electrophoresis (Weinheim)*, 16 (1995) 1175.
- 2891 Nemoto, T., Ota, M., Ohara-Nemoto, Y. and Kaneko, M.: Identification of dimeric structure of proteins by use of the glutathione S-transferase-fusion expression system. *Anal. Biochem.*, 227 (1995) 396-399.
- 2892 Parrado, J., Ayala, A. and Machado, A.: One-step electroblotting of 2- to 100-kDa proteins onto a PVDF membrane. *Appl. Biochem. Biotechnol.*, 53 (1995) 285-292; C.A., 123 (1995) 51524q.
- 2893 Patterson, S.D.: Matrix-assisted laser-desorption/ionization mass spectrometric approaches for the identification of gel-separated proteins in the 5-50 pmol range. *Electrophoresis (Weinheim)*, 16 (1995) 1104-1114.
- 2894 Sanchez, J.-C., Appel, R.D., Golaz, O., Pasquali, C., Ravier, F., Bairoch, A. and Hochstrasser, D.F.: Inside SWISS-2DPAGE database. *Electrophoresis (Weinheim)*, 16 (1995) 1131-1151 - a review with 48 refs.
- 2895 Tomlinson, A.J., Braddock, W.D., Benson, L.M., Oda, R.P. and Naylor, S.: Preliminary investigations of preconcentration-capillary electrophoresis-mass spectrometry. *J. Chromatogr. B*, 669 (1995) 67-73.
- 2896 Wang, H.-P. and Liu, C.-M.: Use of capillary electrophoresis for quantitating the concentration of protein components and of the total protein in fluids. *PCT Int. Appl. WO 95 10,041 (Cl. G01N27/447)*, 13 Apr. 1995, US Appl. 133,514, 07 Oct. 1993; 54 pp.; C.A., 123 (1995) 29040g.
- 2897 Wu, J. and Pawliszyn, J.: Protein analysis by isoelectric focusing in a capillary array with an absorption imaging detector. *J. Chromatogr. B*, 669 (1995) 39-43.
- 2898 Yamamura, T.: (Polyacrylamide gel electrophoresis). *Lipid*, 6 (1995) 65-71; C.A., 123 (1995) 4778r - a review with 7 refs.
- 2899 Zimmermann, C.W., Grus, F.-H. and Dux, R.: Multivariate statistical comparison of autoantibody-repertoires (Western blots) by discriminant analysis. *Electrophoresis (Weinheim)*, 16 (1995) 941-947.
- See also 2671, 2674, 2676, 2699, 2700, 2726, 2741, 2765, 2772, 2774, 2776, 2872, 3019, 3140, 3147, 3293.
- 19b. *Proteins of cells, viruses and subcellular particles*
- 2900 Callaghan, R. and Riordan, J.R.: Collateral sensitivity of multidrug resistant cells to narcotic analgesics is due to effects on the plasma membrane. *Biochim. Biophys. Acta*, 1236 (1995) 155-162.
- 2901 Chan, P.K. and Chan, F.Y.: Nucleophosmin/B23 (NPM) oligomer is a major and stable entity in HeLa cells. *Biochim. Biophys. Acta*, 1262 (1995) 37-42.
- 2902 Dreher, D., Vargas, J.R., Hochstrasser, D.F. and Junod, A.F.: Effects of oxidative stress and Ca²⁺ agonists on molecular chaperones in human umbilical vein endothelial cells. *Electrophoresis (Weinheim)*, 16 (1995) 1205-1214.

- 2903 Flesch, B., Ntambi, E. and Neppert, J.: Biotinylation: a non-radioactive method for the identification of cell surface antigens in immunoprecipitates. *Electrophoresis (Weinheim)*, 16 (1995) 757-762.
- 2904 Freyria, A.-M., Ronzière, M.-C., Boutillon, M.-M. and Herbage, D.: Two-dimensional electrophoresis of intracellular and secreted protein synthesized by fetal bovine chondrocytes in high-density culture. *Electrophoresis (Weinheim)*, 16 (1995) 1268-1272.
- 2905 Haldeman, M.T., Finley, D. and Pickart, C.M.: Dynamics of ubiquitin conjugation during erythroid differentiation *in vitro*. *J. Biol. Chem.*, 270 (1995) 9507-9516.
- 2906 Kato, K., Ito, H., Hasegawa, K., Inaguma, Y., Suzuki, A., Kozawa, O. and Asano, T.: Enhancement of stress-induced synthesis of stress proteins by mastoparan in C6 rat glioma cells. *J. Biochem. (Tokyo)*, 118 (1995) 149-153.
- 2907 Khatypov, R.A., Shirshikova, G.N., Ladygin, V.G. and Shuvalov, V.A.: (The polypeptide composition of thylakoid membranes from *Chlamydomonas reinhardtii* mutants with inactive photosystem II) *Biokhimiya (Moscow)*, 60 (1995) 573-577.
- 2908 Kondo, T., Mihara, K., Inoue, Y., Iijima, M. and Namba, M.: Two-dimensional electrophoretic analysis of down-regulated proteins in human fibroblasts immortalized by treatment with either 4-nitroquinoline 1-oxide or ⁶⁰Co gamma rays. *Electrophoresis (Weinheim)*, 16 (1995) 1067-1073.
- 2909 Konishi, T., Karasaki, Y., Nomoto, M., Ohmori, H., Shibata, K., Abe, T., Shimizu, K., Itoh, H. and Higashi, K.: Induction of heat shock protein 70 and nucleolin and their intracellular distribution during early stage of liver regeneration. *J. Biochem. (Tokyo)*, 117 (1995) 1170-1177.
- 2910 Kwon, H.J., Yoshida, M., Muroya, K., Hattori, S., Nishida, E., Fukui, Y., Beppu, T. and Horimouchi, S.: Morphology of ras-transformed cells becomes apparently normal again with tyrosine kinase inhibitors without a decrease in the ras-GTP complex. *J. Biochem. (Tokyo)*, 118 (1995) 221-228.
- 2911 Lloyds, D. and Hallett, M.B.: Development of oxidase 'priming' in maturing HL60 cells: correlation with protein expression and tyrosine phosphorylation. *Biochim. Biophys. Acta*, 1267 (1995) 65-72.
- 2912 Marks, M.S., Germain, R.N. and Bonifacino, J.S.: Transient aggregation of major histocompatibility complex class II chains during assembly in normal spleen cells. *J. Biol. Chem.*, 270 (1995) 10475-10481.
- 2913 Muguruma, M., Nishimuta, S., Tomisaka, Y., Ito, T. and Matsuura, S.: Organization of the functional domains in membrane cytoskeletal protein talin. *J. Biochem. (Tokyo)*, 117 (1995) 1036-1042.
- 2914 Paulik, M., Morré, D.J. and Morré, D.M.: A 55 kDa protein of transitional endoplasmic reticulum from rat liver binds retinol. *Biochim. Biophys. Acta*, 1266 (1995) 273-277.
- 2915 Römer, I., Jungblut, P., Reik, W., Otto, A. and Klose, J.: A novel strategy to identify maternal and paternal inheritance in the mouse. *Electrophoresis (Weinheim)*, 16 (1995) 823-830.
- 2916 Simonin, D., Diaz, J.-J., Kindbeiter, K., Denoroy, L. and Madjar, J.-J.: Phosphorylation of ribosomal protein L30 after herpes simplex virus type 1 infection. *Electrophoresis (Weinheim)*, 16 (1995) 854-859.
- 2917 Simonin, D., Diaz, J.-J., Kindbeiter, K., Pernas, P. and Madjar, J.-J.: Phosphorylation of herpes simplex virus type 1 Us11 protein is independent of viral genome expression. *Electrophoresis (Weinheim)*, 16 (1995) 1317-1322.
- 2918 Suzuki, K., Yamaguchi, T., Oshizawa, T., Yamamoto, Y., Nishimaki-Mogami, T., Hayakawa, T. and Takahashi, A.: Okadaic acid induces both augmentation and inhibition of opsonized zymosan-stimulated superoxide production by differentiated HL-60 cells. Possible involvement of dephosphorylation of a cytosolic 21K protein in respiratory burst. *Biochim. Biophys. Acta*, 1266 (1995) 261-267.
- 2919 Wasinger, V.C., Cordwell, S.J., Cerpa-Poljak, A., Yan, J.X., Gooley, A.A., Wilkins, M.R., Duncan, M.W., Harris, R., Williams, K.L. and Humphery-Smith, I.: Progress with gene-product mapping of the Mollicutes: *Mycoplasma genitalium*. *Electrophoresis (Weinheim)*, 16 (1995) 1090-1094.

See also 2925, 3042, 3123, 3176, 3208.

19c. *Proteins synthesized by genetic manipulation, monoclonal antibodies*

- 2920 Abumiya, T., Enryoji, K.-i., Kokawa, T., Kamikubo, Y.-i. and Kato, H.: An anti-tissue factor pathway inhibitor (TFPI) monoclonal antibody recognized the third Kunitz domain (K3) of free-form TFPI but not lipoprotein-associated forms in plasma. *J. Biochem. (Tokyo)*, 118 (1995) 178-182.
- 2921 Calero, M., Méndez, E. and Garcia, E.: Expression of the human complex-forming glycoprotein HC (α 1-microglobulin) in *Escherichia coli*. *Biochim. Biophys. Acta*, 1249 (1995) 91-99.
- 2922 Dairaghi, D.J., Shadel, G.S. and Clayton, D.A.: Human mitochondrial transcription factor A and promoter spacing integrity are required for transcription initiation. *Biochim. Biophys. Acta*, 1271 (1995) 127-134.
- 2923 De Zoysa, A., Elstratiou, A., George, R.C., Jahkola, M., Vuopio-Varkila, J., Deshevoi, S., Tseneva, G. and Rikushin, Y.: Molecular epidemiology of *Corynebacterium diphtheriae* from northwestern Russia and surrounding countries studied by using ribotyping and pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 33 (1995) 1080-1083; *C.A.*, 123 (1995) 5056.
- 2924 Masada, R.I., Hague, C., Seid, R., Ho, S., McAlister, S., Pigiet, V. and Starr, C.M.: Fluorophore-assisted-carbohydrate-electrophoresis (FACE) for determining the nature and consistency of recombinant protein glycosylation. *Trends Glycosci. Glyco-technol.*, 7 (1995) 133-147; *C.A.*, 123 (1995) 28851k.
- 2925 Negrini, C., Rivolta, M.N., Kalinec, F. and Kachar, B.: Cloning of an organ of Corti anion exchanger 2 isoform with a truncated C-terminal domain. *Biochim. Biophys. Acta*, 1236 (1995) 207-211.
- 2926 Seifert, H. and Gerner-Smidt, P.: Comparison of ribotyping and pulsed-field gel electrophoresis for molecular typing of *Acinetobacter* isolates. *J. Clin. Microbiol.*, 33 (1995) 1402-1407; *C.A.*, 123 (1995) 5057s.
- 2927 Yu, Y., Terada, K., Nagasaki, A., Takiguchi, M. and Mori, M.: Preparation of recombinant argininosuccinate synthetase and argininosuccinate lyase: expression of the enzymes in rat tissues. *J. Biochem. (Tokyo)*, 117 (1995) 952-957.

See also 2974, 3061, 3167.

19d. Microbial and plant proteins

- 2928 Bergantino, E., Dainese, P., Cerovic, Z., Sechi, S. and Bassi, R.: A post-translational modification of the photosystem II subunit CP20 protects maize from cold stress. *J. Biol. Chem.*, 270 (1995) 8474-8481.
- 2929 Boutillon, C., Wintjes, R., Lippens, G., Drobecq, H. and Tartar, A.: Synthesis, three-dimensional structure and specific ¹⁵N-labelling of the streptococcal protein G B1-domain. *Eur. J. Biochem.*, 231 (1995) 166-180.
- 2930 Cattaneo, T.M.P., Feroldi, A., Toppino, P.M. and Olieman, C.: Sample preparation for selective and sensitive detection of soya proteins in dairy products with chromatographic and electrophoretic techniques. *Neth. Milk Dairy J.*, 48 (1994) 225-234; *C.A.*, 122 (1995) 263757x.
- 2931 Curioni, A., Morel, M.-H., Furegon, L., Redaelli, R. and Dal Belin Peruffo, A.: Purification of wheat glutenin subunits by preparative acid and two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1005-1009.
- 2932 Davidsen, N.B.: Two-dimensional electrophoresis of acidic proteins isolated from ozone-stressed Norway spruce needles (*Picea abies* L. Karst): separation method and image processing. *Electrophoresis (Weinheim)*, 16 (1995) 1305-1311.
- 2933 Jonsson, H., Burtsoff-Asp, C. and Guss, B.: Streptococcal protein MAG - a protein with broad albumin binding specificity. *Biochim. Biophys. Acta*, 1249 (1995) 65-71.
- 2934 Kawaguchi, S.-i. and Kuramitsu, S.: Separation of heat-stable proteins from *Thermus thermophilus* HB8 by two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1060-1066.
- 2935 Latter, G.I., Boutell, T., Monardo, P.J., Kobayashi, R., Fletcher, B., McLaughlin, C.S. and Garrels, J.I.: A *Saccharomyces cerevisiae* Internet protein resource now available. *Electrophoresis (Weinheim)*, 16 (1995) 1170-1174.
- 2936 Leconte, A., Lebrun, P., Nicolas, D. and Seguin, M.: (Use of electrophoresis for the identification of *Hevea* clones). *Plant., Rech., Dev.*, 1 (1994) 28-36; *C.A.*, 123 (1995) 4746d - a review with 9 refs.
- 2937 Lookhart, G.L. and Bean, S.R.: Rapid differentiation of oat cultivars and of rice cultivars by capillary zone electrophoresis. *Cereal Chem.*, 72 (1995) 312-316; *C.A.*, 122 (1995) 310126c.
- 2938 Miura, K., Yokono, T. and Kawaguchi, H.: (Two-dimensional electrophoresis of *Flammulina velutipes*). *Mokuzai Gakkaishi*, 40 (1994) 874-878; *C.A.*, 122 (1995) 310335v.
- 2939 Papeta, N.F., Tsilosani, G.M., Nikolaishvili, N.T. and Kvesitadze, G.I.: (Identification, purification and characterization of xylanase from *Nocardioopsis* sp. inhibiting the growth of some phytopathogenic fungi). *Biokhimiya (Moscow)*, 60 (1995) 578-584.
- 2940 Petersen, A., Becker, W.-M., Moll, H., Blümke, M. and Schlaak, M.: Studies on the carbohydrate moieties of the timothy grass pollen allergen Phl p1. *Electrophoresis (Weinheim)*, 16 (1995) 869-875.
- 2941 Posch, A., van den Berg, B.M. and Görg, A.: Association of protein polymorphism among pepper (*Capsicum annuum* L.) inbred lines with agronomic performance of their crosses. *Electrophoresis (Weinheim)*, 16 (1995) 860-864.
- 2942 Posch, A., van den Berg, B.M., Burg, H.C.J. and Görg, A.: Genetic variability of carrot seed proteins analyzed by one- and two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis (Weinheim)*, 16 (1995) 1312-1316.
- 2943 Posch, A., Weiss, W., Wheeler, C., Dunn, M.J. and Görg, A.: Sequence analysis of wheat grain allergens separated by two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis (Weinheim)*, 16 (1995) 1115-1119.
- 2944 Ranieri, A., Castagna, A., Bini, L. and Soldatini, G.F.: Two-dimensional electrophoresis of thylakoid protein patterns in two wheat cultivars with different sensitivity to sulfur dioxide. *Electrophoresis (Weinheim)*, 16 (1995) 1301-1304.
- 2945 Redaelli, R., Morel, M.-H., Autran, J.-C. and Pogna, N.E.: Genetic analysis of low Mr glutenin subunits fractionated by two-dimensional electrophoresis (A-PAGE x SDS-PAGE). *J. Cereal Sci.*, 21 (1995) 5-13; *C.A.*, 122 (1995) 285881k.
- 2946 Sasek, A. and Cerny, J.: (The model electrophoretic spectra of HMW gliadins and glutenins of common wheat varieties cultivated in the Czech Republic in 1993). *Rostl. Vyroba*, 41 (1995) 163-168; *C.A.*, 123 (1995) 51517q.
- 2947 Werner, W.E.: Ferguson plot analysis of high molecular weight glutenin subunits by capillary electrophoresis. *Cereal Chem.*, 72 (1995) 248-251; *C.A.*, 123 (1995) 8159n.
- 2948 Zivy, M., El Madidi, S. and Thiellement, H.: Distance indices in a comparison between the A, D, I and R genomes of the *Triticeae* tribe. *Electrophoresis (Weinheim)*, 16 (1995) 1295-1300.

See also 2907.

19e. Proteins of blood, serum and blood cells

- 2949 Bläser, J., Triebel, S. and Tschesche, H.: A sandwich enzyme immunoassay for the determination of neutrophil lipocalin in body fluids. *Clin. Chim. Acta*, 235 (1995) 137-145.
- 2950 Bueno, S.M.A., Haupt, K. and Vijayalakshmi, M.A.: Separation of immunoglobulin G from human serum by pseudobioaffinity chromatography using immobilized L-histidine in hollow fibre membranes. *J. Chromatogr. B*, 667 (1995) 57-67.
- 2951 Corsi, D., Galluzzi, L., Crinelli, R. and Magnani, M.: Ubiquitin is conjugated to the cytoskeletal protein α -spectrin in mature erythrocytes. *J. Biol. Chem.*, 270 (1995) 8928-8935.
- 2952 Denton, K.A. and Harris, R.: High-performance capillary electrophoretic separation of human serum albumin using a neutral coated capillary. *J. Chromatogr. A*, 705 (1995) 335-341.
- 2953 Goodarzi, M.T. and Turner, G.A.: Decreased branching, increased fucosylation and changed sialylation of alpha-1-proteinase inhibitor in breast and ovarian cancer. *Clin. Chim. Acta*, 236 (1995) 161-171.
- 2954 Gravel, P., Sanchez, J.-C., Walzer, C., Golaz, O., Hochstrasser, D.F., Balant, L.P., Hughes, G.J., Garcia-Sevilla, J. and Guimon, J.: Human blood platelet protein map established by two-dimensional polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1152-1159.
- 2955 Greco, N.J., Tandon, N.N., Jackson, B. and Jamieson, G.A.: Adenine nucleotide binding and photoincorporation in Glanzmann's thrombasthenia platelets. *Biochim. Biophys. Acta*, 1236 (1995) 142-148.

- 2956 Hoffer, L., Schwinn, H., Biesert, L. and Josic, D.: Improved virus safety and purity of a chromatographically produced Factor IX concentrate by nanofiltration. *J. Chromatogr. B*, 669 (1995) 187-196.
- 2957 Hunt, A.J., Lynch, P.D., Londo, T., Dimond, P., Gordon, N.F., McCormack T., Schutz, A., Percoskie, M., Cao, X., McGrath, J.P., Putney, S. and Hamilton, R.A.: Development and monitoring of purification process for nerve growth factor fusion antibody. *J. Chromatogr. A*, 708 (1995) 61-70.
- 2958 Inoue, J., Yoshida, T., Cho, H. and Shiba, K.: (Improvement of the standard electrophoresis procedure for serum protein fractions. Effect of constant current and constant voltage on cellulose acetate membrane electrophoresis using Separax SP). *Seibutsu Butsuri Kagaku*, 39 (1995) 1-14; C.A., 122 (1995) 285880j.
- 2959 Kato, T., Ogami, K., Shimada, Y., Iwamatsu, A., Sohma, Y., Akahori, H., Horie, K., Kokubo, A., Kudo, Y., Maeda, E., Kobayashi, K., Ohashi, H. et al.: Purification and characterization of thrombopoietin. *J. Biochem. (Tokyo)*, 118 (1995) 229-236.
- 2960 Kobayashi, K., Takahashi, K. and Nagasawa, S.: The role of tyrosine phosphorylation and Ca²⁺ accumulation in Fcγ-receptor-mediated phagocytosis of human neutrophils. *J. Biochem. (Tokyo)*, 117 (1995) 1156-1161.
- 2961 Kroon, D.J., Goltra, S. and Sharma, B.: Analysis of monoclonal antibodies by sodium dodecyl sulfate-capillary gel electrophoresis with special reference to quantitation of half-antibody. *J. Capillary Electrophor.*, 2 (1995) 34-39; C.A., 123 (1995) 7410a.
- 2962 Lehmann, R., Liebich, H., Grübler, G. and Voelter, W.: Capillary electrophoresis of human serum proteins and apolipoproteins. *Electrophoresis (Weinheim)*, 16 (1995) 998-1001.
- 2963 Magi, B., Marzocchi, B., Bini, L., Cellesi, C., Rossolini, A. and Pallini, V.: Two-dimensional electrophoresis of human serum proteins modified by ampicillin during therapeutic treatment. *Electrophoresis (Weinheim)*, 16 (1995) 1190-1192.
- 2964 Marzocchi, B., Magi, B., Bini, L., Cellesi, C., Rossolini, A., Massidda, O. and Pallini, V.: Two-dimensional gel electrophoresis and immunoblotting of human serum albumin modified by reaction with penicillins. *Electrophoresis (Weinheim)*, 16 (1995) 851-853.
- 2965 Matsumoto, N., Kita, K., Kojima, S., Yamamoto, K., Irimura, T., Miyagi, M., Tsunasawa, S. and Toyoshima, S.: Lymphocyte isoforms of mouse p50 LSP1, which are phosphorylated in mitogen-activated T cells, are formed through alternative splicing and phosphorylation. *J. Biochem. (Tokyo)*, 118 (1995) 237-243.
- 2966 Mhatre, R., Nashabeh, W., Schmalzing, D., Yao, X., Fuchs, M., Whitney, D. and Regnier, F.: Purification of antibody Fab fragments by cation-exchange chromatography and pH gradient elution. *J. Chromatogr. A*, 707 (1995) 225-231.
- 2967 Miller, I., Gutleb, A.C. and Gemeiner, M.: Two-dimensional electrophoresis for the study of blood/serum proteins of the otter, an endangered species. *Electrophoresis (Weinheim)*, 16 (1995) 1193-1198.
- 2968 Miller, I., Gutleb, A.C., Kranz, A. and Gemeiner, M.: Forensics on wild animals: differentiation between otter and pheasant blood using electrophoretic methods. *Electrophoresis (Weinheim)*, 16 (1995) 865-868.
- 2969 Okazaki, T., Yanagisawa, Y. and Nagai, T.: (Quantitative analysis of methalbumin by polyacrylamide gel electrophoresis). *Seibutsu Butsuri Kagaku*, 39 (1995) 37-39; C.A., 122 (1995) 262691j.
- 2970 Robinson, M.K., Myrick, J.E., Henderson, L.O., Coles, C.D., Powell, M.K., Orr, G.A. and Lemkin, P.F.: Two-dimensional protein electrophoresis and multiple hypothesis testing to detect potential serum protein biomarkers in children with fetal alcohol syndrome. *Electrophoresis (Weinheim)*, 16 (1995) 1176-1183.
- 2971 Rocha, J., Bohner, J. and Kömpf, J.: Transient bisalbuminemia: separation by isoelectric focusing of human albumin fractions linked to different numbers of benzylpenicilloyl groups. *Electrophoresis (Weinheim)*, 16 (1995) 1031-1033.
- 2972 Tohjo, H., Miyoshi, F., Uchida, E., Niiyama, M., Syuto, B., Moritsu, Y., Ichikawa, S. and Takeuchi, M.: Polyacrylamide gel electrophoretic patterns of chicken serum in acute inflammation induced by intramuscular injection of turpentine. *Poult. Sci.*, 74 (1995) 648-655; C.A., 122 (1995) 285882m.
- 2973 Tu, Y.P. and Yang, F.Y.: Zn²⁺-Mediated domain-domain communication in human erythrocyte Band 3. *J. Biochem. (Tokyo)*, 118 (1995) 161-167.
- 2974 Wohl, T., Klier, H. and Lottspeich, F.: Detection of nonfunctional overexpression gene products using two-dimensional gel electrophoresis with a narrowed pH range. *Electrophoresis (Weinheim)*, 16 (1995) 876-878.
- See also 2835, 2836, 2990, 2999, 3028, 3045, 3060, 3101, 3275.
- 19f. *Structural and muscle proteins*
- 2975 Freundorfer, S., Grupe, G. and Weickmann, D.: Mineral-bound noncollagenous proteins in archaeological human skeletons. *Electrophoresis (Weinheim)*, 16 (1995) 817-819.
- 2976 Fujii, T., Yamana, K., Ogoma, Y. and Kondo, Y.: Interaction of calponin with phospholipids. *J. Biochem. (Tokyo)*, 117 (1995) 999-1003.
- 2977 Gallardo, J.M., Sotelo, C.G., Pineiro, C. and Perez-Martin, R.I.: Use of capillary zone electrophoresis for fish species identification. Differentiation of flatfish species. *J. Agric. Food Chem.*, 43 (1995) 1238-1244; C.A., 122 (1995) 285886r.
- 2978 Hattori, A., Ishii, T., Tatsumi, R. and Takahashi, K.: Changes in the molecular types of connectin and nebulin during development of chicken skeletal muscle. *Biochim. Biophys. Acta*, 1244 (1995) 179-184.
- 2979 Iwata, M., Imamura, Y., Sasaki, T. and Hayashi, T.: Evidence for a short form of α1(IV) as a major polypeptide in bovine lens capsule. *J. Biochem. (Tokyo)*, 117 (1995) 1298-1304.
- 2980 Kovalyov, L.I., Shishkin, S.S., Efimochkin, A.S., Kovalyova, M.A., Ershova, E.S., Egorov, T.A. and Musalyamov, A.K.: The major protein expression profile and two-dimensional protein database of human heart. *Electrophoresis (Weinheim)*, 16 (1995) 1160-1169.
- 2981 Lamande, S.R., Chessler, S.D., Golub, S.B., Byers, P.H., Chan, D., Cole, W.G., Silience, D.O. and Bateman, J.F.: Endoplasmic reticulum-mediated quality control of type I collagen production by cells from osteogenesis imperfecta patients with mutations in the proα1(I) chain carboxyl-terminal propeptide which impair subunit assembly. *J. Biol. Chem.*, 270 (1995) 8642-8649.

- 2982 Miyake, B.: (Analysis of keratins from single hair by electrophoresis using colloidal Coomassie Brilliant Blue staining). *Kagaku Keisatsu Kenkyusho Hokoku, Hokogaku-hen*, 47 (1994) 24-29; *C.A.*, 122 (1995) 258107f.
- 2983 Pleissner, K.-P., Regitz-Zagrosek, V., Weise, C., Neuss, M., Krüde-wagen, B., Söding, P., Buchner, K., Hucho, F., Hildebrandt, A. and Fleck, E.: Chamber-specific expression of human myocardial proteins detected by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 841-850.
- 2984 Price, K.M., Sioman, R., Armstrong, P., Teisner, B. and Grudzinskas, J.G.: The typing of fetal antigen 2 in human amniotic fluid. *Clin. Chim. Acta*, 236 (1995) 181-194.
- 2985 Rehbein, H.: Differentiation of scombroid fish species (tunas, bonitos and mackerels) by isoelectric focusing, titration curve analysis and native polyacrylamide gel electrophoresis of sarco-plasmic proteins. *Electrophoresis (Weinheim)*, 16 (1995) 820-822.
- 2986 Sarmah, S. and Chakraborty, A.: Denaturing electrophoretic pattern of myosin from mithun (*Bos gaurs frontalis*). *Indian J. Anim. Sci.*, 64 (1994) 1203-1204; *C.A.*, 122 (1995) 310110t.
- See also 2705, 2899, 3039, 3276.
- 19g. *Protamines, histones and other nuclear proteins*
- 2987 Bedells, C.H. and Pennington, S.R.: Analysis of growth factor stimulated nucleocytoplasmic protein transport using two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1231-1239.
- 2988 Ibrahim, M.A., Hamed, R.R. and Rasched, I.: Purification and characterization of a novel acid-soluble nuclear protein from developing embryos of the camel tick *Hyalomma dromedarii* (Acarina: Ixodidae). *Biochim. Biophys. Acta*, 1249 (1995) 79-85.
- 2989 Touzet, P., Morin, C., Damerval, C., le Guilloux, M., Zivy, M. and de Vienne, D.: Characterizing allelic proteins for genome mapping in maize. *Electrophoresis (Weinheim)*, 16 (1995) 1289-1294.
- 19h. *Chromoproteins and metalloproteins*
- 2990 Bean, P., Sutphin, M.S., Liu, Y., Anton, R., Reynolds, T.B., Shoenfeld, Y. and Peter, J.B.: Carbohydrate-deficient transferrin and false-positive results for alcohol abuse in primary biliary cirrhosis: differential diagnosis by detection of mitochondrial autoantibodies. *Clin. Chem. (Washington)*, 41 (1995) 858-861.
- 2991 Beattie, J.H. and Richards, M.P.: Analysis of metallothionein isoforms by capillary electrophoresis: optimisation of protein separation conditions using micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 700 (1995) 95-103.
- 2992 Borguet, F., Cornelis, R., Delanghe, J., Lambert, M.-C. and Lameire, N.: Study of the chromium binding in plasma of patients on continuous ambulatory peritoneal dialysis. *Clin. Chim. Acta*, 238 (1995) 71-84.
- 2993 Domokos, G.: Method for the manufacture of highly pure hemoglobin. *Ger. DE 4,412,260 (Cl. C07K14/805)*, 09 Mar. 1995, *Appl.* 09 Apr. 1994; 7 p.; *C.A.*, 122 (1995) 274030f.
- 2994 Fago, A. and Weber, R.E.: The hemoglobin system of the hagfish *Myxine glutinosa*: aggregation state and functional properties. *Biochim. Biophys. Acta*, 1249 (1995) 109-115.
- 2995 Lee, B.H., Hibino, T., Takabe, T., Weisbeek, P.J. and Takabe, T.: Site-directed mutagenetic study on the role of negative patches on silene plastocyanin in the interactions with cytochrome f and photosystem I. *J. Biochem. (Tokyo)*, 117 (1995) 1209-1217.
- 2996 Richards, M.P. and Aagaard, P.J.: Separation of metallothionein isoforms by free-solution capillary electrophoresis in uncoated and surface-modified capillaries. *J. Capillary Electrophor.*, 1 (1994) 90-95; *C.A.*, 122 (1995) 310099w.
- 2997 Richards, M.P. and Beattie, J.H.: Analysis of metalloproteins and metal-binding peptides by capillary electrophoresis. *J. Capillary Electrophor.*, 1 (1994) 196-207; *C.A.*, 122 (1995) 309902w.
- 2998 Richards, M.P. and Beattie, J.H.: Comparison of different techniques for the analysis of metallothionein isoforms by capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 27-37.
- 2999 Van Kamp, G.J., Mulder, K., Kuiper, M. and Wolters, E.C.: Changed transferrin sialylation in Parkinson's disease. *Clin. Chim. Acta*, 235 (1995) 159-167.
- See also 2968.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 3000 Benito, B., Wahl, D., Stuedel, N., Cordier, A. and Steiner, S.: Effects of cyclosporine A on the rat liver and kidney protein pattern, and the influence of vitamin E and C coadministration. *Electrophoresis (Weinheim)*, 16 (1995) 1273-1283.
- 3001 Cossio, G., Sanchez, J.-C., Golaz, O., Wettstein, R. and Hochstrasser, D.F.: Spermatoocytes and round spermatids of rat testis: protein patterns. *Electrophoresis (Weinheim)*, 16 (1995) 1225-1230.
- 3002 Donovan, G.R., Street, M.D. and Baldo, B.A.: Separation of jumper ant (*Myrmecia pilosula*) venom allergens: a novel group of highly basic proteins. *Electrophoresis (Weinheim)*, 16 (1995) 804-810.
- 3003 Giometti, C.S., Tollaksen, S.L., Chubb, C., Williams, C. and Huberman, E.: Analysis of proteins from human breast epithelial cells using two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1215-1224.
- 3004 Inaguma, Y., Hasegawa, K., Goto, S., Ito, H. and Kato, K.: Induction of the synthesis of hsp27 and α B crystallin in tissues of heat-stressed rats and its suppression by ethanol or an α_1 -adrenergic antagonist. *J. Biochem. (Tokyo)*, 117 (1995) 1238-1243.
- 3005 Kanamori, T., Hayakawa, T., Suzuki, M. and Titani, K.: Identification of two 17-kDa rat parotid gland phosphoproteins, subjects for dephosphorylation upon β -adrenergic stimulation, as destrin- and cofilin-like proteins. *J. Biol. Chem.*, 270 (1995) 8061-8067.
- 3006 Kido, A., Kimura, Y. and Oya, M.: Group-specific component subtyping in semen and seminal stains by conventional isoelectric focusing. *Electrophoresis (Weinheim)*, 16 (1995) 1024-1026.
- 3007 Kinghorn, N.M., Norris, C.S., Paterson, G.R. and Otter, D.E.: Comparison of capillary electrophoresis with traditional methods to analyse bovine whey proteins. *J. Chromatogr. A*, 700 (1995) 111-123.

- 3008 Lin, J.D., Huang, C.C., Weng, H.F., Chen, S.C. and Jeng, L.B.: Comparison of membrane proteins from benign and malignant human thyroid tissues by two-dimensional polyacrylamide gel electrophoresis. *J. Chromatogr. B*, 667 (1995) 153-160.
- 3009 Marrakchi, N., Zingali, R.B., Karoui, H., Bon, C. and El Ayeb, M.: Cerastocytin, a new thrombin-like platelet activator from the venom of the Tunisian viper *Cerastes cerastes*. *Biochim. Biophys. Acta*, 1244 (1995) 147-156.
- 3010 Pacakova, V., Stulik, K., Hau, P.T., Jelinek, I., Vins, I. and Sykora, D.: Comparison of high-performance liquid chromatography and capillary electrophoresis for the determination of some bee venom components. *J. Chromatogr. A*, 700 (1995) 187-193.
- 3011 Peterson, G.R., Hill, J.P. and Otter, D.E.: Separation of β -lactoglobulin A, B and C variants of bovine whey using capillary zone electrophoresis. *J. Chromatogr. A*, 700 (1995) 105-110.
- 3012 Prestigiacomo, A.F., Chen, Z. and Stamey, T.A.: A universal calibrator for prostate specific antigen (PSA). *Scand. J. Clin. Lab. Invest., Suppl.*, 55 (1995) 57-59; *C.A.*, 123 (1995) 51376t.
- 3013 Tessmer, U., Quack, T., Donn, F., Leuner, A. and Dernick, R.: Biological activity of prostate-specific antigen isolated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and electroelution. *Electrophoresis (Weinheim)*, 16 (1995) 793-799.
- See also 2815, 2915, 3026.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 3014 Amess, B. and Tolkovsky, A.M.: Programmed cell death in sympathetic neurons: a study by two-dimensional polyacrylamide gel electrophoresis using computer image analysis. *Electrophoresis (Weinheim)*, 16 (1995) 1255-1267.
- 3015 Bresgen, M., Baum, U., Esser, P., Wiedemann, P. and Heimann, K.: (Protein profile of proliferative diabetic retinopathy vitreus. A 2-D electrophoresis). *Ophthalmologe*, 91 (1994) 758-762; *C.A.*, 122 (1995) 262942s.
- 3016 Edwards, R.A., Herrera-Sosa, H., Otto, J. and Bryan, J.: Cloning and expression of a murine fascin homolog from mouse brain. *J. Biol. Chem.*, 270 (1995) 10764-10770.
- 3017 Garcia-Gil, M., Bottai, D., Romano, A., Fineschi, L., Bini, L., Pallini, V. and Brunelli, M.: Repetitive treatment with serotonin modifies protein synthesis and protein phosphorylation in the central nervous system of *Hirudo medicinalis*. *Electrophoresis (Weinheim)*, 16 (1995) 1251-1254.
- 3018 Mattila, K.M. and Frey, H.: Two-dimensional analysis of qualitative and quantitative changes in blood cell proteins in Alzheimer's disease: search for extraneuronal markers. *Appl. Theor. Electrophor.*, 4 (1995) 189-196; *C.A.*, 122 (1995) 310124a.
- See also 2732, 2979.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 3019 Franzén, B., Hirano, T., Okuzawa, K., Uryu, K., Alaiya, A.A., Linder, S. and Auer, G.: Sample preparation of human tumors prior to two-dimensional electrophoresis of proteins. *Electrophoresis (Weinheim)*, 16 (1995) 1087-1089.
- 3020 Lafon, C., Mazars, P., Guerrin, M., Barboule, N., Charcosset, J.-Y. and Valette, A.: Early gene responses associated with transforming growth factor- β 1 growth inhibition and autoinduction in MCF-7 breast adenocarcinoma cells. *Biochim. Biophys. Acta*, 1266 (1995) 288-295.
- 3021 Marton, A., Marko, B., Delbecchi, L. and Bourgaux, P.: Topoisomerase activity associated with polyoma virus large tumor antigen. *Biochim. Biophys. Acta*, 1262 (1995) 59-63.
- 3022 Nawata, S., Tsunaga, N., Numa, F., Tanaka, T., Nakamura, K. and Kato, H.: Serine protease inhibitor activity of recombinant squamous cell carcinoma antigen towards chymotrypsin, as demonstrated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1027-1030.
- 3023 Saijo, S., Watanabe, N. and Kobayashi, Y.: Comparison of reactivity of monoclonal antibody (3F2) to trimeric tumor necrosis factor (TNF- α) with that to monomeric TNF- α . *J. Biochem. (Tokyo)*, 118 (1995) 28-32.
- See also 3008, 3042, 3101.
- 19l. *Specific binding and receptor proteins*
- 3024 Aizawa, H., Sutoh, K., Tsubuki, S., Kawashima, S., Ishii, A. and Yahara, I.: Identification, characterization, and intracellular distribution of cofilin in *Dictyostelium discoideum*. *J. Biol. Chem.*, 270 (1995) 10923-10932.
- 3025 Bose, S., Seetharam, S. and Seetharam, B.: Membrane expression and interactions of human transcobalamin II receptor. *J. Biol. Chem.*, 270 (1995) 8152-8157.
- 3026 Fukui, H., Mizuguchi, H., Liu, Y.Q., Wang, N.P., Hayashi, H., Kangawa, K., Wakamiya, T., Leurs, R., Shiba, T. and Matsuo, H.: Purification and characterization of [3 H]mepyramine (histamine H₁ antagonist)-binding protein from rat liver: a highly homologous protein with cytochrome P450 2D. *J. Biochem. (Tokyo)*, 117 (1995) 993-998.
- 3027 Guignard, F., Muel, J. and Markert, M.: Identification and characterization of a novel human neutrophil protein related to the S100 family. *Biochem. J.*, 309 (1995) 395-401.
- 3028 Harumiya, S., Omori, A., Sugiura, T., Fukumoto, Y., Tachikawa, H. and Fujimoto, D.: EBP-37, a new elastin-binding protein in human plasma: structural similarity to ficolins, transforming growth factor- β 1-binding. *J. Biochem. (Tokyo)*, 117 (1995) 1029-1035.
- 3029 Koch, M., Wahl, R. and Seif, F.J.: Analysis of thyroid stimulating hormone-receptor mutations by temperature-gradient gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 742-743.
- 3030 Lefebvre, P., Gaub, M.-P., Tahayato, A., Rochette-Egly, C. and Formstecher, P.: Protein phosphatases 1 and 2A regulate the transcriptional and DNA binding activities of retinoic acid receptors. *J. Biol. Chem.*, 270 (1995) 10806-10816.
- 3031 Magdolen, V., Rettenberger, P., Lopens, A., Oi, H., Lottspeich, F., Kellermann, J., Creutzburg, S., Goretzki, L., Weidle, U.H., Wilhelm, O., Schmitt, M. and Graeff, H.: Expression of the human urokinase-type plasminogen activator receptor in *E. coli* and Chinese hamster ovary cells: purification of the recombinant proteins and generation of polyclonal antibodies in chicken. *Electrophoresis (Weinheim)*, 16 (1995) 813-816.

- 3032 Morella, K.K., Lai, C., Kumaki, S., Kumaki, N., Wang, Y., Bluman, E.M., Witthuhn, B.A., Ihle, J.N., Giri, J., Gearing, D.P., Cosman, D., Ziegler, S.F. et al.: The action of interleukin-2 receptor subunits defines a new type of signaling mechanism for hematopoietin receptors in hepatic cells and fibroblasts. *J. Biol. Chem.*, 270 (1995) 8298-8310.
- 3033 Offermans, M.T.C., Sonneveld, R.D., Bakker, E., Deutz-Terlouw, P.P., de Geus, B. and Rozing, J.: Denaturing and non-denaturing gel electrophoresis as methods for the detection of junctional diversity in rearranged T cell receptor sequences. *J. Immunol. Methods*, 181 (1995) 101-114; *C.A.*, 122 (1995) 288316r.
- 3034 Suetake, I., Tajima, S. and Asano, A.: A novel DNA binding protein that recognizes the methylated c-Myc binding motif. *J. Biochem. (Tokyo)*, 118 (1995) 244-249.
- 3035 Wang, F., Kan, M., Xu, J., Yan, G. and McKeegan, W.L.: Ligand-specific structural domains in the fibroblast growth factor receptor. *J. Biol. Chem.*, 270 (1995) 10222-10230.
- 3036 Wang, F., Kan, M., Yan, G., Xu, J. and McKeegan, W.L.: Alternately spliced NH₂-terminal immunoglobulin-like loop I in the ectodomain of the fibroblast growth factor (FGF) receptor 1 lowers affinity for both heparin and FGF-1. *J. Biol. Chem.*, 270 (1995) 10231-10235.
- 3037 Yang, B., Yang, B.L. and Goetinck, P.F.: Biotinylated hyaluronic acid as a probe for identifying hyaluronic acid-binding proteins. *Anal. Biochem.*, 228 (1995) 299-306.
- See also 2859, 2933, 3119, 3121, 3122.
- 19m. *Urinary proteins*
- 3038 Kuroda, M.: (Low molecular-weight proteins in urine). *Kensa to Gijutsu*, 22 (1994) 1041-1048; *C.A.*, 122 (1995) 260084q - a review with 9 refs.
- 3039 Shihabi, Z.K.: Myoglobinuria detection by capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 53-58.
- See also 2817.
- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 3040 Grimshaw, J., Trocha-Grimshaw, J., Fisher, W., Kane, A., McCarron, P., Rice, A., Spedding, P.L., Duffy, J. and Mollan, R.A.B.: An investigation of human synovial fluid by capillary electrophoresis. *J. Capillary Electrophor.*, 2 (1995) 40-45; *C.A.*, 122 (1995) 310096t.
- 3041 Lindahl, M., Ståhlborn, B. and Tagesson, C.: Two-dimensional gel electrophoresis of nasal and bronchoalveolar lavage fluids after occupational exposure. *Electrophoresis (Weinheim)*, 16 (1995) 1199-1204.
- 3042 Reid, G.E., Ji, H., Eddes, J.S., Moritz, R.L. and Simpson, R.J.: Nonreducing two-dimensional polyacrylamide gel electrophoretic analysis of human colonic proteins. *Electrophoresis (Weinheim)*, 16 (1995) 1120-1130.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 3043 Wingren, C. and Hansson, U.-B.: A new approach to examine conformational changes occurring upon binding of ligand by biomolecules. *Biochim. Biophys. Acta*, 1244 (1995) 209-215.
- See also 2671.
- 20a. *Oxidoreductases*
- 3044 Boerman, M.H.E.M. and Napoli, J.L.: Characterization of a microsomal retinol dehydrogenase: a short-chain alcohol dehydrogenase with integral and peripheral membrane forms that interacts with holo-CRBP (type I). *Biochemistry*, 34 (1995) 7027-7037.
- 3045 Bonfils, C., Charasse, S., Bonfils, J.-P. and Larroque, C.: Luminescent visualization of low amounts of cytochrome P-450 and hemoproteins by luminal in acrylamide gels. *Anal. Biochem.*, 226 (1995) 302-306.
- 3046 Choi, K.-P., Molnár, I., Yamashita, M. and Murooka, Y.: Purification and characterization of the 3-ketosteroid- Δ^1 -dehydrogenase of *Arthrobacter simplex* produced in *Streptomyces lividans*. *J. Biochem. (Tokyo)*, 117 (1995) 1043-1049.
- 3047 Laeng, P., Bouillon, P., Taupenot, L. and Labourdette, G.: Long-term induction of an aldose reductase protein by basic fibroblast growth factor in rat astrocytes *in vitro*. *Electrophoresis (Weinheim)*, 16 (1995) 1240-1250.
- 3048 Menaya, J., González-Manchón, C., Parrilla, R. and Ayuso, M.S.: Molecular cloning, sequencing and expression of a cDNA encoding a human liver NAD-dependent α -glycerol-3-phosphate dehydrogenase. *Biochim. Biophys. Acta*, 1262 (1995) 91-94.
- 3049 Ono, K., Kawanaka, Y., Izumi, Y., Inui, H., Miyatake, K., Kitaoka, S. and Nakano, Y.: Mitochondrial alcohol dehydrogenase from ethanol-grown *Euglena gracilis*. *J. Biochem. (Tokyo)*, 117 (1995) 1178-1182.
- 3050 Rescigno, A., Sollai, F., Masala, S., Porcu, M.C., Sanjust, E., Rinaldi, A.C., Curreli, N., Grifi, D. and Rinaldi, A.: Purification and characterization of an NAD(P)H:quinone oxidoreductase from *Glycine max* seedlings. *Prepar. Biochem.*, 25 (1995) 57-67.
- 3051 Serebryakova, L.T., Troshina, O.Yu. and Gogotov, I.N.: (Purification and forms of reversible hydrogenase from *Anabaena variabilis* ATCC 29413). *Biokhimiya (Moscow)*, 60 (1995) 508-514.
- 3052 Tordjman, C., Coge, F., Andre, N., Rique, H., Spedding, M. and Bonnet, J.: Characterisation of cyclooxygenase 1 and 2 expression in mouse resident peritoneal macrophages *in vitro*: interactions of non steroidal anti-inflammatory drugs with COX2. *Biochim. Biophys. Acta*, 1256 (1995) 249-256.
- 3053 Xia, J. and Lindahl, P.A.: Decomposition of carbon monoxide dehydrogenase into α metallosubunits and a catalytically-active form consisting primarily of β metallosubunits. *Biochemistry*, 34 (1995) 6037-6042.
- 3054 Zorin, N.A., Pashkova, O.N. and Gogotov, I.N.: (Isolation and characterization of two forms of hydrogenase from *Thiocapsa roseopersicina*). *Biokhimiya (Moscow)*, 60 (1995) 515-522.
- See also 3097.

- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 3055 Goppelt-Struebe, M.: Two step purification of human and murine leukotriene C_4 synthase. *Biochim. Biophys. Acta*, 1256 (1995) 257-261.
- 3056 Hashimoto, W., Suzuki, H., Yamamoto, K. and Kumagai, H.: Effect of site-directed mutations on processing and activity of γ -glutamyltranspeptidase of *Escherichia coli* K-12. *J. Biochem. (Tokyo)*, 118 (1995) 75-80.
- See also 3167.
- 20c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 3057 Babiychuk, E.B., Babiychuk, V.S. and Sobieszek, A.: Modulation of smooth muscle myosin light chain kinase activity by Ca^{2+} /calmodulin-dependent, oligomeric-type modifications. *Biochemistry*, 34 (1995) 6366-6372.
- 3058 Eto, A., Shiki, A., Chikaura, Y., Oka, T. and Nakano, N.I.: Multienzyme control serum (Seraclear-HE) containing human enzymes from established cell lines and other sources. I: Preparation and properties. *Clin. Chem. (Washington)*, 41 (1995) 872-880.
- 3059 Hong, D.-h., Huan, J., Ou, B.-r., Yeh, J.-y., Saido, T.C., Cheeke, P.R. and Forsberg, N.E.: Protein kinase C isoforms in muscle cells and their regulation by phorbol ester and calpain. *Biochim. Biophys. Acta*, 1267 (1995) 45-54.
- 3060 Maeda, H., Inazu, T., Nagai, K., Maruyama, S., Nakagawara, G. and Yamamura, H.: Possible involvement of protein-tyrosine kinases such as p72^{src} in the disc-sphere change response of porcine platelets. *J. Biochem. (Tokyo)*, 117 (1995) 1201-1208.
- 3061 Okajima, T., Goto, S., Tamizawa, K., Tagaya, M., Fukui, T., Shimofuruya, H. and Suzuki, J.: Cloning, sequencing, and expression in *Escherichia coli* of cDNA encoding porcine brain UMP-CMP kinase. *J. Biochem. (Tokyo)*, 117 (1995) 980-986.
- 20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*
- 3062 Abdullah, K., Cromlish, W.A., Payette, P., Laliberté, F., Huang, Z., Street, I. and Kennedy, B.P.: Human cytosolic phospholipase A₂ expressed in insect cells is extensively phosphorylated on Ser-505. *Biochim. Biophys. Acta*, 1244 (1995) 157-164.
- 3063 Bolognese, B., McCord, M. and Marshall, L.A.: Differential regulation of elicited-peritoneal macrophage 14 kDa and 85 kDa phospholipase A₂(s) by transforming growth factor- β . *Biochim. Biophys. Acta*, 1256 (1995) 201-209.
- 3064 Hoshino, T., Kumasaka, K., Kawano, K., Yamagishi, F., Koyama, I., Fujimori-Arai, Y., Nakajima, T. and Komoda, T.: Low serum alkaline phosphatase activity associated with severe Wilson's disease. Is the breakdown of alkaline phosphatase molecules caused by reactive oxygen species? *Clin. Chim. Acta*, 238 (1995) 91-100.
- 3065 Kálmán, F., Ma, S., Fox, R.O. and Horváth, Cs.: Capillary electrophoresis of *S. nuclease* mutants. *J. Chromatogr. A*, 705 (1995) 135-154.
- 3066 Kostka, H., Steinke, M., Gehrisch, S. and Jarosa, W.: (TGGE) A screening method for detecting point mutations in the lipoprotein lipase gene (LPL gene). *Klin. Labor*, 41 (1995) 37-41; C.A., 123 (1995) 26507d.
- 3067 Okano, K. and Kambara, H.: DNA probe assay based on exonuclease III digestion of probes hybridized on target DNA. *Anal. Biochem.*, 228 (1995) 101-108.
- 3068 Otto, V.I., Fried, R., Wiederkehr, F. and Hänsele, E.: Separation of the two most closely related isoenzymes of alkaline phosphatase by two-dimensional electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1284-1288.
- See also 3058.
- 20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*
- 3069 Chinnathambi, S., Lachke, A. and Radhakrishnan, S.: Purification of α -L-arabinofuranosidase using a single-column mini-scale isoelectric focusing unit. *J. Chromatogr. A*, 705 (1995) 400-402.
- 3070 Craig, D., Arriaga, E.A., Banks, P., Zhang, Y., Renborg, A., Palcic, M.M. and Dovichi, N.J.: Fluorescence-based enzymatic assay by capillary electrophoresis laser-induced fluorescence detection for the determination of a few β -galactosidase molecules. *Anal. Biochem.*, 226 (1995) 147-157.
- 3071 Hamada, J.S.: High performance liquid chromatography of *Bacillus circulans* peptidoglutaminase for laboratory and industrial uses. *J. Chromatogr. A*, 702 (1995) 163-172.
- 3072 Kalix, S. and Buchenauer, H.: Direct detection of β -1,3-glucanase in plant extracts by polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1016-1018.
- 3073 Menteur, S., Jestin, L., Risacher, T. and Branlard, G.: Visualization of barley β -glucan degrading isozymes after gel isoelectric focusing. *Electrophoresis (Weinheim)*, 16 (1995) 1019-1023.
- 3074 Nishimura, Y., Kato, K. and Himeno, M.: Biochemical characterization of liver microsomal, Golgi, lysosomal, and serum β -glucuronidases in dibutyl phosphate-treated rats. *J. Biochem. (Tokyo)*, 118 (1995) 56-66.
- 3075 Nishimura, Y., Kato, K., Oda, K., Ishikawa, T., Ikehara, Y. and Himeno, M.: Intracellular sorting of lysosomal β -glucuronidase is altered due to administration of dibutyl phosphate. *J. Biochem. (Tokyo)*, 118 (1995) 46-55.
- 3076 Vandersall, A.S., Cameron, R.G., Nairn III, C.J., Yelonosky, G. and Wodzinski, R.J.: Identification, characterization, and partial purification of glucanase from *Aspergillus niger* (syn. *A. ficuum*) NRRL 3135. *Prepar. Biochem.*, 25 (1995) 29-55.
- See also 2709, 2808.
- 20f. *Other hydrolases*
- 3077 Banerjee, S., Kaplan, H., Yagodnik, C., Breuil, C. and Brown, D.L.: A method for detecting carboxypeptidase activity in the sapstaining fungus, *Ophiostoma piceae*, growing in artificial medium or in wood. *Biotechnol. Tech.*, 9 (1995) 241-246; C.A., 123 (1995) 28286e
- 3078 Bernardo, K., Hurwitz, R., Zenk, T., Desnick, R.J., Ferlinz, K., Schuchman, E.H. and Sandhoff, K.: Purification, characterization, and biosynthesis of human acid ceramidase. *J. Biol. Chem.*, 270 (1995) 11098-11102.

- 3079 Ebisui, C., Tsujinaka, T., Morimoto, T., Fujita, J., Ogawa, A., Ishidoh, K., Kominami, E., Tanaka, K. and Monden, M.: Changes of proteasomes and cathepsins activities and their expression during differentiation of C2C12 myoblasts. *J. Biochem. (Tokyo)*, 117 (1995) 1088-1094.
- 3080 Fridman, R., Toth, M., Pena, D. and Mobashery, S.: Activation of progelatinase B (MMP-9) by gelatinase A (MMP-2). *Cancer Res.*, 55 (1995) 2548-2555.
- 3081 Haginaka, J., Miyano, Y., Saizen, Y., Seyama, C. and Murashima, T.: Separation of enantiomers on a pepsin-bonded column. *J. Chromatogr. A*, 708 (1995) 161-168.
- 3082 Kido, T., Yook, H.-Y. and Ueda, K.: Ligand western blotting for specific detection of active forms of proteases. *Clin. Chim. Acta*, 237 (1995) 31-41.
- 3083 Kitada, S., Shimokata, K., Niidome, T., Ogishima, T. and Ito, A.: A putative metal-binding site in the β subunit of rat mitochondrial processing peptidase is essential for its catalytic activity. *J. Biochem. (Tokyo)*, 117 (1995) 1148-1150.
- 3084 Lafont, P., Guy-Crotte, O., Paulin, C., Galvain, D., Mertani, S., Figarella, C. and Barbier, Y.: A specific immunoradiometric assay of cationic trypsin(ogen) that does not recognize trypsin-alpha-1-proteinase inhibitor complex. *Clin. Chim. Acta*, 235 (1995) 197-206.
- 3085 Raser, K.J., Posner, A. and Wang, K.K.W.: Casein zymography: a method to study μ -calpain, m-calpain, and their inhibitory agents. *Arch. Biochem. Biophys.*, 319 (1995) 211-216.
- 3086 Tyagarajan, K., Chow, D.C., Smolka, A. and Forte, J.G.: Structural interactions between α - and β -subunits of the gastric H,K-ATPase. *Biochim. Biophys. Acta*, 1236 (1995) 105-113.
- 3087 Uchikoba, T., Yonezawa, H. and Kaneda, M.: Cleavage specificity of cucumis, a plant serine protease. *J. Biochem. (Tokyo)*, 117 (1995) 1126-1130.
- 3088 Váradi, A., Molnár, E. and Ashcroft, S.J.H.: Characterisation of endoplasmic reticulum and plasma membrane Ca^{2+} -ATPases in pancreatic β -cells and in islets of Langerhans. *Biochim. Biophys. Acta*, 1236 (1995) 119-127.
- 3089 Yamauchi, Y., Sugimoto, T. and Oji, Y.: A method for the separation and detection of proteases using substrate-containing gel isoelectric focusing electrophoresis. *Biosci., Biotechnol., Biochem.*, 59 (1995) 767-768; *C.A.*, 122 (1995) 309361u.
- 3090 Yoshimoto, Y., Shibukawa, A., Sasagawa, H., Nitta, S. and Nakagawa, T.: Michaelis-Menten analysis of immobilized enzyme by affinity capillary electrophoresis. *J. Pharm. Biomed. Anal.*, 13 (1995) 483-488.
- 3091 Zhang, N. and Jones, B.L.: Characterization of germinated barley endoproteolytic enzymes by two-dimensional gel electrophoresis. *J. Cereal Sci.*, 21 (1995) 145-153; *C.A.*, 123 (1995) 52257s.
- See also 2870, 2871.
- 20g. *Lyases*
- 3092 Kojima, H., Chiba, J., Watanabe, Y. and Numata, O.: Citrate synthase purified from *Tetrahymena* mitochondria is identical with *Tetrahymena* 14-nm filament protein. *J. Biochem. (Tokyo)*, 118 (1995) 189-195.
- 3093 Salerno, C. and Crifo, C.: Microassay of adenylysuccinate by capillary electrophoresis. *Anal. Biochem.*, 226 (1995) 377-379.
- 3094 Zhang, R., Kai, T., Sugimoto, Y., Kusakabe, T., Takasaki, Y., Koga, K. and Hori, K.: *Drosophila melanogaster* aldolase: characterization of the isozymes α , β , and γ generated from a single gene. *J. Biochem. (Tokyo)*, 118 (1995) 183-188.
- See also 2927.
- 20h. *Isomerases*
- 3095 Uchida, K., Kondoh, K. and Matuo, Y.: Recombinant M-, B- and MB-type isozymes of human phosphoglyceric acid mutase: their large-scale production and preparation of polyclonal antibodies specific to M- and B-type isozymes. *Clin. Chim. Acta*, 237 (1995) 43-58.
- 20j. *Complex mixtures and incompletely identified enzymes*
- 3096 Günemann-Schäfer, K. and Kindl, H.: Fatty acid β -oxidation in glyoxysomes. Characterization of a new tetrafunctional protein (MFP III). *Biochim. Biophys. Acta*, 1256 (1995) 181-186.
- 3097 Iwakura, M. and Kokubu, T.: Production of hetero-dimeric dihydrofolate reductase-thymidylate synthase bifunctional enzyme. *J. Biochem. (Tokyo)*, 118 (1995) 67-74.
21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 3098 Cunningham, M.L., Spencer, D.L., Caspary, W.J., Stopper, H. and Phillips, A.: Determination of 5-methylcytosine content in DNA from cells in culture by micellar electrokinetic capillary chromatography. *J. Capillary Electrophor.*, 2 (1995) 30-33; *C.A.*, 122 (1995) 309992a.
- 3099 Dawson, J.R., Nichols, S.C. and Taylor, G.E.: Determination of impurities in a novel analogue of adenosine 5'-triphosphate by capillary electrophoresis. *J. Chromatogr. A*, 700 (1995) 163-172.
- 3100 Gu, J., Liu, Y., Fu, R., Lu, H. and Zhang, Y.: Separation of nucleotides in swine tissues using capillary zone electrophoresis. *J. Beijing Inst. Technol. (Engl. ed.)*, 3 (1994) 155-161; *C.A.*, 123 (1995) 4936r.
- 3101 Lin, S.-B., Huang, S.-S., Choo, K.-B., Chen, P.-J. and Au, L.-C.: Inhibition of α -fetoprotein production in a hepatoma cell line by antisense oligonucleotide analogues. *J. Biochem. (Tokyo)*, 117 (1995) 1100-1104.
- 3102 Rabel, S.R.: Development of analytical methodology utilizing capillary electrophoresis with laser-induced fluorescence detection for the determination of intracellular levels of 6-mercaptopurine metabolites. Avail. *Univ. Microfilms Int.*, Order No. DA9504051, 1994, 174 pp.; *C.A.*, 122 (1995) 281256n.
- 3103 Shihabi, Z.K., Hinsdale, M.E. and Bleyer, A.J.: Xanthine analysis in biological fluids by capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 163-169.
- 3104 Zhang, R., Lu, Z., Zhang, X., Zhao, H., Diasio, R.B., Liu, T., Jiang, Z. and Agrawal, S.: *In vivo* stability and disposition of a self-stabilized oligodeoxynucleotide phosphorothioate in rats. *Clin. Chem. (Washington)*, 41 (1995) 836-843.

See also 2729, 3093, 3173.

21b. Nucleic acids, RNA

- 3105 Adati, N., Ito, T., Koga, C., Kito, K., Sakaki, Y. and Shiohawa, K.: Differential display analysis of gene expression in developing embryos of *Xenopus laevis*. *Biochim. Biophys. Acta*, 1262 (1995) 43-51.
- 3106 DeDionisio, L. and Gryaznov, S.M.: Analysis of a ribonuclease H digestion of N3'→P5' phosphoramidate-RNA duplexes by capillary gel electrophoresis. *J. Chromatogr. B*, 669 (1995) 125-131.
- 3107 Engelmann, S., Ebeling, O. and Schwartz-Albiez, R.: Modulated glycosylation of proteoglycans during differentiation of human B lymphocytes. *Biochim. Biophys. Acta*, 1267 (1995) 6-14.
- 3108 Frigerio, J.-M., Dagorn, J.-C. and Iovanna, J.L.: Cloning, sequencing and expression of the L5, L21, L27a, L28, S5, S9, S10 and S29 human ribosomal protein mRNAs. *Biochim. Biophys. Acta*, 1262 (1995) 64-68.
- 3109 Funai, T. and Ichiyama, A.: Fidelity of translation initiation of mRNA for the precursor of rat mitochondrial serine:pyruvate/alanine:glyoxylate aminotransferase. *J. Biochem. (Tokyo)*, 117 (1995) 1008-1016.
- 3110 Izumi, T., Tajima, S. and Nishikawa, T.: Differential expression of α_1 and α_2 chains of type VI collagen in the upper, middle, and lower dermal fibroblasts *in vitro*. *J. Biochem. (Tokyo)*, 117 (1995) 1004-1007.
- 3111 Kaisakia, P.J., Jerkins, A.A., Goodspeed, D.C. and Steele, R.D.: Cloning and characterization of rat cysteine sulfinic acid decarboxylase. *Biochim. Biophys. Acta*, 1262 (1995) 79-82.
- 3112 Katsivela, E. and Höfle, M.G.: Separation of transfer RNA and 5S ribosomal RNA using capillary electrophoresis. *J. Chromatogr. A*, 700 (1995) 125-136.
- 3113 Kühl, M. and Wedlich, D.: XBU-Cadherin mRNA contains cytoplasmic polyadenylation elements and is polyadenylated during oocyte maturation in *Xenopus laevis*. *Biochim. Biophys. Acta*, 1262 (1995) 95-98.
- 3114 Lübber, B., Fabrizio, P., Kastner, B. and Lührmann, R.: Isolation and characterization of the small nucleolar ribonucleoprotein particle snR30 from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 270 (1995) 11549-11554.
- 3115 Mador, N., Panet, A., Latchman, D. and Steiner, I.: Expression and splicing of the latency-associated transcripts of herpes simplex virus type 1 in neuronal and non-neuronal cell lines. *J. Biochem. (Tokyo)*, 117 (1995) 1288-1297.
- 3116 Mukainaka, Y., Tanaka, K., Hagiwara, T. and Wada, K.: Molecular cloning of two glutamate transporter subtypes from mouse brain. *Biochim. Biophys. Acta*, 1244 (1995) 233-237.
- 3117 Nelson, B.D., Luciakova, K., Li, R. and Betina, S.: The role of thyroid hormone and promoter diversity in the regulation of nuclear encoded mitochondrial proteins. *Biochim. Biophys. Acta*, 1271 (1995) 85-91.
- 3118 Nikawa, J.-i., Murakami, A., Esumi, E. and Hosaka, K.: Cloning and sequence of the SCS2 gene, which can suppress the defect of INO1 expression in an inositol auxotrophic mutant of *Saccharomyces cerevisiae*. *J. Biochem. (Tokyo)*, 118 (1995) 39-45.
- 3119 Obama, H., Tsutsui, J.-i., Ozawa, M., Yoshida, H., Yoshida, Y., Osame, M. and Muramatsu, T.: Midkine (MK) expression in extraembryonic tissues, amniotic fluid, and cerebrospinal fluid during mouse embryogenesis. *J. Biochem. (Tokyo)*, 118 (1995) 88-93.
- 3120 Ohashi, K., Honda, S., Ichinomiya, N., Nakamura, T. and Mizuno, K.: Molecular cloning and *in situ* localization in the brain of rat sky receptor tyrosine kinase. *J. Biochem. (Tokyo)*, 117 (1995) 1267-1275.
- 3121 Ozawa, M.: Cloning of a human homologue of mouse reticulocalbin reveals conservation of structural domains in the novel endoplasmic reticulum resident Ca²⁺-binding protein with multiple EF-hand motifs. *J. Biochem. (Tokyo)*, 117 (1995) 1113-1119.
- 3122 Suzuki, E., Kojima, N., Yoshimura, K., Uyemura, K., Obata, K. and Akagawa, K.: Cloning and sequence analysis of cDNA for a possible DNA-binding protein SE5 in the nervous system. *J. Biochem. (Tokyo)*, 118 (1995) 122-128.
- 3123 Zhang, Y., Wick, D.A., Haas, A.L., Seetharam, B. and Dahms, N.M.: Regulation of lysosomal and ubiquitin degradative pathways in differentiating human intestinal Caco-2 cells. *Biochim. Biophys. Acta*, 1267 (1995) 15-24.
- See also 2674, 2726, 2741, 2810, 2906, 2925, 3004, 3132, 3141, 3167, 3192, 3194, 3200.

21c. Nucleic acids, DNA

- 3124 Barton, B.M., Harding, G.P. and Zuccarelli, A.J.: A general method for detecting and sizing large plasmids. *Anal. Biochem.*, 226 (1995) 235-240.
- 3125 Belenky, A., Smisek, D.L. and Cohen, A.S.: Sequencing of antisense DNA analogues by capillary gel electrophoresis with laser-induced fluorescence detection. *J. Chromatogr. A*, 700 (1995) 137-149.
- 3126 Calvert, R.J., Weghorst, C.M. and Buzard, G.S.: PCR amplification of silver-stained SSCP bands from cold SSCP gels. *BioTechniques*, 18 (1995) 782-786; C.A., 123 (1995) 2127d.
- 3127 Cedervall, B., Kaellman, P. and Dewey, W.C.: Repair of DNA double-strand breaks: errors encountered in the determination of half-life times in pulsed-field gel electrophoresis and neutral filter elution. *Radiat. Res.*, 142 (1995) 23-28; C.A., 122 (1995) 259897a.
- 3128 Chang, H.-T. and Yeung, E.S.: Poly(ethyleneoxide) for high-resolution and high-speed separation of DNA by capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 113-123.
- 3129 Cheng, J., Kasuga, T., Watson, N.D. and Mitchelson, K.R.: Enhanced single-stranded DNA conformation polymorphism analysis by entangled solution capillary electrophoresis. *J. Capillary Electrophor.*, 2 (1995) 24-29; C.A., 122 (1995) 310095s.
- 3130 Chrambach, A., Yarmola, E., Zakharov, S.F. and Garner, M.M.: Commercial automated gel electrophoresis apparatus: application to DNA, band dispersion, nonlinear Ferguson curves, and isolation. *Electrophoresis (Weinheim)*, 16 (1995) 713-718.
- 3131 Courtney, B.C., Williams, K.C., Bing, Q.A. and Schlager, J.J.: Capillary gel electrophoresis as a method to determine ligation efficiency. *Anal. Biochem.*, 228 (1995) 281-286.
- 3132 Dumais, M., Kusukawa, N. and White, H.: Highly derivatized agarose conformational nucleic acid separation. *PCT Int. Appl. WO 95 11,993 (Cl. C12Q1/68)*, 04 May 1995, US Appl. 146,286, 29 Oct. 1993; 40 p p.; C.A., 123 (1995) 51705z.
- 3133 Eid, M.M.A. and Sherwood, J.E.: Pulsed-field gel electrophoresis fingerprinting for identification of *Azospirillum* species. *Curr. Microbiol.*, 30 (1995) 127-131; C.A., 122 (1995) 283331g.

- 3134 Epplen, J.T., Buitkamp, J., Epplen, C., Mäueler, W. and Riess, O.: Indirect DNA gene diagnoses via electrophoresis - an obsolete principle? *Electrophoresis (Weinheim)*, 16 (1995) 683-690.
- 3135 Gelfi, C., Righetti, P.G., Leoncini, F., Brunelli, V., Carrera, P. and Ferrari, M.: CAG triplet analysis in families with androgen insensitivity syndrome by capillary electrophoresis in polymer networks. *J. Chromatogr. A*, 706 (1995) 463-468.
- 3136 Glisic, S., Sunjevaric, I. and Alavantic, D.: Genotyping apolipoprotein B signal peptide insertion/deletion: a comparison of three methods. *Electrophoresis (Weinheim)*, 16 (1995) 899-902.
- 3137 Gurrieri, S.: Molecular mechanisms of DNA motion during agarose gel electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9509786, 1994, 261 pp.; C.A., 123 (1995) 27955k.
- 3138 Heller, C. and Viovy, J.L.: Electrophoretic mobility of DNA in gels. III. Experimental study on band inversion. *Biopolymers*, 35 (1995) 485-492; C.A., 122 (1995) 260350y.
- 3139 Higuchi, Y. and Linn, S.: Purification of all forms of HeLa cell mitochondrial DNA and assessment of damage to it caused by hydrogen peroxide treatment of mitochondria or cells. *J. Biol. Chem.*, 270 (1995) 7950-7956.
- 3140 Hilbert, F., Mayr, B., Lackner, F. and Bauer, F.: The influence of proteins on silver staining of nucleic acids following polyacrylamide gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 903-904.
- 3141 Ide, H., Itoh, H., Tomita, M., Murakumo, Y., Kobayashi, T., Maruyama, H., Osada, Y. and Nawa, Y.: cDNA sequencing and expression of rat mast cell tryptase. *J. Biochem. (Tokyo)*, 118 (1995) 210-215.
- 3142 Johnson, J.M., Weagant, S.D., Jinneman, K.C. and Bryant, J.L.: Use of pulsed-field gel electrophoresis for epidemiological study of *Escherichia coli* O157:H7 during a food-borne outbreak. *Appl. Environ. Microbiol.*, 61 (1995) 2806-2808; C.A., 123 (1995) 51527t.
- 3143 Kadokami, Y. and Lewis, R.V.: Reverse electrophoresis to concentrate DNA fractions. *Anal. Biochem.*, 226 (1995) 193-195.
- 3144 Kikuchi, H., Sugiyama, S., Imajoh-Ohmi, S.: A novel anti-apoptosis serum factor that down-regulates Fas-mediated apoptosis. *J. Biochem. (Tokyo)*, 117 (1995) 936-939.
- 3145 Kim, S., Yoo, H.J., Nam, H.-G. and Hahn, J.H.: Fast capillary-scanning system for detecting fluorescently labelled DNA sequencing fragments separated by capillary gel electrophoresis. *Chromatographia*, 40 (1995) 345-349.
- 3146 Kolchinsky, A. and Funke, R.: Pulsed field gel electrophoresis. *Curr. Innovations Mol. Biol.*, 1 (1995) 69-82; C.A., 123 (1995) 26495y.
- 3147 Kozulic, B.: Gel electrophoresis of DNA and proteins: recent advances in theory and practical applications. *Curr. Innovations Mol. Biol.*, 1 (1995) 51-68; C.A., 123 (1995) 28853n.
- 3148 Larsson, A. and Aakerman, B.: Period times and helix alignment during the cyclic migration of DNA in electrophoresis gels studied with fluorescence microscopy. *Macromolecules*, 28 (1995) 4441-4454; C.A., 122 (1995) 310028x.
- 3149 Lueck, P.C., Helbig, J.H., Hagedorn, H.-J. and Ehret, W.: DNA fingerprinting by pulsed-field gel electrophoresis to investigate a nosocomial pneumonia caused by *Legionella bozemanii* serogroup 1. *Environ. Microbiol.*, 61 (1995) 2759-2761; C.A., 123 (1995) 51526s.
- 3150 Nakashima, H., Yi, M., Ichikawa, N., LeBlond, G.F., Antonarakis, S.E. and Ts'o, P.O.P.: Two-dimensional electrophoresis southern transfer method for detecting human genome variability using a LINE-1 sequence probe. *Anal. Biochem.*, 227 (1995) 319-327.
- 3151 Ng, L.-K., Carballo, M. and Dillon, J.-A.R.: Differentiation of *Neisseria gonorrhoeae* isolates requiring proline, citrulline, and uracil by plasmid content, serotyping, and pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 33 (1995) 1039-1041; C.A., 122 (1995) 285994z.
- 3152 Oh, Y., Yoon, J. and Baek, K.: Isolation and characterization of the gene encoding the *Drosophila melanogaster* transcriptional elongation factor, TFIIS. *Biochim. Biophys. Acta*, 1262 (1995) 99-103.
- 3153 Ozawa, M.: Structure of the gene encoding mouse reticulocalbin, a novel endoplasmic reticulum-resident Ca²⁺-binding protein with multiple EF-hand motifs. *J. Biochem. (Tokyo)*, 118 (1995) 154-160.
- 3154 Ray, T., Mills, A. and Dyson, P.: Tris-dependent oxidative DNA strand scission during electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 888-894.
- 3155 Richter, C., Gogvadze, V., Laffranchi, R., Schlapbach, R., Schweizer, M., Suter, M., Walter, P. and Yaffee, M.: Oxidants in mitochondria: from physiology to diseases. *Biochim. Biophys. Acta*, 1271 (1995) 67-74.
- 3156 Roemling, U., Heuer, T. and Tuemmler, B.: Bacterial genome analysis by pulsed field gel electrophoresis techniques. *Adv. Electrophor.*, 7 (1994) 353-406; C.A., 122 (1995) 283183k - a review with 86 refs.
- 3157 Sanguinetti, C.J., Neto, E.D. and Simpson, A.J.G.: Rapid silver staining and recovery of PCR products separated on polyacrylamide gels. *BioTechniques*, 17 (1994) 914-921; C.A., 122 (1995) 283254j.
- 3158 Semenov, A.N. and Subbotin, A.V.: Instability in the dynamics of long DNA undergoing gel electrophoresis in a strong field. *J. Chem. Phys.*, 102 (1995) 7728-7733; C.A., 123 (1995) 27969t.
- 3159 Sestili, P., Cattabeni, F. and Cantoni, O.: Simultaneous determination of DNA double strand breaks and DNA fragment size in cultured mammalian cells exposed to hydrogen peroxide/histidine or etoposide with CHEF electrophoresis. *Carcinogenesis*, 16 (1995) 703-706; C.A., 122 (1995) 308162t.
- 3160 Sharma, M., Jain, R., Ionescu, E. and Slocum, H.K.: Capillary electrophoretic separation and laser-induced fluorescence detection of the major DNA adducts of cisplatin and carboplatin. *Anal. Biochem.*, 228 (1995) 307-311.
- 3161 Shin, S. and Day, L.A.: Separation and size determination of circular and linear single-stranded DNAs by alkaline agarose gel electrophoresis. *Anal. Biochem.*, 226 (1995) 202-206.
- 3162 Sifakakos, N., Burnett, L., Bennetts, B. and Proos, A.: Nonenzymatic extraction of DNA from blood collected into serum separator tubes. *Clin. Chem. (Washington)*, 41 (1995) 1045-1046.
- 3163 Skov, M.N., Pedersen, K. and Larsen, J.L.: Comparison of pulsed-field gel electrophoresis, ribotyping, and plasmid profiling for typing of *Vibrio anguillarum* serovar O1. *Appl. Environ. Microbiol.*, 61 (1995) 1540-1545; C.A., 122 (1995) 260346b.
- 3164 Slater, G.W., Desruisseaux, C., Villeneuve, C., Guo, H.L. and Drouin, G.: Trapping gel electrophoresis of end-labeled DNA: an analytical model for mobility and diffusion. *Electrophoresis (Weinheim)*, 16 (1995) 704-712.

- 3165 Stellwagen, J. and Stellwagen N.C.: Effect of spike pulses on the orientation of the agarose gel matrix. *Electrophoresis (Weinheim)*, 16 (1995) 700-703.
- 3166 Stellwagen, N.C.: Use of polyacrylamide gel electrophoresis to detect structural variations in kilobase-sized DNAs. *Electrophoresis (Weinheim)*, 16 (1995) 691-699.
- 3167 Tajima, S., Tsuda, H., Wakabayashi, N., Asano, A., Mizuno, S. and Nishimori, K.: Isolation and expression of a chicken DNA methyltransferase cDNA. *J. Biochem. (Tokyo)*, 117 (1995) 1050-1057.
- 3168 Tsen, H.-Y., Yu, G.-K. and Lin, I.-T.: Plasmid profiles and pulsed-field gel electrophoresis for type A enterotoxigenic *Staphylococcus aureus* isolated from foods. *J. Food Prot.*, 58 (1995) 147-153; C.A., 122 (1995) 286357f.
- 3169 Wang, J., Delgado, D.A. and Deen, D.F.: Use of pulsed-field gel electrophoresis to investigate factors influencing the measurement of DNA double-strand breaks in human brain tumor specimens. *Int. J. Radiat. Biol.*, 67 (1995) 153-160; C.A., 122 (1995) 285584r.
- 3170 Wang, Z., Liu, Y. and Tian, P.: (An improved method for recovering DNA from agarose gel). *Weishengwuxue Tongbao*, 21 (1994) 120-122; C.A., 122 (1995) 260353b.
- 3171 Wöhl, T., Brecht, M., Lottspeich, F. and Ammer, H.: The use of genomic DNA probes for in-gel hybridization. *Electrophoresis (Weinheim)*, 16 (1995) 739-741.
- See also 2674, 2699, 2726, 2741, 2761, 2915, 3021, 3079, 3105, 3194.
- 21d *Structural studies on RNA and RNA mapping*
- 3172 Fujii, H.: Base sequencing apparatus. U.S. US 5,419,825 (Cl. 204-299R; C25B7/00), 30 May 1995, US Appl. 737,416, 29 Jul. 1991; 19 pp.; C.A., 123 (1995) 49248j.
- 21e *Structural studies on DNA and DNA mapping*
- 3173 Baba, Y., Tomisaki, R. and Tshako, M.: Three-dimensional electropherogram for the separation of oligodeoxynucleotides and DNA restriction fragments using capillary gel electrophoresis with a photodiode array detector. *J. Liq. Chromatogr.*, 18 (1995) 1317-1324.
- 3174 Bory, C., Chantin, C. and Bozon, D.: Capillary electrophoretic analysis of DNA restriction fragments and PCR products for polymorphism and mutation studies in cystic fibrosis and Gaucher's disease. *J. Pharm. Biomed. Anal.*, 13 (1995) 511-514.
- 3175 Brewer, L.R., Davidson, C., Balch, J. and Carrano, A.: Three dimensional imaging of DNA fragments during electrophoresis using a confocal detector. *Proc. SPIE-Int. Soc. Opt. Eng.*, 2386 (1995) 86-96; C.A., 123 (1995) 28760e.
- 3176 Busch, U., Teufel, C.H., Boehmer, R., Wilske, B. and Preac-Mursic, V.: Molecular characterization of *Borrelia burgdorferi sensu lato* strains by pulsed-field gel electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 744-747.
- 3177 Butler, J.M., McCord, B.R., Jung, J.M., Lee, J.A., Budowle, B. and Allen, R.O.: Application of dual internal standards for precise sizing of polymerase chain reaction products using capillary electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 974-980.
- 3178 Debuire, B., Sol, O., Lemoine, A. and May, E.: Nonisotopic competitive RT-PCR assay to measure MDR1 gene expression. *Clin. Chem. (Washington)*, 41 (1995) 819-825.
- 3179 Dürr, C., Hinney, A., Luckenbach, C. and Ritter, H.: Detection of two hypervariable (ATTT)n loci in the human genome. *Electrophoresis (Weinheim)*, 16 (1995) 719-721.
- 3180 Evans, M.D., Wolfe, J.T., Perrett, D., Lunec, J. and Herbert, K.E.: Analysis of internucleosomal DNA fragmentation in apoptotic thymocytes by dynamic sieving capillary electrophoresis. *J. Chromatogr. A*, 700 (1995) 151-162.
- 3181 Feussner, G., Lohrmann, J. and Dörmeyer, J.: Rapid and reliable identification of human apolipoprotein E1 (Gly₁₂₇→Asp, Arg₁₅₈→Cys) variant. *Clin. Chem. (Washington)*, 41 (1995) 1043-1045.
- 3182 Fung, E.N. and Yeung, E.S.: High-speed DNA sequencing by using mixed poly(ethylene oxide) solutions in uncoated capillary columns. *Anal. Chem.*, 67 (1995) 1913-1919.
- 3183 Gelfi, C., Leoncini, F., Righetti, P.G., Cremonesi, L., di Blasio, A.M., Carniti, C. and Vignali, M.: Separation and quantitation of reverse transcriptase polymerase chain reaction fragments of basic fibroblast growth factor by capillary electrophoresis in polymer networks. *Electrophoresis (Weinheim)*, 16 (1995) 780-783.
- 3184 Gombocz, E. and Cortez, E.: Separation, real-time migration monitoring and selective zone retrieval using a computer controlled system for automated analysis. *Appl. Theor. Electrophor.*, 4 (1995) 197-209; C.A., 122 (1995) 310243p.
- 3185 Hernandez, A., Uhrberg, M., Enczmann, J., Witt, I., Reitsma, P.H. and Wernet, P.: Rapid identification of gene defects in protein C deficiency by temperature gradient gel electrophoresis. *Blood Coagulation Fibrinolysis*, 6 (1995) 23-30; C.A., 123 (1995) 48665f.
- 3186 Jarrett, H.W.: Preparation of cDNA-silica using reverse transcriptase and its DNA sequence determination. *J. Chromatogr. A*, 708 (1995) 13-18.
- 3187 Jikuya, H., Aoyama, Y., Nagai, N. and Fujiwake, H.: (Analysis of the parameters for reading of long DNA sequences). *Shimadzu Hyoron*, 51 (1994, Pub. 1995) 281-285; C.A., 122 (1995) 283226b - review with 3 refs.
- 3188 Liang, P.: Analysis of dystrophin gene mutations by pulsed field gel electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9419248, 1993, 147 pp.; C.A., 123 (1995) 2020p.
- 3189 Lu, M., Funsch, B., Wiese, M. and Roggendorf, M.: Analysis of hepatitis C virus quasispecies populations by temperature gradient gel electrophoresis. *J. Gen. Virol.*, 76 (1995) 881-887; C.A., 123 (1995) 48661b.
- 3190 Lu, X. and Yeung, E.S.: Optimization of excitation and detection geometry for multiplexed capillary array electrophoresis of DNA fragments. *Appl. Spectrosc.*, 49 (1995) 605-609; C.A., 123 (1995) 48671e.
- 3191 Macdonald, R. and Kalmakoff, J.: Comparison of pulsed-field gel electrophoresis DNA fingerprints of field isolates of the entomopathogen *Bacillus popilliae*. *Appl. Environ. Microbiol.*, 61 (1995) 2446-2449; C.A., 123 (1995) 2143f.
- 3192 Maeda, M., Asahara, S., Nishi, T., Mushiake, S., Oka, T., Shimada, S., Chiba, T., Tohyama, M. and Futai, M.: The rat intrinsic factor gene: its 5'-upstream region and chief cell-specific transcription. *J. Biochem. (Tokyo)*, 117 (1995) 1305-1311.

- 3193 Menke, M.A.O.H., Tiemann, M., Vogelsang, D., Boie, C. and Parwaresch, R.: Temperature gradient gel electrophoresis for analysis of a polymerase chain reaction-based diagnostic clonality assay in the early stages of cutaneous T-cell lymphomas. *Electrophoresis (Weinheim)*, 16 (1995) 733-738.
- 3194 Nakagawa, C.W., Mutoh, N. and Hayashi, Y.: Transcriptional regulation of catalase gene in the fission yeast *Schizosaccharomyces pombe*: molecular cloning of the catalase gene and Northern blot analyses of the transcript. *J. Biochem. (Tokyo)*, 118 (1995) 109-116.
- 3195 Oto, M., Suehiro, T. and Yuasa, Y.: DNA hybridization analysis of PCR products by non-gel sieving capillary electrophoresis. *PCR Methods Appl.*, 4 (1995) 303-304; *C.A.*, 123 (1995) 2096t.
- 3196 Reymer, P.W.A., Groenemeyer, B.E., van de Burg, R. and Kastelein, J.J.P.: Apolipoprotein E genotyping on agarose gels. *Clin. Chem. (Washington)*, 41 (1995) 1046-1047.
- 3197 Schlötterer, C.: Temperature-gradient gel electrophoresis as a screening tool for polymorphisms in multigene families. *Electrophoresis (Weinheim)*, 16 (1995) 722-728.
- 3198 Schuetz, W.: (Capillary electrophoresis of DNA restriction fragments and PCR products with laser-induced fluorescence detection (LIF)). *Bioforum*, 17 (1994) 510-512; *C.A.*, 122 (1995) 260021s - a review with 6 refs.
- 3199 Schwartz, D.W.M., Jungl, E.M., Krenek, O.R. and Mayr, W.R.: Simple and rapid typing for VNTR polymorphisms using high resolution electrophoresis of PCR products on rehydratable polyacrylamide gels. *Adv. Forensic Haemogenet.*, 5 (1994) 170-172; *C.A.*, 122 (1995) 257242j.
- 3200 Sekiguchi, K., Yokota, C., Asashima, M., Kaname, T., Fan, Q.-W., Muramatsu, T. and Kadomatsu, K.: Restricted expression of *Xenopus* midkine gene during early development. *J. Biochem. (Tokyo)*, 118 (1995) 94-100.
- 3201 Serwer, P., Hayes, S.J., Watson, R.H. and Khan, S.A.: Gel electrophoretic analysis of bacteriophage assembly intermediates in bacteriophage plaques. *Appl. Theor. Electrophor.*, 4 (1995) 211-217; *C.A.*, 122 (1995) 310125b.
- 3202 Sestini, R., Orlando, C., Zentilin, L., Lami, D., Gelmini, S., Pincani, P., Giacca, M. and Pazzagli, M.: Gene amplification for c-erbB-2, c-myc, epidermal growth factor receptor, int-2, and N-myc measured by quantitative PCR with a multiple competitor template. *Clin. Chem. (Washington)*, 41 (1995) 826-832.
- 3203 Slater, G.W. and Mayer, P.: Electrophoretic resolution versus fluctuations of the lateral dimensions of a capillary. *Electrophoresis (Weinheim)*, 16 (1995) 771-779.
- 3204 Sugie, H., Sugie, Y., Ito, M., Fukuda, T., Nonaka, I. and Igarashi, Y.: Genetic analysis of Japanese patients with myophosphorylase deficiency (McArdle's disease): single-codon deletion in exon 17 is the predominant mutation. *Clin. Chim. Acta*, 236 (1995) 81-86.
- 3205 Tiemann, M., Menke, M.A.O.H., Asbeck, R., Wacker, H.-H. and Parwaresch, R.: Temperature gradient gel electrophoresis for analysis of clonal evolution in non-Hodgkin's lymphoma of the thyroid. *Electrophoresis (Weinheim)*, 16 (1995) 729-732.
- 3206 Trott, D.J., Masters, A.M., Carson, J.M., Ellis, T.M. and Hampson, D.J.: Genetic analysis of *Dermatophilus* spp. using multilocus enzyme electrophoresis. *Zentralbl. Bakteriol.*, 282 (1995) 24-34; *C.A.*, 123 (1995) 2035x.
- 3207 Yamanaka, H.I., Kirii, Y. and Ohmoto, H.: An improved phage display antibody cloning system using newly designed PCR primers optimized for Pfu DNA polymerase. *J. Biochem. (Tokyo)*, 117 (1995) 1218-1227.
- 3208 Yang, C., Maruyama, S., Yanagi, S., Wang, X., Takata, M., Kurosaki, T. and Yamamura, H.: Syk and Lyn are involved in radiation-induced signaling, but inactivation of Syk or Lyn alone is not sufficient to prevent radiation-induced apoptosis. *J. Biochem. (Tokyo)*, 118 (1995) 33-38.
- See also 2726, 2922, 3125, 3172.
- 21f. *Complex mixtures of nucleic acids and their fragments*
- 3209 Cann, J.R., Pfenninger, O. and Pettijohn, D.E.: Theory of the mobility-shift assay of nonspecific protein-DNA complexes governed by conditional probabilities: the HU:DNA complex. *Electrophoresis (Weinheim)*, 16 (1995) 881-887.
- 3210 Chang, H.-T. and Chrambach, A.: Dispersion coefficients of a protein and DNA fragment in polyacrylamide gel electrophoresis as a function of parameters defining the effective gel pore size and particle size. *Electrophoresis (Weinheim)*, 16 (1995) 895-898.
22. ALKALOIDS
- 3211 Bjornsdottir, I. and Hansen, S.H.: Determination of opium alkaloids in opium by capillary electrophoresis. *J. Pharm. Biomed. Anal.*, 13 (1995) 687-693.
- 3212 Chicharro, M., Zapardiel, A., Bermejo, E., Perez-Lopez, J.A. and Hernandez, L.: Direct determination of ephedrine alkaloids and epinephrine in human urine by capillary zone electrophoresis. *J. Liq. Chromatogr.*, 18 (1995) 1363-1381.
- 3213 Flurer, C.L., Lin, L.A., Satzger, R.D. and Wolnik, K.A.: Determination of ephedrine compounds in nutritional supplements by cyclodextrin-modified capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 133-139.
- 3214 Lirne, M.W., Jones, A.D., Morin, D., Segall, H.J. and Wilson, D.W.: Biliary excretion of pyrrolic metabolites of [¹⁴C]monocrotaline in the rat. *Drug Metab. Disp.*, 23 (1995) 422-429.
- 3215 Yang, S.S. and Smetena, I.: Evaluation of capillary electrophoresis for the analysis of nicotine and selected minor alkaloids from Tobacco. *Chromatographia*, 40 (1995) 375-378.
23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN
- 23d. *Pyridine derivatives*
- 3216 McKillop, A.G., Smith, R.M., Rowe, R.C. and Wren, S.A.C.: Separation and identification of the Z and E isomers of 2-(3-pentenyl)pyridine by capillary electrophoresis and nuclear magnetic resonance spectroscopy. *J. Chromatogr. A*, 700 (1995) 69-72.
- See also 2739.

23e. Other N-heterocyclic compounds

- 3217 Chankvetadze, B., Endresz, G. and Blaschke, G.: Enantiomeric resolution of chiral imidazole derivatives using capillary electrophoresis with cyclodextrin-type buffer modifiers. *J. Chromatogr. A*, 700 (1995) 43-49.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 3218 Furuta, R. and Doi, T.: Enantiomeric separation of a thiazole derivative by high-performance liquid chromatography and micellar electrokinetic chromatography. *J. Chromatogr. A*, 708 (1995) 245-251.
- 3219 Shah, P.A. and Quinones, L.: Validation of a micellar electrokinetic capillary chromatography (MECC) method for the determination of *p*-toluenesulfonic acid impurity in a pharmaceutical intermediate. *J. Liq. Chromatogr.*, 18 (1995) 1349-1362.
- 3220 Vogt, C., Heinig, K., Langer, B., Mattusch, J. and Werner, G.: Determination of linear alkylbenzenesulfonates by high-performance liquid chromatography and capillary zone electrophoresis. *Fresenius J. Anal. Chem.*, 352 (1995) 508-514.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- See 2845, 2853, 2856, 2860, 2916, 2960, 2965, 3005, 3017, 3060, 3099, 3307.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 3221 Han, F., Fasching, J.L. and Brown, P.R.: Speciation of organotin compounds by capillary electrophoresis using indirect ultraviolet absorbance detection. *J. Chromatogr. B*, 669 (1995) 103-112.
- 3222 Li, K., Li, S.F.Y. and Lee, H.K.: Determination of organolead and organotin compounds in water samples by micellar electrokinetic chromatography. *J. Liq. Chromatogr.*, 18 (1995) 1325-1347.

See also 3160.

26b. Boranes, silanes and related non-metallic compounds

See 3292.

26c. Coordination compounds

See 2745, 3267.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 3223 Lee, J.J., Moon, D.C. and Lee, K.-J.: (Analysis of water-soluble vitamins in pharmaceutical products by capillary electrophoresis). *Yakhak Hoechi*, 38 (1994) 38-45; *C.A.*, 122 (1995) 223003s.

28. ANTIBIOTICS

- 3224 Cutting, J.H., Kiessling, W.M., Bond, F.L., McCarron, J.E., Kreuzer, K.S., Hurlbut, J.A. and Sofos, J.N.: Agarose gel electrophoretic detection of six β -lactam antibiotic residues in milk. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 663-667.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 3225 Harumi, T., Kitamura, Y. and Kuzuhara, S.: Analysis of pesticides and pigments by electrophoresis. *Jpn. Kokai Tokkyo Koho JP 07 55,769* [95 55,769] (Cl. G01N27/447), 03 Mar. 1995, Appl. 93/222,766, 17 Aug. 1993; 5 pp.; *C.A.*, 122 (1995) 313410b.
- 3226 Süsse, H. and Müller, H.: Application of micellar electrokinetic capillary chromatography to the analysis of pesticides. *Fresenius J. Anal. Chem.*, 352 (1995) 470-473.

29e. Herbicides

- 3227 Dinelli, G., Vicari, A. and Brandolini, V.: Detection and quantitation of sulfonylurea herbicides in soil at the ppb level by capillary electrophoresis. *J. Chromatogr. A*, 700 (1995) 201-207.
- 3228 Krynitsky, A.J. and Swineford, D.M.: Determination of sulfonylurea herbicides in grains by capillary electrophoresis. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1091-1096.

See also 3270.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 3230 Tapley, K.N.: Capillary electrophoretic analysis of the reactions of bifunctional reactive dyes under various conditions including a study of the analysis of the traditionally difficult to analyse phthalocyanine dyes. *J. Chromatogr. A*, 706 (1995) 555-562.
- 3231 Thompson, C.O. and Trenerry, V.C.: Determination of synthetic colours in confectionery and cordials by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 704 (1995) 195-201.

See also 2715.

- 30b. *Chloroplast and other natural pigments*
- See 3225.
31. PLASTICS AND THEIR INTERMEDIATES
- 3232 Huff, B.V.O. and McIntire, G.R.: Determination of the electrophoretic mobility of polystyrene particles by capillary electrophoresis. *J. Microcolumn Sep.*, 6 (1994) 591-594; C.A., 122 (1995) 241061p.
- 3233 Ono, M., Tanaka, C. and Yoshinaga, K.: (The analysis of water-soluble component in latex by capillary electrophoresis). *Kami Parupu Kenkyu Happyokai Koen Yoshishu*, 61 (1994) 96-99; C.A., 122 (1995) 316665z.
- 3234 Schulz, S.F. and Sticher, H.: Surface charge densities and electrophoretic mobilities of aqueous colloidal suspensions of latex spheres with different ionizable groups. *Prog. Colloid Polym. Sci.*, 97 (1994) 85-88; C.A., 122 (1995) 249174y.
- See also 2705.
32. DRUG ANALYSIS
- 32a. *Drug analysis, general techniques*
- 3235 Filbey, S.D. and Altria, K.D.: Robustness testing of a capillary electrophoresis method for the determination of potassium content in the potassium salt of an acidic drug. *J. Capillary Electrophor.*, 1 (1994) 190-195; C.A., 122 (1995) 299249x.
- 3236 Henion, J.: Coupled techniques for problem solving in pharmaceutical analysis: liquid chromatography/mass spectrometry and capillary electrophoresis/mass spectrometry using atmospheric pressure ionization. In: Crommelin, D.J.A., Midha, K.K. and Nagi, T. (Editors), *Top. Pharm. Sci. 1993, Proc. Int. Congr. Pharm. Sci. F.I.P., 53rd, 1993*, Medpharm. Sci. Pub., Stuttgart, 1994, pp. 297-313; C.A., 122 (1995) 222975y.
- 3237 Li, S.F.Y., Ng, C.L. and Ong, C.P.: Pharmaceutical analysis by capillary electrophoresis. *Adv. Chromatogr. (N.Y.)*, 35 (1995) 199-257; C.A., 122 (1995) 274187n - a review with 125 refs.
- 3238 Lloyd, D.K.: Determination of some drugs by capillary electrophoresis with direct injection of biological fluids. *Methodol. Surv. Bioanal. Drugs*, 23(Biofluid and Tissue Analysis for Drugs, Including Hypolipidaemics) (1994) 41-50; C.A., 122 (1995) 255317p.
- 3239 Lloyd, D.K.: Pharmaceutical and biomedical applications of capillary electrophoresis. *Prog. Pharm. Biomed. Anal.*, 1 (1994) 3-40; C.A., 122 (1995) 222971u - a review with refs.
- 3240 Smith, J.T. and Vinjamoori, D.V.: Rapid determination of logarithmic partition coefficients between *n*-octanol and water using micellar electrokinetic capillary chromatography. *J. Chromatogr. B*, 669 (1995) 59-66.
- 3241 Terabe, S. and Nishi, H.: Capillary electrophoresis and its contribution in the pharmaceutical industry. In: Crommelin, D.J.A., Midha, K.K. and Nagi, T. (Editors), *Top. Pharm. Sci. 1993, Proc. Int. Congr. Pharm. Sci. F.I.P., 53rd, 1993*, Medpharm. Sci. Pub., Stuttgart, 1994, pp. 319-331; C.A., 122 (1995) 222970t - a review with 66 refs.
- 3242 Tomlinson, A.J., Benson, L.M. and Naylor, S.: Online capillary electrophoresis-mass spectrometry for the analysis of drug metabolite mixtures: practical considerations. *J. Capillary Electrophor.*, 1 (1994) 127-135; C.A., 122 (1995) 281265q.
- 3243 Weinmann, W., Logemann, E. and Werp, J.: Drug analysis by capillary electrophoresis. In: Mueller, R.K. (Editor), *Contrib. Forensic Toxicol., Proc. Int. Meet. Int. Assoc. Forensic Toxicol., 31st 1993*, MOLINAPress, Leipzig, 1994, pp. 494-498; C.A., 123 (1995) 2880u.
- 3244 Zhang, C.-X. and Thormann, W.: Determination of drug levels in human serum by micellar electrokinetic capillary chromatography with direct sample injection using different quantitation strategies. *J. Capillary Electrophor.*, 1 (1994) 208-218; C.A., 122 (1995) 281264p.
- See also 2671, 2762, 2767.
- 32b. *Antirheumatics and antiinflammatory drugs*
- 3245 Bechet, I., Fillet, M., Hubert, P. and Crommen, J.: Quantitative analysis of non-steroidal anti-inflammatory drugs by capillary zone electrophoresis. *J. Pharm. Biomed. Anal.*, 13 (1995) 497-503.
- 3246 Perrett, D. and Ross, G.A.: Rapid determination of drugs in biofluids by capillary electrophoresis. Measurement of antipyrine in saliva for pharmacokinetic studies. *J. Chromatogr. A*, 700 (1995) 179-186.
- See also 2770.
- 32c. *Autonomic and cardiovascular drugs*
- 3247 Aumatell, A. and Wells, R.J.: Determination of a cardiac antiarrhythmic, tricyclic antipsychotics and antidepressants in human and animal urine by micellar electrokinetic capillary chromatography using a bile salt. *J. Chromatogr. B*, 669 (1995) 331-344.
- 3248 Brettnall, A.E. and Clarke, G.S.: Investigation and optimisation of the use of micellar electrokinetic chromatography for the analysis of six cardiovascular drugs. *J. Chromatogr. A*, 700 (1995) 173-178.
- 3249 Lalljie, S.P.D., Vindevogel, J. and Sandra, P.: Capillary electrophoresis as a fast screening method for the determination of clenbuterol in urine. *J. Capillary Electrophor.*, 1 (1994) 241-245; C.A., 122 (1995) 305722q.
- 3250 Luo, G., Wang, Y., Ewing, A.G. and Strein, T.G.: Chiral separation of a novel calcium antagonist using capillary electrophoresis with cyclodextrins and diode array detection. *J. Capillary Electrophor.*, 1 (1994) 175-180; C.A., 122 (1995) 299210c.
- 3251 Matchett, M.W., Branch, S.K. and Jefferies, T.M.: Application of modified cyclodextrins in capillary electrophoresis for enantiomeric resolution of propranolol and analogues. *J. Chromatogr. A*, 705 (1995) 351-361.
- 3252 Sadecka, J. and Polonsky, J.: Determination of some β -blockers by capillary isotachopheresis in human serum. *Pharmazie*, 50 (1995) 434-435.
- 3253 Wang, C.J., Tian, Z., Watkins, R., Cook, J. and Lin, C.: Determination of a cyclic guanine monophosphate phosphodiesterase inhibitor (SCH 51866) in rat serum using capillary zone electrophoresis. *J. Chromatogr. B*, 669 (1995) 141-148.

See also 3212.

32d. *Central nervous system drugs*

- 3254 Boonkerd, S., Detaevnier, M.R., Michotte, Y. and Vindevoegel, J.: Suppression of chiral recognition of 3-hydroxy-1,4-benzodiazepines during micellar electrokinetic capillary chromatography with bile salts. *J. Chromatogr. A*, 704 (1995) 238-241.
- 3255 Ferslew, K.E., Hagardorn, A.N. and McCormick, W.F.: Application of micellar electrokinetic capillary chromatography to forensic analysis of barbiturates in biological fluids. *J. Forensic Sci.*, 40 (1995) 245-249; *C.A.*, 122 (1995) 284045d.
- 3256 Garcia, L.L., Shihabi, Z.K. and Oles, K.: Determination of gabapentin in serum by capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 157-162.
- 3257 Hansen, S.H., Bjornsdottir, I. and Tjornelund, J.: Separation of basic drug substances of very similar structure using micellar electrokinetic chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 489-495.
- 3258 Hiraoka, A., Miura, I., Akai, J., Tominaga, I., Hattori, M., Machida, S. and Arato, T.: Capillary electrophoresis as an aid in therapeutic monitoring of the serum anti-epileptic drug concentrations. *Seibutsu Shiryō Bunseki*, 17 (1994) 274-279; *C.A.*, 122 (1995) 281274s.
- 3259 Hufschmid, E., Theurillat, R., Martin, U. and Thormann, W.: Exploration of the metabolism of dihydrocodeine via determination of its metabolites in human urine using micellar electrokinetic capillary chromatography. *J. Chromatogr. B*, 668 (1995) 159-170.
- 3260 Lehotay, J. and Cizmarik, J.: Correlation between capacity factor and relative step height of zones H from isotachopheresis and local anaesthetic activity in the series of piperidinoethyl esters of 2-, 3-alkoxyphenylcarbamic acids. *Pharmazie*, 50 (1995) 433-434.
- 3261 Piperaki, S., Penn, S.G. and Goodall, D.M.: Systematic approach to treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography. II. A study of the enantiomeric separation of fluoxetine and norfluoxetine. *J. Chromatogr. A*, 700 (1995) 59-67.
- 3262 Quaglia, M.G., Farina, A., Bossu, E. and Dell'Aquila, C.: Analysis of non-benzodiazepinic anxiolytic agents by capillary zone electrophoresis. *J. Pharm. Biomed. Anal.*, 13 (1995) 505-509.
- 3263 Trenerry, V.C., Robertson, J. and Wells, R.J.: Analysis of illicit amphetamine seizures by capillary electrophoresis. *J. Chromatogr. A*, 708 (1995) 169-176.

See also 2756, 2768, 2782, 3247.

32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*

- 3264 Hsu, L.C., Constable, D.J.C., Orvos, D.R. and Hannah, R.E.: Comparison of high-performance liquid chromatography and capillary zone electrophoresis in penciclovir biodegradation kinetic studies. *J. Chromatogr. B*, 669 (1995) 85-92.

See also 2728.

32f. *Cytostatics*

- 3265 Anigbogu, V.C., Copper, C.L. and Sepaniak, M.J.: Separation of stereoisomers of aminoglutethimide using three capillary electrophoretic techniques. *J. Chromatogr. A*, 705 (1995) 343-349.

See also 3102.

32g. *Other drug categories*

- 3266 Abushoffa, A.M. and Clark, B.J.: Resolution of the enantiomers of oxamniquine by capillary electrophoresis and high-performance liquid chromatography with cyclodextrins and heparin as chiral selectors. *J. Chromatogr. A*, 700 (1995) 51-58.
- 3267 Bullock, J.: Assay for the dianhydride of diethylenetriaminepentaacetic acid and its major degradation products by capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 149-155.
- 3268 Iwagami, S., Sawabe, Y. and Nakagawa, T.: Determination of components in eye lotion by capillary electrophoresis. *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 85-91; *C.A.*, 122 (1995) 222974x.
- 3269 Luksa, J. and Josic, Dj.: Determination of cimetidine in human plasma by free capillary zone electrophoresis. *J. Chromatogr. B*, 667 (1995) 321-327.

See also 3259.

32h. *Toxicological and forensic applications*

- 3270 Dinelli, G., Vicari, A. and Bonetti, A.: Separation of sulfonylurea metabolites in water by capillary electrophoresis. *J. Chromatogr. A*, 700 (1995) 195-200.
- 3271 Lurie, I.S., Chan, K.C., Spratley, T.K., Casale, J.F. and Issaq, H.J.: Separation and detection of acidic and neutral impurities in illicit heroin via capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 3-13.
- 3272 Northrop, D.M., McCord, B.R. and Butler, J.M.: Forensic applications of capillary electrophoresis. *J. Capillary Electrophor.*, 1 (1994) 158-168; *C.A.*, 122 (1995) 283954n - a review with 78 refs.
- 3273 Sheu, S.-J. and Chen, H.-R.: Simultaneous determination of twelve constituents of I-tzu-tang, a Chinese herbal preparation, by high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr. A*, 704 (1995) 141-148.

See also 3211, 3243, 3255, 3263.

32i. *Plant extracts*

- 3274 Sheu, S.-J. and Lu, C.-F.: Determination of six bioactive components of Hsia-cheng-chi-tang by capillary electrophoresis. *J. High Resolut. Chromatogr.*, 18 (1995) 269-270.

See also 3213, 3276.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

- 3275 Jellum, E.: Capillary electrophoresis for medical diagnosis. *J. Capillary Electrophor.*, 1 (1994) 97-105; C.A., 122 (1995) 310100q.

See also 3295.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

- See 2805, 2828, 2953, 2984, 2990, 2992, 2999, 3040, 3064, 3082, 3084, 3103, 3162, 3174, 3178, 3181, 3196, 3202, 3204, 3286.

34. FOOD ANALYSIS

34b. Complex mixtures (single compounds by cross-reference only)

- 3276 Huang, T.-S., Marshall, M.R. and Wei, C.-I.: Identification of red snapper (*Lutjanus campechanus*) using electrophoretic techniques. *J. Food Sci.*, 60 (1995) 279-283; C.A., 122 (1995) 263806n.
- 3277 Huang, T.S.: Identification of red snapper (*Lutjanus campechanus*) using electrophoretic techniques and monoclonal antibodies. Avail. *Univ. Microfilms Int.*, Order No. DA9505775, 1993, 157 p.; C.A., 122 (1995) 329313h.
- 3278 Thompson, C.O., Trenerry, V.C. and Kemmery, B.: Determination of cyclamate in low joule foods by capillary zone electrophoresis with indirect ultraviolet detection. *J. Chromatogr. A*, 704 (1995) 203-210.

See also 2786, 2930, 2945, 2977, 3224, 3225, 3304.

34c. Organoleptically important compounds (flavors, odors, volatiles)

- 3279 Liu, J. and Li, S.F.Y.: Separation and determination of Stevia sweeteners by capillary electrophoresis and high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 1703-1719.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

- 3280 Schmitt, P. and Kettrup, A.: (Capillary electrophoresis in environmental analysis. An example of application in the pesticide and humic substance analysis). *GIT Fachz. Lab.*, 38 (1994) 1312-1318; C.A., 122 (1995) 298136w - a review with 37 refs.

35b. Air pollution (complex mixtures; single compounds by cross-reference only)

See 3289, 3300.

- 35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

- 3281 Takeda, S., Kanda, A., Wakida, S.-i., Yamane, M., Kawahara, A. and Higashi, K.: (Analysis of environmental pollutants by electrokinetic chromatography, (XI). Analysis of colored wastewater containing dyestuffs). *Kankyo Kagaku*, 4 (1994) 524-525; C.A., 122 (1995) 273567z.

See also 2783, 2839, 3222, 3296, 3305, 3306.

35d. Soil pollution (complex mixtures; single compounds by cross-reference only)

See 3227.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 3282 Altria, K.D., Gill, I., Howells, J.S., Luscombe, C.N. and Williams, R.Z.: Trace analysis of detergent residues by capillary electrophoresis. *Chromatographia*, 40 (1995) 527-531.

36c. Complex mixtures, technical products and unidentified compounds

See 2731.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 3283 Jensen, A.E., Cheville, N.F., Ewalt, D.R., Payeur, J.B. and Thoen, C.O.: Application of pulsed-field gel electrophoresis for differentiation of vaccine strain RB51 from field isolates of *Brucella abortus* from cattle, bison, and elk. *Am. J. Vet. Res.*, 56 (1995) 308-312; C.A., 122 (1995) 260332u.
- 3284 Radko, S.P., Sokoloff, A.V., Garner, M.M. and Chrambach, A.: Capillary electrophoresis of rat liver microsomes in polymer solutions. *Electrophoresis (Weinheim)*, 16 (1995) 981-992.
- 3285 Sonohara, R., Muramatsu, N., Ohshima, H. and Kondo, T.: Difference in surface properties between *Escherichia coli* and *Staphylococcus aureus* as revealed by electrophoretic mobility measurements. *Biophys. Chem.*, 55 (1995) 273-277; C.A., 123 (1995) 52146e.

38. INORGANIC COMPOUNDS

38a. Cations

- 3286 Che, P., Xu, J., Shi, H. and Ma, Y.: Quantitative determination of serum iron in human blood by high-performance capillary electrophoresis. *J. Chromatogr. B*, 669 (1995) 45-51.
- 3287 Cheng, K., Zhao, Z., Garrick, R., Nordmeyer, F.R., Lee, M.L. and Lamb, J.D.: Separation of metal cations by electrophoresis in a positively charged coated capillary. *J. Chromatogr. A*, 706 (1995) 517-526.

- 3288 Colburn, B.A., Starnes, S.D., Sepaniak, M.J. and Hinton, R.: Quantitative aspects of rare earth metal determinations using capillary electrophoresis with indirect absorbance detection. *Sep. Sci. Technol.*, 30 (1995) 1511-1527; C.A., 123 (1995) 46911j.
- 3289 Dabek-Zlotorzynska, E. and Dlouhy, J.F.: Application of capillary electrophoresis in atmospheric aerosol analysis: determination of cations. *J. Chromatogr. A*, 706 (1995) 527-534.
- 3290 Francois, C., Morin, P. and Dreux, M.: Effect of the concentration of 18-crown-6 added to the electrolyte upon the separation of ammonium, alkali and alkaline-earth cations by capillary electrophoresis. *J. Chromatogr. A*, 706 (1995) 535-553.
- 3291 Li, Q. and Yeung, E.S.: Contamination control in capillary electrophoresis and quantitative measurement of potassium and sodium in single human erythrocytes. *J. Capillary Electrophor.*, 1 (1994) 55-61; C.A., 122 (1995) 285883n.
- 3292 Lin, L., Wang, J. and Caruso, J.: Arsenic speciation using capillary zone electrophoresis with indirect ultraviolet detection. *J. Chromatogr. Sci.*, 33 (1995) 177-180.
- 3293 Messerschmidt, J., Alt, F. and Tölg, G.: Detection of platinum species in plant material. *Electrophoresis (Weinheim)*, 16 (1995) 800-803.
- 3294 Oshurkova, O.V. and Gorshkov, A.I.: (Separation of the ionic components of salts of alkali and alkaline-earth metals and halogens by capillary isotachopheresis). *Ukr. Khim. Zh.*, 60 (1994) 366-371; C.A., 123 (1995) 24723x.
- 3295 Takatsu, A., Eyama, S. and Uchiumi, A.: Determination of aluminium in serum by capillary zone electrophoresis with laser-induced fluorescence detection. *Chromatographia*, 40 (1995) 125-128.
- 3296 Wang, T. and Li, S.F.Y.: Migration behaviour of alkali and alkaline-earth metal ion-EDTA complexes and quantitative analysis of magnesium in real samples by capillary electrophoresis with indirect ultraviolet detection. *J. Chromatogr. A*, 707 (1995) 343-353.
- 3297 Yang, Q., Zhuang, Y., Smeyers-Verbeke, J. and Massart, D.L.: Interpretation of migration behaviour of inorganic cations in capillary ion electrophoresis based on an equilibrium model. *J. Chromatogr. A*, 706 (1995) 503-515.
- 3298 Zhang, H.-W., Jia, L. and Hu, Z.-D.: Determination of palladium(II) as a chloro complex by capillary zone electrophoresis. *J. Chromatogr. A*, 704 (1995) 242-246.
- 3299 Benz, N.J.: Liquid chromatographic method for determination of water in soils and the optimization of anion separators by capillary zone electrophoresis. Avail. *Univ. Microfilms Int.*, Order No. DA9503530, 1994, 125 p.; C.A., 122 (1995) 329330m.
- 3300 Dabek-Zlotorzynska, E., Dlouhy, J.F., Houle, N., Piechowski, M. and Ritchie, S.: Comparison of capillary zone electrophoresis with ion chromatography and standard photometric methods for the determination of inorganic anions in atmospheric aerosols. *J. Chromatogr. A*, 706 (1995) 469-478.
- 3301 Debus, G.: (Capillary ion electrophoresis with conductivity detection). *GIT Fachz. Lab.*, 39 (1995) 38-44; C.A., 122 (1995) 305350s.
- 3302 Haddad, P.R., Harakuwe, A.H. and Buchberger, W.: Separation of inorganic and organic anionic components of Bayer liquor by capillary zone electrophoresis. I. Optimisation of resolution with electrolyte-containing surfactant mixtures. *J. Chromatogr. A*, 706 (1995) 571-578.
- 3303 Hettiarachchi, K., Ha, Y., Tran, T. and Cheung, A.P.: Application of HPLC and CZE to the analysis of polyoxometalates. *J. Pharm. Biomed. Anal.*, 13 (1995) 515-523.
- 3304 Jimidar, M., Hartmann, C., Cousement, N. and Massart, D.L.: Determination of nitrate and nitrite in vegetables by capillary electrophoresis with indirect detection. *J. Chromatogr. A*, 706 (1995) 479-492.
- 3305 Liu, J.: (Application of capillary electrophoresis technique in the analysis of common anions in drinking water). *Fenxi Huaxue*, 23 (1995) 365; C.A., 122 (1995) 247795c.
- 3306 Nguyen, H.M.L., Tamisier-Karolak, S.L., Czok, M., Laugier, R. and Cardot, P.: Analysis of anions in rain water by capillary electrophoresis. *Analisis*, 23 (1995) 82-87; C.A., 123 (1995) 17298j.
- 3307 Shamsi, S.A. and Danielson, N.D.: Ribonucleotide electrolytes for capillary electrophoresis of polyphosphates and polyphosphonates with indirect photometric detection. *Anal. Chem.*, 67 (1995) 1845-1852.
- 3308 Ueda, T., Maekawa, T., Sadamitsu, D., Oshita, S., Ogino, K. and Nakamura, K.: The determination of nitrite and nitrate in human blood plasma by capillary zone electrophoresis. *Electrophoresis (Weinheim)*, 16 (1995) 1002-1004.
- 3309 Weiss, J. and Hauffe, D.: (Capillary electrophoresis of inorganic anions and organic acids with suppressed conductivity detection). *GIT Spez. Chromatogr.*, 14 (1994) 57-63; C.A., 122 (1995) 329218f - a review with 12 refs.

See also 2750, 3294.

38b. Anions