



Bibliography Section

Liquid Column Chromatography

1. REVIEWS AND BOOKS

- 1 Backer, A.: The chromatographic analysis of refined and synthetic waxes. *J. Chromatogr. Libr.*, 56 (1995) 55-93; C.A., 123 (1995) 147207q - a review with 112 refs.
- 2 Conway, W.D. and Petroski, R.J. (Editors): *Modern Countercurrent Chromatography. (Developed from a Symposium sponsored by the Division of Agricultural and Food Chemistry at the 206th National Meeting of the American Chemical Society Chicago, Illinois, August 22-27, 1993)*. In: ACS Symp. Ser., Vol. 593, ACS, Washington, 1995, 238 p.; C.A., 123 (1995) 124099s.
- 3 Jacob, L.: (Biochromatography). *CLB, Chem. Labor. Biotech.*, 45 (1994) 427-430; C.A., 123 (1995) 164231k - a review without references.
- 4 Rowland, S.J. and Revill, A.T.: Chromatography in petroleum geochemistry. *J. Chromatogr. Libr.*, 56 (1995) 127-141; C.A., 122 (1995) 318119y - a review with 103 refs.
- 5 Scott, R.P.W.: *Chromatography Techniques: Separation Methods for Industry and Academia* In: *Chromatographic Science Series*, Vol. 70, Dekker, New York, 1995, 400 p.; C.A., 123 (1995) 187285v.
- 6 Serwe, M. and Meyer, H.E.: Microseparation techniques I: high performance liquid chromatography. In: Kellner, R., Lottspeich, F. and Meyer, H.E. (Editors), *Microcharact. Proteins*, VCH, Weinheim, 1994, pp. 29-45; C.A., 123 (1995) 106750t - a review with 18 refs.
- 7 Siponi, A. and Verga, G.R.: The O-FID and its applications in petroleum product analysis. *J. Chromatogr. Libr.*, 56 (1995) 143-158; C.A., 122 (1995) 318120s - a review with 25 refs.

See also 18, 20, 36, 37, 38, 43, 47, 48, 52, 94, 97, 99, 105, 106, 107, 124, 131, 134, 137, 139, 142, 144, 145, 151, 153, 158, 161, 162, 165, 167, 169, 170, 171, 177, 180, 206, 211, 232, 254, 256, 267, 278, 280, 282, 284, 287, 288, 295, 299, 308, 309, 312, 434, 436, 440, 631, 654, 673, 687, 703, 759, 775, 777, 778, 792, 945, 951, 952, 953, 955, 961, 987.

2. FUNDAMENTALS, THEORY AND GENERAL

- 2a. General
 - 8 Culley, T. and Dolan, J.W.: Mobile-phase proportioning problems and standard operating procedures. *LC-GC Int.*, 8 (1995) 381-383.
 - 9 Dolan, J.W.: Internal standard problems, buffer precipitation, and column storage. *LC-GC Int.*, 8 (1995) 560-564.
 - 10 Dolan, J.W.: International liquid chromatography problems. *LC-GC Int.*, 8 (1995) 324-327.
 - 11 Eimer, T., Unger, K.K. and Tsuda, T.: Pressurised flow electro-chromatography with reversed phase capillary columns. *Fresenius J. Anal. Chem.*, 352 (1995) 649-653.
 - 12 Hayashi, Y. and Matsuda, R.: Stochastic utility of internal standard method in liquid chromatography. *Anal. Sci.*, 11 (1995) 389-400; C.A., 123 (1995) 159753c.
 - 13 Kirkland, J.J.: Experiences with HPLC-grade solvents. *J. Chromatogr. A*, 715 (1995) 199-200.
 - 14 Majors, R.E.: Current trends in HPLC column usage - a comparative study between the United States and Europe. *LC-GC Int.*, 8 (1995) 368-374.
 - 15 McDowall, R.D.: The right sample in the right tube? *LC-GC Int.*, 8 (1995) 384-385.
 - 16 Webster, G.: Comprehensive and correct chromatography documentation-made easy! *LC-GC Int.*, 8 (1995) 588-592.
 - 17 Willis, B.: Analytical chemistry instrumentation and systems: a view to the future. *Int. Lab.*, 25, No. 4 (1995) 11-17.
 - 18 Zhao, G., Mao, Q., Tang, B. and Shao, X.: (Application of factor analysis in chromatography). *Fenxi Huaxue*, 23 (1995) 224-230; C.A., 123 (1995) 73801e - a review with 68 refs.

For additional information see C.A.:
123 (1995) 187062v.

See also 20, 91, 161.

2b. Thermodynamics and theoretical relationships

- 19 Gagnon, P., Grund, E. and Lindbäck, T.: Media selection and rapid development of binding conditions for high-performance, hydrophobic-interaction chromatography (HIC). *LC-GC Int.*, 8 (1995) 502-510.

- 20 Hwang, Y.-L.: Wave propagation in mass-transfer processes: from chromatography to distillation. *Ind. Eng. Chem. Res.*, 34 (1995) 2849-2864; C.A., 123 (1995) 86834t - a review with many refs.
- 21 Kalinichev, A.: Effects of nonlinearity in the theory of multi-component chromatography. *Ind. Eng. Chem. Res.*, 34 (1995) 2625-2633; C.A., 123 (1995) 86867f.
- 22 Kazakevich, Y.V. and McNair, H.M.: Study of the excess adsorption of the eluent components on different reversed-phase adsorbents. *J. Chromatogr. Sci.*, 33 (1995) 321-327.
- 23 Nikolaropoulos, S.A., Arvanitopoulou, E. and Katsanos, N.A.: Diffusion coefficients in liquids by HPLC and the respective calibrations factors. *Fresenius J. Anal. Chem.*, 352 (1995) 629-642.
- 24 Rowe, K., Bowlin, D., Zou, M. and Davis, J.H.: Application of 2-D statistical theory of overlap to three separation types: 2-D thin-layer chromatography, 2-D gas chromatography, and liquid chromatography/capillary electrophoresis. *Anal. Chem.*, 67 (1995) 2994-3003.
- 25 Velayudhan, A. and Horvath, C.: Characterizing and compensating for enrichment effect in displacement chromatography with multicomponent carriers. *Ind. Eng. Chem. Res.*, 34 (1995) 2789-2795; C.A., 123 (1995) 60003w.

See also 28, 94, 279, 435, 792.

2c. Relationship between structure and chromatographic behaviour

- 26 Liu, H., Ong, S., Glunz, L. and Pidgeon, C.: Predicting drug-membrane interactions by HPLC: structural requirements of chromatographic surfaces. *Anal. Chem.*, 67 (1995) 3550-3557.
- 27 Luco, J.M., Yamin, L.J. and Ferretti, H.F.: Molecular topology and quantum chemical descriptors in the study of reversed-phase liquid chromatography. Hydrogen-bonding behavior of chalcones and flavanones. *J. Pharm. Sci.*, 84 (1995) 903-908.

See also 76, 160.

2d. Measurement of physico-chemical and related values

- 28 Glod, B.K., Haddad, P.R. and Alexander, P.W.: Retention volume of acidic compounds in ion-exclusion chromatography. *Acta Chromatogr.*, 3 (1994) 23-36; C.A., 123 (1995) 131634r.
- 29 Stanley, B.J. and Guiochon, G.: Calculation of adsorption energy distributions of silica samples using nonlinear chromatography. *Langmuir*, 11 (1995) 1735-1743; C.A., 123 (1995) 94167t.

For additional information see C.A.:

- 123 (1995) 57334z, 84885m, 86511k, 131653w, 159828f, 159829g.

See also 26, 93, 234, 255, 756, 758, 761, 767, 769, 770, 771, 772, 773, 774, 775, 776, 777, 780, 781, 829, 845, 897, 966.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 30 Chandler, H.M.: Interrupted-flow assay device. *PCT Int. Appl.* WO 95 16,208 (Cl. GO1N33/558), 15 Jun. 1995, US Appl. 163,341, 07 Dec. 1993; 76 pp.; C.A., 123 (1995) 164647a.
- 31 Chen, J. and Pawliszyn, J.B.: Solid phase microextraction coupled to high-performance liquid chromatography. *Anal. Chem.*, 67 (1995) 2530-2533.
- 32 Dolan, J.W.: Pump seal problems. *LC-GC Int.*, 8 (1995) 490-493.
- 33 Peichung, Lu. and Xiangmin, Z.: Intelligent optimization of multi-column system of chromatography. *Fresenius J. Anal. Chem.*, 352 (1995) 633-638.
- 34 Pfeiffer, W.: Method and device for the separation of dissolved and undissolved mixtures. *PCT Int. Appl.* WO 95 14,220 (Cl. GO1N1/18), 26 May 1995, DE Appl. 4,339,166, 16 Nov. 1993; 30 pp.; C.A., 123 (1995) 164643w.
- 35 Stevenson, R.: The world of separation science: PITTCON® '95. *Int. Lab.*, 25, No. 5 (1995) 8B-8N.

For additional information see C.A.:

- 123 (1995) 73720c, 73874f.

See also 56, 69, 84, 439.

3b. Detectors and detection reagents

- 36 Anonymous: (New detectors in liquid chromatography of carbohydrates). *Kem-Kemi*, 21 (1994) 694-697; C.A., 123 (1995) 101659k - a review with 12 refs.
- 37 Brykina, G.D., Marchenko, D.Yu. and Shpigun, O.A.: Solid-phase spectrophotometry. *J. Anal. Chem.*, 50 (1995) 440-446; C.A., 123 (1995) 186943w - a review with 54 refs.
- 38 Calokerinos, A.C., Deftereos, N.T. and Baeyens, W.R.G.: Chemiluminescence in drug assay. *J. Pharm. Biomed. Anal.*, 13 (1995) 1063-1071 - a review with 122 refs.
- 39 Casella, I.G.: Platinum/glassy carbon electrode as detector for liquid chromatographic determination of hydroxyl-containing compounds. *Anal. Chim. Acta*, 311 (1995) 37-46.
- 40 Chen, P., Nie, L. and Yao, S.-Z.: Application of a series piezoelectric sensor as a detector in ion chromatography. *J. Chromatogr. Sci.*, 33 (1995) 268-272.
- 41 Chen, P., Nie, L.-H., Yao, S.-Z. and Fan, X.-X.: A frequency-voltage conversion circuit for piezoelectric quartz crystal detector for liquid chromatography. *Instrum. Sci. Technol.*, 23 (1995) 137-145; C.A., 123 (1995) 159511x.
- 42 De Backer, B. and Nagels, L.J.: Trends in the coupling of hydrodynamic analysis techniques (LC, CE, and FIA) with electrochemical methods. *LC-GC Int.*, 8 (1995) 498-501.
- 43 Hayakawa, K.: (Development of photometric detection in liquid chromatography). *Yakugaku Kenkyu no Shinpo* 1994, 11 (1995) 127-136; C.A., 123 (1995) 73807m - a review with 39 refs.
- 44 Johnston, T.E.: Chromatographic frequency domain fluorescence modulation lifetime errors caused by photomultiplier baseline offsets. *Anal. Chem.*, 67 (1995) 2835-2841.

- 45 Kou, H.S., Wu, H.L., Chen, S.H. and Wu, S.M.: Chemically removable derivatization reagent for chromatography. II. 2-(1-Naphthyl)ethyl 2-[1-(4-benzyl)-piperazyl]ethanesulfonate dihydrochloride. *J. Liq. Chromatogr.*, 18 (1995) 2323-2335.
- 46 Nunes, J.A., Berniolles, S. and Tong, W.G.: Sensitive laser wave-mixing detection methods for biomedical applications. *Proc. SPIE-Int. Soc. Opt. Eng.*, 2388(Advances in Fluorescence Sensing Technology II) (1995) 205-212; C.A., 123 (1995) 137868s.
- 47 Ryerson, T.B. and Sievers, R.E.: Sulphur-selective chemiluminescence detection in chromatography. *Chem. Anal. (N.Y.)*, 131 (1995) 1-33; C.A., 123 (1995) 187337p - a review with 81 refs.
- 48 Scheer, H.: Diode array detection in liquid chromatography. *Methods Enzymol.*, 246 (1995) 749-758; C.A., 123 (1995) 106784g - a review with 10 refs.
- 49 Suddaby, K.G., Sanaye, R.A., O'Driscoll, K.F. and Rudin, A.: Eliminating lag time estimation in multidetector size-exclusion chromatography. Calibrating each detector independently. *Adv. Chem. Ser.*, 247 (1995) 79-91; C.A., 123 (1995) 170823w.
- 50 Tran, C.D. and Lu, J.: Characterization of the acousto-optic tunable filter for the ultraviolet and visible regions and development of an acousto-optic tunable filter based rapid scanning detector for high-performance liquid chromatography. *Anal. Chim. Acta*, 314 (1995) 57-66.
- 51 Verity, I.: Solvent purity and the individual requirements of different HPLC detectors. *LC-GC Int.*, 8 (1995) 652-655.
- 52 Vigdergauz, M.S. and Bulanova, A.V.: (Obtaining and interpretation of multielement chromatographic spectra). *Izv. Vyssh. Uchebn. Zaved. Khim., Khim. Tekhnol.*, 37 (1994) 92-97; C.A., 123 (1995) 101657h - a review with 10 refs.
- 53 Zhou, J., Zhang, L. and Wang, E.: Enhanced amperometric detector for local anesthetics in liquid chromatography with metal-oxide dispersed glassy carbon electrodes. *Electroanalysis*, 5 (1993) 295-301; C.A., 123 (1995) 123293p.
- For additional information see C.A.:
123 (1995) 73748t, 106885r, 138144q.
- See also 182, 183, 202, 207, 238, 277, 281, 286, 315, 336, 341, 353, 356, 364, 373, 656, 657, 662, 672, 683, 761, 765, 770, 772, 773, 780, 820.
- 3c. Sorbents and columns, packing procedures
- 54 Agarwal, R. and Gupta, M.N.: Evaluation of gluteraldehyde-modified chitosan as a matrix for hydrophobic interaction chromatography. *Anal. Chim. Acta*, 313 (1995) 253-257.
- 55 Boguszewski, B., Lodziowski, R. and Suprynowicz, Z.: Method column filling for high efficiency liquid chromatography columns. *Pol. PL 159,827* (Cl. B01D15/08), 29 Jan. 1993, Appl. 282,181, 23 Oct. 1987; 3 p.; C.A., 123 (1995) 174148c.
- 56 Boss, K.-S., Rudolphi, A., Vielhauer, S., Walfort, A., Lubda, D. and Eisenbeiss, F.: Alkyl-diol silica (ADS): restricted access precolumn packings for direct injection and coupled-column chromatography of biofluids. *Fresenius J. Anal. Chem.*, 352 (1995) 684-690.
- 57 Danielson, N.D., Wangsa, J. and Shamsi, S.A.: Synthesis and characterization of a polymeric fluorocarbon-diamine reversed phase weak anion exchange silica HPLC column packing. *J. Liq. Chromatogr.*, 18 (1995) 2579-2592.
- 58 Denizli, A. and Piskin, E.: Heparin-immobilized polyhydroxyethylmethacrylate microbeads for cholesterol removal: a preliminary report. *J. Chromatogr. B*, 670 (1995) 157-161.
- 59 Friebe, S., Gebauer, S., Krauss, G.J., Goerner, G. and Krueger, J.: HPLC on calixarene bonded silica gels. I. Characterization and applications of the *p*-tert.-butyl-calix[4]arene bonded material. *J. Chromatogr. Sci.*, 33 (1995) 281-284.
- 60 Fujiki, M.: Branched organosilanes and their polymers. *Eur. Pat. Appl. EP 612,756* (Cl. C07F7/12) 31 Aug. 1994, JP Appl. 93/56,321, 23 Feb. 1993; 66 p.; C.A., 122 (1995) 266369h.
- 61 Fujima, H., Kitamura, H., Nakamura, C., Kitagawa, H., Wada, H. and Hakinaka, J.: (Chromatographic properties of a pepsin-conjugated column and optimization of enantiomeric resolution). *Kuomatogurafi*, 15 (1994) 196-197; C.A., 123 (1995) 101694t.
- 62 Gawdzik, B. and Matynia, T.: Highly crosslinked porous copolymer of methacrylic ester of *p,p'*-dihydroxyphenylpropane diglycidyl ether and divinylbenzene as new HPLC packing. *Chem. Anal. (Warsaw)*, 40 (1995) 103-109; C.A., 123 (1995) 114467w.
- 63 Guo, Y. and Colón, L.A.: A stationary phase for open tubular liquid chromatography and electrochromatography using sol-gel technology. *Anal. Chem.*, 67 (1995) 2511-2516.
- 64 Hennion, M.-C., Coquart, V., Guenu, S. and Sella, C.: Retention behaviour of polar compounds using porous graphitic carbon with water-rich mobile phases. *J. Chromatogr. A*, 712 (1995) 287-301.
- 65 Ibrahim, H. and Aubry, A.-F.: Development of melanin-based high-performance liquid chromatography stationary phase and its use in the study of drug-melanin binding interactions. *Anal. Biochem.*, 229 (1995) 272-277.
- 66 Jungbauer, A. and Lettner, H.P.: Liquid chromatography column designed for *in situ* chemical sterilisation. *U.S. Pat. 5,423,982* (Cl. 210-198.2; B01D15/08), 13 Jun. 1995, Appl. 251,847, 31 May 1994; 16 p.; C.A., 123 (1995) 59882a.
- 67 Kamimori, H. and Konishi, M.: (Evaluation and application of liquid chromatographic columns coated with acylcarnitines for enantiomeric separations). *Kuomatogurafi*, 15 (1994) 198-199; C.A., 123 (1995) 101695u.
- 68 Krepelka, I.: PERLOZA - products made of regenerated beaded cellulose. In: Kennedy, J.F., Philips, G.O. and Williams, P.A. (Editors), *Cellulosics*, Horwood, Chichester, 1993, pp. 9-16; C.A., 123 (1995) 59174w.
- 69 Majors, R.E.: New chromatography columns and accessories at the 1995 Pittsburgh Conference, Part II. *LC-GC Int.*, 8 (1995) 312-323.
- 70 Marme, S. and Unger, K.: (Zeolites in adsorption separation, purification, and enrichment of valuable organic products in preparative column liquid chromatography). *Chem.-Ing.-Tech.*, 67 (1995) 580-582; C.A., 123 (1995) 173778w - a review with 8 refs.
- 71 Murakami, R., Hachisako, H., Yamada, K. and Motozato, Y.: Properties of poly(vinyl alcohol) silica hybrid gel particles. *J. Mater. Sci. Lett.*, 14 (1995) 937-938; C.A., 123 (1995) 171801z.
- 72 O'Gara, J.E., Alden, B.A., Walter, T.H., Petersen, J.S., Niederländer, C.L. and Neue, V.D.: Simple preparation of a C₈ HPLC stationary phase with an internal polar functional group. *Anal. Chem.*, 67 (1995) 3809-3813.

- 73 Omi, S., Katami, K., Taguchi, T., Kaneko, K. and Iso, M.: Synthesis and applications of porous SPG (Shirasu Porous Glass) microspheres. *Macromol. Symp.*, 92 (1995) 309-320; C.A., 123 (1995) 58008h.
- 74 Park, W.K.: Self-regenerating ion exchange column chromatography. U.S. US 5,419,819 (Cl. 204-157.2; B01D5/00), 30 May 1995, Appl. 214,261, 17 Mar. 1994; 7 p.; C.A., 123 (1995) 173752h.
- 75 Peseck, J.J., Matyska, M.T., Hemphällä, H. and Christensen, P.: High performance liquid chromatographic characterization of diol bonded phases synthesized via a hydride intermediate. *J. Liq. Chromatogr.*, 18 (1995) 2507-2526.
- 76 Sander, L.C. and Wise, S.A.: Influence of stationary phase chemistry on shape recognition in liquid chromatography. *Anal. Chem.*, 67 (1995) 3284-3292.
- 77 Szabo, G. and Fulop, N.: Studies on the stabilization of phenyl phases for reversed-phase liquid chromatography. In: Kalasz, H., Ettre, L.S. and Pick, J. (Editors), *New Approaches Chromatogr. '93 [Pap. Budapest Chromatogr. Conf.]* 1990, Fekete Sas Kiado, Budapest, 1993, pp. 89-102; C.A., 123 (1995) 131629t.
- 78 Vissers, J.P.C., Chervet, J.-P., van Paassen, P.N.H. and Salzmann, J.-P.: The stability of packed capillary LC columns in routine analyses. *LC-GC Int.*, 8 (1995) 648-651.
- 79 Weber, T.P., Jackson, P.T. and Carr, P.W.: Chromatographic evaluation of porous carbon-clad zirconia microparticles. *Anal. Chem.*, 67 (1995) 3042-3050.
- 80 Zhang, Y., Aimoto, S., Lu, L., Yang, Q. and Lundahl, P.: Immobilized liposome chromatography for analysis of interactions between lipid bilayers and peptides. *Anal. Biochem.*, 229 (1995) 291-298.
- 81 Zühlke, J., Knopp, D. and Niessner, R.: Sol-gel glass as a new support matrix in immunoaffinity chromatography. *Fresenius J. Anal. Chem.*, 352 (1995) 654-659.

For additional information see C.A.:

- 123 (1995) 60451r, 73878k, 101639d, 101640x, 101748p, 114216p, 159825c, 159826d, 187436v, 187437w, 187440s.

See also 14, 26, 92, 128, 131, 133, 135, 137, 139, 140, 141, 157, 163, 166, 214, 350, 393, 401, 442, 443, 444, 459, 490, 517, 614, 729, 754, 783, 833, 975.

3d. Quantitative analysis

See 101, 822.

3e. Preparative scale chromatography

82 Sarker, M. and Guiochon, G.: Study of the operating conditions of axial compression columns for preparative chromatography. *J. Chromatogr. A*, 709 (1995) 227-239.

See also 70, 83, 133, 148, 152, 169, 193, 194, 314, 475, 950.

3f. Programmed temperature, pressure, vapors, gradients

83 Velayudhan, A. and Ladish, M.R.: Effect of modulator sorption on gradient shape in ion-exchange chromatography. *Ind. Eng. Chem. Res.*, 34 (1995) 2805-2810; C.A., 123 (1995) 59997s.

See also 931.

4. SPECIAL TECHNIQUES

4a. Automation

- 84 Dolan, J.W. and Culley, T.: Testing autosampler performance. *LC-GC Int.*, 8 (1995) 628-630.
- 85 Holland, D.A. and Jorgenson, J.W.: Separation of nanoliter samples of biological amines by a comprehensive two-dimensional microcolumn liquid chromatography system. *Anal. Chem.*, 67 (1995) 3275-3283.

See also 88, 390, 391, 525, 767, 795, 851.

4b. Computerization and modelling

- 86 Altepeter, B.J.: (HPLC/mass spectrometry for reliable identification of components). *LaborPraxis*, 19 (1995) 90-92; C.A., 123 (1995) 73812j.
- 87 Dyson, N.: Modern data handling methods [in chromatography]. *J. Chromatogr. Libr.*, 56 (1995) 375-399; C.A., 123 (1995) 131232h.
- 88 Galushko, S.V. and Kamenchuk, A.A.: Fast optimization of the multi-segment gradient profile in reversed-phase high performance liquid chromatography using the Monte-Carlo method. *LC-GC Int.*, 8 (1995) 581-587.
- 89 Galushko, S.V., Kamenchuk, A.A. and Pit, G.L.: Software for method development in reversed-phase liquid chromatography. *Int. Lab.*, 25, No. 4 (1995) 14C-14H.
- 90 Gobburu, J.V.S., Shelver, W.L. and Shelver, W.H.: Application of artificial neural networks in the optimization of HPLC mobile-phase parameters. *J. Liq. Chromatogr.*, 18 (1995) 1957-1972.
- 91 Huber, L. and Thomas, M.: Validating computer-controlled analytical systems for pharmaceutical laboratories. *LC-GC Int.*, 8 (1995) 572-580.
- 92 Hyun, M.H., Jin, J.S., Na, M.S. and Jyung, K.K.: Chiral recognition models for the liquid chromatographic resolution of enantiomers on (S)-Naproxen-derived chiral stationary phase bearing both α -acid and -basic sites. *Bull. Korean Chem. Soc.*, 16 (1995) 344-348; C.A., 123 (1995) 18072t.
- 93 Jackson, C. and Yan, W.W.: Computer simulation study of multidetector size-exclusion chromatography. Flory-Schulz molecular weight distribution. *Adv. Chem. Ser.*, 247 (1995) 69-77; C.A., 123 (1995) 170822v.
- 94 Kowalska, T.: (Modelling of retention processes in liquid chromatography). *Zesz. Nauk. Univ. Opolskiego: Chem.*, 17 (1994) 5-18; C.A., 123 (1995) 159548q.
- 95 Lesellier, E., Saint Martin, P. and Tchapla, A.: Problems occurring in an attribution step and the benefits of software flexibility during a separation optimization process. *LC-GC Int.*, 8 (1995) 386-395.
- 96 McDowall, R.D.: Chromatographic data-acquisition systems and LIMS: is integration possible? *LC-GC Int.*, 8 (1995) 566-570.
- 97 Noble, D.: Chromatography data systems. *Anal. Chem.*, 67 (1995) 617A-620A.

- 98 Pietrogrande, M.C., Dondi, F., Felinger, A. and Davis, J.M.: Statistical study of peak overlapping in multicomponent chromatograms: importance of the retention pattern. *Chemom. Intell. Lab. Syst.*, 28 (1995) 239-258; C.A., 123 (1995) 159776n.
- 99 Rieger, A.: (A new type of operating system for advanced chromatography data systems). *GIT Fachz. Lab.*, 39 (1995) 450-452; C.A., 123 (1995) 187344p - a review without refs.
- 100 Schlegelmilch, F.: (Software for chromatography). *LaborPraxis*, 19 (1995) 94-96; C.A., 123 (1995) 73485e.
- 101 Tamisier-Karolak, S.L., Tod, M., Bonnardel, P., Czok, M. and Cardot, P.: Daily validation procedure of chromatographic assay using gaussoexponential modelling. *J. Pharm. Biomed. Anal.*, 13 (1995) 959-970.
- 102 Tondeur, D.: Paradigms and paradoxes in modelling adsorption and chromatographic separations. *Ind. Eng. Chem. Res.*, 34 (1995) 2782-2788; C.A., 123 (1995) 60020z.
- 103 Zhang, H., Huang, P. and Ye, Z.: (Continuous annular ion exchange chromatography). *Gaoxiao Huaxue Gongcheng Xuebao*, 8 (1994) 383-388; C.A., 123 (1995) 59333x.
- See also 16, 18, 24, 29, 33, 44, 111, 113, 121, 149, 301, 784, 790, 794, 963, 969.
- 4c. Combination with other physico-chemical techniques (MS, IR etc.)
- 104 Adachi, T., Nemoto, M. and Ito, Y.: Method of increasing the sensitivity of liquid chromatography-atmospheric pressure chemical ionization mass spectrometry using a semi-micro column. *J. Chromatogr. A*, 715 (1995) 13-18.
- 105 Barth, H.G.: Hyphenated polymer separation techniques. Present and future role. *Adv. Chem. Ser.*, 247 (1995) 3-11; C.A., 123 (1995) 170715n - a review with 20 refs.
- 106 Byrdy, F.A. and Caruso, J.A.: Selective chromatographic detection by plasma mass spectrometry. *Chem. Anal. (N.Y.)*, 131 (1995) 171-207; C.A., 123 (1995) 187338q - a review with 97 refs.
- 107 Dawkins, J.V.: Compositional heterogeneity of copolymers by coupled techniques with chromatographic columns and multidetectors. *Adv. Chem. Ser.*, 247 (1995) 197-209; C.A., 123 (1995) 170716p - a review with 25 refs.
- 108 Dobberstein, P. and Muenster, H.: Application of a new atmospheric pressure ionization source for double focusing sector instruments. *J. Chromatogr. A*, 712 (1995) 3-15.
- 109 Doerge, D.R. and Bajic, S.: Isotope dilution LC/MS using a particle beam interface. *Prog. Ind. Microbiol.*, 32 (1995) 37-50; C.A., 123 (1995) 162796z.
- 110 Honing, M., Barceló, D., Jager, M.E., Slobodnik, J., van Baar, B.L.M. and Brinkman, U.A.T.: Effect of ion source pressure on ion formation of carbamates in particle-beam chemical-ionization mass spectrometry. *J. Chromatogr. A*, 712 (1995) 21-30.
- 111 Huang, S.K. and Garza, N.R.: Optimization of the Hewlett-Packard particle-beam liquid chromatography-mass spectrometry interface by statistical experimental design. *J. Am. Soc. Mass Spectrom.*, 6 (1995) 507-512; C.A., 123 (1995) 187359x.
- 112 Kanai, M., Ingendoh, A., Seta, K., Nakayama, H., Isobe, T., Okuyama, T.: (Off-line HPLC-MALDI/TOF-MS system for protein mapping, peptide mapping and structural analysis of oligosaccharide chain of calreticulin). *Kuromatogurai*, 16 (1995) 23-30; C.A., 123 (1995) 164571w.
- 113 McGuire, J.N., Proefke, M.L., Conway, W.D. and Rinehart, K.L.: On-line fast atom bombardment mass spectrometric detection in high-speed countercurrent chromatography through a moving belt interface. *ACS Symp. Ser.*, 593 (1995) 129-142; C.A., 123 (1995) 187391b.
- 114 Nagra, D.S. and Li, L.: Liquid chromatography-time-of-flight mass spectrometry with continuous-flow matrix-assisted laser desorption ionization. *J. Chromatogr. A*, 711 (1995) 235-245.
- 115 Nojima, K., Fujimaki, S., Hertsens, R.C. and Morita, T.: Application of liquid chromatography-atmospheric pressure chemical ionization mass spectrometry to a sector mass spectrometer. *J. Chromatogr. A*, 712 (1995) 17-19.
- 116 Schmeer, K., Behnke, B. and Bayer, E.: Capillary electrochromatography - electrospray mass spectrometry: a microanalysis technique. *Anal. Chem.*, 67 (1995) 3656-3658.
- 117 Señoráns, F.J., Herráiz, M. and Tabera, J.: On-line reversed-phase liquid chromatography-capillary gas chromatography using a programmed temperature vaporizer as interface. *J. High Resolut. Chromatogr.*, 18 (1995) 433-438.
- 118 Simonsick, W.J., Jr. and Prokai, L.: Size-exclusion chromatography with electrospray mass spectrometric detection. *Adv. Chem. Ser.*, 247 (1995) 41-56; C.A., 123 (1995) 170820t.
- 119 Suzuki, K.T., Yoneda, S., Itoh, M. and Ohmichi, M.: Enriched stable isotopes of elements used as tracers: methods of presenting high-performance liquid chromatographic-inductively coupled argon plasma mass spectrometric data. *J. Chromatogr. B*, 670 (1995) 63-71.
- 120 Van der Hoeven, R.A.M., Buscher, B.A.P., Tjaden, U.R. and van der Greef, J.: Performance of an electrospray-interfaced thermospray ion source in hyphenated techniques. *J. Chromatogr. A*, 712 (1995) 211-218.
- 121 Visentini, J., Kwong, E.C., Carrier, A., Zidarov, D. and Bertrand, M.J.: Comparison of softwares used for the detection of analytes present at low levels in liquid chromatographic-mass spectrometric experiments. *J. Chromatogr. A*, 712 (1995) 31-43.
- See also 24, 86, 146, 212, 213, 218, 234, 256, 263, 284, 299, 310, 312, 316, 318, 357, 366, 368, 370, 372, 381, 390, 391, 407, 424, 427, 439, 441, 482, 492, 526, 630, 660, 666, 667, 669, 687, 688, 735, 737, 738, 740, 748, 751, 753, 757, 761, 769, 770, 771, 772, 774, 780, 796, 799, 810, 815, 823, 827, 837, 846, 850, 864, 865, 881, 883, 899, 913, 920, 930, 951, 956, 969, 970, 971, 989.
- 4d. Affinity chromatography (advances)
- 122 Boeg-Hansen, T.C. and Davis, W.C.: Preparation and use of immunoaffinity columns with monoclonal antibodies without purification from ascites and tissue-culture medium. *Methods Mol. Biol. (Totowa)*, 45(Monoclonal Antibody Protocols) (1995) 189-193; C.A., 123 (1995) 164313p.

- 123 Brassfield, A.L.: Antigen purification by monoclonal antibody immunoaffinity chromatography. *Methods Mol. Biol. (Totowa)*, 45(Monoclonal Antibody Protocols) (1995) 195-203; C.A., 123 (1995) 164314q.
- 124 Ogawa, H. and Matsumoto, I.: (Fundamentals of affinity chromatography). *Bunseki*, (1995) 430-439; C.A., 123 (1995) 131604f - a review with 16 refs.
- 125 Rasmussen, J.K., Curry, R.A. and Walker, M.M.: Heparin functional supports. *PCT Int. Appl. WO 95 05,400* (Cl. C08B37/10), 23 Feb. 1995, US Appl. 109,341, 19 Aug. 1993; 23 pp.; C.A., 123 (1995) 138147t.
- For additional information see C.A.:
123 (1995) 107247w.
- See also 58, 143, 275, 293, 294, 295, 383, 393, 433, 437, 462, 505, 529, 563, 569, 783.
- 4f. Trace analysis and preseparation techniques**
- 126 Ashraf-Khorassani, M., Barzegar, M. and Yamini, Y.: On-line coupling of supercritical fluid extraction with high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 472-476.
- 127 Cai, Y., Cabañas, M., Fernández-Turiel, J.L., Abalos, M. and Bayona, J.M.: On-line preconcentration of selenium(IV) and selenium(VI) in aqueous matrices followed by liquid chromatography-inductively coupled plasma mass spectrometry determination. *Anal. Chim. Acta*, 314 (1995) 183-192.
- 128 Hosoya, K., Kageyama, Y., Yoshizako, K., Kimata, K., Araki, T. and Tanaka, N.: Uniform-sized polymer-based separation media prepared using vinyl methacrylate as a cross-linking agent. Possible powerful adsorbent for solid-phase extraction of halogenated organic solvents in an aqueous environment. *J. Chromatogr. A*, 711 (1995) 247-255.
- 129 Majors, R.E.: The use of membranes in sample preparation. *LC-GC Int.*, 8 (1995) 375-380.
- 130 Wiley, S.P. and Leazenbee, L.: Autofiltration of pharmaceutical HPLC samples. *Int. Lab.*, 25, No. 4 (1995) 14J-14P.
- See also 155, 162, 184, 729, 734, 741, 853, 940, 969, 974, 981, 999, 1000.
- 4g. Enantiomers, separation**
- 131 Allenmark, S.: Separation of enantiomers by protein-based chiral phases. In: Subramanian, G. (Editor), *Pract. Approach Chiral Sep. Liq. Chromatogr.*, VCH, Weinheim, 1994, pp. 183-216; C.A., 123 (1995) 142942r - a review with 85 refs.
- 132 Bidlo-Igly, M., Abraham, G., Solyom, S. and Mueller, J.: Chiral separation os some 1-azabicyclo[3.3.1]non-3-ene derivatives of Chiralcel of stationary phase. *ACH-Models Chem.*, 131 (1994) 819-828; C.A., 123 (1995) 65962t.
- 133 Felix, G., Descorts, V., Kopaciewicz, W. and Coryell, B.: Evaluation of a commercial BSA chiral stationary phase in semi-preparative chromatography. *LC-GC Int.*, 8 (1995) 396-398.
- 134 Furuta, R., Okamoto, M., Takahashi, K. and Doi, T.: (Chiral separation by chromatography. Recent progress and future trends). *Sumitomo Kagaku (Osaka)*, (1995) 87-100; C.A., 123 (1995) 137711k - a review with 50 refs.
- 135 Haginaka, J., Seyama, C. and Kanasugi, N.: The absence of chiral recognition ability in ovomucoid: ovoglycoprotein - bonded HPLC stationary phases for chiral recognition. *Anal. Chem.*, 67 (1995) 2539-2547.
- 136 Husain, N.: Spectroscopic and chromatographic studies of guest binding to cyclodextrins and proteins. Avail. *Univ. Microfilms Int.*, Order No. DA9505697, 1994, 244 p.; C.A., 123 (1995) 131609m.
- 137 Kinkel, J.N.: Optically active polyacrylamide/silica composites and related packings and their application in chiral separations. In: Subramanian, G. (Editor), *Pract. Approach Chiral Sep. Liq. Chromatogr.*, VCH, Weinheim, 1994, pp. 217-277; C.A., 123 (1995) 114212j - a review with 152 refs.
- 138 Ma, Y. and Ito, Y.: Chiral separation by high-speed countercurrent chromatography. *Anal. Chem.*, 67 (1995) 3069-3074.
- 139 Nimura, N.: (Optical resolution; chromatographic technique). *Bunseki*, (1994) 119-126; C.A., 123 (1995) 180046j - a review with 80 refs.
- 140 Ooi, T., Kitahara, H., Matsushita, Y. and Kisu, N.: Preparation of support-bound tartaric acid-amino acid monoamide derivative as reagent for chirality recognition and resolving agent for chromatography. *Jpn. Kokai Tokkyo Koho JP 06,298,672* [94,298,672] (Cl. C07B57/00), 25 Oct. 1994, Appl. 93/89,956, 16 Apr. 1993; 11 pp.; C.A., 123 (1995) 144624n.
- 141 Pinkerton, T.C., Howe, W.J., Ulrich, E.L., Comiskey, J.P., Haginaka, J., Murashima, T., Walkenhorst, W.F., Westler, W.M. and Markley, J.L.: Protein binding chiral discrimination of HPLC stationary phases made with whole, fragmented, and third domain turkey ovomucoid. *Anal. Chem.*, 67 (1995) 2354-2367.
- 142 Serban, M. and Ostafe, V.: (Physical method for immobilizing biologically active substances). *Stud. Cercet. Biochim.*, 37 (1994) 99-129; C.A., 123 (1995) 136618y - a review with 103 refs.
- For additional information see C.A.:
123 (1995) 73827t.
- See also 60, 61, 67, 92, 172, 250, 272, 330, 340, 348, 350, 354, 359, 360, 361, 363, 384, 633, 638, 680, 730, 785, 787, 807, 809, 812, 821, 827, 830, 843, 849, 858, 859, 877, 879, 880, 909, 914, 939, 942.
- 4h. Other special techniques**
- 143 Beeskow, T.C., Kusharyoto, W., Anspach, F.B., Kröner, K.H. and Deckwer, W.-D.: Surface modification of microporous polyamide membranes with hydroxyethyl cellulose and their application as affinity membranes. *J. Chromatogr. A*, 715 (1995) 49-65.
- 144 Berthod, A.: Separation with a liquid stationary phase: the countercurrent chromatography technique. *Instrum. Sci. Technol.*, 23 (1995) 75-89; C.A., 123 (1995) 187340j - a review with 16 refs.
- 145 Conway, W.D.: Overview of countercurrent chromatography. *ACS Symp. Ser.*, 593 (1995) 1-14; C.A., 123 (1995) 187346r - a review with 39 refs.
- 146 Dekkers, S.E.G., Tjaden, U.R. and van der Greef, J.: Development of an instrumental configuration for pseudo-electrochromatography-electrospray mass spectrometry. *J. Chromatogr. A*, 712 (1995) 201-209.

- 147 Fung, Y.S. and Dao, K.L.: Oxygen bomb combustion ion chromatography for elemental analysis of heteroatoms in fuel and wastes development. *Anal. Chim. Acta*, 315 (1995) 347-355.
- 148 Hsu, S.T.: Design and dynamics of large-scale chromatographic processes (recycle chromatography purification). Avail. *Univ. Microfilms Int.*, Order No. DA9512990, 1994, 151 p.; C.A., 123 (1995) 147601p.
- 149 Levin, S., Nudelman, R., Reschigian, P. and Pasti, L.: Simulation of fractograms of fat emulsions in power-programmed sedimentation field-flow fractionation (SdFFF). *J. Pharm. Biomed. Anal.*, 13 (1995) 869-877.
- 150 Shiundu, P.M. and Giddings, J.C.: Influence of bulk and surface composition on the retention of colloidal particles in thermal field-flow fractionation. *J. Chromatogr. A*, 715 (1995) 117-126.
- 151 Tamura, A. and Masujima, T.: (Development of chromatovideoscopy. It's principle and application). *Jasco Rep.*, 37 (1995) 1-6; C.A., 123 (1995) 73797h - a review with 21 refs.
- 152 Weisz, A., Andruszewski, D., Shinomiya, K. and Ito, Y.: Preparative separation of components of commercial 4,5,6,7-tetrachlorofluorescein chromatography. *ACS Symp. Ser.*, 593 (1995) 203-217; C.A., 123 (1995) 85913f.

For additional information see C.A.:
123 (1995) 117294y, 117302z.

See also 2, 25, 66, 85, 103, 113, 116, 138, 279, 320, 442, 490, 693, 759.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. Cyclic hydrocarbons, fullerenes

- 153 Gadzala, R.M. and Buszewski, B.: Properties and determination of polycyclic aromatic hydrocarbons (PAHs) using chromatographic methods. *Pol. J. Environ. Stud.*, 4 (1995) 5-15; C.A., 123 (1995) 159730t - a review with 79 refs.
- 154 Hadjimohammadi, M.R. and Fatemi, M.H.: Separation and improvement in detection of polycyclic aromatic hydrocarbons by reverse-phase high performance liquid chromatography using micellar mobile phase and fluorescence detector. *J. Liq. Chromatogr.*, 18 (1995) 2569-2578.
- 155 Hoefer, F., Jensen, D., Ezzel, J. and Richeter, B.: (Accelerated solvent extraction of polycyclic aromatic hydrocarbons (PAHs) from solid samples with subsequent HPLC analysis). *GIT Spez. Chromatogr.*, 15 (1995) 68-71; C.A., 123 (1995) 178544b.
- 156 Höner, A., Arnold, M., Hüser, N. and Kleiböhmer, W.: Monitoring polycyclic aromatic hydrocarbons in waste gases. *J. Chromatogr. A*, 710 (1995) 129-137.
- 157 Kimata, K., Hirose, T., Moriuchi, K., Hosoya, K., Araki, T. and Tanaka, N.: High-capacity stationary phases containing heavy atoms for HPLC separation of fullerenes. *Anal. Chem.*, 67 (1995) 2556-2561.
- 158 Lee, H.K.: Recent applications of gas and high-performance liquid chromatographic techniques to the analysis of polycyclic aromatic hydrocarbons in airborne particulates. *J. Chromatogr. A*, 710 (1995) 79-92 - a review with 106 refs.

- 159 Lintelmann, J., Waadt, K., Sauerbrey, R. and Kettrup, A.: On-site solid phase extraction and coupled-column high performance liquid chromatographic determination of six polycyclic aromatic hydrocarbons in water. *Fresenius J. Anal. Chem.*, 352 (1995) 735-742.
- 160 Lopez-Lopez, D., Rubio-Barroso, S. and Polo-Diez, L.M.: Study of PAH's separation and phase-solute interaction by micellar liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2397-2425.
- 161 Neal, A.C.: HPLC and column liquid chromatography. *J. Chromatogr. Libr.*, 56 (1995) 347-374; C.A., 123 (1995) 13135f - a review with 72 refs.
- 162 Peltonen, K. and Kuljukka, T.: Air sampling and analysis of polycyclic aromatic hydrocarbons. *J. Chromatogr. A*, 710 (1995) 93-108 - a review with 131 refs.
- 163 Saito, Y., Ohta, H., Nagashima, H., Itoh, K., Jinno, K., Okamoto, M., Chen, Y.-L., Luehr, G. and Archer, J.: Separation of fullerenes with novel stationary phases in microcolumn high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 1897-1908.
- 164 Shklyav, S.A., Sold, N.V., Simskaya, N.R. and Gerich, S.V.: (Analysis of a series of polycyclic aromatic hydrocarbons on a Miikkrom microcolumn chromatograph). *Vest. Khar'k. Univ.*, 378 (1993) 113-117; C.A., 123 (1995) 101726e.
- 165 Spitzer, T. and Takeuchi, T.: Determination of benzanthrone in environmental samples. *J. Chromatogr. A*, 710 (1995) 109-116 - a review with 77 refs.
- 166 Xiao, J. and Meyerhoff, M.E.: High-performance liquid chromatography of C₆₀, C₇₀, and higher fullerenes on tetraphenylporphyrin-silica stationary phases using strong mobile phase solvents. *J. Chromatogr. A*, 715 (1995) 19-29.

For additional information see C.A.:
123 (1995) 131667d, 159787s.

See also 31, 64, 75, 76, 126, 128, 652, 957.

5c. Halogen derivatives

See 957.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

- 167 Davidson, R.R., Glover, C.J., Burr, B.J. and Bullin, J.A.: Size exclusion chromatography of asphalts. *Chromatogr. Sci. Ser.*, 69 (1995) 211-247; C.A., 123 (1995) 60743n - a review with 112 refs.
- 168 Dhole, V.R. and Ghosal, G.K.: Quality assurance of petrol by HPLC. *J. Liq. Chromatogr.*, 18 (1995) 2475-2488.
- 169 Mascherpa, A.: (Liquid chromatographic techniques). *Riv. Combust.*, 48 (1994) 442-449; C.A., 123 (1995) 174531x - a review.
- 170 Melda, K.J. and Clemons, J.M.: Reformulated gasoline measurements by process chromatography. *Proc. Annu ISA Anal. Div. Symp.*, 28 (1995) 65-71; C.A., 123 (1995) 117671u - a review with 5 refs.

- 171 Partl, M.N., Hugener, M. and Raab, C.: Quality change and structural damage of polymer-bitumen binders during road construction. *Mater. Struct.*, 28 (1995) 112-113; C.A., 123 (1995) 151045c - a review with no refs.

For additional information see C.A.:
123 (1995) 87933m.

See also 1, 4, 7.

6. ALCOHOLS

- 172 Kim, J.-H., Nishida, Y., Ohru, H. and Meguro, H.: Highly sensitive high-performance liquid chromatographic method to discriminate enantiomeric monoacylglycerols based on fluorescent chiral derivatization with (S)-(+)-2-tert.-butyl-2-methyl-1,3-benzodioxole-4-carboxylic acid. *J. Chromatogr. A*, 709 (1995) 375-380.

See also 39, 64, 662, 947.

7. PHENOLS

- 173 Achilli, G., Cellerino, G.P. and D'Erl, Melzi G.V.: Simultaneous determination of 35 phenolic compounds in distilled beverages using HPLC with a Coulometric Electrode Array System (CEAS). In: Cantagrel, R. (Editor), *Symp. Sci. Int. Cognac*, 1st 1992, Tec & Doc-Lavoisier, Paris, 1993, pp. 484-489; C.A., 123 (1995) 167856d.
- 174 Andreoni, N. and Fiorentini, R.: (Determination of phenolic compounds in virgin olive oil). *Riv. Ital. Sostanze Grasse*, 72 (1995) 163-164; C.A., 123 (1995) 167882j.
- 175 Baiocchi, C., Marengo, E., Roggero, M.A., Giacosa, D., Giorcelli, A. and Toccori, S.: Chromatographic and chemometric investigation of the chemical defence mechanism of poplar tree genotypes against a bark fungine parasite. *J. Chromatogr. A*, 715 (1995) 95-104.
- 176 Baiocchi, C., Roggero, M.A., Giacosa, D. and Marengo, E.: HPLC separation and determination of complex mixtures of phenolic compounds on different stationary phases. An efficient general screening method for the presence of phenols in water. *J. Chromatogr. Sci.*, 33 (1995) 338-346.
- 177 Baranowska, I.: (Chromatographic and spectrophotometric methods used in the determination of phenols). *Chemik*, 47 (1994) 350-352; C.A., 123 (1995) 101658j - a review with 50 refs.
- 178 Barroso, C.G., Galvez, M.C., Guillen, D.A. and Perez-Bustamante, J.A.: Analytical characterization of polyphenolic compounds in vinegars by HPLC. *Bull. Liaison - Groupe Polyphenols*, 16(Pt. 2) (1992) 69-73; C.A., 123 (1995) 167858f.
- 179 Bianchi, J., Cros, E. and Pontillon, J.: (Role of phenolic compounds in the development of the soluble coloring matter during cocoa alkalization). *Bull. Liaison - Groupe Polyphenols*, 16(Pt. 2) (1992) 92-95; C.A., 123 (1995) 167982s.

- 180 Bosch Reig, F., Campins Falcó, P. and Verdú Andrés, J.: Preconcentration, sample clean-up, and HPLC determination of phenol and its chloro, methyl, and nitro derivatives in biological samples. *J. Liq. Chromatogr.*, 18 (1995) 2229-2241 - a review with 44 refs.

- 181 Galceran, M.T. and Moyano, E.: Determination of hydroxy-substituted polycyclic aromatic hydrocarbons by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. A*, 715 (1995) 41-48.

- 182 Kou, H.-S., Wu, H.-L., Wang, Y.-F., Chen, S.-H. and Wu, S.-S.: Chemically removable derivatization reagent for liquid chromatography. I. 2-(N-Phthalimido)ethyl 2-(dimethylamino)ethanesulfonate. *J. Chromatogr. A*, 710 (1995) 267-272.

- 183 McCreedy, T. and Fielden, P.R.: Amperometric detector for high-performance liquid chromatography, featuring a glassy carbon working electrode array in the wall-jet configuration. *Analyst (Cambridge)*, 120 (1995) 2343-2346.

- 184 Puig, D. and Barceló, D.: Comparative study of on-line solid phase extraction followed by UV and electrochemical detection in liquid chromatography for the determination of priority phenols in river water samples. *Anal. Chim. Acta*, 311 (1995) 63-69.

- 185 Revenga, J., Rodríguez, F. and Tijero, J.: Determination of 9,10-dihydroxyanthracene and anthraquinone in Kraft pulping liquors by high-performance liquid chromatography. *J. Chromatogr. A*, 712 (1995) 372-377.

- 186 Roume, C. and Jacquet, M.: (Development of chromatographic method for studying complex polyphenols: comparison of wines undergoing various treatments). *Bull. Liaison - Groupe Polyphenols*, 16(Pt. 2) (1992) 217-221; C.A., 123 (1995) 167864c.

- 187 Rozbeh, M. and Hurtubise, R.J.: The liquid chromatographic separation of metabolites of benzo[a]pyrene with γ -cyclodextrin as a mobile phase additive. *J. Liq. Chromatogr.*, 18 (1995) 1909-1931.

See also 64, 331, 788, 966.

8. SUBSTANCES CONTAINING HETERO CYCLIC OXYGEN

8a. Flavonoids

- 188 Greenham, J., Williams, C. and Harborne, J.B.: Identification of lipophilic flavonols by a combination of chromatographic and spectral techniques. *Phytochem. Anal.*, 6 (1995) 211-217; C.A., 123 (1995) 106906y.

- 189 Han, J. and Cai, G.: (RP-HPLC method for the determination of chlorogenic acid and gardenoside in *Artemisia capillaris* compound injection). *Zhongguo Yaoxue Zazhi*, 30 (1995) 166-167; C.A., 123 (1995) 123289s.

- 190 Liu, O., Bonness, M.S., Liu, M., Seradge, E., Dixon, R.A. and Mabry, T.J.: Enzymes of B-ring-deoxy flavonoid biosynthesis in elicited cell cultures of "old man" cactus (*Cephalocereus senilis*). *Arch. Biochem. Biophys.*, 321 (1995) 397-404.

- 191 Pietta, P.G., Mauri, P.L., Simonetti, P. and Testolin, G.: HPLC and MEKC determination of major flavonoids in selected food pools. *Fresenius J. Anal. Chem.*, 352 (1995) 788-792.

- 192 Repcak, M. and Martonfi, P.: The variability pattern of apigenin glucosides in *Chamomilla recutita* diploid and tetraploid cultivars. *Pharmazie*, 50 (1995) 696-699.
- 193 Rubio, B. and Diaz, A.M.: Flavonoid glycosides in leaves and flowers from *Scolymus hispanicus*. *Pharmazie*, 50 (1995) 629-631.
- 194 Villaescusa, L., Diaz, A.M. and Bartolome, C.: Methoxylated flavonoids from *Jasonia glutinosa*, D.C. and their relation with other species of *Jasonia*. *Pharmazie*, 50 (1995) 639-640.

For additional information see C.A.:
123 (1995) 164320p.

See also 27, 942.

8b. Aflatoxins and other mycotoxins

- 195 Joshua, H.: Analysis of aflatoxins in naturally contaminated corn by HPLC, post-column photochemical derivatization, and fluorescence detection. *Am. Lab. (Shelton)*, 27 (1995) 36J-36M; C.A., 123 (1995) 167866g.
- 196 Kussak, A., Andersson, B. and Andersson, K.: Immunoaffinity column clean-up for the high-performance liquid chromatographic determination of aflatoxins B₁, B₂, G₁, G₂, M₁ and Q₁ in urine. *J. Chromatogr. B*, 672 (1995) 253-259.
- 197 Naseem, S.M., Pace, J.G. and Wannemacher, R.W., Jr.: A high-performance liquid chromatographic method for determining [³H]T-2 and its metabolites in biological fluids of the cynomolgus monkey. *J. Anal. Toxicol.*, 19 (1995) 151-156.
- 198 Wang, G., He, C., Lu, C., Zhao, J. and Wang, Y.: (Determination of aflatoxin B₁, B₂, G₁ and G₂ in peanut and corn with immunoaffinity column cleanup and high performance liquid chromatography). *Sepu*, 13 (1995) 239-246; C.A., 123 (1995) 167903s.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 199 Chen, C.-T. and Sheu, S.-J.: Separation of coumarins by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 710 (1995) 323-329.
- 200 Walsh, J.S., Patanella, J.E., Halm, K.A. and Facchine, K.L.: An improved HPLC assay for the assessment of liver slice metabolic viability using 7-ethoxycoumarin. *Drug Metab. Disp.*, 23 (1995) 869-874.
- 201 Wawrzynowicz, T. and Waksmundzka-Hajnos, M.: Influence of various chromatographic parameter on the separation of natural mixtures under column overloading conditions. *Fresenius J. Anal. Chem.*, 352 (1995) 779-782.

See also 27, 203, 324, 355, 786, 931, 947.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 202 Bousquet, E., Tirendi, S., Prezzavento, O. and Tateo, F.: 2,5-Dihydroxybenzohydrazide as electroactive labeling reagent for aliphatic aldehydes by high performance liquid chromatography with electrochemical and ultraviolet detection. *J. Liq. Chromatogr.*, 18 (1995) 1933-1945.

- 203 Diaz, A., Munguia, O., Farina, J. and Llabres, M.: Lactoyl-lactic acid determination of DL-lactide as starting material in the synthesis of polylactic acid. *Drug Dev. Ind. Pharm.*, 21 (1995) 1697-1703; C.A., 123 (1995) 153030t.
- 204 Fukunaga, K., Takama, K. and Suzuki, T.: High-performance liquid chromatographic determination of plasma malondialdehyde level without a solvent extraction procedure. *Anal. Biochem.*, 230 (1995) 20-23.
- 205 Lindahl, R. and Levin, J.-O.: Laboratory validation of a diffusive sampler for the determination of glutaraldehyde in air. *J. Chromatogr. A*, 710 (1995) 175-180.
- 206 Nishikawa, H. and Sakai, T.: Derivatization and chromatographic determination of aldehydes in gaseous and air samples. *J. Chromatogr. A*, 710 (1995) 159-165 - a review with 59 refs.
- 207 Nuijens, M.J., Zomer, M., Mank, A.J.G., Gooijer, C., Velthorst, N.H. and Hofstraat, J.W.: A cyanine fluorophore with a hydrazide functionality as labelling reagent for aldehydes in liquid chromatography. *Anal. Chim. Acta*, 311 (1995) 47-55.
- 208 Pina, E., Sousa, A.T. and Brojo, A.P.: HPLC method development and validation for formaldehyde in enteric coating of hard gelatin capsules. *J. Liq. Chromatogr.*, 18 (1995) 2683-2693.
- 209 Poon, W.W., Marbois, B.N., Faull, K.F. and Clarke, C.F.: 3-Hexaprenyl-4-hydroxybezoic acid forms a predominant intermediate pool in ubiquinone biosynthesis in *Saccharomyces cerevisiae*. *Arch. Biochem. Biophys.*, 320 (1995) 305-314.
- 210 Robison, T.W., Forman, H.J. and Thomas, M.J.: Release of aldehydes from rat alveolar macrophages exposed *in vitro* to low concentrations of nitrogen dioxide. *Biochim. Biophys. Acta*, 1256 (1995) 334-340.

See also 165, 181, 185, 187, 280, 541, 650, 922.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

- 211 Bruno, A.E. and Krattiger, B.: On-column refractive index detection of carbohydrates separated by HPLC and CE. *J. Chromatogr. Libr.*, 58 (1995) 431-446; C.A., 123 (1995) 131595d - a review with 29 refs.
- 212 Johansson, L., Karlsson, H. and Karlsson, K.-A.: Separation and detection of 4-hexadecylaniline maltooligosaccharide derivatives with packed capillary liquid chromatography-frit fast atom bombardment-mass spectrometry. *J. Chromatogr. A*, 712 (1995) 149-154.
- 213 Kohler, M. and Leary, J.A.: LC/MS/MS of carbohydrates with postcolumn addition of metal chlorides using a triaxial electrospray probe. *Anal. Chem.*, 67 (1995) 3501-3508.
- 214 Kugimiya, A., Matsui, J., Takeuchi, T., Yano, K., Muguruma, H., Elgersma, A.V. and Karube, I.: Recognition of sialic acid using molecularly imprinted polymer. *Anal. Lett.*, 28 (1995) 2317-2323.
- 215 Langemeier, J.M. and Rogers, D.E.: Rapid method for sugar analysis of doughs and baked products. *Cereal Chem.*, 72 (1995) 349-351; C.A., 123 (1995) 110349m.

- 216 Manzella, S.M., Dharmesh, S.M., Beranek, M.C., Swanson, P. and Baenziger, J.U.: Evolutionary conservation of the sulfated oligosaccharides on vertebrate glycoprotein hormones that control circulatory half-life. *J. Biol. Chem.*, 270 (1995) 21665-21671.
- 217 Murata, K., Murata, A. and Yoshida, K.: High-performance liquid chromatographic identification of eight constitutional disaccharides from heparan sulfate isomers digested with heparitinases. *J. Chromatogr. B*, 670 (1995) 3-10.
- 218 Nakayama, H., Matsuoka, K., Shinkai, F., Isobe, T., Kanai, M., Seta, K. and Okuyama, T.: (Analysis of structure and heterogeneity of oligosaccharides in calreticulin using HPLC-ESI/TSQMS). *Kuromatogurafi*, 16 (1995) 13-21; C.A., 123 (1995) 164570v.
- 219 Nesterowicz, J., Pierzynowska-Korniak, G. and Zadernowski, R.: Application of uridine to the determination of glucopyranosides in faba bean (*Vicia faba*) seeds by high pressure liquid chromatography (HPLC), short report. *Pol. J. Food Nutr. Sci.*, 4 (1995) 82-87; C.A., 123 (1995) 167872f.
- 220 Omichi, K. and Hase, S.: Determination of the linkage positions of reducing-end residues of oligosaccharides by partial periodate oxidation of pyridylaminated derivatives. *Anal. Biochem.*, 227 (1995) 404-407.
- 221 Puanglarp, N., Oxley, D., Currie, G.J., Bacic, A., Craik, D.J. and Yellowless, D.: Structure of the N-linked oligosaccharides from tridacnin, a lectin found in the haemolymph of the giant clam *Hipposus hippocampus*. *Eur. J. Biochem.*, 232 (1995) 873-880.
- 222 Rohrer, J.S.: Separation of asparagine-linked oligosaccharides by high-pH anion-exchange chromatography with pulsed amperometric detection: empirical relationships between oligosaccharide structure and chromatographic retention. *Glycobiology*, 5 (1995) 359-360; C.A., 123 (1995) 106911w.
- 223 Shirane, K., Furukawa, K., Fukuchi, K., Yamazaki, H., Tsuji, M. and Okamoto, Y.: Effects of N-3554S, a polypropenyl phosphate, on B16-F10 mouse melanoma cells. *Biochim. Biophys. Acta*, 1268 (1995) 285-292.
- 224 Sugahara, K., Ohkita, Y., Shibata, Y., Yoshida, K. and Ikegami, A.: Structural studies on the hexasaccharide alditois isolated from the carbohydrate-protein linkage region of dermatan sulfate proteoglycans of bovine aorta. Demonstration of iduronic acid-containing components. *J. Biol. Chem.*, 270 (1995) 7204-7212.
- 225 Thompson, M.: The potential application of ion exclusion chromatography for additional sucrose recovery from molasses. *Proc. Annu. Congr.-S. Afr. Sugar Technol. Assoc.*, 68 (1994) 105-108; C.A., 123 (1995) 173282e.
- 226 Torto, N., Buttler, T., Gorton, L., Marko-Varga, G., Stalbrand, H. and Tjerneld, F.: Monitoring of enzymatic hydrolysis of ivory nut mannan using on-line microdialysis sampling and anion-exchange chromatography with integrated pulsed electrochemical detection. *Anal. Chim. Acta*, 313 (1995) 15-24.
- 227 Wilson, I.B.H., Webberley, M.C., Revers, L. and Flitsch, S.L.: Dolichol is not a necessary moiety for lipid-linked oligosaccharide substrates of the mannosyltransferases involved in *in vitro* N-linked oligosaccharide assembly. *Biochem. J.*, 310 (1995) 909-916.
- See also 36, 103, 231, 235, 245, 582.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 228 Anumula, K.R.: Rapid quantitative determination of sialic acids in glycoproteins by high-performance liquid chromatography with a sensitive fluorescence detection. *Anal. Biochem.*, 230 (1995) 24-30.
- 229 Armstrong, D.C. and Johns, M.R.: Improved molecular weight analysis of streptococcal hyaluronic acid by size exclusion chromatography. *Biotechnol. Tech.*, 9 (1995) 491-496; C.A., 123 (1995) 137805u.
- 230 Bigge, J.C., Patel, T.P., Bruce, J.A., Goulding, P.N., Charles, S.M. and Parekh, R.B.: Nonselective and efficient fluorescent labeling of glycans using 2-amino benzamide and anthranilic acid. *Anal. Biochem.*, 230 (1995) 229-238.
- 231 Boudart, G., de Champ-Guillaume, G., Lafitte, C., Ricart, G., Barthe, J.-P., Mazau, F. and Esquerre-Tugayé, M.-T.: Elicitors and suppressors of hydroxyproline-rich glycoprotein accumulation are solubilized from plant cell walls by endopolagalacturonase. *Eur. J. Biochem.*, 232 (1995) 449-457.
- 232 Chuang, J.-Y.: Size exclusion chromatography of starch. *Chromatogr. Sci. Ser.*, 69 (1995) 381-408; C.A., 123 (1995) 173262y - a review with 69 refs.
- 233 Dall'Amico, R., Montini, G., Pisanello, L., Piovesan, G., Bottaro, S., Cracco, A.T., Zucchello, G. and Zucchello, F.: Determination of inulin in plasma and urine by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 672 (1995) 155-159.
- 234 Hanselmann, R., Buchard, W., Lemmes, R. and Schwengers, D.: Characterization of DEAE-dextran by means of light scattering and combined size-exclusion chromatography/low-angle laser light scattering/viscometry. *Macromol. Chem. Phys.*, 196 (1995) 2259-2275; C.A., 123 (1995) 86447u.
- 235 Lortat-Jacob, H., Turnbull, J.E. and Grimaud, J.-A.: Molecular organization of the interferon γ -binding domain in heparan sulphate. *Biochem. J.*, 310 (1995) 497-505.
- 236 McConville, M.J., Schnur, L.F., Jaffe, C. and Schneider, P.: Structure of *Leishmania* lipophosphoglycan: inter- and intra-specific polymorphism in Old World species. *Biochem. J.*, 310 (1995) 807-818.
- 237 Moseley, R., Waddington, R., Evans, P., Halliwell, B. and Embrey, G.: The chemical modification of glycosaminoglycan structure by oxygen-derived species *in vitro*. *Biochim. Biophys. Acta*, 1244 (1995) 245-252.
- 238 Sakamoto, H.: Colloidal particle- or microparticle-labeled reagent and heparin for immunoassay. *Jpn. Kokai Tokkyo Koho JP 07,151,754 [95,151,754] (Cl. GO1N33/531)*, 16 Jun. 1995, Appl. 93/323,391, 30 Nov. 1993; 10 pp.; C.A., 123 (1995) 138168a.
- 239 Sheng, G.-Q., Kresbach, G., Spiro, M.J. and Spiro, R.G.: Evaluation of the cell specificity and sulfate dependence of glomerular extracellular matrix proteoglycan synthesis. *Arch. Biochem. Biophys.*, 321 (1995) 83-93.
- 240 Thomas, S.S., Plenkiewicz, J., Ison, E.R., Bois, M., Zou, W., Szarek, W.A. and Kisilevsky, R.: Influence of monosaccharide derivatives on liver cell glycosaminoglycan synthesis: 3-deoxy-D-xylo-hexose (3-deoxy-D-galactose) and methyl (methyl 4-chloro-4-deoxy- β -D-galactopyranosid) uronate. *Biochim. Biophys. Acta*, 1272 (1995) 37-48.

For additional information see C.A.:
123 (1995) 110361j, 137768j.

- 241 Varelas, J.B., Kollar, J., Huynh, T.D. and Hering, T.M.: Expression and characterization of a single recombinant proteoglycan tandem repeat domain of link protein that binds zinc and hyaluronate. *Arch. Biochem. Biophys.*, 321 (1995) 21-30.

See also 217.

10c. Glycoproteins and their constituents

- 242 Barry, F.P., Rosenberg, L.C., Gaw, J.U., Gaw, J.U., Koob, T.J. and Neame, P.J.: N- and O-linked keratan sulfate on the hyaluronan binding region of aggrecan from mature and immature bovine cartilage. *J. Biol. Chem.*, 270 (1995) 20516-20524.
- 243 Grabenhorst, E., Hoffmann, A., Nimtz, H., Zettlmeissl, G. and Conradt, H.S.: Construction of stable BHK-21 cells expressing human secretory glycoproteins and human Gal(β1-4)GlcNAc-R α2,6-sialyltransferase. α2,6-Linked NeuAc is preferentially attached to the Gal(β1-4)GlcNAc(β1-2)Man(α1-3)-branch of dian- tennary oligosaccharides from secreted recombinant β-trace proteins. *Eur. J. Biochem.*, 232 (1995) 718-725.
- 244 Hauffe, D.: (Analysis of glycoproteins). *GIT Fachz. Lab.*, 38 (1994) 1220-1224; *C.A.*, 123 (1995) 112718e.
- 245 Kishino, S., Nomura, A., Sugawara, M., Iseki, K., Kakinoki, S., Kitabatake, A. and Miyazaki, K.: Purification method for α-1-acid glycoprotein with subsequent high-performance liquid chromatographic determination of monosaccharides in plasma of healthy subjects and patients with renal insufficiency. *J. Chromatogr. B*, 672 (1995) 199-205.
- 246 Koike, T., Titani, K., Suzuki, M., Beppu, H., Kuzuya, H., Maruta, K., Shimpo, K. and Fujita, K.: The complete amino acid sequence of a mannose-binding lectin from "Kidachi Aleo" (*Aloe arborescens* Miller var. *natalensis* Berger). *Biochem. Biophys. Res. Commun.*, 214 (1995) 163-170.
- 247 Kramerov, A.A., Arbatsky, N.P., Rozovsky, Ya.M., Pochechueva, T.I., Mikhaleva, E.A., Druzhinina, T.N., Gvozdev, V.A. and Shibaev, V.N.: (The glycoprotein from *Drosophila melanogaster* cultured cells contains O-glycosidically linked Gal(β1-3)GalNAc disaccharide units). *Biokhimiya (Moscow)*, 60 (1995) 805-812.
- 248 Srinivasan, S.R., Xu, J.-H., Vijayagopal, P., Radhakrishnamurthy, B. and Berenson, G.S.: Low-density lipoprotein binding affinity of arterial chondroitin sulfate proteoglycan variants modulates cholesterol ester accumulation in macrophages. *Biochim. Biophys. Acta*, 1272 (1995) 61-67.
- 249 Tagaki, J., Aoyama, T., Ueki, S., Ohba, H., Saito, Y. and Lorand, L.: Identification of factor-XIIa-reactive glutamyl residues in the propolyptide of bovine von Willebrand factor. *Eur. J. Biochem.*, 232 (1995) 773-777.
- For additional information see *C.A.*:
123 (1995) 137995f.
- See also 221, 228, 426, 494, 517, 661.
11. ORGANIC ACIDS AND LIPIDS
- 11a. *Organic acids and simple esters*
- 250 Asami, M. and Nakamura, K.-i.: Enantiomeric separation and recognition mechanism of 2-arylpropionic acid derivatives by high-performance liquid chromatography using a chiral column. *Chirality*, 7 (1995) 28-33; *C.A.*, 123 (1995) 169176f.
- 251 Briscoe, J.F.: The determination of formic and acetic acid in chondritic meteorites using exclusion chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9514869, 1994, 160 p.; *C.A.*, 123 (1995) 159786r.
- 252 Carnacho, M., Godessart, N., Antón, R., García, M. and Vila, L.: Interleukin-1 enhances the ability of cultured human umbilical vein endothelial cells to oxidize linoleic acid. *J. Biol. Chem.*, 270 (1995) 17279-17286.
- 253 Gondo, K. and Ito, S.: Acid pretreatment and chromatographic analysis for mevalonic acid determination. *Jpn. Kokai Tokkyo Koho JP 07,159,400 [95,159,400]* (Cl. GO1N33/52), 23 Jun. 1995, Appl. 93/310,538, 10 Dec. 1993; 13 pp.; *C.A.*, 123 (1995) 164659f.
- 254 Gutnikov, G.: Fatty acid profiles of lipid samples. *J. Chromatogr. B*, 671 (1995) 71-89 - a review with 218 refs.
- 255 Helia, O., Paulikova, I., Svajdlenka, E., Devinsky, F., Lacko, I. and Olaszova, E.: The hydrolytic activity of microsomal esterases: Organ dependence and species variability. *Pharmazie*, 50 (1995) 705-706.
- 256 Kuksis, A. and Myher, J.J.: Application of tandem mass spectrometry for the analysis of long-chain carboxylic acids. *J. Chromatogr. B*, 671 (1995) 35-70 - a review with 116 refs.
- 257 Macchia, L., Hamberg, M., Kumlin, M., Butterfield, J.H. and Haeggström, J.Z.: Arachidonic acid metabolism in the human mast cell line HMC-1: 5-lipoxygenase gene expression and biosynthesis of thromboxane. *Biochim. Biophys. Acta*, 1257 (1995) 58-74.
- 258 Mank, A.J.G., Beekman, M.C., Velthorst, N.H., Brinkman, U.A.T., Lingeman, H. and Gooijer, C.: Column liquid chromatography with diode laser-induced fluorescence detection of carboxylic acids after pre-column derivatisation. *Anal. Chim. Acta*, 315 (1995) 209-216.
- 259 Nagabhushana, K.S. and Ravindranath, B.: Efficient medium-scale chromatographic group separation of anacardic acids from solvent-extracted cashew nut (*Anacardium occidentale*) shell liquid. *J. Agric. Food Chem.*, 43 (1995) 2381-2383; *C.A.*, 123 (1995) 147267j.
- 260 Prieur, C., Larroque, M. and Brun, S.: Determination of residual methacrylic acid as a food contaminant potentially released by packaging materials. *Sci. Aliments.*, 15 (1995) 75-81; *C.A.*, 123 (1995) 142066h.
- 261 Röschinger, W., Millington, D.S., Gage, D.A., Huang, Z.-H., Iwamoto, T., Yano, S., Packman, S., Johnston, K., Berry, S.A. and Sweetman, L.: 3-Hydroxyisovalerylcarnitine in patients with deficiency of 3-methylcrotonyl CoA carboxylase. *Clin. Chim. Acta*, 240 (1995) 35-51.
- 262 Shen, Y., Sangiah, S. and Ye, M.Y.: Determination of hydroxyl radical formation in the testes of cadmium-treated mice by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2217-2228.

- 263** Sidelmann, U., Gavaghan, C., Carles, H.A.J., Farrant, R.D., Lindon, J.C., Wilson, I.D. and Nicholson, J.K.: Identification of the positional isomers of 2-fluorobenzoic acid 1-O-acyl glucuronide by directly coupled HPLC-NMR. *Anal. Chem.*, 67 (1995) 3401-3404.
- 264** Sohlenius, A.-K., Wigren, J., Bäckström, K., Andersson, K. and DePierre, J.W.: Synergistic induction of acyl-CoA oxidase activity, an indicator of peroxisome proliferation, by arachidonic acid and retinoic acid in Morris hepatoma 7800C1 cells. *Biochim. Biophys. Acta*, 1258 (1995) 257-264.
- 265** Tapia, M.I., Ochoa de Alda, J.A.G., Llama, M.J. and Serra, J.L.: Determination of 2-oxoglutarate in the presence of citrate and/or isocitrate after ion-exclusion high-performance liquid chromatography. *Anal. Lett.*, 28 (1995) 1959-1971.
- 266** Teichert, J. and Preiss, R.: Determination of lipoic acid in human plasma by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 672 (1995) 277-281.
- 267** Toyo'oka, T.: Use of derivatization to improve the chromatographic properties and detection selectivity of physiologically important carboxylic acids. *J. Chromatogr. B*, 671 (1995) 91-112 - a review with 127 refs.
- 268** VanRollins, M., Kochanek, P.M., Evans, R.W., Schiding, J.K. and Nemoto, E.M.: Optimization of epoxyeicosatrienoic acid syntheses to test their effects on cerebral blood flow *in vivo*. *Biochim. Biophys. Acta*, 1256 (1995) 263-274.
- 269** Wheelan, P. and Murphy, R.C.: Metabolism of 6-trans-isomers of leukotriene B₄ in cultured hepatoma cells and in human polymorphonuclear leukocytes. Identification of a Δ⁶-reductase metabolic pathway. *J. Biol. Chem.*, 270 (1995) 19845-19852.
- 270** Zdrobovchenko, G.M., Veremeychenko, S.N. and Solyanik, I.P.: (The fatty acid content and components of the hydrophilic moiety of lipid A of *Pseudomonas syringae* lipopolysaccharides). *Biokhimiya (Moscow)*, 60 (1995) 1039-1047.
- 271** Zosmer, A., Rendell, N.B., Taylor, G.W., Elder, M.G. and Sullivan, M.H.F.: Formation and metabolism of 14,15-epoxyeicosatrienoic acid by human reproductive tissues. *Biochim. Biophys. Acta*, 1258 (1995) 234-240.
- For additional information see C.A.:
123 (1995) 65930f, 137786p.
- See also 57, 64, 175, 189, 203, 209, 326, 532, 545, 649.
- 11b. Prostaglandins**
- 272** Demin, P., Reynaud, D. and Pace-Asciak, C.R.: High-performance liquid chromatographic separation of fluorescent esters of hepxolin enantiomers on a chiral stationary phase. *J. Chromatogr. B*, 672 (1995) 282-289.
- See also 271.
- 11c. Lipids and their constituents**
- 273** Aitzetmueller, K.: HPLC of triglycerides: identification of palm oils and palm oil fractions. Survey by a working party of the DGF, 142th report: German standard methods for investigation of fats fatty products, tensides, and related materials, 109th report: analysis of fats XXXII. *Fat. Sci. Technol.*, 97 (1995) 127-131; C.A., 123 (1995) 173325w.
- 274** Al-Rashood, K.A., Abdel-Moety, E.M., Rauf, A., Abou-Shaab, R.R. and Al-Khamis, K.I.: Triacylglycerols-profiling by high performance liquid chromatography: a tool for detection of pork fat (lard) in processed foods. *J. Liq. Chromatogr.*, 18 (1995) 2661-2673.
- 275** Anspach, F.B. and Hilbeck, O.: Removal of endotoxins by affinity sorbents. *J. Chromatogr. A*, 711 (1995) 81-92.
- 276** Bernhard, W., Linck, M.H. and Creutzburg, H.: Separation of the principal phospholipid classes by high-pressure liquid chromatography. *Ger. DE 4,409,987 (Cl. GO1N30/88)*, 06 Jul. 1995, Appl. 23 Mar. 1994; 13 pp.; C.A., 123 (1995) 164658e.
- 277** Bünger, H. and Pison, U.: Quantitative analysis of pulmonary surfactant phospholipids by high-performance liquid chromatography and light-scattering detection. *J. Chromatogr. B*, 672 (1995) 25-31.
- 278** Dobson, G., Christie, W.W. and Nikolova-Damyanova, B.: Silver ion chromatography of lipids and fatty acids. *J. Chromatogr. B*, 671 (1995) 197-222 - a review with 230 refs.
- 279** Evans, D.M., Williams, K.P., Parsons, G. and Jindal, S.: A tandem-column chromatographic method for studying the interaction between ligands and their targets: lipopolysaccharide as a model. *Anal. Biochem.*, 229 (1995) 42-47.
- 280** Kinter, M.: Analytical technologies for lipid oxidation products analysis. *J. Chromatogr. B*, 671 (1995) 223-236 - a review with 81 refs.
- 281** Korytowski, W., Geiger, P.G. and Girotti, A.W.: High-performance liquid chromatography with mercury cathode electrochemical detection: application to lipid hydroperoxide analysis. *J. Chromatogr. B*, 670 (1995) 189-197.
- 282** Lipp, M.: Review of methods for the analysis of triglycerides in milk fat: application for studies of milk quality and adulteration. *Food Chem.*, 54 (1995) 213-221; C.A., 123 (1995) 141978v - a review with 125 refs.
- 283** Lotze, A., Stroud, C.Y. and Soldin, S.J.: Serial lecithin/sphingomyelin ratios and surfactant/albumin ratios in tracheal aspirates from term infants with respiratory failure receiving extracorporeal membrane oxygenation. *Clin. Chem. (Washington)*, 41 (1995) 1182-1188.
- 284** Myher, J.J. and Kuksis, A.: General strategies in chromatographic analysis of lipids. *J. Chromatogr. B*, 671 (1995) 3-33 - a review with 171 refs.
- 285** Nogueira, J.M.F., Pereira, J.L.C. and Sandra, P.: Chromatographic methods for the analysis of crude tall-oil. *J. High Resolut. Chromatogr.*, 18 (1995) 425-432.
- 286** Parlesak, A. and Bode, C.: Lipopolysaccharide determination by reversed-phase high-performance liquid chromatography after fluorescence labeling. *J. Chromatogr. A*, 711 (1995) 277-288.
- 287** Ruiz-Gutiérrez, V. and Barron, L.J.R.: Methods for the analysis of triacylglycerols. *J. Chromatogr. B*, 671 (1995) 133-168 - a review with 327 refs.

288 Singh, A.K. and Jiang, Y.: Quantitative chromatographic analysis of inositol phospholipids and related compounds. *J. Chromatogr. B*, 671 (1995) 255-280 - a review with 130 refs.

289 Torti, M., Ramaschi, G., Montserrat, N., Sinigaglia, F., Balduini, C., Plantavid, M., Breton, M., Chap, H. and Mauro, G.: Evidence for a glycoprotein IIb-IIIa- and aggregation-independent mechanism of phosphatidylinositol 3',4'-bisphosphate synthesis in human platelets. *J. Biol. Chem.*, 270 (1995) 113179-113185.

See also 1, 80, 149, 256, 259, 270, 309.

11d. Lipoproteins and their constituents

290 Bjørkerud, S. and Bjørkerud, B.: Growth-stimulating effect of lipoproteins on human arterial smooth-muscle cells and lung fibroblasts is due to Apo B-containing lipoproteins, type LDL and VLDL, and requires LDL receptors. *Biochim. Biophys. Acta*, 1268 (1995) 237-247.

291 Greenspan, P., Ryu, B.-H., Mao, F. and Gutman, R.L.: Association of negatively-charged phospholipids with low-density lipoprotein (LDL) increases its uptake and the deposition of cholesteryl ester by macrophages. *Biochim. Biophys. Acta*, 1257 (1995) 257-264.

292 Hida, T., Hayashi, K., Yukishige, S., Tanida, S., Kawamura, N. and Harada, S.: Synthesis and biological activities of TAN-1511 analogues. *J. Antibiot.*, 48 (1995) 589-603.

293 McCormick, S.P.A., Day, W.A. and George, P.M.: Characterization of lipoproteins containing a truncated form of apolipoprotein B, apolipoprotein B32. *Biochim. Biophys. Acta*, 1258 (1995) 49-56.

294 Rothwell, T.C., Kananna, V.S., Jin, F.-Y., Koren, E., Foley, T. and Kashyap, M.L.: Characterization of a monoclonal antibody (HB-22) and development of an ELISA for human apolipoprotein A-I. *Clin. Chem. (Washington)*, 41 (1995) 1150-1158.

295 Tadey, T. and Purdy, W.C.: Chromatographic techniques for the isolation and purification of lipoproteins. *J. Chromatogr. B*, 671 (1995) 237-253 - a review with 105 refs.

296 Takizawa, M., Hida, T., Horiguchi, T., Hiramoto, A., Harada, S. and Tanida, S.: TAN-1511 A, B and C, microbial lipopeptides with G-CSF and GM-CSF inducing activity. *J. Antibiot.*, 48 (1995) 579-588.

12. ORGANIC PEROXIDES

See 280, 281.

13. STEROIDS

13a. General techniques

297 Fenske, M.: Rapid and efficient method for extraction and separation of glucocorticosteroids and sex steroids from urines. *J. Chromatogr. B*, 670 (1995) 373-375.

298 Hill, M., Lapčík, O., Hampl, R., Starka, L. and Putz, Z.: Radioimmunoassay of three deoxycorticoids in human plasma following HPLC separation. *Steroids*, 60 (1995) 615-620; *C.A.*, 123 (1995) 160996r.

299 Volin, P.: High-performance liquid chromatographic analysis of corticosteroids. *J. Chromatogr. B*, 671 (1995) 319-340 - a review with 162 refs.

13b. Pregnane and androstane derivatives

300 De Orsi, D., Gagliardi, L., Chimenti, F. and Tonelli, D.: HPLC determination of beclomethasone dipropionate and its degradation products in bulk drug and pharmaceutical formulations. *Anal. Lett.*, 28 (1995) 1655-1663.

301 Gau, Y.S., Sun, S.W. and Chen, R.R.-L.: Optimization of high-performance liquid chromatographic separation for progestogenic, estrogenic, and androgenic steroids using factorial design. *J. Liq. Chromatogr.*, 18 (1995) 2373-2382.

302 Kurosawa, S., Yoshimura, T., Kurosawa, T., Chiba, H., Kobayashi, K., Koike, T. and Tohma, M.: Simultaneous determination of 18-oxygenated corticosteroids by high-performance liquid chromatography with fluorescence detection. *J. Liq. Chromatogr.*, 18 (1995) 2383-2396.

303 Lovgren, U., Johansson, M., Kronqvist, K. and Edholm, L-E.: Biocompatible sample pretreatment for immunochemical techniques using micellar liquid chromatography for separation of corticosteroids. *J. Chromatogr. B*, 672 (1995) 33-44.

304 Milad, M.A., Ludwig, E.A., Anné, S., Middleton, E., Jr. and Jusko, W.J.: Pharmacodynamic model for joint exogenous and endogenous corticosteroid suppression of lymphocyte trafficking. *J. Pharmacokin. Biopharm.*, 22 (1994) 469-480.

305 Okumura, T., Nakajima, Y., Takamatsu, T. and Matsuoka, M.: Column-switching high-performance liquid chromatographic system with a laser-induced fluorimetric detector for direct, automated assay of salivary cortisol. *J. Chromatogr. B*, 670 (1995) 11-20.

306 Sanwald, P., Blankson, E.A., Duléry, B.D., Schoun, J., Huebert, N.D. and Dow, J.: Isocratic high-performance liquid chromatographic method for the separation of testosterone metabolites. *J. Chromatogr. B*, 672 (1995) 207-215.

307 Singh, S.K., Verma, N. and Gupta, R.C.: Sensitive high-performance liquid chromatographic assay method for the determination of guggulsterone in serum. *J. Chromatogr. B*, 670 (1995) 173-176.

308 Venturelli, E., Cavalleri, A. and Secreto, G.: Methods for urinary testosterone analysis. *J. Chromatogr. B*, 671 (1995) 363-380 - a review with 85 refs.

See also 799.

13c. Estrogens

See 297, 301.

13d. Sterols

309 Hoving, E.B.: Chromatographic methods in the analysis of cholesterol and related lipids. *J. Chromatogr. B*, 671 (1995) 341-362 - a review with 113 refs.

310 Lanuzza, F., Micali, G. and Calabro, G.: (Sterol analysis of olive oil by transesterification and HPLC-HRGC coupling). *Riv. Ital. Sostanze Grasse*, 72 (1995) 105-109; *C.A.*, 123 (1995) 142069m.

- 311 Payne, D.W., Shackleton, C., Toms, H., Ben-Shlomo, I., Kol, S., deMoura, M., Strauss, J.F. and Adashi, E.Y.: A novel nonhepatic hydroxycholesterol 7α -hydroxylase that is markedly stimulated by interleukin-1 β . Characterization in the immature rat ovary. *J. Biol. Chem.*, 270 (1995) 18888-18896.

See also 58, 117, 209.

13e. *Bile acids and alcohols*

- 312 Scalia, S.: Bile acid separation. *J. Chromatogr. B*, 671 (1995) 299-317 - a review with 92 refs.

14. STEROID GLYCOSIDES AND SAPONINS

- 313 Kelly, K.L., Kimball, B.A. and Johnston, J.J.: Quantitation of digitoxin, digoxin, and their metabolites by high-performance liquid chromatography using pulsed amperometric detection. *J. Chromatogr. A*, 711 (1995) 289-295.
 314 Mansouri, L. and Henry, M.: An ontogenetic study on the saponin contents of ivy. *Pharmazie*, 50 (1995) 642-643.
 315 Park, M.K., Kim, B.K., Park, J.H., Shin, Y.G. and Cho, K.H.: High performance liquid chromatographic determination of ginsenosides using photoreduction fluorescence detection. *J. Liq. Chromatogr.*, 18 (1995) 2077-2088.
 316 Wolfender, J.-L., Hostettmann, K., Abe, F., Nagao, T., Okabe, H. and Yamauchi, T.: Liquid chromatography combined with thermospray and continuous-flow fast atom bombardment mass spectrometry of glycosides in crude plant extracts. *J. Chromatogr. A*, 712 (1995) 155-168.

See also 822.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. *Terpenes*

- 317 Beutler, J.A., Kashman, Y., Tischler, M., Cardellina, J.H., II, Gray, G.N., Currens, M.J., Wall, M.E., Wani, M.C. and Boyd, M.R.: A reinvestigation of *Maprouena* triterpenes. *J. Natural Prod.*, 58 (1995) 1039-1046.
 318 Cremin, P., Donnelly, D.M.X., Wolfender, J.-L. and Hostettmann, K.: Liquid chromatographic-thermospray mass spectrometric analysis of sesquiterpenes of *Armillaria* (Eumycota: Basidiomycotina) species. *J. Chromatogr. A*, 710 (1995) 273-285.
 319 Debrunner, B., Neuenschwander, M. and Brenneisen, R.: Sesquiterpenes of *Petasites hybridus* (L.) G.M. et Sch.: distribution of sesquiterpenes over plant organs. *Pharm. Acta Helv.*, 70 (1995) 167-173; C.A., 123 (1995) 179609p.
 320 Du, Q., Xiong, X. and Ito, Y.: Separation of bioactive quadri-terpenic acids from the fruit of *Ligustrum lucidum* Ait by high-speed countercurrent. *J. Liq. Chromatogr.*, 18 (1995) 1997-2004.
 321 Garlaschelli, L., Vidari, G., Virtuani, M., Vita-Finzi, P. and Mellerio, G.: The structures of new lanostane triterpenes from the fruiting bodies of *Hebeloma senscens*. *J. Natural Prod.*, 58 (1995) 992-1002.

- 322 Lauren, D.R., Jensen, D.J. and Douglas, J.A.: Analysis of taxol, 10-deacetylbaicatin III and related compounds in *Taxus baccata*. *J. Chromatogr. A*, 712 (1995) 303-309.

- 323 Shibota, H., Sawa, Y., Oka, T.-a., Sonoke, S., Kim, K.K. and Yoshioka, M.: Steviol and steviol-glycoside: glucosyltransferase activities in *Stevia rebaudiana* Bertoni - purification and partial characterization. *Arch. Biochem. Biophys.*, 321 (1995) 390-396.

See also 314, 893, 896, 944.

15c. *Bitter substances*

- 324 Ohnishi, S., Fukamiya, N., Okano, M., Tagahara, K. and Lee, K.-H.: Bruceosides D, E, and F, three new cytotoxic quassinoïd glucosides from *Brucea javanica*. *J. Natural Prod.*, 58 (1995) 1032-1038.

16. NITRO AND NITROSO COMPOUNDS

- 325 Bouvier, E.S.P. and Oehrle, S.A.: Analysis and identification of nitroaromatic and nitramine explosives in water using HPLC and photodiode-array detection. *LC-GC Int.*, 8 (1995) 338-348.
 326 Li, H., Hu, Q., Yang, X. and Zhou, Z.: Direct determination of 4-nitrobenzoyl chloride by high-performance liquid chromatography based on silanophilic interaction. *J. Chromatogr. A*, 710 (1995) 347-350.

See also 180.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. *Amines and polyamines*

- 327 Bagni, N., Mengoli, M. and Matsuzaki, S.: Quantifying polyamines in *Agrobacterium rhizogenes* strains and in Ri plasmid transformed cells. *Methods Mol. Biol. (Totowa)*, 44(Agrobacterium Protocols) (1995) 223-236; C.A., 123 (1995) 164326v.
 328 Brown, K.K., Teass, A.W., Simon, S. and Ward, E.M.: A biological monitoring method for a o-toluidine and aniline in urine using high performance liquid chromatography with electrochemical detection. *Appl. Occup. Environ. Hyg.*, 10 (1995) 557-565; C.A., 123 (1995) 151557w.
 329 Golloch, A., Hagen, J. and Orth, K.: (Aromatic amines in waters). *LaborPraxis*, 19 (1995) 58-63; C.A., 123 (1995) 178868k.
 330 Hyun, M.H., Min, C.S., Cho, Y.J. and Na, M.S.: An improved chiral recognition model for resolving N-acyl- α -arylalkylamines on Pirkle-type π -acidic chiral stationary phases. *J. Liq. Chromatogr.*, 18 (1995) 2527-2541.
 331 Rao, R.N., Bhavani, T. and Husain, S.: Monitoring the levels of 2-amino-4-nitrophenol in industrial process effluents by high performance liquid chromatography. *J. High Resolut. Chromatogr.*, 18 (1995) 449-451.
 332 Wehr, J.B.: Purification of plant polyamines with anion-exchange column clean-up prior to high-performance liquid chromatographic analysis. *J. Chromatogr. A*, 709 (1995) 241-247.

For additional information see C.A.:
123 (1995) 106886s.

See also 59, 64, 350, 358.

17b. Catecholamines and their metabolites

- 333 Clauzing, P., Gough, B., Holson, R.R., Slikker, W., Jr. and Bowyer, J.F.: Amphetamine levels in brain microdialysate, caudate/putamen, substantia nigra and plasma after dosage that produces either behavioral or neurotoxic effects. *J. Pharmacol. Exp. Ther.*, 274 (1995) 614-621.
- 334 Ghindilis, A.L., Michael, N. and Markower, A.: A new sensitive and simple method for detection of catecholamines from adrenal chromaffin cells. *Pharmazie*, 50 (1995) 599-600.
- 335 Huang, T., Yang, L., Gitzen, J., Kissinger, P.T., Vreeke, M. and Heller, A.: Detection of basal acetylcholine in rat brain microdialysate. *J. Chromatogr. B*, 670 (1995) 323-327.
- 336 Jensen, T.B. and Marley, P.D.: Development of an assay for histamine using automated high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 670 (1995) 199-207.
- 337 Kema, I.P., Schoots, C.J.F., Gidding, C.E.M., Okken, A., Aarnoudse, J.G. and Muskiet, F.A.J.: N-Methyladrenaline: age-dependent urinary excretion, perinatal organ content and relation with 'classical' catecholamines. *Clin. Chim. Acta*, 238 (1995) 137-150.
- 338 Kreiss, D.S. and Lucki, I.: Effects of acute and repeated administration of antidepressant drugs on extracellular levels of 5-hydroxytryptamine measured *in vivo*. *J. Pharmacol. Exp. Ther.*, 274 (1995) 866-876.
- 339 Kumar, A.M., Kumar, M., Fernandez, B.J., Goodkin, K., Schneiderman, N. and Eisdorfer, C.: An isocratic HPLC-ECD assay of urinary normetanephrine and metanephrine. *J. Liq. Chromatogr.*, 18 (1995) 2257-2269.
- 340 Kusu, F., Matsumoto, K. and Takamura, K.: Direct separation and determination of synephrine enantiomers by high-performance liquid chromatography with electrochemical detection. *Chem. Pharm. Bull.*, 43 (1995) 1158-1161.
- 341 Niwa, O., Tabei, H., Solomon, B.P., Xie, F. and Kissinger, P.T.: Improved detection limit for catecholamines using liquid chromatography-electrochemistry with a carbon interdigitated array microelectrode. *J. Chromatogr. B*, 670 (1995) 21-28.
- 342 Pugsley, T.A., Davis, M.D., Akunne, H.C., Cooke, L.W., Whetzel, S.Z., MacKenzie, R.G., Shih, Y.-H., van Leeuwen, D.H., Demattos, S.B., et al.: CI-1007, a dopamine partial agonist and potential antipsychotic agent. I. Neurochemical effects. *J. Pharmacol. Exp. Ther.*, 274 (1995) 898-911.
- 343 Sato, M., Nakano, T., Takeuchi, M., Kumagai, T., Kanno, N., Nagahisa, E. and Sato, Y.: Specific determination of histamine in fish by high-performance liquid chromatography after diazo coupling. *Biosci., Biotechnol., Biochem.*, 59 (1995) 1208-1210; C.A., 123 (1995) 142039b.
- 344 Tarrach, F.: (Determination of low quantities of histamine in wine and sparkling wine). *Dtsch. Lebensm.-Rundsch.*, 91 (1995) 73-75; C.A., 123 (1995) 167915x.
- 345 Uchiyama, S., Ito, N. and Hasebe, Y.: Chemically amplified detection of catecholamines by liquid chromatography using an L-ascorbic acid carrier. *Electroanalysis*, 7 (1995) 731-733; C.A., 123 (1995) 160992m.
- 346 Yamazaki, T., Akiyama, T. and Shindo, T.: Routine high-performance liquid chromatographic determination of myocardial interstitial norepinephrine. *J. Chromatogr. B*, 670 (1995) 328-331.
- See also 42, 358, 432.
- 17c. Urea and guanidine derivatives**
- 347 Shintani, H.: Solid phase extraction (SPE) of blood urea compared with liquid-liquid extraction regarding artifact formation. *J. Liq. Chromatogr.*, 18 (1995) 2167-2174.
- 17d. Other amine derivatives and amides (excl. peptides)**
- See 778.
- 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS**
- 18a. Amino acids and their derivatives**
- 348 Brückner, H., Westhauser, T. and Godel, H.: Liquid chromatographic determination of D- and L-amino acids by derivatization with o-phthalaldehyde and N-isobutyryl-L-cysteine. Applications with reference to the analysis of peptidic antibiotics, toxins, drugs and pharmaceutically used amino acids. *J. Chromatogr. A*, 711 (1995) 201-215.
- 349 Carney, J.R. and Rinehart, K.L.: Biosynthesis of brominated tyrosine metabolites by *Aplysina fistularis*. *J. Natural Prod.*, 58 (1995) 971-985.
- 350 Chen, C.-C. and Lin, C.-E.: HPLC separation of enantiomers of amino acids and amino alcohols on ionically bonded chiral stationary phases consisting of cyanuric chloride with amino acid and dialkylamine substituents. *J. Chromatogr. Sci.*, 33 (1995) 229-235.
- 351 Csapó, J., Csapó-Kiss, Z., Csordás, E., Martin, T.G., Folestad, S., Tivesten, A. and Némethy, S.: Rapid method for determination of diaminopimelic acid using ion exchange column chromatography. *Anal. Lett.*, 28 (1995) 2049-2061.
- 352 Diez, S., Zhou, P., Daunert, S. and Valiente, M.: Immobilized soft-metal affinity system for amino acids based on an 8-hydroxyquinoline-Pd(II) complex; characterization using glycine as a model. *Anal. Chim. Acta*, 315 (1995) 339-345.
- 353 Engström, A., Andersson, P.E., Josefsson, B. and Pfeffer, W.D.: Determination of 2-(9-anthryl)ethyl chloroformate-labelled amino acids by capillary electrophoresis and liquid chromatography with absorbance or fluorescence detection. *Anal. Chem.*, 67 (1995) 3018-3022.
- 354 Goodnough, D.B., Lutz, M.P. and Wood, P.L.: Separation and quantification of D- and L-phosphoserine in rat brain using Na-(2,4-dinitro-5-fluorophenyl)-L-alaninamide (Marfey's reagent) by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 672 (1995) 290-294.
- 355 Henle, T., Zehetner, G. and Klostermeyer, H.: Fast and sensitive determination of furosine [in food]. *Z. Lebensm.-Unters. Forsch.*, 200 (1995) 235-237; C.A., 123 (1995) 167911t.

- 356 Huddleston, M.J., Annan, R.S., Bean, M.F. and Carr, S.A.: Selective detection of Thr-, Ser-, and Tyr-phosphopeptides in complex digests by electrospray LC-MS. In: Crabb, J.W. (Editor), *Tech. Protein Chem. V, [Pap. Symp. Protein Soc.], 7th 1993*, Academic, San Diego, 1994, pp. 123-130; C.A., 123 (1995) 138058q.
- 357 Ito, T., Kidouchi, K., Sugiyama, N., Kajita, M., Chiba, T., Niwa, T. and Wada, Y.: Liquid chromatographic-atmospheric pressure chemical ionization mass spectrometric analysis of glycine conjugates and urinary isovalerylglycine in isovaleric acidemia. *J. Chromatogr. B*, 670 (1995) 317-322.
- 358 Krause, I., Bockhardt, A., Neckermann, H., Henle, T. and Klostermeyer, H.: Simultaneous determination of amino acids and biogenic amines by reversed-phase high-performance liquid chromatography of the dabsyl derivatives. *J. Chromatogr. A*, 715 (1995) 67-79.
- 359 Lin, S. and Maddox, N.J.: Direct chromatographic separation of 2-(2-amino-1,3-thiazol-4-yl)methylglycine and its methyl ester enantiomers using a chiral crown ether column. *J. Liq. Chromatogr.*, 18 (1995) 1947-1955.
- 360 Peter, A., Somlai, Cs. and Penke, B.: High-performance liquid chromatographic method for the separation of the optical isomers of γ,γ' -di-tert.-butyl-D,L- γ -carboxyglutamic acid and D,L- γ -carboxyglutamic acid. *J. Chromatogr. A*, 710 (1995) 297-302.
- 361 Rizzi, A.M., Cladrowa-Runge, S., Jonsson, H. and Osla, S.: Enantiomeric resolution of derivatized DL-amino acids by high-performance liquid chromatography using a β -cyclodextrin chiral stationary phase: a comparison between derivatization labels. *J. Chromatogr. A*, 710 (1995) 287-295.
- 362 Scaloni, A., Simmaco, M. and Bossa, F.: D-L-Amino acid analysis using automated precolumn derivatization with 1-fluoro-2,4-dinitrophenyl-5-alanine amide. *Amino Acids*, 8 (1995) 305-313; C.A., 123 (1995) 137793p.
- 363 Spencer, B.J. and Purdy, W.C.: High-performance liquid chromatographic separation of derivatized amino acids enantiomers using modified β -cyclodextrin as a mobile phase additive. *Anal. Lett.*, 28 (1995) 1865-1881.
- 364 Takeuchi, T. and Miwa, T.: Signal enhancement of dansyl amino acids by fluorimetric detection with a packed flow cell in microcolumn liquid chromatography. *Anal. Chim. Acta*, 311 (1995) 231-236.
- 365 Vázquez-Ortiz, F.A., Caire, G., Higuera-Ciapara, I. and Hernández, G.: High performance liquid chromatographic determination of free amino acids in shrimp. *J. Liq. Chromatogr.*, 18 (1995) 2059-2068.
- 366 Wu, N., Webb, A., Peck, T.L. and Sweedler, J.V.: On-line NMR detection of amino acids and peptides in microbore LC. *Anal. Chem.*, 67 (1995) 3101-3107.
- See also 80, 427, 640, 672, 787, 968.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 367 Anjuere, F., Layer, A., Cerottini, J.-C., Servis, C. and Luescher, I.F.: Synthesis of a radioiodinated photoreactive MAGE1 peptide derivative and photoaffinity labeling of cell-associated human leukocyte antigen-A1 molecules. *Anal. Biochem.*, 229 (1995) 61-67.
- 368 Apffel, A., Fischer, S., Goldberg, G., Goodley, P.C. and Kuhlmann, F.E.: Enhanced sensitivity for peptide mapping with electrospray liquid chromatography-mass spectrometry in the presence of signal suppression due to trifluoroacetic acid-containing mobile phases. *J. Chromatogr. A*, 712 (1995) 177-190.
- 369 Authier, F., Mort, J.S., Bell, A.W., Posner, B.I. and Bergeron, J.J.M.: Proteolysis of glucagon within hepatic endosomes by membrane-associated cathepsins B and D. *J. Biol. Chem.*, 270 (1995) 15798-15807.
- 370 Bateman, K.P., Thibault, P., Douglas, D.J. and White, R.L.: Mass spectral analyses of microcysts from toxic cyanobacteria using on-line chromatographic and electrophoretic separations. *J. Chromatogr. A*, 712 (1995) 253-268.
- 371 Bolhagen, R., Schmiedberger, M. and Grell, E.: High-performance liquid chromatographic purification of extremely hydrophobic peptides: transmembrane segments. *J. Chromatogr. A*, 711 (1995) 181-186.
- 372 Brudel, M., Kertscher, U., Schröder, D., Melzig, M.F. and Mehlis, B.: Liquid chromatographic-mass spectrometric studies on the enzymatic degradation of β -endorphin by endothelial cells. *J. Chromatogr. A*, 712 (1995) 169-175.
- 373 Chen, J.-G. and Weber, S.G.: Detection of bioactive oligopeptides after microbore HPLC with electrochemical detection of their Cu(II) complexes: effect of operating parameters on sensitivity and selectivity. *Anal. Chem.*, 67 (1995) 3596-3604.
- 374 Chen, M., Inestrosa, N.C., Ross, G.S. and Fernandez, H.L.: Platelets are the primary source of amyloid β -peptide in human blood. *Biochem. Biophys. Res. Commun.*, 213 (1995) 96-103.
- 375 DiSalvo, J., Bayne, M.L., Conn, G., Kwok, P.W., Trivedi, P.G., Soderman, D.D., Palisi, T.M., Sullivan, K.A. and Thomas, K.A.: Purification and characterization of a naturally occurring vascular endothelial growth factor•placenta growth factor heterodimer. *J. Biol. Chem.*, 270 (1995) 7717-7723.
- 376 Duve, H., Thorpe, A., Scott, A.G., Johnsen, A.H., Rehfeld, J.F., Hines, E. and East, P.D.: The sulfakinins of the fly *Caliphora vomitoria*. Peptide isolation, gene cloning and expression studies. *Eur. J. Biochem.*, 232 (1995) 633-640.
- 377 Eswaran, D., Ali, M.S., Shenoy, B.C., Korotchkina, L.G., Roche, T.E. and Patel, M.S.: Arginine-239 in the beta subunit is at or near the active site of bovine pyruvate dehydrogenase. *Biochim. Biophys. Acta*, 1252 (1995) 203-208.
- 378 Griko, Y.V. and Kapanadze, M.D.: Purification and characterization of human pancreatic polypeptide expressed in *E. coli*. *Biochem. Biophys. Res. Commun.*, 213 (1995) 239-248.
- 379 Hashimoto, T., Kurosawa, K. and Sakura, N.: Structure-activity relationships of neuromedin U. II. Highly potent analogs substituted at the N-terminus of neuromedin U-8. *Chem. Pharm. Bull.*, 43 (1995) 1154-1157.
- 380 Ichiki, Y., Kitamura, K., Kangawa, K., Kawamoto, M., Matsuo, H. and Eto, T.: Distribution and characterization of immunoreactive adrenomedullin in porcine tissue, and isolation of adrenomedullin[26-52] and adrenomedullin[34-52] from porcine duodenum. *J. Biochem. (Tokyo)*, 118 (1995) 765-770.
- 381 Kobayashi, N., Kanai, M., Seta, K. and Nakamura, K.-i.: Quantitative analysis of synthetic human calcitonin by liquid chromatography-mass spectrometry. *J. Chromatogr. B*, 672 (1995) 17-23.

- 382 Kojima, N., Kawabata, S.-i., Makinose, Y., Nishino, N. and Iwanaga, S.: Substrate specificity of rabbit liver metalloendopeptidase and its new fluorogenic peptide substrates. *J. Biochem. (Tokyo)*, 118 (1995) 855-861.
- 383 Lakhiai, H., Muller, D. and Jozefowicz, J.: Interaction mechanisms between insulin and N-acetylneurameric acid in affinity chromatography. *J. Chromatogr. A*, 711 (1995) 93-103.
- 384 Liu, Y.-M., Miao, J.-R. and Toyo'oka, T.: Enantiometric separation of di- and tri-peptides with chiral fluorescence labeling reagents by liquid chromatography. *Anal. Chim. Acta*, 314 (1995) 169-173.
- 385 Lorenz, M.W., Kellner, R. and Hoffmann, K.H.: A family of neuropeptides that inhibit juvenile hormone biosynthesis in the cricket, *Gryllus bimaculatus*. *J. Biol. Chem.*, 270 (1995) 21103-21108.
- 386 Luo, J.-L., Hammarqvist, F., Cotgreave, I.A., Lind, C., Andersson, K. and Werner, J.: Determination of intracellular glutathione in human skeletal muscle by reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 670 (1995) 29-36.
- 387 Ma, N.-T., Wang, Y.-F., Roe, B.A. and Harrison, R.G.: Purification of an L-asparaginase-atrial natriuretic peptide fusion protein expressed in *Escherichia coli*. *Biotechnol. Bioeng.*, 47 (1995) 483-491; C.A., 123 (1995) 110188h.
- 388 McGwire, G.B. and Skidgel, R.A.: Extracellular conversion of epidermal growth factor (EGF) to des-Arg⁵³-EGF by carboxypeptidase M. *J. Biol. Chem.*, 270 (1995) 17154-17158.
- 389 Moore, A.J., Loadman, P.M., Devine, D.A. and Bibby, M.C.: Extraction of the synthetic lytic peptide, cecropin B, from biological fluid and analysis using reversed-phase high-performance liquid chromatography. *J. Chromatogr. B*, 672 (1995) 225-231.
- 390 Moore, A.W., Jr. and Jorgenson, J.W.: Comprehensive three-dimensional separation of peptides using size exclusion chromatography/reversed phase liquid chromatography/optically gated capillary zone electrophoresis. *Anal. Chem.*, 67 (1995) 3456-3463.
- 391 Moore, A.W., Jr. and Jorgenson, J.W.: Rapid comprehensive two-dimensional separations of peptides via RPCL - optically gated capillary zone electrophoresis. *Anal. Chem.*, 67 (1995) 3448-3455.
- 392 Ohguro, H. and Palczewski, K.: Separation of phospho- and non-phosphopeptides using reverse phase column chromatography. *FEBS Lett.*, 368 (1995) 452-454; C.A., 123 (1995) 137808x.
- 393 Ono, S., Yamamoto, Y., Sakai, Y., Yoshimura, T., Shimasaki, C. and Tsukurimichi, E.: Property of chromatography gel on immobilized amphipathic β -structural peptide (II). *Pept. Chem.*, 32nd (1994, Pub. 1995) 117-120; C.A., 123 (1995) 137773g.
- 394 Pridal, L., Ingwersen, S.H., Larsen, F.S., Holst, J.J., Adelhorst, K. and Kirk, O.: Comparison of sandwich enzyme-linked immunoadsorbent assay and radioimmunoassay for determination of exogenous glucagon-like peptide-1(7-36)amide in plasma. *J. Pharm. Biomed. Anal.*, 13 (1995) 841-850.
- 395 Purcell, A.W., Aguilar, M.I. and Hearn, M.T.W.: Conformational effects in reversed-phase high-performance liquid chromatography of polypeptides. I. Resolution of insulin variants. *J. Chromatogr. A*, 711 (1995) 61-70.
- 396 Purcell, A.W., Aguilar, M.I. and Hearn, M.T.W.: Conformational effects in reversed-phase high-performance liquid chromatography of polypeptides. II. The role of insulin A and B chains in the chromatographic behaviour of insulin. *J. Chromatogr. A*, 711 (1995) 71-79.
- 397 Rowley, D.R., Dang, T.D., Larsen, M., Gerdes, M.J., McBride, L. and Lu, B.: Purification of a novel protein (ps20) from urogenital sinus mesenchymal cells with growth inhibitory properties *in vitro*. *J. Biol. Chem.*, 270 (1995) 22058-22065.
- 398 Sakura, N., Kurosawa, K. and Hashimoto, T.: Structure-activity relationships of neuromedin U. I. Contractile activity of dog neuromedin U-related peptides on isolated chicken crop smooth muscle. *Chem. Pharm. Bull.*, 43 (1995) 1148-1153.
- 399 Shimura, K. and Kasai, K.-i.: Fluorescence-labeled peptides as isoelectric point (pI) markers in capillary isoelectric focusing with fluorescence detection. *Electrophoresis (Weinheim)*, 16 (1995) 1479-1484.
- 400 Stevens, J. and Dixon, H.B.F.: The removal of 2-oxoacyl residues from the N-terminus of peptides and cystatin in non-denaturing conditions. *Biochim. Biophys. Acta*, 1252 (1995) 195-202.
- 401 Sun, L. and Carr, P.W.: Mixed-mode retention of peptides on phosphate-modified polybutadiene-coated zirconia. *Anal. Chem.*, 67 (1995) 2517-2523.
- 402 Surendran, N., Ugwu, S.O., Sterling, E.J. and Blanchard, J.: High-performance liquid chromatographic assay for Melanotan-1 ((NLe⁴-DPhen)⁷) α -melanocyte-stimulating hormone) in biological matrices. *J. Chromatogr. B*, 670 (1995) 235-242.
- 403 Suzawa, Y., Hiraoka, B.Y., Harada, M. and Deguchi, T.: High-performance liquid chromatographic determination of prolyl-carboxypeptidase activity in monkey kidney. *J. Chromatogr. B*, 670 (1995) 152-156.
- 404 Veenstra, J.A. and Lambrou, G.: Isolation of a novel RFamide peptide from the mudgut of the american cockroach, *Periplaneta americana*. *Biochem. Biophys. Res. Commun.*, 213 (1995) 519-524.
- 405 Yamaguchi, H., Nakazato, M., Miyazato, M., Kangawa, K., Matsuo, H. and Matsukura, S.: Two novel rat guanylin molecules, guanylin-94 and guanylin-16, do not increase cyclic GMP production in T84 cells. *Biochem. Biophys. Res. Commun.*, 214 (1995) 1204-1210.
- 406 Yan, C.C. and Huxtable, R.J.: Fluorimetric determination of monobromobimane and o-phthalaldehyde adducts of γ -glutamylcysteine and glutathione: application to assay of γ -glutamylcysteinyl synthetase activity and glutathione concentration in liver. *J. Chromatogr. B*, 672 (1995) 217-224.
- 407 Yates, J.R., III, Eng, J.K. and McCormack, A.L.: Mining genomes: correlating tandem mass spectra of modified and unmodified peptides to sequences in nucleotide databases. *Anal. Chem.*, 67 (1995) 3202-3210.

For additional information see C.A.:
123 (1995) 179383k.

See also 59, 80, 85, 116, 210, 366, 506, 511, 823, 867.

18c. Elucidation of structure of proteins and enzymes

- 408 Bendixen, E., Harpel, P.C. and Sottrup-Jensen, L.: Location of the major ϵ -(glutamyl)lysyl cross-linking site in transglutaminase-modified human plasminogen. *J. Biol. Chem.*, 270 (1995) 17929-17933.
- 409 Bertrand, R., Derancourt, J. and Kassab, R.: Production and properties of skeletal myosin subfragment 1 selectively labeled with fluorescein at lysine-553 proximal to the strong actin-binding site. *Biochemistry*, 34 (1995) 9500-9507.
- 410 Casey, E.B., Zhao, H. and Abraham, E.C.: Role of glycine 1 and lysine 2 in the glycation of bovine γ B-crystallin. Site-directed mutagenesis of lysine to threonine. *J. Biol. Chem.*, 270 (1995) 20781-20786.
- 411 Chang, A., Condon, R., Neumann, G.M. and Polya, G.M.: Purification and characterization of a heat-stable wheat substrate for wheat embryo calcium-dependent protein kinase. *Biochim. Biophys. Acta*, 1244 (1995) 317-324.
- 412 Chang, J.: Size of hirudin sequence required to fold into an active core domain. *Biochemistry*, 34 (1995) 11160-11166.
- 413 Chang, L.-S., Lin, J., Kuo, K.-W., Lin, S.-R. and Chang, C.C.: Characterization of epitopes in native and unfolded cobrotoxin: evidence of an immunodominant C-terminal region related to the production of precipitating and non-precipitating antibodies against cobrotoxin. *J. Biochem. (Tokyo)*, 118 (1995) 686-692.
- 414 Fan, N., Rank, K.B., Leone, J.W., Heinrikson, R.L., Bannow, C.A., Smith, C.W., Evans, D.B., Poppe, S.M., Tarpley, W.G., Rothrock, D.J. et al.: The differential processing of homodimers of reverse transcriptases from human immunodeficiency viruses type 1 and 2 is a consequence of the distinct specificities of the viral proteases. *J. Biol. Chem.*, 270 (1995) 13573-13579.
- 415 Hayashi, K., Izu, H., Oda, K., Fukuhara, K.-i., Matsuo, M., Takano, R. and Hara, S.: The primary structure of pepstatin-insensitive carboxyl proteinase produced by *Pseudomonas* sp. No. 101. *J. Biochem. (Tokyo)*, 118 (1995) 738-744.
- 416 Hiraike, N., Sohma, H., Kuroki, Y. and Akino, T.: Epitope mapping for monoclonal antibody against human surfactant protein A (SP-A) that alters receptor binding of SP-A and the SP-A-dependent regulation of phospholipid secretion by alveolar type II cells. *Biochim. Biophys. Acta*, 1257 (1995) 214-222.
- 417 Le Coeur, C., Zhao, Q., Garreau, I., Sannier, F., Maurice, M., Durand, P. and Piot, J.M.: Analytical peptide mapping of a complex yellowfin tuna myoglobin peptic hydrolysate by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2353-2371.
- 418 Maste, M.C.L., Rinia, H.A., Brands, C.M.J., Egmond, M.R. and Norde, W.: Inactivation of a subtilisin in colloidal systems. *Biochim. Biophys. Acta*, 1252 (1995) 261-268.
- 419 Mino, Y.: Protein chemotaxonomy of genus *Datura*. IV. Amino acid sequence of *Datura* ferredoxins depends not on the species but the section of *Datura* plants from which it comes. *Chem. Pharm. Bull.*, 43 (1995) 1186-1189.
- 420 Miyagi, M., Sakiyama, F., Kato, I. and Tsunasawa, S.: Complete covalent structure of porcine liver acylamino acid-releasing enzyme and identification of its active site serine residue. *J. Biochem. (Tokyo)*, 118 (1995) 771-779.
- 421 Nagamine-Natsuka, Y., Norioka, S. and Sakiyama, F.: Molecular cloning, nucleotide sequence, and expression of the gene encoding a trypsin-like protease from *Streptomyces erythraeus*. *J. Biochem. (Tokyo)*, 118 (1995) 338-346.
- 422 Naidoo, N., Cooperman, B.S., Wang, Z., Liu, X. and Rubin, H.: Identification of lysines within α 1-antichymotrypsin important for DNA binding. An unusual combination of DNA-binding elements. *J. Biol. Chem.*, 270 (1995) 14548-14555.
- 423 Ni, R., Tomita, Y., Tokunaga, F., Liang, T.J., Noda, C., Ichihara, A. and Tanaka, K.: Molecular cloning of two types of cDNA encoding subunit RC6-I of rat proteasomes. *Biochim. Biophys. Acta*, 1264 (1995) 45-52.
- 424 Qian, M.G. and Lubman, D.M.: Analysis of tryptic digests using microbore HPLC with an ion trap storage/reflection time-of flight detector. *Anal. Chem.*, 67 (1995) 2870-2877.
- 425 Qian, Z.-Y., Jollès, P., Migliore-Samour, D., Schoentgen, F. and Fiat, A.-M.: Sheep κ -casein peptides inhibit platelet aggregation. *Biochim. Biophys. Acta*, 1244 (1995) 411-417.
- 426 Saito, T., Kawabata, S., Hirata, M. and Iwanaga, S.: A novel type of limulus lectin-L6. Purification, primary structure, and antibacterial activity. *J. Biol. Chem.*, 270 (1995) 14493-14499.
- 427 Weigt, C., Korte, H., von Strandmann, R.P., Hengstenberg, W. and Meyer, H.E.: Identification of phosphocysteine by electrospray mass spectrometry combined with Edman degradation. *J. Chromatogr. A*, 712 (1995) 141-147.
- 428 Wu, J. and Eyre, D.R.: Structural analysis of cross-linking domains in cartilage type XI collagen. Insights on polymeric assembly. *J. Biol. Chem.*, 270 (1995) 18865-18870.
- 429 Yamaguchi, T. and Masada, M.: Comparative studies on cysteine synthase isozymes from spinach leaves. *Biochim. Biophys. Acta*, 1251 (1995) 91-98.
- 430 Yuan, C. and Borchardt, R.T.: Photoaffinity labeling of human placental S-adenosylhomocysteine hydrolase with [2- 3 H]8-acoxy-adenosine. *J. Biol. Chem.*, 270 (1995) 16140-16146.

See also 368, 369, 377, 445, 469, 498.

19. PROTEINS

19a. General techniques

- 431 Antia, F.D., Fellegvari, I. and Horvath, C.: Displacement of proteins in hydrophobic interaction chromatography. *Ind. Eng. Chem. Res.*, 34 (1995) 2796-2804; C.A., 123 (1995) 106889v.
- 432 Burke, W.J., Schmitt, C.A., Li, S.W., Gillespie, K.N., Park, D.H. and Joh, T.H.: Simultaneous measurement of enzyme activity, protein, mRNA, and metabolites in small samples. *Anal. Biochem.*, 230 (1995) 345-348.
- 433 Frantzen, F., Grimsrud, K., Heggli, D.-E. and Sundrehagen, E.: Protein-boronic acid conjugates and their binding to low-molecular-mass *cis*-diols and glycated hemoglobin. *J. Chromatogr. B*, 670 (1995) 37-45.
- 434 Gonzalez-Lara, R. and Gonzalez, L.M.: Reversed-phase HPLC: determination of variety musts by SPE/RP-HPLC analysis of their soluble proteins. *Mater. Eng.*, 9 (1995) 217-221; C.A., 123 (1995) 106761x - a review with 11 refs.

- 435 Heeter, G.A. and Liapis, A.I.: Perfusion chromatography: performance of periodic countercurrent column operation and its comparison with fixed-bed operation. *J. Chromatogr. A*, 711 (1995) 3-21.
- 436 Hill, C.R.: Immunoaffinity purification with monoclonal antibodies. In: Birch, J.R. and Lennox, E.S. (Editors), *Monodonal Antibodies: Princ. Appl.*, Wiley-Liss, New York, 1995, pp. 121-136; C.A., 123 (1995) 106760w - a review with 75 refs.
- 437 Huang, P.Y. and Carbonell, R.G.: Affinity purification of proteins using ligands derived from peptide libraries. *Biotechnol. Bioeng.*, 47 (1995) 288-297; C.A., 123 (1995) 106897w.
- 438 Krook, M., Lindblad, C., Birnbaum, S., Naess, H., Eriksen, J.A. and Mosbach, K.: Selection of peptides with surface affinity for α -chymotrypsin using a phage display library. *J. Chromatogr. A*, 711 (1995) 119-128.
- 439 Lemmo, A.V.: Instrumentation for two dimensional liquid chromatography-capillary zone electrophoresis for the separation of protein mixtures (liquid chromatography capillary zone electrophoresis, protein separation). Avail. *Univ. Microfilms Int.*, Order No. DA9523069, 1994, 325 p.; C.A., 123 (1995) 187063w.
- 440 Lundell, N.: A peptide in a haystack - strategies for a biochromatographer. *LC-GC Int.*, 8 (1995) 636-647 - a review with 109 refs.
- 441 Osborn, B.L. and Abramson, F.P.: Stable-isotope dilution studies of an intact protein using high-performance liquid chromatography/chemical reaction interface mass spectrometry. *Anal. Biochem.*, 229 (1995) 347-350.
- 442 Ovalle, R.: Nonideal size-exclusion chromatography with zwitterions causes a pI-dependent elution of proteins from glycerol propylsilane-modified silica. *Anal. Biochem.*, 229 (1995) 1-7.
- 443 Sun, L. and Carr, P.W.: Chromatography of proteins using polybutadiene-coated zirconia. *Anal. Chem.*, 67 (1995) 3717-3721.
- 444 Xie, J., Aguilar, M.-I. and Hearn, M.T.W.: Studies on the adsorption capacities of proteins with a tentacle-type ion exchanger and their relationship to the stoichiometric retention parameter Z_c^1 . *J. Chromatogr. A*, 711 (1995) 43-52.
- For additional information see C.A.:
123 (1995) 13772f.
- See also 112, 143, 395, 396, 948.
- 19b. Proteins of cells, viruses and subcellular particles**

- 445 Dumont-Misopein, A., Lavergne, J.-P. and Reboud, J.-P.: Properties of elongation factor-2 fragments obtained by partial proteolysis. *Biochim. Biophys. Acta*, 1263 (1995) 221-227.
- 446 Eliseikina, I.A., Agalarov, S.Ch., Muranova, T.A., Markova, L.F., Kashparov, I.A., Avlyakulov, N.K. and Garber, M.B.: (Preparative isolation of proteins from ribosomal 30S subparticles of *Thermus thermophilus* under non-denaturing conditions). *Biokhimiya (Moscow)*, 60 (1995) 1720-1730.
- 447 Gupta, A., Mal, T.K., Jayasuryan, N. and Chauhan, V.S.: Assignment of disulphide bonds in the X protein (HBx) of hepatitis B virus. *Biochem. Biophys. Res. Commun.*, 212 (1995) 919-924.
- 448 Itoh, H., Kobayashi, R., Wakui, H., Komatsuda, A., Ohtani, H., Miura, A.B., Otaka, M., Masamune, O., Andoh, H., Koyama, K., Sato, Y. and Tashima, Y.: Mammalian 60-kDa stress protein (chaperonin homolog). Identification, biochemical properties, and localization. *J. Biol. Chem.*, 270 (1995) 13429-13435.

- 449 Kalasz, H. and Báthori, M.: Chromatographic techniques in studies of the calcium channels and of calcium antagonists in physiological cells. *TrAC*, 14 (1995) 442-449.
- 450 Lambert, W.J. and Stamper, G.F.: Development of an analytical reversed-phase high-performance liquid chromatography assay for transforming growth factor β 3. *J. Chromatogr. A*, 709 (1995) 249-256.
- 451 Löster, K., Baum, O., Hofmann, W. and Reutter, W.: Characterization of molecular aggregates of $\alpha_1\beta_1$ -integrin and other rat liver membrane proteins by combination of size-exclusion chromatography and chemical cross-linking. *J. Chromatogr. A*, 711 (1995) 187-199.
- 452 Quadros, E.V. and Jacobsen, D.W.: The dynamics of cobalamin utilization in L-1210 mouse leukemia cells: a model of cellular cobalamin metabolism. *Biochim. Biophys. Acta*, 1244 (1995) 395-403.
- 453 Tateishi, S., Mori, S., Sugano, T., Hori, N., Ohtsuka, E. and Yamaizumi, M.: Separation of protein factors that correct the defects in the seven complementation groups of xeroderma pigmentosum cells. *J. Biochem. (Tokyo)*, 118 (1995) 819-824.

For additional information see C.A.:
123 (1995) 102746y, 187379d.

See also 456.

- 19c. *Proteins synthesized by genetic manipulation, monodonal antibodies*
- 454 Anspach, F.B., Spille, H. and Rinas, U.: Purification of recombinant human basic fibroblast growth factor: stability of selective sorbents under cleaning in place conditions. *J. Chromatogr. A*, 711 (1995) 129-139.
- 455 Denisov, A.A., Torsky, S.P., Nicolaeva, O.G. and Vologantseva, I.Ya.: (Recombinant human lymphotoxin: synthesis in *Escherichia coli* cells, purification and some properties). *Biokhimiya (Moscow)*, 60 (1995) 1403-1410.
- 456 Gill, D.S., Roush, D.J., Shick, K.A. and Willson, R.C.: Microcalorimetric characterization of the anion-exchange adsorption of recombinant cytochrome b_5 and its surface-charge mutants. *J. Chromatogr. A*, 715 (1995) 81-93.
- 457 Kroher, M., Raffioni, S. and Steele, R.E.: Single step purification of biologically active recombinant rat basic fibroblast growth factor by immobilized metal affinity chromatography. *Biochim. Biophys. Acta*, 1250 (1995) 29-34.
- 458 Nitta, I., Ueda, T., Nojima, T. and Watanabe, K.: Template-dependent polypeptide synthesis in a factor- and energy-free translation system promoted by pyridine. *J. Biochem. (Tokyo)*, 118 (1995) 841-849.
- 459 Serres, A., Muller, D. and Jozefowicz, J.: Purification of monoclonal antibodies on dextran-coated silica support grafted by thiophilic ligand. *J. Chromatogr. A*, 711 (1995) 151-157.
- 460 Toyoguchi, T., Yamaguchi, K., Imajoh-Ohmi, S., Kato, N., Kusunoki, M., Kageyama, H., Sakiyama, S., Nagasawa, S., Moriya, H. and Sakiyama, H.: Purification and characterization of recombinant hamster tissue complement C1s. *Biochim. Biophys. Acta*, 1250 (1995) 90-96.

- 461 Tsai, P.K., Burke, C.J., Irwin, J.W., Bruner, M.W., Tung, J.-S., Hollis, G.F., Mark, G.E., Kessler, II, J.A., Boots, L.J., Conley, A.J. and Middaugh, C.R.: Enhancing the avidity of a human recombinant anti-HIV-1 monoclonal antibody through oligomerization. *J. Pharm. Sci.*, 84 (1995) 866-870.

For additional information see C.A.:
123 (1995) 103819m, 141040h.

See also 387, 421, 423, 438, 465.

19d. Microbial and plant proteins

- 462 Blennow, A., Surin, B.P., Ehring, H., McLennan, N.F. and Spangfort, M.D.: Isolation and biochemical characterization of highly purified *Escherichia coli* molecular chaperone Cpn60 (GroEL) by affinity chromatography and urea-induced monomerization. *Biochim. Biophys. Acta*, 1252 (1995) 69-78.

- 463 Dai, R., Yamazaki, T., Yamazaki, I. and Song, P.-S.: Initial spectroscopic characterization of the ciliate photoreceptor stenotrin. *Biochim. Biophys. Acta*, 1231 (1995) 58-68.

- 464 Heesche-Wagner, K., Schwarz, T., Bartholmes, P. and Kauffmann, M.: Purification of unstable proteins from *Halobacterium salinarium* crude cell extracts: combined cell disruption and desalting by a hollow-fiber membrane module as an access to perform ion-exchange chromatography. *J. Chromatogr. A*, 711 (1995) 175-179.

- 465 Hengen, P.N.: Purification of His-Tag fusion proteins from *Escherichia coli*. *Trends Biochem. Sci.*, 20 (1995) 285-286; C.A., 123 (1995) 106900s.

- 466 Jones, M.L. and Kurzban, G.P.: Noncooperativity of biotin binding to tetrameric streptavidin. *Biochemistry*, 34 (1995) 11750-11756.

- 467 Lichko, L.P.: (H^+ -ATPase and H^+ -pyrophosphatase of the yeast vacuolar membrane). *Biokhimiya (Moscow)*, 60 (1995) 851-863.

- 468 Ling, J., Liu, W.-y. and Wang, T.P.: Simultaneous existence of two types of ribosome-inactivating proteins in the seeds of *Cinnamomum camphora* - characterization of the enzymatic activities of these cytotoxic proteins. *Biochim. Biophys. Acta*, 1252 (1995) 15-22.

- 469 Shimoji, H., Iimura, Y. and Obata, T.: Molecular cloning of CWP1: a gene encoding a *Saccharomyces cerevisiae* cell wall protein solubilized with *Rarobacter faecitabidus* protease I. *J. Biochem. (Tokyo)*, 118 (1995) 302-311.

- 470 Welte, C., Nickel, R. and Wild, A.: Three-dimensional crystallization of the light-harvesting complex from *Mantonella squamata* (Prasinophyceae) requires an adequate purification procedure. *Biochim. Biophys. Acta*, 1231 (1995) 265-274.

See also 514.

19e. Proteins of blood, serum and blood cells

- 471 Agadir, A., Cornic, M., Jerome, M., Menot, M.L., Cambier, N., Gaub, M.P., Gourmel, B., Lefebvre, P., Degos, L. and Chomienne, C.: Characterization of nuclear retinoic acid binding activity in sensitive leukemic cell lines: cell specific uptake of ATRA and RAR α protein modulation. *Biochem. Biophys. Res. Commun.*, 213 (1995) 112-122.

- 472 Bjerrum, O.J., Bjerrum, M.J. and Heegaard, N.H.H.: Electrophoretic and chromatographic differentiation of two forms of albumin in equilibrium at neutral pH: New screening techniques for determination of ligand binding to albumin. *Electrophoresis (Weinheim)*, 16 (1995) 1401-1407.

- 473 Chen, Z., Prestigiacomo, A. and Stamey, T.A.: Purification and characterization of prostate-specific antigen (PSA) complexed to α_1 -antichymotrypsin: potential reference material for international standardization of PSA immunoassays. *Clin. Chem. (Washington)*, 41 (1995) 1273-1282.

- 474 Jacobs, D.B. and Munch, D.C.: Method for modulating specific activity of interferon alpha. *U.S. US 5,434,249* (Cl. 530-351; C07K17/02), 18 Jul. 1995, Appl. 889,136, 27 May 1992; 5 pp.; C.A., 123 (1995) 167611v.

- 475 Leibl, H., Tomasits, R. and Mannhalter, J.W.: Isolation of human serum IgA using thiophilic adsorption chromatography. *Protein Expression Purif.*, 6 (1995) 408-410; C.A., 123 (1995) 141056t.

- 476 Luo, D., Mah, N., Krantz, M., Wilde, K., Wisharat, D., Zhang, Y., Jacobs, F. and Martin, L.: V1-Linker-Vh orientation-dependent expression of single chain Fv containing an engineered disulfide-stabilized bond in the framework regions. *J. Biochem. (Tokyo)*, 118 (1995) 825-831.

- 477 Sakamoto, Y., Kitamura, K., Madison, J., Watkins, S., Laurell, C.-B., Nomura, M., Higashiyama, T. and Putnam, F.W.: Structural study of the glycosylated and unglycosylated forms of a genetic variant of human serum albumin (63 Asp \rightarrow Asn). *Biochim. Biophys. Acta*, 1252 (1995) 209-216.

For additional information see C.A.:
123 (1995) 65811t, 137777m.

See also 19, 416, 455, 459, 460, 493.

19f. Structural and muscle proteins

- 478 Honda, H., Tamura, T., Hatori, K. and Matsuno, K.: Decorating actin filaments with troponin T-I complexes and acceleration of their sliding movement on myosin molecules. *Biochim. Biophys. Acta*, 1251 (1995) 43-47.

- 479 Manzella, S., Ananth, S., Oegema, T.R., Rodén, L., Rosenberg, L.C. and Meezan, E.: Inhibition of glycogenin-catalyzed glycosyl and xylosyl transfer by cytidine 5'-diphosphate and related compounds. *Arch. Biochem. Biophys.*, 320 (1995) 361-368.

- 480 Postnikova, G.B. and Tselikova, S.V.: (Study of ground squirrel myoglobin. Redox reaction between ground squirrel oxymyoglobin and ferricytochrome c). *Biokhimiya (Moscow)*, 60 (1995) 1121-1129.

- 481 Resing, K.A., Johnson, R.S. and Walsh, K.A.: Mass spectrometric analysis of 21 phosphorylation sites in the internal repeat of rat profilaggrin, precursor of an intermediate filament associated protein. *Biochemistry*, 34 (1995) 9477-9487.

- 482 Thulin, C.D. and Walsh, K.A.: Identification of the amino terminus of human filaggrin using differential LC/MS techniques: implications for profilaggrin processing. *Biochemistry*, 34 (1995) 8687-8692.

See also 409, 417, 428.

19g. *Protamines, histones and other nuclear proteins*

- 483 Berger, R.G., Hoffmann, R., Zeppezauer, M., Wagner-Reeder, W., Maljers, L., Ingendoh, A. and Hillenkamp, F.: Separation and characterisation of bovine histone H1 subtypes by combined ion-exchange and reversed-phase chromatography and mass spectrometry. *J. Chromatogr. A.*, 711 (1995) 159-165.

See also 445.

19h. *Chromoproteins and metalloproteins*

- 484 Apostolova, M., Bontchev, P.R., Nachev, Ch. and Sirakova, I.: Analytical determination of metallothioneins in rat liver with respect of their relation to arterial hypertension. *Anal. Lab.*, 4 (1995) 35-37; C.A., 123 (1995) 138096a.
- 485 Bonander, N., Karlsson, B.G. and Vännngård, T.: Disruption of the disulfide bridge in azurin from *Pseudomonas aeruginosa*. *Biochim. Biophys. Acta*, 1251 (1995) 48-54.
- 486 Kildibekov, N.A., Omarov, R.T., Shvetsov, A.A. and L'vov, N.P.: Isolation and purification of the protein containing a molybdenum cofactor from pea *Pisum sativum* L. seeds). *Biokhimiya (Moscow)*, 60 (1995) 1411-1418.
- 487 Koh, T.S., Elgar, D.F. and Ayers, J.S.: Citrate derivatives of cellulose and agarose for the purification of bovine lactoferrin. *Bull. Singapore Natl. Inst. Chem.*, 22 (1994) 75-84; C.A., 123 (1995) 137776k.
- 488 Kouzaki, N., Kawashima, H., Chung, M.C.-M. and Shimizu, S.: Purification and characterization of two forms of cytochrome b from an arachidonic acid-producing fungus, *Mortierella hygrophila*. *Biochim. Biophys. Acta*, 1256 (1995) 319-326.
- 489 Moreau, S., Davies, M.J. and Puppo, A.: Reaction of ferric leghemoglobin with H₂O₂: formation of heme-protein cross-links and dimeric species. *Biochim. Biophys. Acta*, 1251 (1995) 17-22.
- 490 Müller-Schulte, D. and Brunner, H.: Novel magnetic microspheres on the basis of poly(vinyl alcohol) as affinity medium for quantitative detection of glycated haemoglobin. *J. Chromatogr. A*, 711 (1995) 53-60.
- 491 Ohtaki, N. and Inoue, S.: (Chromatographic behavior of transferrins on calcium metaphosphate column). *Kuromatogurafu*, 15 (1994) 282-283; C.A., 123 (1995) 101696v.
- 492 Sannolo, N., Carbone, V., Ferranti, P., Fiume, I., Mamone, G. and Malorni, A.: Mass spectrometric analysis of haemoglobin adducts formed by methyl bromide *in vitro*. *J. Chromatogr. B*, 670 (1995) 349-353.

See also 433.

19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*

- 493 Basheer, A.R., El-Asmar, M.F. and Soslau, G.: Characterization of a potent platelet aggregation inducer from *Cerastes cerastes* (Egyptian sand viper) venom. *Biochim. Biophys. Acta*, 1250 (1995) 97-109.
- 494 Calvete, J.J., Mann, K., Schäfer, W., Sanz, L., Reinert, M., Nessau, S., Raïda, M. and Töpfer-Petersen, E.: Amino acid sequence of HSP-1, a major protein of stallion seminal plasma: effect of glycosylation on its heparin- and gelatin-binding capabilities. *Biochem. J.*, 310 (1995) 615-622.

- 495 Calvete, J.J., Reinert, M., Sanz, L. and Töpfer-Petersen, E.: Effect of glycosylation on the heparin-binding capability of boar and stallion seminal plasma proteins. *J. Chromatogr. A*, 711 (1995) 167-173.

- 496 Drakopoulou, E., Cotton, J., Virelizier, H., Bernardi, E., Schoofs, A.R., Partiseti, M., Choquet, D., Gurrola, G., Possani, L.D. and Vita, C.: Chemical synthesis, structural and functional characterisation of noxiustoxin, a powerful blocker of lymphocyte voltage-dependent K⁺ channels. *Biochem. Biophys. Res. Commun.*, 213 (1995) 901-907.

- 497 Laing, G.D., Kamiguti, A.S., Wilkinson, M.C., Lowe, G.M. and Theakston, R.D.G.: Characterisation of a purified phospholipase A₂ from the venom of the Papuan black snake (*Pseudechis papuanus*). *Biochim. Biophys. Acta*, 1250 (1995) 137-143.

- 498 Lin, S.-R., Chi, S.-H., Chang, L.-S., Kuo, K.-W. and Chang, C.-C.: Chemical modification of cationic groups of a novel α-neurotoxin (Oh-4) from king cobra (*Ophiophagus hannah*) venom. *J. Biochem. (Tokyo)*, 118 (1995) 297-301.

- 499 Macleod, A. and Ozimek, L.: Separation of β-lactoglobulin A by biospecific subunit exchange affinity chromatography. *Milchwissenschaft*, 50 (1995) 303-307; C.A., 123 (1995) 142038a.

- 500 Nagao, Y.: Expression of c-mos protein in cultured rat spermatogenic cells and evidence that rat testicular 43 kDa c-mos protein has an elongated amino acid sequence at the N-terminus. *Biochim. Biophys. Acta*, 1245 (1995) 130-143.

- 501 Visser, S., Slanger, C.J., Lagerwerf, F.M., van Dongen, W.D. and Haverkamp, J.: Identification of a new genetic variant of bovine β-casein using reversed-phase high-performance liquid chromatography and mass spectrometric analysis. *J. Chromatogr. A*, 711 (1995) 141-150.

See also 413, 425, 487.

19j. *Proteins of brain, cerebrospinal fluid and eye*

- 502 Carver, J.A., Guerreiro, N., Nicholls, K.A. and Truscott, R.J.W.: On the interaction of α-crystallin with unfolded proteins. *Biochim. Biophys. Acta*, 1252 (1995) 251-260.
- 503 Kantorow, M., Horwitz, J., van Boekel, M.A.M., de Jong, W.W. and Piatigorsky, J.: Conversion from oligomers to tetramers enhances autop phosphorylation by lens αA-crystallin. Specificity between αA- and αB-crystallin subunits. *J. Biol. Chem.*, 270 (1995) 17215-17220.
- 504 Medvedev, A.E., Kamyshanskaya, N.S., Halket, J., Glover, V. and Sandler, M.: (Brain endogenous monoamine oxidase a inhibitor (tribulin A): purification and identification). *Biokhimiya (Moscow)*, 60 (1995) 659-667.

See also 410.

19l. *Specific binding and receptor proteins*

- 505 Ax, R.L., Miller, D.J. and Winer, M.A.: Detection of heparin-binding seminal plasma proteins. U.S. US 5,434,139 (Cl. 514-21; A61K37/02), 18 Jul. 1995, US Appl. 86,954, 28 Jul. 1989; 26 pp.; C.A., 123 (1995) 164660z.

- 506 Banks, M., Gruber, P., Proudfoot, A.E.I., Arod, C.Y., Allot, B., Bernard, A.R., Sebille, E., McKinnon, M., Wells, T.N.C. and Solari, R.: Soluble interleukin-5 receptor α -chain binding assays: use for screening and analysis of interleukin-5 mutants. *Anal. Biochem.*, 230 (1995) 321-328.
- 507 Cao, J., Ramos-Martinez, J.I. and Villamarín, J.A.: Characterization of a cAMP-binding protein from the bivalve mollusc *Mytilus galloprovincialis*. *Eur. J. Biochem.*, 232 (1995) 664-670.
- 508 Christopher, G.K. and Sundermann, C.A.: Isolation and partial characterization of the insulin binding sites of *Tetrahymena pyriformis*. *Biochem. Biophys. Res. Commun.*, 212 (1995) 515-523.
- 509 Crosby, J., Sherman, D.H., Bibb, M.J., Revill, W.P., Hopwood, D.A. and Simpson, T.J.: Polyketide synthase acyl carrier proteins from *Streptomyces*: expression in *Escherichia coli*, purification and partial characterisation. *Biochim. Biophys. Acta*, 1251 (1995) 32-42.
- 510 De Vries, K.J., Heinrichs, A.A.J., Cunningham, E., Brunink, F., Westerman, J., Somerharju, P.J., Cockcroft, S., Wirtz, K.W.A. and Snoek, G.T.: An isoform of the phosphatidylinositol-transfer protein transfers sphingomyelin and is associated with the Golgi system. *Biochem. J.*, 310 (1995) 643-649.
- 511 Haniu, M., Talvenheimo, J., Le, J., Katta, V., Welcher, A. and Rohde, M.F.: Extracellular domain of neurotrophin receptor trkB: disulfide structure, N-glycosylation sites, and ligand binding. *Arch. Biochem. Biophys.*, 322 (1995) 256-264.
- 512 Kang, S.-H., Xu, X., Heidenreich, O., Gryaznov, S. and Nerenberg, M.: A new affinity purification procedure for DNA-binding proteins using bromoacetyl agarose. *Nucleic Acids Res.*, 23 (1995) 2344-2345; C.A., 123 (1995) 164312n.
- 513 Karthikyan, N. and Thampan, R.V.: The nuclear estrogen receptor R-II of the goat uterus: distinct possibility that the R-II is the deglycosylated form of the nonactivated estrogen receptor (naER). *Arch. Biochem. Biophys.*, 321 (1995) 442-452.
- 514 Liang, O.D., Flock, J.-I. and Wadström, T.: Isolation and characterisation of a vitronectin-binding surface protein from *Staphylococcus aureus*. *Biochim. Biophys. Acta*, 1250 (1995) 110-116.
- 515 O'Keefe, D.O. and Paiva, A.M.: Assay for recombinant hepatitis B surface antigen using reversed-phase high-performance liquid chromatography. *Anal. Biochem.*, 230 (1995) 48-54.
- 516 Okamoto, S., Nakamura, K., Nihira, T. and Yamada, Y.: *Virginiae* butanolide binding protein from *Streptomyces virginiae*. Evidence that VbrA is not the *virginiae* butanolide binding protein and reidentification of the true binding protein. *J. Biol. Chem.*, 270 (1995) 12319-12326.
- 517 Petrescu, S., Branza-Nichita, N., Mita-Lazar, M., Petrescu, A.-J. and Motas, C.: Immunoaffinity chromatography on antibodies immobilized on nitrocellulose powder. *Anal. Biochem.*, 229 (1995) 299-303.
- 518 Pshezhetsky, A.V., Vinogradova, M.V., Elsliger, M.-A., El-Zein, F., Svedas, V.K. and Potier, M.: Continuous spectrophotometric assay of human lysosomal cathepsin A/protective protein in normal and galactosialidosis cells. *Anal. Biochem.*, 230 (1995) 303-307.
- 519 Wimalawansa, S.J.: Purification and biochemical characterization of neuropeptide Y₂ receptor. *J. Biol. Chem.*, 270 (1995) 18523-18530.
- See also 426, 521, 661.
- 19n. Other proteins (incl. proteinous inhibitors of enzymic activity)
- 520 De Filippis, V., Vindigni, A., Altichieri, L. and Fontana, A.: Core domain of hirudin from the leech *Hirudinaria manillensis*: chemical synthesis, purification, and characterization of a Trp₃ analog of fragment 1-47. *Biochemistry*, 34 (1995) 9552-9564.
- 521 Hashimoto, E., Takeuchi, F., Tanaka, Y. and Yamamura, H.: Purification and phosphorylation of a Mr 25,000 protein, an effective phosphate acceptor for casein kinase II and protein kinase C, detected in the cytosolic fraction of *Xenopus laevis* oocytes. *J. Biochem. (Tokyo)*, 118 (1995) 453-460.
- 522 Hatano, K.-i., Kojima, M., Tanokura, M. and Takahashi, K.: Primary structure, sequence-specific ¹N-NMR assignments and secondary structure in solution of bromelain inhibitor VI from pineapple stem. *Eur. J. Biochem.*, 232 (1995) 335-343.
- 523 Jung, H.I., Kim, S.I., Ha, K., Joe, C.O. and Kang, K.W.: Isolation and characterization of guamerin, a new human leukocyte elastase inhibitor from *Hirundo nigropurpurea*. *J. Biol. Chem.*, 270 (1995) 13879-13884.
- 524 Ohba, Y., Goto, Y., Kimura, Y., Suzuki, F., Hisa, T., Takahashi, K. and Takigawa, M.: Purification of an angiogenesis inhibitor from culture medium conditioned by a human chondrosarcoma-derived chondrocytic cell line, HCS-2/8. *Biochim. Biophys. Acta*, 1245 (1995) 1-8.
- 525 Pazhanisamy, S., Stuver, C.M. and Livingston, D.J.: Automation of a high-performance liquid chromatography-based enzyme assay: evaluation of inhibition constants for human immunodeficiency virus-1 protease inhibitors. *Anal. Biochem.*, 229 (1995) 48-53.
- See also 412, 422, 457, 479.
- ## 20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 526 Hsieh, F.Y.L., Tong, X., Wachs, T., Ganem, B. and Henion, J.: Kinetic monitoring of enzymatic reactions in real time by quantitative high-performance liquid chromatography-mass spectrometry. *Anal. Biochem.*, 229 (1995) 20-25.
- See also 432.
- ### 20a. Oxidoreductases
- 527 Barber, M.J., van Valkenburgh, H., Trimboli, A.J., Pollock, V.V., Neame, P.J. and Bastian, N.R.: The amino acid sequence of *Rhodobacter sphaeroides* dimethyl sulfoxide reductase. *Arch. Biochem. Biophys.*, 320 (1995) 266-275.
- 528 Buters, J.T.M., Shou, M., Hardwick, J.P., Korzekwa, K.R. and Gonzalez, F.J.: cDNA-directed expression of human cytochrome P450 CYP1A1 using baculovirus. Purification, dependency on NADPH-P450 oxidoreductase, and reconstitution of catalytic properties without purification. *Drug Metab. Disp.*, 23 (1995) 696-701.
- 529 Galaev, I.Yu., Arvidsson, P. and Mattiasson, B.: Polymer displacement in dye-affinity chromatography. *J. Chromatogr. A*, 710 (1995) 259-266.
- 530 Gasparyan, V.K.: (Flavin-containing polyamine oxidase of the liver: spectral characteristics and localization). *Biokhimiya (Moscow)*, 60 (1995) 1632-1636.

- 531 Hubatsch, I., Maurer, P., Engel, D. and Adolph, H.W.: Preparation and characterization of isozymes and isoforms of horse liver alcohol dehydrogenase. *J. Chromatogr. A.*, 711 (1995) 105-112.
- 532 Husseini, H., Mitrovic, V. and Schlepper, M.: Determination of monoamine oxidase B activity by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B.*, 672 (1995) 138-142.
- 533 Imaoka, S., Hiroi, T., Tamura, Y., Yamazaki, H., Shimada, T., Komori, M., Degawa, M. and Funae, Y.: Mutagenic activation of 3-methoxy-4-aminoazobenzene by mouse renal cytochrome P450 CYP4B1: cloning and characterization of mouse CYP4B1. *Arch. Biochem. Biophys.*, 321 (1995) 255-262.
- 534 Kempf, A.C., Zanger, U.M. and Meyer, U.A.: Truncated human P450 2D6: expression in *Escherichia coli*, Ni²⁺-chelate affinity purification, and characterization of solubility and aggregation. *Arch. Biochem. Biophys.*, 321 (1995) 277-288.
- 535 Mahmoud, Y.A.-G., Abu El Souod, S.M. and Niehaus, W.G.: Purification and characterization of malate dehydrogenase from *Cryptococcus neoformans*. *Arch. Biochem. Biophys.*, 322 (1995) 69-75.
- 536 Montero, C., Duley, J.A., Fairbanks, L.D., McBride, M.B., Micheli, V., Cant, A.J. and Morgan, G.: Demonstration of induction of erythrocyte inosine monophosphate dehydrogenase activity in Ribavirin-treated patients using a high performance liquid chromatography linked method. *Clin. Chim. Acta*, 238 (1995) 169-178.
- 537 Morgunov, I.G. and Il'chenko, A.P.: (Isolation, purification and some properties of glycerol dehydrogenase from *Candida valida* yeast). *Biokhimiya (Moscow)*, 60 (1995) 883-890.
- 538 Regdel, D., Schewe, T. and Kuehn, H.: (Comparative characteristics of lipoxygenase isoenzymes from green pea seeds). *Biokhimiya (Moscow)*, 60 (1995) 953-961.
- 539 Rifkind, A.B., Lee, C., Chang, T.K.H. and Waxman, D.J.: Arachidonic acid metabolism of human cytochrome P450s 2C8, 2C9, 2E1, and 1A2: regioselective oxygenation and evidence for a role for CYP2C enzymes in arachidonic acid epoxidation in human liver microsomes. *Arch. Biochem. Biophys.*, 320 (1995) 380-389.
- 540 Spaldin, V., Madden, S., Adams, D.A., Edwards, R.J., Davies, D.S. and Park, B.K.: Determination of human hepatic cytochrome P4501A2 activity *in vitro* use of tacrine as an isozyme-specific probe. *Drug Metab. Disp.*, 23 (1995) 929-934.
- 541 Steinebach, V., Groen, B.W., Wijmenga, S.S., Niessen, W.M.A., Jongejan, J.A. and Duine, J.A.: Identification of topaquinone, as illustrated for pig kidney diamine oxidase and *Escherichia coli* amine oxidase. *Anal. Biochem.*, 230 (1995) 159-166.
- 542 Sueyoshi, T., Park, L.J., Moore, R., Juvonen, R.O. and Negishi, M.: Molecular engineering of microsomal P450 2a-4 to a stable, water-soluble enzyme. *Arch. Biochem. Biophys.*, 322 (1995) 265-271.
- 543 Varlamov, V.P., Lopatin, S.A., Ilyina, A.V., Bannikova, G.E., Chlenov, M.A., Vasiyarov, G.G., Mitrochkin, A.A., Morgunova, E.Yu., Pierrot, M. and Reglier, M.: New approaches to chromatographic purification of bovine dopamine-β-hydroxylase. *J. Chromatogr. A.*, 711 (1995) 113-118.
- 544 Yamazaki, T., Kominami, S., Nishida, N. and Takemori, S.: Incorporation of bovine adrenal 3β-hydroxy-5-ene steroid dehydrogenase/Δ⁵-Δ⁴ isomerase into phospholipid vesicles. *Biochim. Biophys. Acta*, 1257 (1995) 293-296.
- 545 Yeum, K.-J., Lee-Kim, Y.C., Yoon, S., Lee, K.Y., Park, I.S., Lee, K.S., Kim, B.S., Tang, G., Russell, R.M. and Krinsky, N.I.: Similar metabolites formed from β-carotene by human gastric mucosal homogenates, lipoxygenase, or linoleic acid hydroperoxide. *Arch. Biochem. Biophys.*, 321 (1995) 167-174.
- 546 Zeng, Y., Weiss, C., Yao, T.-T., Huang, J., Siconolfi-Baez, L., Hsu, P. and Rushbrook, J.I.: Isocitrate dehydrogenase from bovine heart: primary structure of subunit 3/4. *Biochem. J.*, 310 (1995) 507-516.
- 20b. *Transferases (excl. E.C. 2.7.-.)*
- 547 Attucci, S., Aitken, S.M., Gulick, P.J. and Ibrahim, R.K.: Farnesyl pyrophosphate synthase from white lupin: molecular cloning, expression, and purification of the expressed protein. *Arch. Biochem. Biophys.*, 321 (1995) 493-500.
- 548 Balducci, E., Orsomando, G., Polzonetti, V., Vita, A., Emanuelli, M., Raffaelli, N., Ruggieri, S., Magni, G. and Natalini, P.: NMN adenyltransferase from bull testis: purification and properties. *Biochem. J.*, 310 (1995) 395-400.
- 549 Broadway, N.M. and Saggerson, E.D.: Solubilization and separation of two distinct carnitine acyltransferases from hepatic microsomes: characterization of the malonyl-CoA-sensitive enzyme. *Biochem. J.*, 310 (1995) 989-995.
- 550 Ciesla, J., Golos, B., Dzik, J.M., Pawelczak, K., Kempny, M., Makowski, M., Bretner, M., Kulikowski, T., Machnicka, B., Rzeszotarska, B. and Rode, W.: Thymidylate synthases from *Hymenolepis diminuta* and regenerating rat liver: purification, properties, and inhibition by substrate and cofactor analogues. *Biochim. Biophys. Acta*, 1249 (1995) 127-136.
- 551 Eccleston, V.S. and Harwood, J.L.: Solubilisation, partial purification and properties of acyl-CoA: glycerol-3-phosphate acyltransferase from avocado (*Persea americana*) fruit mesocarp. *Biochim. Biophys. Acta*, 1257 (1995) 1-10.
- 552 Filippov, M.Yu., Solovyeva, O.N. and Kochetov, G.A.: (Some properties of multiple forms of baker's yeast transketolase). *Biokhimiya (Moscow)*, 60 (1995) 1089-1094.
- 553 Hooper, L.V., Hindsgaul, O. and Baenziger, J.U.: Purification and characterization of the GalNAc-4-sulfotransferase responsible for sulfation of GalNAcβ1,4GlcNAc-bearing. *J. Biol. Chem.*, 270 (1995) 16327-16332.
- 554 Kim, S., Chung, S. and Steinert, P.M.: Highly active soluble processed forms of the transglutaminase I enzyme in epidermal keratinocytes. *J. Biol. Chem.*, 270 (1995) 18026-18035.
- 555 Masutani, H. and Kimura, H.: Purification and characterization of secretory-type GDP-L-fucose:β-D-galactoside 2-α-L-fucosyltransferase from human gastric mucosa. *J. Biochem. (Tokyo)*, 118 (1995) 541-545.
- 556 Rammesmayer, G., Pichorner, H., Adams, P., Jensen, R.G. and Bohnert, H.J.: Characterization of IMT1, myo-inositol O-methyltransferase, from *Mesembryanthemum crystallinum*. *Arch. Biochem. Biophys.*, 322 (1995) 183-188.
- 557 Rouimi, P., Debrauwer, L. and Tulliez, J.: Electrospray ionization-mass spectrometry as a tool for characterization of glutathione S-transferase isozymes. *Anal. Biochem.*, 229 (1995) 304-312.
- 558 Schadewaldt, P., Hummel, W., Wendel, U. and Adelmeyer, F.: Enzymatic method for determination of branched-chain amino acid aminotransferase activity. *Anal. Biochem.*, 230 (1995) 199-204.

- 559 Tollefson, T.O. and Hutchison, III, C.A.: Mammalian DNA (cytosine-5)-methyltransferase expressed in *Escherichia coli*, purified and characterized. *J. Biol. Chem.*, 270 (1995) 18543-18550.
- 560 Vogt, A., Sun, J., Qian, Y., Tan-Chiu, E., Hamilton, A.D. and Sebti, S.M.: Burkitt lymphoma Daudi cells contain two distinct farnesyltransferases with different divalent cation requirements. *Biochemistry*, 34 (1995) 12398-12403.
- 561 Wildersten, M. and Mannervik, B.: Glutathione transferases with novel active sites isolated by phage display from a library of random mutants. *J. Mol. Biol.*, 250 (1995) 115-122; C.A., 123 (1995) 163976v.
- 562 Wragg, S., Hagen, F.K. and Tabak, L.A.: Kinetic analysis of a recombinant UDP-N-acetyl-D-galactosamine: polypeptide N-acetylgalactosaminyltransferase. *J. Biol. Chem.*, 270 (1995) 16947-16954.
- 563 Yoshida, K.-i., Sumi, S., Honda, M., Hosoya, Y., Yano, M., Arai, K. and Ueda, Y.: Serial lectin affinity chromatography demonstrates altered asparagine-linked sugar chain structures of γ -glutamyltransferase in human renal cell carcinoma. *J. Chromatogr. B*, 672 (1995) 45-51.
- 20c. Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)
- 564 Benner, G.E., Dennis, P.B. and Masaracchia, R.A.: Activation of an S6/H4 kinase (PAK 65) from human placenta by intramolecular and intermolecular autophosphorylation. *J. Biol. Chem.*, 270 (1995) 21121-21128.
- 565 Jiang, Y., Zhang, S.-J., Wu, S.-M. and Lee, M.Y.W.T.: Immunoaffinity purification of DNA polymerase δ . *Arch. Biochem. Biophys.*, 320 (1995) 297-304.
- 566 Kitagawa, M., Mukai, H., Shibata, H. and Ono, Y.: Purification and characterization of a fatty acid-activated protein kinase (PKN) from rat testis. *Biochem. J.*, 310 (1995) 657-664.
- 567 Minor, T., Osswald, B., Kraus, T.W., July, N., Isselhard, W. and Klar, E.: Determination of plasma activities of purine nucleoside phosphorylase by high-performance liquid chromatography: estimates of nonparenchymal cell injury after porcine liver transplantation. *J. Chromatogr. B*, 670 (1995) 332-336.
- 568 Trigon, S. and Morange, M.: Different carboxyl-terminal domain kinase activities are induced by heat-shock and arsenite. Characterization of their substrate specificity, separation by Mono Q chromatography, and comparison with the mitogen-activated protein kinases. *J. Biol. Chem.*, 270 (1995) 13091-13098.
- See also 414, 526.
- 20d. Hydrolases, acting on ester bonds (E.C. 3.1.-.-)
- 569 Abdullah, K.M., Leger, S., Perrier, H., Cromlish, W.A., Kennedy, B. and Gresser, M.: Purification of baculovirus-overexpression cytosolic phospholipase A₂ using a single-step affinity column chromatography. *Protein Expression Purif.*, 6 (1995) 291-297; C.A., 123 (1995) 105962b.
- 570 Brown, H.A., Gutowski, S., Kahn, R.A. and Sternweis, P.C.: Partial purification and characterization of Arf-sensitive phospholipase D from porcine brain. *J. Biol. Chem.*, 270 (1995) 14935-14943.
- 571 De Caro, J., Ferrato, F., Verger, R. and de Caro, A.: Purification and molecular characterization of lamb pregastric lipase. *Biochim. Biophys. Acta*, 1252 (1995) 321-329.
- 572 Deprez, P.N., Signorelli, J. and Inestrosa, N.C.: Effect of protamine on the solubilization of collagen-tailed acetylcholinesterase: potential heparin-binding consensus sequences in the tail of the enzyme. *Biochim. Biophys. Acta*, 1252 (1995) 53-58.
- 573 Hara, S., Kudo, I., Komatani, T., Takahashi, K., Nakatani, Y., Natori, Y., Ohshima, M. and Inoue, K.: Detection and purification of two 14 kDa phospholipase A₂ isoforms in rat kidney: their role in eicosanoid synthesis. *Biochim. Biophys. Acta*, 1257 (1995) 11-17.
- 574 Kireyeva, N.N., Bobruskin, I.D. and Severin, S.E.: (cAMP-Specific cyclic nucleotide phosphodiesterase from particulate fraction of human brain). *Biokhimiya (Moscow)*, 60 (1995) 694-708.
- 575 Mirbod, F., Banno, Y., Ghannoum, M.A., Ibrahim, A.S., Nakashima, S., Kitajima, Y., Cole, G.T. and Nozawa, Y.: Purification and characterization of lysophospholipase-transacylase (h-LPTA) from a highly virulent strain of *Candida albicans*. *Biochim. Biophys. Acta*, 1257 (1995) 181-188.
- 576 Moulard, M., Condemine, G., Nasser, W. and Robert-Baudouy, J.: Purification and characterization of the nuclease NucM of *Erwinia chrysanthemi*. *Biochim. Biophys. Acta*, 1262 (1995) 133-138.
- 577 Ncube, I., Gitlesen, T., Adlercreutz, P., Read, J.S. and Mattiasson, B.: Fatty acid selectivity of a lipase purified from *Vernonia galamensis* seed. *Biochim. Biophys. Acta*, 1257 (1995) 149-156.
- 578 Vanderlaan, M., Lotti, R., Siek, G., King, D. and Goldstein, M.: Perfusion immunoassay for acetylcholinesterase: analyte detection based on intrinsic activity. *J. Chromatogr. A*, 711 (1995) 23-31.
- 579 Zelinskaya, N.V., Kovalevskaya, N.P., Matvienko, N.N., Zheleznyaya, L.A. and Matvienko, N.I.: (A site-specific endonuclease and methylase from the thermophilic strain of *Bacillus* species IS4). *Biokhimiya (Moscow)*, 60 (1995) 1435-1449.
- 20e. Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)
- 580 Ito, Y., Nakamura, M., Hotani, T. and Imoto, T.: Insect lysozyme from house fly (*Musca domestica*) larvae: possible digestive function based on sequence and enzymatic properties. *J. Biochem. (Tokyo)*, 118 (1995) 546-551.
- 581 Leah, R., Kigel, J., Svendsen, I. and Mundy, J.: Biochemical and molecular characterization of a barley seed β -glucosidase. *J. Biol. Chem.*, 270 (1995) 15789-15797.
- 582 Takahashi, N., Lee, K.B., Nakagawa, H., Tsukamoto, Y., Kawamura, Y., Li, Y.-T. and Lee, Y.-C.: Enzymatic sialylation of N-linked oligosaccharides using an α -(2,3)-specific *trans*-sialidase from *Trypanosoma cruzi*: structural identification using a three-dimensional elution mapping technique. *Anal. Biochem.*, 230 (1995) 333-342.
- For additional information see C.A.:
123 (1995) 106082b.

20f. Other hydrolases

- 583 Asakura, M., Nagahashi, Y., Hamada, M., Kawai, M., Kadowayashi, K., Narahara, M., Nakagawa, S., Kawai, Y., Hama, T. and Miyake, M.: Purification and properties of β -citryl-L-glutamate-hydrolysing enzyme from rat testis particulate. *Biochim. Biophys. Acta*, 1250 (1995) 35-42.
- 584 Colige, A., Beschin, A., Samyn, B., Goebels, Y., van Beeumen, J.J., Nusgens, B.V. and Lapierre, C.M.: Characterization and partial amino acid sequencing of a 107-kDa procollagen I N-proteinase purified by affinity chromatography on immobilized type XIV collagen. *J. Biol. Chem.*, 270 (1995) 16724-16730.
- 585 Dekeyser, P.M., Buttle, D.J., Devreese, B., van Beeumen, J., Demester, J. and Lauwers, A.: Kinetic constants for the hydrolysis of aggrecan by the papaya proteinases and their relevance for chemonucleolysis. *Arch. Biochem. Biophys.*, 320 (1995) 375-379.
- 586 Jeohn, G., Serizawa, S., Iwamatsu, A. and Takahashi, K.: Isolation and characterization of gastric trypsin from the microsomal fraction of porcine gastric antral mucosa. *J. Biol. Chem.*, 270 (1995) 14748-14755.
- 587 Kabakov, V.E., Klyachko, N.L. and Levashov, A.V.: (Penicillin acylase from *Escherichia coli*: catalytically active subunits). *Biokhimiya (Moscow)*, 60 (1995) 791-797.
- 588 Kawada, A., Hara, K., Hiruma, M., Noguchi, H. and Ishibashi, A.: Rat epidermal cathepsin L-like proteinase: purification and some hydrolytic properties toward filaggrin and synthetic substrates. *J. Biochem. (Tokyo)*, 118 (1995) 332-337.
- 589 Lee, K.B., Matsuoka, K., Nishimura, S.-i. and Lee, Y.C.: A new approach to assay endo-type carbohydrates: bifluorescent-labeled substrates for glycoamidases and ceramide glycanases. *Anal. Biochem.*, 230 (1995) 31-36.
- 590 Lorenz, B., Batel, R., Bachinski, N., Müller, W.E.G. and Schröder, H.C.: Purification and characterization of two exopolyphosphatases from the marine sponge *Tethya lyncurium*. *Biochim. Biophys. Acta*, 1245 (1995) 17-28.
- 591 Molinari, M., Maki, M. and Carafoli, E.: Purification of μ -calpain by a novel affinity chromatography approach. New insights into the mechanism of the interaction of the protease with targets. *J. Biol. Chem.*, 270 (1995) 14576-14581.
- 592 Okamoto, K., Yu, H., Misumi, Y., Ikebara, Y. and Yamamoto, K.: Isolation and sequencing of two sDNA clones encoding rat spleen cathepsin E and analysis of the activation of purified procathepsin E. *Arch. Biochem. Biophys.*, 322 (1995) 103-111.
- 593 Pilote, L., McKercher, G., Thibeault, D. and Lamarre, D.: Enzymatic characterization of purified recombinant human renin. *Biochem. Cell Biol.*, 73 (1995) 163-170.
- 594 Shibatani, T. and Ward, W.F.: Sodium dodecyl sulfate (SDS) activation of the 20S proteasome in rat liver. *Arch. Biochem. Biophys.*, 321 (1995) 160-166.
- 595 Szilagyi, C.M., Sarfati, P., Pradayrol, L. and Morisset, J.: Purification, characterization and substrate specificity of rat pancreatic elastase II. *Biochim. Biophys. Acta*, 1251 (1995) 55-65.
- 596 Woo, S., Lee, J.I., Park, I.K., Yoo, Y.J., Cho, C.M., Kang, M., Ha, D.B., Tanaka, K. and Chung, C.H.: Multiple ubiquitin C-terminal hydrolases from chick skeletal muscle. *J. Biol. Chem.*, 270 (1995) 18766-18773.

- 597 Yada, Y., Higuchi, K. and Imokawa, G.: Purification and biochemical characterization of membrane-bound epidermal ceramidases from guinea pig skin. *J. Biol. Chem.*, 270 (1995) 12677-12684.

See also 403, 408, 415, 421, 423, 430.

20g. Lyases

- 598 Baron, C.B., Ozaki, S., Watanabe, Y., Hirata, M., LaBelle, E.F. and Coburn, R.F.: Inositol 1,4,5-trisphosphate binding to porcine tracheal smooth muscle aldolase. *J. Biol. Chem.*, 270 (1995) 20459-20465.
- 599 Nazaryan, K.B., Simonyan, S.Z., Kazaryan, B.A., Perez, P. and Clement F.: (Purification of human brain enolase and phosphoglycerate mutase and formation of the bienzymatic complex). *Biokhimiya (Moscow)*, 60 (1995) 746-753.
- 600 Savage, T.J., Ichii, H., Hume, S.D., Little, D.B. and Croteau, R.: Monoterpene synthase from gymnosperms and angiosperms: stereospecificity and inactivation by cysteinyl- and arginyl-directed modifying reagents. *Arch. Biochem. Biophys.*, 320 (1995) 257-265.

See also 429.

20h. Isomerases

- 601 Hirtzlin, J., Färber, P.M., Franklin, R.M. and Bell, A.: Molecular and biochemical characterization of a *Plasmodium falciparum* cyclophilin containing a cleavable signal sequence. *Eur. J. Biochem.*, 232 (1995) 765-772.
- 602 Riedel, B., Ueland, P.M. and Svardal, A.M.: Fully automated assay for cobalamin-dependent methylmalonyl CoA mutase. *Clin. Chem. (Washington)*, 41 (1995) 1164-1170.

20i. Ligases

- 603 Kuusksalu, A., Pihlak, A., Müller, W.E.G. and Kelve, M.: The (2'-5')oligoadenylate synthetase is present in the lowest multicellular organisms, the marine sponges. Demonstration of the existence and identification of its reaction products. *Eur. J. Biochem.*, 232 (1995) 351-357.

20j. Complex mixtures and incompletely identified enzymes

- 604 Kastelyanos, O.F., Ermolova, O.V., Sinitsyn, A.P., Popova, N.N., Okunev, O.N., Kerns, G. and Kude, E.: (A scheme for purification of *Penicillium verruculosum* cellulase complex enzymes. Study of biochemical properties and specificity of the enzyme). *Biokhimiya (Moscow)*, 60 (1995) 925-943.
- 605 Wang, P., Royer, M. and Houtz, R.L.: Affinity purification of ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit ϵ N-methyltransferase. *Protein Expression Purif.*, 6 (1995) 528-536; C.A., 123 (1995) 136822k.

See also 420.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 606 Ausserer, W.A. and Biros, M.L.: High-resolution analysis and purification of synthetic oligonucleotides with strong anion-exchange HPLC. *BioTechniques*, 19 (1995) 136-139; C.A., 123 (1995) 106893s.
- 607 Buchko, G.W., Wagner, J.R., Cadet, J., Raoul, S. and Weinfield, M.: Methylene blue-mediated photooxidation of 7,8-dihydro-8-oxo-2'-deoxyguanosine. *Biochim. Biophys. Acta*, 1263 (1995) 17-24.
- 608 De Bellis, G., Invernizzi, L., Debernardi, S., Pergolizzi, R. and Luzzana, M.: Anion-exchange HPLC analysis of biotinylated oligonucleotides. *BioTechniques*, 19 (1995) 230-234; C.A., 123 (1995) 137807w.
- 609 Hagmar, P., Bailey, M., Tong, G., Haralambidis, J., Sawyer, W. and Davidson, B.E.: Synthesis and characterisation of fluorescent oligonucleotides. Effect of internal labelling on protein recognition. *Biochim. Biophys. Acta*, 1244 (1995) 259-268.
- 610 Jacobsson, K.-G., Jurealm, M. and Lindberg, J.E.: Separating hippuric acid from samples of bovine urine provides an improved method to determine allantoin and other purine derivatives. *Swed. J. Agric. Res.*, 25 (1995) 33-36; C.A., 123 (1995) 106894t.
- 611 Keuzenkamp-Jansen, C.W., de Abreu, R.A., Bokkerink, J.P.M. and Trijbels, J.M.F.: Determination of extracellular and intracellular thiopurines and methylthiopurines by high-performance liquid chromatography. *J. Chromatogr. B*, 672 (1995) 53-61.
- 612 Li, K.-M., Todorovic, R., Rogan, E.G., Cavalier, E.L., Ariese, F., Suh, M., Jankowiak, R. and Small, G.J.: Identification and quantitation of dibenzo[a,l]pyrene-DNA adducts formed by rat liver microsomes *in vitro*: preponderance of depurinating adducts. *Biochemistry*, 34 (1995) 8034-8049.
- 613 Liu, Z., Li, T. and Wang, E.: Simultaneous determination of guanine, uric acid, hypoxanthine and xanthine in human plasma by reversed-phase high-performance liquid chromatography with amperometric detection. *Analyst (Cambridge)*, 120 (1995) 2181-2184.
- 614 McNeff, C. and Carr, P.W.: High-performance anion-exchange chromatography of oligonucleotides and oligodeoxynucleotides on quaternized polyethylenimide-coated zirconia. *Anal. Chem.*, 67 (1995) 2350-2353.
- 615 Palmisano, F., Rotunno, T., La Sorsa, M., Zambonin, G.C. and Abbate, I.: Simultaneous determination of pseudouridine, neopterine and creatinine in urin by ion-pair high-performance liquid chromatography with in-series ultraviolet and fluorescence detection. *Analyst (Cambridge)*, 120 (1995) 2185-2189.
- 616 Pezennec, S., Berger, G., Andrianambintsoa, S., Radziszewski, N., Girault, G., Balmiche, J.M. and Bäuerlein, E.: Kinetics of ATP hydrolysis by the F₁-ATPase from *Bacillus PS3*: a reappraisal of the effects of ATP and Mg²⁺. *Biochim. Biophys. Acta*, 1231 (1995) 98-110.
- 617 Sumi, S., Kidouchi, K., Ohba, S. and Wada, Y.: Automated determination of hypoxanthine and xanthine in urine by high-performance liquid chromatography with column switching. *J. Chromatogr. B*, 670 (1995) 376-378.
- 618 Sumi, S., Kidouchi, K., Ohba, S. and Wada, Y.: Automated screening system for purine and pyrimidine metabolism disorders using high-performance liquid chromatography. *J. Chromatogr. B*, 672 (1995) 233-239.
- 619 Wise, C. and Fullerton, F.: Analytical procedure for determination of S-adenosylmethionine, S-adenosylhomocysteine, and S-adenosylethionine in same isocratic HPLC run, with a procedure for preparation and analysis of the analog S-adenosylhomocysteine sulfoxide. *J. Liq. Chromatogr.*, 18 (1995) 1005-1017.
- 620 Xia, Y. and Zweier, J.L.: Substrate control of free radical generation from xanthine oxidase in the postischemic heart. *J. Biol. Chem.*, 270 (1995) 18797-18803.
- 621 Xu, Y.-K., Kaufman, F.R., Donnell, G.N., Giudici, T., Alfi, O. and Ng, W.G.: HPLC analysis of uridine diphosphate sugars: decreased concentrations of uridine diphosphate galactose in erythrocytes and cultured skin fibroblasts from classical galactosemia patients. *Clin. Chim. Acta*, 240 (1995) 21-33.
- 622 Zhan, D.-J., Herreno-Saenz, D., Chiu, L.-H., von Tungeln, L.S., Wu, Y.-S., Lewtas, J. and Fu, P.P.: Separation of ³²P-labeled 3',5'-bisphosphate nucleotides of polycyclic aromatic hydrocarbon anti-diol-epoxides and derivatives. *J. Chromatogr. A*, 710 (1995) 149-157.
- 623 Zuo, S., Boorstein, R.J., Cunningham, R.P. and Teebor, G.W.: Comparison of the effects of UV irradiation on 5-methyl-substituted and unsubstituted pyrimidines in alternating pyrimidine-purine sequences in DNA. *Biochemistry*, 34 (1995) 11582-11590.
- For additional information see C.A.:
123 (1995) 179615n.
- See also 59, 567, 900.
- 21b. *Nucleic acids, RNA*
- 624 Kanduc, D., Bracalelo, A., Mosavi, A. and Farber, E.: Isolation of rat liver tRNA^{Ala}, tRNA^{Gly}, tRNA^{His}, and tRNA^{Ileu} isoacceptor species by HPLC. *Biochem. Arch.*, 11 (1995) 67-72; C.A., 123 (1995) 106968v.
- 625 Kanduc, D., Bracalelo, A. and Mosavi, A.: Isolation of rat liver tRNA^{Asp}, tRNA^{Cys}, tRNA^{Tyr}, and tRNA^{Val} isoacceptor species by HPLC. *Biochem. Arch.*, 11 (1995) 121-127; C.A., 123 (1995) 106892r.
- 626 Van Hille, B., Lohri, A., Reuter, J. and Herrmann, R.: Nonradioactive quantification of mdr1 mRNA by polymerase chain reaction amplification coupled with HPLC. *Clin. Chem. (Washington)*, 41 (1995) 1087-1093.
- See also 432.
- 21c. *Nucleic acids, DNA*
- 627 Vaca, C.E., Fang, J.-L., Mutanen, M. and Valsta, L.: ³²P-Postlabeling determination of DNA adducts of malonaldehyde in humans: total white blood cells and breast tissue. *Carcinogenesis*, 16 (1995) 1847-1851; C.A., 123 (1995) 164487y.

21d. Structural studies on RNA and RNA mapping

- 628 Kowalak, J.A., Bruenger, E. and McCloskey, J.A.: Posttranscriptional modification of the central loop of domain V in *Escherichia coli* 23 S ribosomal RNA. *J. Biol. Chem.*, 270 (1995) 17758-17764.

21e. Structural studies on DNA and DNA mapping

- 629 Chernov, A.V., Zheleznyaya, L.A. and Matvienko, N.I.: (Site-specific endonuclease BspKT8 from the thermophilic strain of *Bacillus species* KT8). *Biokhimiya (Moscow)*, 60 (1995) 1318-1325.

21f. Complex mixtures of nucleic acids and their fragments

- 630 Chaudhary, A.K., Nokubo, M., Oglesby, T.D., Marnett, L.J. and Blair, I.A.: Characterization of endogenous DNA adducts by liquid chromatography/electrospray ionization tandem mass spectrometry. *J. Mass Spectrom.*, 30 (1995) 1157-1166; C.A., 123 (1995) 162835m.

See also 512.

22. ALKALOIDS

- 631 Clauwaert, K., Lambert, W. and de Leenheer, A.: High performance liquid chromatographic determination of cocaine and its main metabolites in biological samples: a review. *J. Liq. Chromatogr.*, 18 (1995) 2097-2114 - a review with 52 refs.
- 632 De Andrade, J.B., Pinheiro, H.L.C., Lopes, A., Martins, S., Amorin, A.M.M. and Branda, A.M.: (Determination of caffeine in beverages by high-performance liquid chromatography (HPLC)). *Quim. Nova*, 18 (1995) 379-381; C.A., 123 (1995) 110346h.
- 633 Deng, Y., Maruyama, W., Dostert, P., Takahashi, T., Kawai, M. and Naoi, M.: Determination of the (R)- and (S)-enantiomers of salsolinol and N-methylsalsolinol by use of a chiral high-performance liquid chromatographic column. *J. Chromatogr. B*, 670 (1995) 47-54.
- 634 Galle, K., Bladt, S. and Wagner, H.: (Mahonia: TLC and HPLC analysis of the bark of *Mahonia aquifolium* and related plants). *Dtsch. Apoth. Ztg.*, 134 (1994) 35-44; C.A., 123 (1995) 153029z.
- 635 Johnston, J.J., Petty, E.E., Okuno, I. and Nolte, D.L.: Comparison of reversed-phase and ion-pair chromatography for the determination of strychnine in animal tissues. *J. Chromatogr. B*, 670 (1995) 73-79.
- 636 Muztar, J., Chari, G., Bhat, R., Ramarao, S. and Vidyasagar, D.: A high-performance liquid chromatographic procedure for the separation of cocaine and some of its metabolites from acepromazine, ketamine, and atropine from serum. *J. Liq. Chromatogr.*, 18 (1995) 2635-2645.
- 637 Pacifici, R.: Determination of morphine and its 3- and 6-glucuronides in serum of heroin addicts by liquid chromatography and mass-spectrometry. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) A177.

- 638 Shao, G., Wang, D.-S., Wu, F., Chen, S.-J. and Luo, X.: Separation and determination of (1)-ephedrine and (d)-pseudoephedrine in plasma by high-performance liquid chromatography with fluorescence detection. *J. Liq. Chromatogr.*, 18 (1995) 2133-2145.

- 639 Sukbuntherng, J., Walters, A., Chow, H.-H. and Mayersohn, M.: Quantitative determination of cocaine, cocaethylene (ethylococaine), and metabolites in plasma and urine by high-performance liquid chromatography. *J. Pharm. Sci.*, 84 (1995) 799-804.

For additional information see C.A.:

123 (1995) 123307w.

See also 935, 940.

23. OTHER SUBSTANCES CONTAINING HETERO CYCLIC NITROGEN

23a. Porphyrins and other pyrroles

- 640 Zamora, R. and Hidalgo, F.J.: Linoleic acid oxidation in the presence of amino compounds produces pyrroles by carbonyl amine reactions. *Biochim. Biophys. Acta*, 1258 (1995) 319-327.

23b. Bile pigments

- 641 Burnett, J.R., Lim, C.W., Mahoney, G.N. and Crooke, M.J.: A method for the separation of delta bilirubin using Cibacron Blue affinity chromatography. *Clin. Chim. Acta*, 239 (1995) 37-46.
- 642 Giulian, J.-M., Dalmasso, C., Millet, V., Unal, D. and Charrel, M.: Influence of protoisomers in bilirubin determination of Kodak Ektachem and Hitachi Analysers in neonatal specimens study of the contribution of structural and configurational isomers. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 503-512.
- 643 Hirota, K.: Urinary excretion of alpha- and beta-isomers of biliverdin-IX in humans. *Biol. Pharm. Bull.*, 18 (1995) 481-484; C.A., 123 (1995) 137811t.
- 644 Nakayama, K.: Differences between enzymatic and diazo methods for measuring direct bilirubin. *Eur. J. Clin. Chem. Clin. Biochem.*, 33 (1995) 513-517.
- 645 Yamaguchi, T., Horio, F., Hashizume, T., Tanaka, M., Ikeda, S., Kakinuma, A. and Nakajima, H.: Bilirubin is oxidized in rats treated with endotoxin and acts as a physiological antioxidant synergistically with ascorbic acid *in vivo*. *Biochem. Biophys. Res. Commun.*, 214 (1995) 11-19.

23c. Indole derivatives and plant hormones (gibberellins)

- 646 Li, J., Chen, G., Webster, J.M. and Czyzewska, E.: Antimicrobial metabolites from a bacterial symbiont. *J. Natural Prod.*, 58 (1995) 1081-1086.
- 647 Stresser, D.M., Williams, D.E., Griffin, D.A. and Bailey, G.S.: Mechanisms of tumor modulation by indole-3-carbinol. Disposition and excretion in male Fischer 344 rats. *Drug Metab. Disp.*, 23 (1995) 965-975.

- 648 Winder, A.J., Odh, G., Rosengren, E. and Rorsman, H.: Fibroblasts co-expressing tyrosinase and the b-protein synthesise both eumelanin and phaeomelanin. *Biochim. Biophys. Acta*, 1268 (1995) 300-310.

See also 181.

23d. Pyridine derivatives

- 649 Cordi, A., Lacoste, J.M., Dudault, J., Espinal, J., Boulanger, M., Broux, O., Husson, B., Volland, J.P. and Mahieu, J.P.: Synthesis of 1,2-diacyl-3-nicotinoyl glycerol derivatives and evaluation of their acute effects on plasma lipids in the rat. *Arzneim.-Forsch.*, 45 (1995) 997-1001.
- 650 Gorbunov, A.A., Kostik, E.I. and Gagarin, S.F.: (Determination of impurity amounts of α -pyridil in α -pyridinol by high-performance liquid chromatography). *Zavod. Lab.*, 61 (1995) 15-16; C.A., 123 (1995) 131714s.

See also 652.

23e. Other N-heterocyclic compounds

- 651 Alam, M.K., Sasaki, M., Watanabe, T. and Maeyama, K.: Simultaneous determination of histamine and N-methylhistamine by high-performance liquid chromatography-chemiluminescence coupled with immobilized diamine oxidase. *Anal. Biochem.*, 229 (1995) 26-34.
- 652 Carlsson, H. and Östman, C.: Retention mechanisms of polycyclic aromatic nitrogen heterocyclics on bonded amino phases in normal-phase liquid chromatography. *J. Chromatogr. A*, 715 (1995) 31-39.
- 653 Mierzwka, R., Patel, M. and Chu, M.: HPLC separation of two novel diketopiperazine isomers on polyvinyl alcohol functionalized silica gel. *J. Liq. Chromatogr.*, 18 (1995) 2019-2027.
- 654 Motohashi, N., Meyer, R., Molnár, J., Párkányi, C. and Fang, X.: Chromatographic determination of benz[c]acridines and related compounds in airborne carcinogens. *J. Chromatogr. A*, 710 (1995) 117-128 - a review with 61 refs.

See also 796, 946.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 655 Altenbach, B. and Giger, W.: Determination of benzene- and naphthalenesulfonates in wastewater by solid-phase extraction with graphitized carbon black and ion-pair liquid chromatography with UV detection. *Anal. Chem.*, 67 (1995) 2325-2333.
- 656 Cole, D.E.C., Evrovski, J. and Pirone, R.: Urinary thiosulfate determined by suppressed ion chromatography with conductometric detection. *J. Chromatogr. B*, 672 (1995) 149-154.
- 657 Kleinman, W.A. and Richie, J.P., Jr.: Determination of thiols and disulfides using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 672 (1995) 73-80.

- 658 Lee, B.H., Bertram, B., Rajca, A., Frank, N., Schmezer, P. and Wiessler, M.: Inhibitory effects of mixed disulfides from disulfiram on the metabolism and genotoxicity of N-nitrosodiethylamine. *Arzneim.-Forsch.*, 45 (1995) 1024-1028.

- 659 Stolcova, M., Kaszonyi, A., Hronec, M. and Liptaj, T.: Reversed-phase high-performance liquid chromatographic method for the quantitative determination of alkylbis(2-benzothiazolylsulphen)amides. *J. Chromatogr. A*, 710 (1995) 351-356.

- 660 Volmer, D., Wilkes, J.G. and Keven, K.: Liquid chromatography/mass spectrometry multiresidue determination of sulfonylureas after online trace enrichment. *Rapid Commun. Mass Spectrom.*, 9 (1995) 767-771; C.A., 123 (1995) 105031x.

For additional information see C.A.:
123 (1995) 179072h.

See also 47, 126, 266, 787, 968.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 661 Alvarez, V., Parodi, A.J. and Couso, R.: Characterization of the mannose 6-phosphate-dependent pathway of lysosomal enzyme routing in an invertebrate. *Biochem. J.*, 310 (1995) 589-595.
- 662 Guse, A.H., Goldwich, A., Weber, K. and Mayr, G.W.: Non-radioactive, isomer-specific inositol phosphate mass determinations: high-performance liquid chromatography-micro-metal-dye detection strongly improves speed and sensitivity of analyses from cells and micro-enzyme assays. *J. Chromatogr. B*, 672 (1995) 189-198.
- 663 Mizuno, H., Morita, M. and Akanuma, H.: Phosphorylation of 1,5-anhydro-D-glucitol in mammalian cells. *J. Biochem. (Tokyo)*, 118 (1995) 411-417.

See also 236, 288, 354, 356, 392, 411, 427, 567, 958, 1004.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

- 664 Compañó, R., Granados, M., Leal, C. and Prat, M.D.: Liquid chromatographic determination of triphenyltin and tributyltin using fluorimetric detection. *Anal. Chim. Acta*, 314 (1995) 175-182.
- 665 Falter, R. and Schöler, H.F.: Determination of mercury species in natural waters at picogram level with on-line RP C18 preconcentration and HPLC-UV-PCO-CVAAS. *Fresenius J. Anal. Chem.*, 353 (1995) 34-38.
- 666 Sato, K., Kohri, M. and Okochi, H.: (The speciation of organotin compounds in seawater by micellar LC/ICP-MS). *Bunseki Kagaku*, 44 (1995) 561-568; C.A., 123 (1995) 122479s.

See also 1002.

26b. Boranes, silanes and related non-metallic compounds

See 60.

26c. Coordination compounds

- 667 Ehrsson, H.C., Wallin, I.B., Andersson, A.S. and Edlund, P.O.: Cisplatin, transplatin, and their hydrated complexes: separation and identification using porous graphitic carbon and electrospray ionization mass spectrometry. *Anal. Chem.*, 67 (1995) 3608-3611.
- 668 Lee, C.-H., Kang, C.-H., Kim, E.-K. and Lee, W.: Separation and elution behaviors of some metal-2-hydroxyarylazopyrazolone chelates by reversed phase high performance liquid chromatography (I). *Anal. Sci. Technol.*, 7 (1994) 103-114; *C.A.*, 123 (1995) 159759j.
- 669 Mazzucotelli, A., Bavastello, V., Magi, E., Rivaro, P. and Tomba, C.: Analysis of gadolinium polyaminopolycarboxylic complexes by HPLC-ultrasonic nebulizer-ICP-AES hyphenated technique. *Anal. Proc.*, 32 (1995) 165-167; *C.A.*, 123 (1995) 187265p.
- 670 Nagaosa, Y. and Mizuyuki, T.: Determination of Cu(II) and Ni(II) as chelates with butyl xanthate by liquid chromatography with electrochemical and spectrophotometric detection. *Anal. Chim. Acta*, 311 (1995) 225-229.
- 671 Shapovalova, E.N., Mishenina, I.V., Bol'shova, T.A. and Shpigun, O.A.: (Ion-pair high-performance liquid chromatography of 4-(5-bromo-2-thiazolylazo)resorcinol and its chelates). *Vestn. Mosk. Univ., Ser. 2: Khim.*, 36 (1995) 46-50; *C.A.*, 123 (1995) 73544y.
- 672 Ye, L. and Lucy, C.A.: Chromatographic determination of metal-EDTA complexes with postcolumn detection using fluorescent ternary complexes. *Anal. Chem.*, 67 (1995) 2534-2538.

See also 979, 991.

27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)
- 673 Aizetmueller, K.: (HPLC of vitamin A-esters in margarine). *Fat Sci. Technol.*, 96 (1994) 279-285; *C.A.*, 123 (1995) 141901q - a review with 36 refs.
- 674 Blanco Gomis, D., Escotet Arias, V., Fidalgo Alvarez, L.E. and Gutierrez Alvarez, D.: Determination of fat-soluble vitamins by capillary liquid chromatography in bovine blood plasma. *Anal. Chim. Acta*, 315 (1995) 177-181.
- 675 Delgado-Zamarreño, M.M., Sanchez-Perez, A., Gomez-Perez, M.C. and Hernandez-Mendez, J.: Automatic determination of liposoluble vitamins in butter and margarine using Triton X-100 aqueous micellar solution by liquid chromatography with electrochemical detection. *Anal. Chim. Acta*, 315 (1995) 201-208.
- 676 Derguini, F., Nakanishi, K., Hämmерling, U., Chu, R., Eppinger, T., Levi, E. and Buck, J.: 13,14-Dihydroxy-retinol, a new bioactive retinol metabolite. *J. Biol. Chem.*, 270 (1995) 18875-18880.
- 677 Dilworth, F.J., Scott, I., Green, A., Strugnell, S., Guo, Y., Roberts, E.A., Kremer, R., Calverley, M.J., Makin, H.I.J. and Jones, G.: Different mechanisms of hydroxylation site selection by liver and kidney cytochrome P450 species (CYP27 and CYP24) involved in vitamin D metabolism. *J. Biol. Chem.*, 270 (1995) 16766-16774.

- 678 Fie, M., Zee, J.A. and Amiot, J.: (Separation and determination of B₁₂ vitamers in dairy products by a high-performance liquid chromatography-radioassay method). *Sci. Aliments*, 14 (1994) 763-775; *C.A.*, 123 (1995) 110321w.
- 679 Hibiya, M., Teradaira, R., Sugimoto, T., Fujita, K. and Nagatsu, T.: Simultaneous determination of N²-(3-aminopropyl)biopterin (oncopterin), biopterin and neopterin by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. B*, 672 (1995) 143-148.
- 680 Kuo B.-S., Mathews, B.S., Stuhler, J.D., Carrel, B.K., Ho, J. and Rose, J.Q.: Disposition and cardioselectivity of MDL 74,405, a vitamin E-like free radical scavenger, in rats and dogs after intravenous infusion. *Drug Metab. Disp.*, 23 (1995) 757-764.
- 681 Kurlandsky, S.B., Gamble, M.V., Ramakrishnan, R. and Blaner, W.S.: Plasma delivery of retinoic acid to tissues in the rat. *J. Biol. Chem.*, 270 (1995) 17850-17857.
- 682 Lykkesfeldt, J., Loft, S. and Poulsen, H.E.: Determination of ascorbic acid and dehydroascorbic acid in plasma by high-performance liquid chromatography with coulometric detection - are there reliable biomarkers of oxidative stress? *Anal. Biochem.*, 229 (1995) 329-335.
- 683 MacCrehan, W.A. and Schönberger, E.: Determination of vitamin K₁ in serum using catalytic-reduction liquid chromatography with fluorescence detection. *J. Chromatogr. B*, 670 (1995) 209-217.
- 684 Sass, J.O., Masgrau, E., Saurat, J.-H. and Nau, H.: Metabolism of oral 9-cis-retinoic acid in the human. Identification of 9-cis-retinoyl-β-glucuronide and 9-cis-4-oxo-retinoyl-β-glucuronide as urinary metabolites. *Drug Metab. Disp.*, 23 (1995) 887-891.
- 685 Shimada, K., Sugaya, K., Kaji, H., Nakatani, I., Mitamura, K. and Tsutsumi, N.: Syntheses and enzymatic hydrolysis of 25-hydroxyvitamin D. *Chem. Pharm. Bull.*, 43 (1995) 1379-1384.
- 686 Shimizu, M., Iwasaki, Y., Ishida, H. and Yamada, S.: Determination of 25-hydroxyvitamin D₃ in human plasma using a non-radioactive tetranorvitamin D analogue as an internal standard. *J. Chromatogr. B*, 672 (1995) 63-71.
- 687 Wyss, R.: Chromatographic and electrophoretic analysis of biomedically important retinoids. *J. Chromatogr. B*, 671 (1995) 381-425 - a review with 313 refs.
- 688 Yeung, B., Vourou, P., Siu-Caldera, M.-L. and Reddy, G.S.: Study of the metabolism of 1,25-dihydroxy-16-ene-vitamin D₃ in rat kidney using online HPLC-electrospray tandem mass spectrometry. *Proc. Workshop Vitam. D*, 9th(Vitamin D) (1994) 196-197; *C.A.*, 123 (1995) 164367j.
- 689 Zarzycki, P.K., Kowalski, P., Nowakowska, J. and Lamparczyk, H.: High-performance liquid chromatographic and capillary electrophoretic determination of free nicotinic acid in human plasma and separation of its metabolites by capillary electrophoresis. *J. Chromatogr. A*, 709 (1995) 203-208.

See also 40, 452, 1001.

28. ANTIBIOTICS

- 690 Abdel-Moety, E.M., Al-Rashood, K.A., Al-Deeb, O.A. and Khattab, N.A.: Stability indicating HPLC method for determination of ampicillin and dicloxacillin in bulk form, admixtures, and capsules. *Sci. Pharm.*, 63 (1995) 7-15; C.A., 123 (1995) 179619s.
- 691 Al-Angary, A.A.: A simple high-performance liquid chromatographic assay for loracarbef in human plasma. *Anal. Lett.*, 28 (1995) 1819-1831.
- 692 Al-Rashood, K.A.: Simultaneous determination of ampicillin and dicloxacillin in pharmaceutical formulations by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2457-2465.
- 693 Chandra, A. and Nair, M.G.: Azalomycin F complex from *Streptomyces hygroscopicus*, MSU/MN-4-75B. *J. Antibiot.*, 48 (1995) 896-898.
- 694 Dabbs, E.R., Yazawa, K., Tanaka, Y., Mikami, Y., Miyaji, M., Andersen, S.J., Morisaki, N., Iwasaki, S., Shida, O., Takagi, H. and Kadokawa, K.: Rifampicin inactivation by *Bacillus* species. *J. Antibiot.*, 48 (1995) 815-819.
- 695 Fekete, J., Romvári, Z., Szepesi, I. and Morovján, G.: Liquid chromatographic determination of the antibiotic fumagillin in fish meat samples. *J. Chromatogr. A*, 712 (1995) 378-381.
- 696 Fujii, N., Katsuyama, T., Kobayashi, E., Hara, M. and Nakano, H.: The clecarmycins, new antitumor antibiotics produced by *Streptomyces*: fermentation, isolation and biological properties. *J. Antibiot.*, 48 (1995) 768-772.
- 697 Gau, C.-S., Chiang, Y.-Y. and Hong, W.-h.: HPLC assay of amoxicillin, aztreonam and imipenem. *Chin. Pharm. J. (Taipei)*, 47 (1995) 157-170; C.A., 123 (1995) 179621m.
- 698 Haagsma, N., Scherpenisse, P., Simmonds, R.J., Wood, S.A. and Rees, S.A.: High-performance liquid chromatographic determination of spectinomycin in swine, calf and chicken plasma using post-column derivatization. *J. Chromatogr. B*, 672 (1995) 165-171.
- 699 Hansen, E.B., Jr., Dooley, K.L. and Thompson, H.C., Jr.: High-performance liquid chromatographic analysis of the antituberculosis drugs aconzamide and isoniazid. *J. Chromatogr. B*, 670 (1995) 259-266.
- 700 Hormazábal, V. and Yndestad, M.: Detection of benzylpenicillin in milk by HPLC. *J. Liq. Chromatogr.*, 18 (1995) 2467-2474.
- 701 Hormazábal, V. and Yndestad, M.: Determination of dihydrostreptomycin sulfate in milk by HPLC using ion-pair and postcolumn derivatization. *J. Liq. Chromatogr.*, 18 (1995) 2695-2702.
- 702 Hosokawa, N., Naganawa, H., Iinuma, H., Hamada, M., Takeuchi, T., Kanbe, T. and Hori, M.: Thiazinotrienomycins, new ansamycin group antibiotics. *J. Antibiot.*, 48 (1995) 471-478.
- 703 Ikeda, H. and Omura, S.: Control of avermectin biosynthesis in *Streptomyces avermitillii* for the selective production of a useful component. *J. Antibiot.*, 48 (1995) 549-462.
- 704 Jeong, T.-S., Kim, S.U., Son, K.-H., Kwon, B.-M., Kim, Y.-K., Choi, M.-U. and Bok, S.-H.: GERI-BP001 compounds, new inhibitors of acyl-CoA:cholesterol acyltransferase from *Aspergillus fumigatus* F37. I. Production, isolation, and physico-chemical and biological properties. *J. Antibiot.*, 48 (1995) 751-756.
- 705 Kakeya, H., Takahashi, I., Okada, G., Isono, K. and Osada, H.: Epolactaene, a novel neuritogenic compound in human neuroblastoma cells, produced by a marine fungus. *J. Antibiot.*, 48 (1995) 733-735.
- 706 Lhoest, G., Verbeeck, R.K., Maton, N., Muthelet, P. and Latinne, D.: The *in vitro* immunosuppressive activity of the C15-demethylated metabolite of FK-506 is governed by ring- and open-chain tautomerism effects. *J. Pharmacol. Exp. Ther.*, 274 (1995) 622-626.
- 707 Li, L., Miles, M.V., Hall, W. and Carson, S.W.: An improved micromethod for vancomycin determination by high-performance liquid chromatography. *Ther. Drug Monit.*, 17 (1995) 366-370; C.A., 123 (1995) 131876w.
- 708 Matsuzak, K., Ikeda, H., Masuma, R., Tanaka, H. and Omura, S.: Isochromophilones I and II, novel inhibitors against gp120-CD4 binding produced by *Penicillium multicolor* FO-2338. I. Screening, taxonomy, fermentation, isolation and biological activity. *J. Antibiot.*, 48 (1995) 703-707.
- 709 McBride, J.H. and Kim, S.S.: Analysis of whole-blood cyclosporin G by liquid chromatography in renal transplant recipients. *J. Clin. Lab. Anal.*, 9 (1995) 238-242; C.A., 123 (1995) 159950q.
- 710 McCloud, T.G., Burns, M.P., Majadly, F.D., Muschik, G.M., Miller, D.A., Poole, K.K., Roach, J.M., Ross, J.T. and Lebherz, W.B., III: Production of brefeldin-A. *J. Ind. Microbiol.*, 15 (1995) 5-9; C.A., 123 (1995) 167649p.
- 711 Morino, T., Shimada, K.-i., Nakatani, A., Nishikawa, K., Harada, T. and Saito, S.: Medelamines, novel anticancer agents which cancel RAS2^{val19} induced heat shock sensitivity of yeast. *J. Antibiot.*, 48 (1995) 904-906.
- 712 Murata, H., Ohama, I., Harada, K.-i., Suzuki, M., Ikemoto, T., Shibuya, T., Haneishi, T., Torikata, A., Itezono, Y. and Nakayama, N.: Structural elucidation of aculeximycin IV. Absolute structure of aculeximycin, belonging to a new class of macrolide antibiotics. *J. Antibiot.*, 48 (1995) 850-862.
- 713 Murata, H., Suzuki, K., Tabayashi, T., Hattori, C., Takada, Y., Harada, K.-i., Suzuki, M., Ikemoto, T., Shibuya, T., Haneishi, T., Torikata, A., et al.: Structural elucidation of aculeximycin III. Planar structure of aculeximycin, belonging to a new class of macrolide antibiotics. *J. Antibiot.*, 48 (1995) 838-849.
- 714 Naidong, W., Hua, S., Roets, E. and Hoogmartens, J.: Assay and purity control of minocycline by thin-layer chromatography using UV and fluorescence densitometry - A comparison with liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 905-910.
- 715 Najjar, T.A., Al-Dhuwailie, A.A. and Tekle, A.: Comparison of high-performance liquid chromatography with fluorescence polarization immunoassay for the analysis of vancomycin in patients with chronic renal failure. *J. Chromatogr. B*, 672 (1995) 295-299.
- 716 Paesen, J., Cypres, W., Pauwels, K., Roets, E. and Hoogmartens, J.: Study of the stability of tylosin A in aqueous solutions. *J. Pharm. Biomed. Anal.*, 13 (1995) 1153-1159.
- 717 Paesen, J., Solie, A., Roets, E. and Hoogmartens, J.: Liquid chromatography of josamicin on poly(styren-divinylbenzene). *Fresenius J. Anal. Chem.*, 352 (1995) 797-800.
- 718 Pelyvas, I., Madi-Puskas, M., Toth, Z.G., Varga, Z., Hornyak, M., Batta, G. and Sztaricskai, F.: Synthesis of new pseudodisaccharide aminoglycoside antibiotics from carbohydrates. *J. Antibiot.*, 48 (1995) 683-695.

- 719 Peyrin, E., Guillaume, Y. and Guinchard, C.: High-performance liquid chromatographic determination of tazobactam by precolumn derivatization. *J. Chromatogr. B*, 672 (1995) 160-164.
- 720 Rashid, M.A., Gustafson, K.R., Cardellina, J.H., II and Boyd, M.R.: Mycalolides D and E, new cytotoxic macrolides from a collection of the stony coral *Tubastrea faulkneri*. *J. Natural Prod.*, 58 (1995) 1120-1125.
- 721 Ren, K. and Yu, C.: (Determination of cefuroxime in human serum by HPLC). *Zhongguo Yiyuan Yaoxue Zazhi*, 15 (1995) 103-105; *C.A.*, 123 (1995) 101917t.
- 722 Santos, M., García, E., López, F.G., Lanao, J.M. and Domínguez-Gil, A.: Determination of netilmicin in plasma by HPLC. *J. Pharm. Biomed. Anal.*, 13 (1995) 1059-1062.
- 723 Shakoor, O., Taylor, R.B. and Moody, R.R.: Analysis of amoxicillin in capsules and oral suspensions by high-performance liquid chromatography. *Analyst (Cambridge)*, 120 (1995) 2191-2194.
- 724 Tomoda, H., Tabata, N., Yang, D.-J., Takayanagi, H. and Omura, S.: Terpendoles, novel ACAT inhibitors produced by *Albophoma yamanashiensis* III. Production, isolation and structure elucidation of new compounds. *J. Antibiot.*, 48 (1995) 793-804.
- 725 Tsuchiya, K., Kobayashi, S., Harada, T., Kurokawa, T., Nakagawa, T., Shimada, N. and Kobayashi, K.: Gualamycin, a novel acaricide produced by *Streptomyces* sp. NK11687. I. Taxonomy, production, isolation, and preliminary characterization. *J. Antibiot.*, 48 (1995) 626-629.
- 726 Van der Pyl, D., Cans, P., Debernard, J.J., Herman, F., Lelievre, Y., Tahraoui, L., Vuilhorgne, M. and Leboul, J.: RPR113228, a novel farnesyl-protein transferase inhibitors produced by *Chrysosporium lobatum*. *J. Antibiot.*, 48 (1995) 736-737.
- 727 Venkateshwaran, T.G. and Stewart, J.T.: Determination of metronidazole in vaginal tissue by high-performance liquid chromatography using solid-phase extraction. *J. Chromatogr. B*, 672 (1995) 300-304.
- 728 Yano, Y., Hiraoka, A. and Oguma, T.: Enhancement of tetracycline binding to rat renal brush border membrane by vancomycin. *J. Pharmacol. Exp. Ther.*, 274 (1995) 695-699.

For additional information see *C.A.*:
123 (1995) 131879z.

See also 292, 296, 348.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 729 Pichon, V., Chen, L., Guenu, S. and Hennion, M.-C.: Comparison of sorbents for the solid-phase extraction of the highly polar degradation products of atrazine (including ammeline, ammelide and cyanuric acid). *J. Chromatogr. A*, 711 (1995) 257-267.

See also 108, 110, 731.

29b. Chlorinated insecticides

- 730 Buser, H.-R. and Müller, M.D.: Isomer-selective and enantioselective determination of DDT and related compounds using chiral high-resolution gas chromatography/mass spectrometry and chiral high-performance liquid chromatography. *Anal. Chem.*, 67 (1995) 2691-2698.
- 731 Specht, W., Petz, S. and Gilsbach, W.: Gas-chromatographic determination of pesticides residues after clean-up by gel-permeation chromatography and mini-silica gel column chromatography. 6. Communication. *Fresenius J. Anal. Chem.*, 353 (1995) 183-190.
- 29c. Phosphorus insecticides
- 732 Ameno, K., Fuke, C., Ameno, S., Kinoshita, H., Shirakawa, Y. and Ijiri, I.: HPLC quantitation of fenitrothion and 3-methyl-4-nitrophenol in biological fluids after prepacked cartridge extraction and its application to a poisoning case. *J. Liq. Chromatogr.*, 18 (1995) 2123-2131.
- 733 Barceló, D., Lacorte, S. and Marty, J.L.: Validation of an enzymatic biosensor with liquid chromatography for pesticide monitoring. *TrAC*, 14 (1995) 334-340.
- 734 García Pinto, C., Pérez Pavón, J.L. and Moreno Cordero, B.: Cloud point preconcentration and high-performance liquid chromatographic determination of organophosphorus pesticides with dual electrochemical detection. *Anal. Chem.*, 67 (1995) 2606-2612.
- 735 Lacorte, S. and Barceló, D.: Determination of organophosphorus pesticides and their transformation products in river waters by automated on-line solid-phase extraction followed by thermospray liquid chromatography-mass spectrometry. *J. Chromatogr. A*, 712 (1995) 103-112.

See also 731.

29d. Carbamates

See 110, 743.

29e. Herbicides

- 736 Ameno, K., Fuke, C., Ameno, S., Kinoshita, H. and Ijiri, I.: Simultaneous quantitation of diquat and its two metabolites in serum and urine by ion-paired HPLC. *J. Liq. Chromatogr.*, 18 (1995) 2115-2121.
- 737 Barnes, K.A., Startin, J.R., Thorpe, S.A., Reynolds, S.L. and Fussell, R.J.: Determination of the pesticide diflubenzuron in mushrooms by high-performance liquid chromatography-atmospheric pressure chemical ionisation mass spectrometry. *J. Chromatogr. A*, 712 (1995) 85-93.
- 738 Brede, C. and Lundanes, E.: Identification of diflubenzuron by packed-capillary supercritical fluid chromatography-mass spectrometry with electron-capture negative ionization. *J. Chromatogr. A*, 712 (1995) 95-101.
- 739 Dugay, J. and Hennion, M.-C.: Evaluation of the performance of analytical procedures for the trace-level determination of aminotriazole in drinking waters. *TrAC*, 14 (1995) 407-414.

740 Molina, C., Durand, G. and Barceló, D.: Trace determination of herbicides in estuarine waters by liquid chromatography-high-flow pneumatically assisted electrospray mass spectrometry. *J. Chromatogr. A*, 712 (1995) 113-122.

741 Pichon, V., Chen, L., Hennion, M.-C., Daniel, R., Martel, A., le Goffic, F., Abian, J. and Barceló, D.: Preparation and evaluation of immunosorbents for selective trace enrichment of phenylurea and triazine herbicides in environmental waters. *Anal. Chem.*, 67 (1995) 2451-2460.

See also 729.

29f. Fungicides

742 Brandsteterova, E., Kubalec, P., Rady, A. and Krcmery, V.: Determination of intraconazole and its metabolite in plasma using SPE-HPLC. *Pharmazie*, 50 (1995) 597-599.

743 García Sánchez, F., Navas Diaz, A. and Pareja, A.G.: Normal-phase liquid chromatography on amino-bonded-phase column of fluorescence detected pesticides. *J. Liq. Chromatogr.*, 18 (1995) 2543-2558.

744 Will, W.: Determination of vinclozolin metabolites in human urine by high-performance liquid chromatography and electrochemical detection. *Fresenius J. Anal. Chem.*, 353 (1995) 215-218.

29g. Other types of pesticides and various agrochemicals

745 Mesmer, M.Z. and Satzger, R.D.: Determination of brodifacoum in commercial rodenticides by using high-performance liquid chromatography with fluorescence detection. *Analyst (Cambridge)*, 120 (1995) 2195-2197.

See also 725, 737, 738, 786.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

746 Andrisano, V., Cavrini, V., Summer, P. and Passuti, S.: Determination of impurities in oxidation hair dyes as raw materials by liquid chromatography (HPLC). *Int. J. Cosmet. Sci.*, 17 (1995) 53-60; C.A., 123 (1995) 92838p.

747 Chivukula, M., Spadaro, J.T. and Renganathan, V.: Lignin peroxidase-catalyzed oxidation of sulfonated azo dyes generates novel sulfophenyl hydroperoxides. *Biochemistry*, 34 (1995) 7765-7772.

748 Turnipseed, S.B., Roybal, J.E., Rupp, H.S., Hurlbut, J.A. and Long, A.R.: Particle beam liquid chromatography-mass spectrometry of triphenylmethane dyes: application to confirmation of malachite green in incurred catfish tissue. *J. Chromatogr. B*, 670 (1995) 55-62.

See also 152.

30b. Chloroplast and other natural pigments

749 Eijkelhoff, C. and Dekker, J.P.: Determination of the pigment stoichiometry of the photochemical reaction center of Photosystem II. *Biochim. Biophys. Acta*, 1231 (1995) 21-28.

750 Elbergali, A.K., Brereton, R.C. and Rahmani, A.: Resolvability indices and variable selection in factor analysis of diode-array high-performance liquid chromatography: Development of approach for multi-peak clusters and application to mixtures of chlorophyll degradation products. *Analyst (Cambridge)*, 120 (1995) 2207-2216.

751 Khachik, F., Englert, G., Beecher, G.R. and Smith, J.C., Jr.: Isolation, structural elucidation, and partial synthesis of lutein dehydration products in extracts from human plasma. *J. Chromatogr. B*, 670 (1995) 219-233.

752 Laznik, B. and Kovac, M.: The determination of higher plant photosynthetic pigments. *Acta Pharm. (Zagreb)*, 45 (1995) 227-229; C.A., 123 (1995) 106898x.

753 Stradi, R., Celentano, G. and Nava, D.: Separation and identification of carotenoids in bird's plumage by high-performance liquid chromatography-diode array detection. *J. Chromatogr. B*, 670 (1995) 337-348.

754 Wirth, H.-J. and Hearn, M.T.W.: Adsorbents for the removal of humic acid from surface water based on modified porous zirconia and silica. *J. Chromatogr. A*, 711 (1995) 223-233.

See also 463, 545, 687.

31. PLASTICS AND THEIR INTERMEDIATES

755 Azuma, C., Dias, M.L. and Mano, E.B.: Size exclusion behavior of polymers in amide solvents. I. Elution characteristics of acrylonitrile polymers in N,N-dimethylformamide. *Polym. Bull. (Berlin)*, 34 (1995) 585-592; C.A., 123 (1995) 57097z.

756 Azuma, C., Dias, M.L. and Mano, E.B.: Size exclusion behavior of polymers in amide solvents. II. Molecular weight determination of acrylonitrile polymers in N,N-dimethylformamide. *Polym. Bull. (Berlin)*, 34 (1995) 593-598; C.A., 123 (1995) 113345t.

757 Barnes, K.A., Damant, A.P., Startin, J.R. and Castle, L.: Qualitative liquid chromatographic-atmospheric-pressure chemical-ionisation mass spectrometric analysis of polyethylene terephthalate oligomers. *J. Chromatogr. A*, 712 (1995) 191-199.

758 Bodzek, M. and Korus, I.: (Use of gel-permeation chromatography (GPC) for the characterization of polymer membranes). *Polimery (Warsaw)*, 40 (1995) 164-169; C.A., 123 (1995) 171876a.

759 Bos, J. and Tijssen, R.: Hydrodynamic chromatography of polymers. *J. Chromatogr. Libr.*, 56 (1995) 95-126; C.A., 123 (1995) 145053n - a review with 47 refs.

760 Chubarova, E.V. and Nestorov, V.V.: (Correlation of solvent quality with degradation of high-molecular-weight polystyrene during chromatography on porous and nonporous silicate beads). *Vysokomol. Soedin., Ser. A, Ser. B*, 36 (1994) 1372-1378; C.A., 123 (1995) 84525u.

- 761 Degoulet, C., Nicolai, T., Durand, D. and Busnel, J.P.: Characterization of polydisperse solutions of branched poly(methyl methacrylate) using size exclusion chromatography with online multiangle light scattering and viscosity detection. *Macromolecules*, 28 (1995) 6819-6824; C.A., 123 (1995) 170948r.
- 762 Fedorov, E.K. and Kedik, C.A.: (Study of molecular weight and chemical composition heterogeneity of random copolymer of N-vinylpyrrolidone and 2-methyl-5-vinylpyridine by liquid chromatography method). *Vysokomol. Soedin. Ser. A Ser. B*, 36 (1994) 1548-1552; C.A., 123 (1995) 84526v.
- 763 Grubisic-Gallot, Z., Gallot, Y. and Sedlacek, J.: Study of polystyrene-block-poly(methyl methacrylate) micelles by size exclusion chromatography/low angle laser light scattering. Influence of copolymer composition and molecular weight. *J. Liq. Chromatogr.*, 18 (1995) 2291-2307.
- 764 Hunkeler, D.J., Janco, M., Guryanova, V.V. and Berek, D.: Limiting conditions in the liquid chromatography of polymers. *Adv. Chem. Ser.*, 247 (1995) 13-19; C.A., 123 (1995) 170818y.
- 765 Jackson, C. and Barth, H.G.: Concerns regarding the practice of multiple detector size-exclusion chromatography. *Adv. Chem. Ser.*, 247 (1995) 59-68; C.A., 123 (1995) 170821u.
- 766 Kawai, T., Akashima, M. and Teramachi, S.: Compositional fraction of poly(methyl methacrylate)-graft-polydimethylsiloxane by reversed-phase high-performance liquid chromatography. *Polymer*, 36 (1995) 2851-2852; C.A., 123 (1995) 84523s.
- 767 Kilz, P., Krueger, R.-P., Much, H. and Schulz, G.: Two-dimensional chromatography for the deformulaion of complex copolymers. *Adv. Chem. Ser.*, 247(Chromatographic Characterization of Polymers) (1995) 223-241; C.A., 123 (1995) 170829c.
- 768 Krol, P. and Chrzaszcz, R.: (Chemical analysis of urethane oligomers prepared from toluene 2,4-diisocyanate (2,4-TDI) and 1,4-butanediol). *Polimery*, 39 (1994) 683-688; C.A., 123 (1995) 57496d.
- 769 Lee, H.C. and Chang, T.: On-line determination of dn/dc for size exclusion chromatography coupled with a light scattering detector. *Bull. Korean Chem. Soc.*, 16 (1995) 640-643; C.A., 123 (1995) 145206q.
- 770 Lesec, J., Millequant, M., Patin, M. and Teyssie, P.: Star-branched polymers in multidetection gel-permeation chromatography. *Adv. Chem. Ser.*, 247 (1995) 167-179; C.A., 123 (1995) 170827a.
- 771 Lloyd, P.M., Suddaby, K.G., Varney, J.E., Scrivener, E., Derrick, P.J. and Haddleton, D.M.: A comparison between matrix-assisted laser desorption/ionization time-of-flight mass spectrometry and size exclusion chromatography in the mass characterization of synthetic polymers with narrow molecular-mass distribution: poly(methyl methacrylate) and poly(styrene). *Eur. Mass Spectrom.*, 1 (1995) 293-300; C.A., 123 (1995) 170831x.
- 772 Meehan, E. and O'Donohue, S.: Characterization of block copolymers using size-exclusion chromatography with multiple detectors. *Adv. Chem. Ser.*, 247 (1995) 243-251; C.A., 123 (1995) 170830w.
- 773 Mourey, T.H. and Cool, H.: Size-exclusion chromatography with light-scattering detection at two angles. *Adv. Chem. Ser.*, 247 (1995) 123-140; C.A., 123 (1995) 170826z.
- 774 Nagy, D.J.: Characterization of poly(vinyl alcohol) using SEC multiangle laser light scattering. *Am. Lab. (Shelton)*, 27 (1995) 47J, 47L, 47N, 47P, 47R, 47T, 47V; C.A., 123 (1995) 113633k.
- 775 Rudin, A.: Determination of molecular weight distributions of copolymers by size exclusion chromatography. *Chromatogr. Sci. Ser.*, 69 (1995) 147-159; C.A., 123 (1995) 170700d - a review with 48 refs.
- 776 Sokolov, M.I., Klinskikh, A.F., Granovskaya, G.L.: (Use of gel permeation chromatography to determine the molecular weight distribution of carboxy-containing nitrile rubbers). *Kauch. Rezina*, (1994) 43-45; C.A., 123 (1995) 172224a.
- 777 Trathnigg, B.: Determination of MWD and chemical composition of polymers by chromatographic techniques. *Prog. Polym. Sci.*, 20 (1995) 615-650; C.A., 123 (1995) 57002q - a review with 161 refs.
- 778 Wang, P.J.: Size exclusion chromatography of polyamides, polyesters and fluoropolymers. *Chromatogr. Sci. Ser.*, 69 (1995) 161-183; C.A., 123 (1995) 170701e - a review with 44 refs.
- 779 Watanabe, S., Tanabe, M., Jikei, M., Kakimoto, M.-A. and Imai, Y.: Characterization of ethylene-containing polyamide using size exclusion chromatography with light scattering detector. *Polym. J. (Tokyo)*, 27 (1995) 762-765; C.A., 123 (1995) 84524t.
- 780 Willis, J.N. and Wheeler, L.: Use of gel permeation chromatography-Fourier transform infrared spectrometry interface for polymer analysis. *Adv. Chem. Ser.*, 247 (1995) 253-263; C.A., 123 (1995) 170965u.
- 781 Wu, C., Zhang, Y., Yan, X. and Cheng, R.: (A new method to calibrate size exclusion chromatography). *Gaofenzi Xuebao*, (1995) 343-348; C.A., 123 (1995) 113350r.

For additional information see C.A.:

123 (1995) 84528x, 170773e.

See also 49, 93, 105, 107, 118, 171, 967.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 782 Biagi, G.L., Recanatini, M., Barbaro, A.M., Guerra, M.C., Sapone, A., Borea, P.A. and Pietrogrande, M.C.: Chromatographic parameters as lipophilicity indices for QSAR studies. In: Kalasz, H., Ettre, L.S. and Pick, J. (Editors), *New Approaches Chromatogr. '93, (Pap. Budapest Chromatogr. Conf.)*, 1990, Fekete Sas Kiado, Budapest, 1993, pp. 215-234; C.A., 123 (1995) 102008j.
- 783 Brekkan, E., Lu, L. and Lundahl, P.: Properties of immobilized-liposome-chromatographic supports for interaction analysis. *J. Chromatogr. A*, 711 (1995) 33-42.
- 784 Chaminade, P., Feraud, S., Baillet, A. and Ferrier, D.: (Validation of an HPLC analytical method. Ruggedness test and method validation). *S.T.P. Pharma Prat.*, 5 (1995) 17-35; C.A., 123 (1995) 153013q.
- 785 Goto, J.: Sensitive chiral derivatization reagents for separation of enantiomeric drugs on liquid chromatography. In: Crommeliin D.J.A., Midha, K.K. and Nagi, T. (Editors), *Top. Pharm. Sci. 1993, Proc. Int. Congr. Pharm. Sci. F.I.P., 53rd 1993*, Medpharm Sci. Pub., Stuttgart, 1994, pp. 247-252; C.A., 123 (1995) 131742z.

- 786 Lang, D. and Böcker, R.: Highly sensitive and specific high-performance liquid chromatographic analysis of 7-hydroxywarfarin, a marker for human cytochrome P-4502C9 activity. *J. Chromatogr. B*, 672 (1995) 305-309.
- 787 Leroy, P., Bellucci, L. and Nicolas, A.: Chiral derivation for separation of racemic amino and thiol drugs by liquid chromatography and capillary electrophoresis. *Chirality*, 7 (1995) 235-242; C.A., 123 (1995) 123298u.
- 788 Makovskaya, V., Dean, J.R., Tomlinson, W.R. and Comber, M.: Octanol-water partition coefficients of substituted phenols and their correlation with molecular descriptors. *Anal. Chim. Acta*, 315 (1995) 193-200.
- 789 Makovskaya, V., Dean, J.R., Tomlinson, W.R., Hitchen, S.M. and Comber, M.: Determination of octanol-water partition coefficients using gradient liquid chromatography. *Anal. Chim. Acta*, 315 (1995) 183-192.
- 790 Ong, T.-H., Wong, Y.-S., Woo, S.-O. and Ng, S.: Factorial designs for optimization in high-performance liquid chromatography of pharmaceuticals. *Bull. Singapore Natl. Inst. Chem.*, 22 (1994) 19-29; C.A., 123 (1995) 123274h.
- 791 Tsai, R.-S., Carrupt, P.-A. and Testa, B.: Measurement of partition coefficient using centrifugal partition chromatography. Method development and application to the determination of solute properties. *ACS Symp. Ser.*, 593(Modern Countercurrent Chromatography) (1995) 143-154; C.A., 123 (1995) 101899p.
- 792 Valko, K.: Retention prediction of pharmaceutical compounds. *J. Chromatogr. Libr.*, 57 (1995) 47-92; C.A., 123 (1995) 934008 - a review with 103 refs.
- 793 Virlichie, J.L. and Ayache, A.: (Application of a model to the study of the ruggedness study to the validation of an HPLC analysis method). *S.T.P. Pharma Prat.*, 5 (1995) 37-48; C.A., 123 (1995) 153053x.
- 794 Virlichie, J.L. and Ayache, A.: A ruggedness test model and its application for HPLC method validation. *S.T.P. Pharma Prat.*, 5 (1995) 49-60; C.A., 123 (1995) 153014r.
- 795 Wickman, A., Otto, D. and Bender, C.: Automation of sample preparation HPLC analysis of pharmaceutical dosage forms. *Am. Lab. (Shelton)*, 27 (1995) 27-31; C.A., 123 (1995) 123273g.
- For additional information see C.A.:
123 (1995) 159966z.
- See also 38, 130, 133, 300, 348.
- 32b. Antirheumatics and antiinflammatory drugs**
- 796 Herron, W.J., Eadie, J. and Penman, A.D.: Estimation of ranolazine and eleven Phase I metabolites in human plasma by liquid chromatography-atmospheric pressure chemical ionisation mass spectrometry with selected-ion monitoring. *J. Chromatogr. A*, 712 (1995) 55-60.
- 797 Ibrahim, F.B.: Quantitative determination of oxaprozin and several of its related compounds by high-performance reversed-phase liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2621-2633.
- 798 Jang, E.-J., Lee, Y.-J., Park, M.-G. and Shim, C.-K.: HPLC assay of 6-methoxy-2-naphthylacetic acid, a major metabolite of nabumetone, in human serum. *Anal. Lett.*, 28 (1995) 2379-2389.
- 799 Koupai-Abyzani, M.R., Yu, N., Esaw, B. and Laviolle, B.: Determination of triamcinolone acetonide in equine serum and urine by liquid chromatography-atmospheric pressure ionization mass spectrometry. *J. Anal. Toxicol.*, 19 (1995) 182-186.
- 800 Leroy, P., Baudrit, O., Blanchin, M.-D., Nicolas, A. and Fabre, H.: Electrochemical oxidation of CBS-113 A, a new anti-inflammatory drug: applicability to liquid chromatography-electrochemical detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 1133-1140.
- 801 Li, K., Zhao, F.-L., Yuan, Y.-S. and Tan, L.: Determination of diclofenac sodium in human plasma by reversed-phase liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2205-2216.
- 802 Mason, J.L. and Hobbs, G.J.: A rapid high performance liquid chromatographic assay for the measurement of diclofenac in human plasma. *J. Liq. Chromatogr.*, 18 (1995) 2045-2058.
- 803 Niopas, I. and Georgarakis, M.: Determination of tolafenamic acid in human plasma by HPLC. *J. Liq. Chromatogr.*, 18 (1995) 2675-2682.
- 804 Shinde, V.M., Desai, B.S. and Tendolkar, N.M.: Simultaneous determination of paracetamol, diclofenac sodium and chlorzoxazone by HPLC from tablet. *Indian J. Pharm. Sci.*, 57 (1995) 35-37; C.A., 123 (1995) 93433q.
- 805 Sochor, J., Klimes, J., Sedlacek, J. and Zahradnick, M.: Determination of ibuprofen in erythrocytes and plasma by high performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 899-903.
- 806 Sochor, J., Klimes, J., Sedlacek, J. and Zahradnick, M.: High-performance liquid chromatographic analysis of kebuzone and its metabolites in the samples of erythrocytes, plasma, and whole blood. *J. Liq. Chromatogr.*, 18 (1995) 2147-2166.
- 807 Van Overbeke, A., Baeyens, W. and Dewaele, C.: Comparative study on the enantiomeric separation of several non-steroidal anti-inflammatory drugs on two cellulose-based chiral stationary phases. *J. Liq. Chromatogr.*, 18 (1995) 2427-2443.

See also 853.

32c. Autonomic and cardiovascular drugs

- 808 Al-Dhawailie, A.A.: High performance liquid chromatographic determination of amiodarone in plasma and tissues. *Anal. Lett.*, 28 (1995) 2391-2400.
- 809 Böhm, R., Ellrich, R. and Koytchev, R.: Quantitation of R- and S-propafenone and of the main metabolite in plasma. *Pharmazie*, 50 (1995) 542-545.
- 810 Cannell, R.J.P., Knaggs, A.R., Dawson, M.J., Manchee, G.R., Eddershaw, P.J., Waterhouse, I., Sutherland, D.R., Bowers, G.D. and Sidebottom, P.J.: Microbial biotransformation of the angiotensin II antagonist GR117289 by *Streptomyces rimosus* to identify a mammalian metabolite. *Drug Metab. Disp.*, 23 (1995) 724-729.
- 811 Carignan, G., Carrier, K., Laganière, S. and Lessard, M.: Simultaneous determination of diltiazem and quinidine in human plasma by liquid chromatography. *J. Chromatogr. B*, 672 (1995) 261-269.

- 812 Christensen, E.B., Hansen, S.H. and Rasmussen, S.N.: Assay of tocainide enantiomers in plasma by solid-phase extraction and indirect chiral high-performance liquid chromatography after derivatization with (-)-menthyl chloroformate. *J. Chromatogr. B*, 670 (1995) 243-249.
- 813 Dovletoglou, A., Thomas, S.M., Berwick, L., Ellison, D.K. and Twy, P.C.: Development of practical HPLC methods for analysis and quality assessment of the novel carbonic anhydrase inhibitor MK-0507 and the acetamidosulfonamide intermediate. *J. Liq. Chromatogr.*, 18 (1995) 2337-2352.
- 814 El Walily, A.F.M., Belal, S.F., Heaba, E.A. and El Kersh, A.: Simultaneous determination of enalapril maleate and hydrochlorothiazide by first-derivative ultraviolet spectrophotometry and high-performance liquid chromatography. *J. Pharm. Biomed. Anal.*, 13 (1995) 851-856.
- 815 Gilbert, J.D., Greber, T.F., Ellis, J.D., Barrish, A., Olah, T.V., Fernández-Metzler, C., Yuan, A.S. and Burke, C.J.: The development and cross-validation of methods based on radioimmunoassay and LC/MS-MS for the quantification of the Class III antiarrhythmic agent, MK-0499, in human plasma and urine. *J. Pharm. Biomed. Anal.*, 13 (1995) 937-950.
- 816 Gonzalez, A.G., Herrador, M.A. and Asuero, A.G.: Determination of β -adrenoceptor blocking agents in tablets using reversed-phase high-performance liquid chromatography. *Int. J. Pharm.*, 123 (1995) 1149-151; *C.A.*, 123 (1995) 123397t.
- 817 Gumieniczek, A., Przyborowski, L. and Plizga, J.: Determination of dilazep dihydrochloride in tablets by HPLC. *Pharmazie*, 50 (1995) 541-542.
- 818 Heath, T.G., Massad, D.D., Carroll, J.I., Mathews, B.S., Chang, J., Scott, D.O., Kuo, B.-S. and Toren, P.C.: Quantification of a dual angiotensin I-converting enzyme-neutral endopeptidase inhibitor and the active thiol metabolite in dog plasma by high-performance liquid chromatography with ultraviolet absorbance detection. *J. Chromatogr. B*, 670 (1995) 91-101.
- 819 Herrmann, R. and Kleinboesem, C.H.: Nachweis der Bioäquivalenz einer neuen Sotalololhydrochlorid-Zubereitung im Vergleich zu einer Standardformulierung. *Arzneim.-Forsch.*, 45 (1995) 849-852.
- 820 Holeman, J.A. and Danielson, N.D.: Microbore liquid chromatography of tertiary amine anticholinergic pharmaceuticals with tris(2,2'-bipyridine)ruthenium(III) chemiluminescence detection. *J. Chromatogr. Sci.*, 33 (1995) 297-302.
- 821 Hooper, W.D. and Baker, P.V.: Enantioselective analysis of sotalol in plasma by reversed-phase high-performance liquid chromatography using diastereomeric derivatives. *J. Chromatogr. B*, 672 (1995) 89-96.
- 822 Ivanovic, D., Medenica, M., Radulovic, D. and Djugumovic, Z.: Pharmaceutical analysis of β -methyldigoxin in dosage forms using RP-HPLC. *Drug. Dev. Ind. Pharm.*, 21 (1995) 1789-1797; *C.A.*, 123 (1995) 93411f.
- 823 Lausecker, B. and Hopfgartner, G.: Determination of an endothelin receptor antagonist in human plasma by narrow-bore liquid chromatography and ionspray tandem mass spectrometry. *J. Chromatogr. A*, 712 (1995) 75-83.
- 824 Lee, J.W., Naidong, W., Johnson, T., Dzerk, A., Miyabayashi, T. and Motohashi, M.: Development and validation of column-switching high-performance liquid chromatographic methods for the determination of a potent All receptor antagonist, TCV-116, and its metabolites in human serum and urine. *J. Chromatogr. B*, 670 (1995) 287-298.
- 825 Leuenberger, P.M., Ha, H.R., Pletscher, W., Meier, P.J. and Sticher, O.: Measurement of nimodipine metabolism in rat liver microsomes by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2243-2255.
- 826 Michelitsch, A., Reiner, J., Schubert-Zsilavecz, M. and Likussar, W.: 2,2'-Bis(3-isobutyloxycarbonyl-5-methoxycarbonyl-2,6-dimethyl-4-pyridyl)-azobenzene-N,N'-dioxide: a new degradation product of nisoldipine by UV light. *Pharmazie*, 50 (1995) 548-549.
- 827 Mück, W.M.: Enantiospecific determination of nimodipine in human plasma by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. A*, 712 (1995) 45-53.
- 828 Nakabeppe, H., Nakajima, A., Kamikawai, Y. and Shinozaki, Y.: Identification of efonidipine hydrochloride metabolites in rats. *Arzneim.-Forsch.*, 45 (1995) 766-770.
- 829 Oertel, R. and Richter, K.: Die Metaboliten des Beta-Rezeptorenblockers Talinolol beim Menschen - Zusammenhang zwischen Struktur, Polarität und ausgeschiedener Menge. *Pharmazie*, 50 (1995) 637-638.
- 830 Piperaki, S., Tsantili-Kakoulidou, A. and Parissi-Poulou, M.: Solvent selectivity in chiral chromatography using a β -cyclodextrin-bonded phase. *Chirality*, 7 (1995) 257-266; *C.A.*, 123 (1995) 123399v.
- 831 Rathinavelu, A. and Malave, A.: High-performance liquid chromatography using electrochemical detection for the determination of prazosin in biological samples. *J. Chromatogr. B*, 670 (1995) 177-182.
- 832 Sastry, M.K., Chakrabarti, P.K., Nanda, P.K. and Naik, S.R.: An improved method for separation of diltiazem and related substances by HPLC. *Indian J. Pharm. Sci.*, 57 (1995) 61-65; *C.A.*, 123 (1995) 65943n.
- 833 Simmons, B.R. and Stewart, J.T.: HPLC determination of atenolol in human serum on underivatized silica using solid phase extraction. *Anal. Lett.*, 28 (1995) 2017-2026.
- 834 Stockis, A., Deroubaix, X., Jeanbaptiste, B., Lins, R., Allemon, A.M. and Laufen, H.: Relative bioavailability of carbinoxamine and phenylephedrine from a retard capsule after single and repeated dose administration in healthy subjects. *Arzneim.-Forsch.*, 45 (1995) 1009-1012.
- 835 Taft, D.R. and Sweeney, K.R.: The influence of protein binding on the elimination of acetazolamide by the isolated perfused rat kidney: evidence of albumin-mediated tubular secretion. *J. Pharmacol. Exp. Ther.*, 274 (1995) 752-760.
- 836 Valladao, Mendes de Sousa, D., Zuanon Netto, J. and Ionashiro, M.: (Separation and identification of diuretic drugs). *Edectica Quin.*, 19 (1994) 57-65; *C.A.*, 122 (1995) 248495s.
- 837 Van Rhijn, J.A., O'Keeffe, M., Heskamp, H.H. and Collins, S.: Rapid analysis of β -agonists in urine by thermospray tandem mass spectrometry. *J. Chromatogr. A*, 712 (1995) 67-73.
- 838 Vistelle, R., Lamiable, D., Morin, E., Trenque, T. and Kaltenbach, M.: Urinary excretion of cicletanine in the rat. Stereochemical aspects. *Drug Metab. Disp.*, 23 (1995) 988-992.

For additional information see *C.A.*:
123 (1995) 153031u.

See also 843.

32d. Central nervous system drugs

- 839 Al-Obaidy, S.S., Po, A.L.W., McKiernan, P.J., Glasgow, J.F.T. and Millership, J.: Assay of paracetamol and its metabolites in urine, plasma and saliva of children with chronic liver disease. *J. Pharm. Biomed. Anal.*, 13 (1995) 1033-1039.
- 840 Andersin, R. and Tammilehto, S.: Photochemical decomposition of midazolam. IV. Study of pH-dependent stability by high-performance liquid chromatography. *Int. J. Pharm.*, 123 (1995) 229-235; C.A., 123 (1995) 152722h.
- 841 Ashforth, E.I.L., Carlile, D.J., Chenery, R. and Houston, J.B.: Prediction of *in vivo* disposition from *in vitro* systems: clearance of phenytoin and tolbutamide using rat hepatic microsomal and hepatocyte data. *J. Pharmacol. Exp. Ther.*, 274 (1995) 761-766.
- 842 Atmaca, S. and Tatar, S.: UV- and VIS-spectrophotometric and high-performance liquid chromatographic determinations of fluvoxamine maleate and its tablets. *Acta Pharm. Turc.*, 37 (1995) 33-37; C.A., 123 (1995) 123309y.
- 843 Boulton, D.W. and Fawcett, J.P.: Determination of salbutamol enantiomers in human plasma and urine by chiral high-performance liquid chromatography. *J. Chromatogr. B*, 672 (1995) 103-109.
- 844 Brewster, M.E., Anderson, W.R., Estes, K.S., Bodor, N.S., Pop, E. and Panetta, J.A.: Tissue distribution of LY231617, an antioxidant with neuroprotectant activity, in the rat. *J. Pharm. Sci.*, 84 (1995) 791-798.
- 845 Castanier, L., Rubat, C., Coudert, P., Albussion, E., Chopineau, J. and Couquelet, J.: Synthesis of new serotonergic 2-substituted 2,4-diaryl pyridazin-3-ones. *Arzneim.-Forsch.*, 45 (1995) 947-951.
- 846 Chavez, C.M., Constanzer, M.L. and Matuszewski, B.K.: Determination of pirenzepine in human plasma using liquid chromatography with tandem mass spectrometric detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 1179-1184.
- 847 Costa Queiroz, R.H., Lanchote, V.L., Bonato, P.S. and de Carvalho, D.: Simultaneous HPLC analysis of tricyclic antidepressants and metabolites in plasma samples. *Pharm. Acta Helv.*, 70 (1995) 181-186; C.A., 123 (1995) 159994g.
- 848 Crowley, J.R. and Hollenberg, P.F.: Mechanism-based inactivation of rat liver cytochrome P4502B1 by phencyclidine and its oxidative product, the iminium ion. *Drug Metab. Disp.*, 23 (1995) 786-793.
- 849 Fitòs, I., Visy, J., Simonyi, M. and Hermansson, J.: Separation of enantiomers of benzodiazepines on the Chiral-AGP column. *J. Chromatogr. A*, 709 (1995) 265-273.
- 850 Fujimaki, Y., Sudo, K. and Hakusui, H.: *In vitro* metabolism of nefiracetam by liver microsomes from rats, dogs and monkeys. *Arzneim.-Forsch.*, 45 (1995) 748-752.
- 851 García-Encina, G., Farrán, R., Puig, S., Serafini, M.T. and Martínez, L.: Automated high-performance liquid chromatographic assay for lesopitron, a novel anxiolytic, in human plasma using on-line solid-phase extraction. *J. Chromatogr. B*, 670 (1995) 103-110.
- 852 Gurley, B.J., Marx, M. and Olsen, K.: Phenytoin free fraction determination: comparison of an improved direct serum injection high-performance liquid chromatographic method to ultrafiltration coupled with fluorescence polarization immunoassay. *J. Chromatogr. B*, 670 (1995) 358-364.
- 853 Jimenez-Carmona, M.M., Tena, M.T. and Luque de Castro, M.D.: Ion-pair-supercritical fluid extraction of clenbuterol from food samples. *J. Chromatogr. A*, 711 (1995) 269-276.
- 854 Langen, M.C.J. and Ros, J.J.W.: Adsorption of tricyclic antidepressants to the barrier of serum separator tubes. *Ziekenhuisfarmacie*, 11 (1995) 64-66; C.A., 123 (1995) 101901h.
- 855 Law, M.Y.L. and Moody, D.E.: Simultaneous quantitation of amphetamine and 4'-hydroxyamphetamine by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2029-2043.
- 856 Lehotay, J., Cizmarik, J. and Brandsteterova, E.: Study of local anesthetics. Part 128: HPLC study of the degradation of heptacaine and carbisocaine in urine, human and rabbit serums. *Acta Pharm. Hung.*, 65 (1995) 95-96; C.A., 123 (1995) 159972y.
- 857 Ma, C.-K., Mok, C.-S. and Hon, P.-K.: Determination of benzhexol hydrochloride and ethambutol hydrochloride tablets by liquid chromatography. *Anal. Chim. Acta*, 314 (1995) 77-85.
- 858 Mather, L.E., Rutten, A.J. and Plummer, J.L.: Pharmacokinetics of bupivacaine enantiomers in sheep: influence of dosage regimen and study design. *J. Pharmacokin. Biopharm.*, 22 (1994) 481-498.
- 859 Meza-Toledo, S.E., Ortega-Gonzalo, C., Juárez-Carbalal, E. and Carvajal-Sandoval, G.: Stereoselective anticonvulsant activity of the enantiomers of (\pm)-2-hydroxy-2-phenylbutyramide. *Arzneim.-Forsch.*, 45 (1995) 756-759.
- 860 Molins Legua, C., Campiñs Falcó, P. and Sevillano Cabeza, A.: Amphetamine and methamphetamine determination in urine by reversed-phase high-performance liquid chromatography with sodium 1,2-naphthoquinone 4-sulfonate as derivatizing agent and solid-phase extraction for sample clean-up. *J. Chromatogr. B*, 672 (1995) 81-88.
- 861 Papadoyannis, I.N., Zouzou, A.C. and Samanidou, V.F.: Solid-phase extraction study and RP-HPLC analysis of lamotrigine in human biological fluids and in antiepileptic tablet formulations. *J. Liq. Chromatogr.*, 18 (1995) 2593-2609.
- 862 Ponder, G.W. and Stewart, J.T.: A liquid chromatographic method for the determination of promethazine enantiomers in human urine and serum using solid-phase extraction and fluorescence detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 1161-1166.
- 863 Rodrigues, A.D., Mulford, D.J., Lee, R.D., Surber, B.W., Kukulka, M.J., Ferrero, J.L., Thomas, S.B., Shet, M.S. and Estabrook, R.W.: *In vitro* metabolism of terfenadine by a purified recombinant fusion protein containing cytochrome P4503A4 and NADPH-P450 reductase. Comparison to human liver microsomes and precision-cut liver tissue slices. *Drug Metab. Disp.*, 23 (1995) 765-775.
- 864 Rozman, E., Galceran, M.T., Anglada, L. and Albet, C.: Investigation of the metabolism of ebrotidine in human urine by liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *Drug Metab. Disp.*, 23 (1995) 976-981.
- 865 Segarra, V., Carrera, F., Fabregas, J.-L. and Claramunt, J.: Degradation profile and identification of the major degradation products of dobutropride under several conditions by GC/MS and HPLC-particle beam/MS. *J. Pharm. Biomed. Anal.*, 13 (1995) 987-993.
- 866 Swanson, S.P., Catlow, J., Pohland, R.C., Chay, S.H. and Johnson, T.: Disposition of the opioid antagonist, LY255582, in rats and dogs. *Drug Metab. Disp.*, 23 (1995) 916-921.

- 867 Taki, Y., Sakane, T., Nadai, T., Sezaki, H., Amidon, G.L., Langguth, P. and Yamashita, S.: Gastrointestinal absorption of peptide drug: Quantitative evaluation of the degradation and the permeation of metkephamid in rat small intestine. *J. Pharmacol. Exp. Ther.*, 274 (1995) 373-377.
- 868 Tan, H., Yin, Z. and Cai, M.: (Bioavailability of acetaminophen in ten healthy male volunteers by reversed-phase HPLC with electrochemical detection). *Zhongguo Yiyuan Yaoyue Zaishi*, 15 (1995) 51-53; *C.A.*, 123 (1995) 122927m.
- 869 Van Belle, K., Verfaillie, I., Ebinger, G. and Michotte, Y.: Liquid chromatographic assay using a microcolumn coupled to a U-shaped optical cell for high-sensitivity ultraviolet absorbance detection of oxcarbazepine and its major metabolite in microdialysates. *J. Chromatogr. B*, 672 (1995) 97-102.
- 870 Venkateshwaran, T.G., King, D.T. and Stewart, J.T.: HPLC determination of ondansetron-atropine and ondansetron-glycopyrrolate mixtures in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 18 (1995) 2647-2659.
- 871 Walker, R.B. and Kanfer, I.: Sensitive high-performance liquid chromatographic determination of cyclizine and its demethylated metabolite, norcyclizine, in biological fluids using coulometric detection. *J. Chromatogr. B*, 672 (1995) 172-177.
- 872 Walters, D.L., Strong, C.R., Green, S.V. and Curtis, M.A.: Development and validation of a gradient reversed-phase high-performance liquid chromatographic assay for S(-)-2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0923) from a transdermal delivery system. *J. Chromatogr. B*, 670 (1995) 299-307.
- 873 Waters, S.M., Konkoy, C.S. and Davis, T.P.: Neuropeptide metabolism on intact, regional brain slices: effect of dopaminergic agents on substance P, cholecystokinin and Met-enkephalin degradation. *J. Pharmacol. Exp. Ther.*, 274 (1995) 783-789.
- 874 Watt, A.P., O'Connor, D., Thomas, S., Herbert, R.H. and Matassa, V.G.: *In vitro* studies of the oxidative metabolism of L-737, 415, a CS-cycloalkylamine-1,4-benzodiazepin-2-one CCK₈ receptor antagonist. *J. Liq. Chromatogr.*, 18 (1995) 2271-2282.
- 875 Woodland, J.M. and Elhardt, W.J.: Characterization of an O-glucuronide metabolite of the CCK-B antagonist LY288513 and its plasma levels in mouse, rat, and dog. *Drug Metab. Disp.*, 23 (1995) 910-915.
- 876 Yamashita, S., Furuno, K., Kawasaki, H., Gomita, Y., Yoshinaga, H., Yamatogi, Y. and Ohtahara, S.: Simple and rapid analysis of lamotrigine, a novel antiepileptic, in human serum by high-performance liquid chromatography using a solid-phase extraction technique. *J. Chromatogr. B*, 670 (1995) 354-357.
- For additional information see *C.A.*:
123 (1995) 153035y.
- See also 53, 65, 342, 804, 830, 939.
- 32e. *Chemotherapeutics (exc. cytostatics and antibiotics)*
- 877 Brocks, D.R., Dennis, M.J. and Schaefer, W.H.: A liquid chromatographic assay for the stereospecific quantitative analysis of halofantrine in human plasma. *J. Pharm. Biomed. Anal.*, 13 (1995) 911-918.
- 878 Chouini-Lalanne, N., Malet-Martino, M.-C., Gilard, V., Ader, J.-C. and Martino, R.: Structural determination of a glucuronide conjugate of flucytosine in humans. *Drug Metab. Disp.*, 23 (1995) 813-817.
- 879 Karle, J.M., Olmeda, R., Freeman, S.G. and Schroeder, A.C.: Quantification of the individual enantiomer plasma concentrations of the candidate antimalarial agent N⁴-[2,6-dimethoxy-4-methyl-5-[(3-trifluoromethyl)phenoxy]-8-quinolinyl]-1,4-pentane diamine (WR 238,605). *J. Chromatogr. B*, 670 (1995) 251-257.
- 880 Lim, B.-G., Ching, C.-B., Tan, R.B.H. and Ng, S.-C.: Recovery of (-)-praziquantel from racemic mixtures by continuous chromatography and crystallization. *Chem. Eng. Sci.*, 50 (1995) 2289-2298; *C.A.*, 123 (1995) 93448y.
- 881 McCracken, R.J., Blanchflower, W.J., Rowan, C., McCoy, A.M. and Kennedy, D.G.: Determination of furazolidone in porcine tissue using thermospray liquid chromatography-mass spectrometry and study of the pharmacokinetics and stability of its residues. *Analyst (Cambridge)*, 120 (1995) 2347-2351.
- 882 Medicott, N.J., Tucker, I.G., Holborow, D.W. and Rathbone, M.J.: Liquid chromatographic analysis of chlorhexidine at specific sites in the saliva film after application of a tooth-bonded delivery system. *J. Liq. Chromatogr.*, 18 (1995) 2193-2204.
- 883 Potts, W., van Horn, R., Anderson, K., Blake, T., Graver, E., Joseph, G., Dreyer, G., Shu, A., Heys, R. and Fong, K.-L.: Characterization of the metabolites of the peptidomimetic human immunodeficiency virus type 1 protease inhibitor SK&F 107461 in rats using liquid chromatography/mass spectrometry. *Drug Metab. Disp.*, 23 (1995) 799-805.
- 884 Prueksaritanont, T., Balani, S.K., Dwyer, L.M., Ellsi, J.D., Kauffman, L.R., Varga, S.L., Pitzenberger, S.M. and Theoharides, A.D.: Species differences in the metabolism of a potent HIV-1 reverse transcriptase inhibitor L-738,372. *in vivo* and *in vitro* studies in rats, dogs, monkeys, and humans. *Drug Metab. Disp.*, 23 (1995) 688-695.
- 885 Simó-Alfonso, E.F., Ramis-Ramos, G., García-Alvarez-Coque, M.C. and Esteve-Romero, J.S.: Determination of sulphonamides in human urine by azo dye precolumn derivatization and micellar liquid chromatography. *J. Chromatogr. B*, 670 (1995) 183-187.
- 886 Tendolkar, N.M., Desai, B.S., Gandhi, J.S. and Shinde, V.M.: Simultaneous determination of tinidazole and furazolidine in suspension by HPTLC and HPLC. *Anal. Lett.*, 28 (1995) 1641-1653.
- 887 Vree, T.B., Schoondermark-van de Ven, E., Verwey-van Wissen, C.P.W.G.M., Baars, A.M., Swolfs, A., van Galen, P.M. and Amatdjais-Groenen, H.: Isolation, identification and determination of sulfadiazine and its hydroxy metabolites and conjugates from man and Rhesus monkey by high-performance liquid chromatography. *J. Chromatogr. B*, 670 (1995) 111-123.
- 888 Ye, L., Stewart, J.T. and Zhang, H.: A comparison of disc and cartridge solid-phase extraction for the LC determination of rifampin and 25-desacetylrifampin in human serum. *J. Pharm. Biomed. Anal.*, 13 (1995) 1185-1188.
- See also 646, 699, 727, 728, 742.

32f. Cytostatics

- 889 Aboagye, E.O., Graham, M.A., Lewis, A.D., Workman, P., Kelson, A.B. and Tracy, M.: Development and validation of a solid-phase extraction and high-performance liquid chromatographic assay for a novel fluorinated 2-nitroimidazole hypoxia probe (SR-4554) in Balb/c mouse plasma. *J. Chromatogr. B*, 672 (1995) 125-132.
- 890 Augey, V., Cociglio, M., Galtier, M., Yearoo, R., Pinsani, V. and Bressolle, F.: High-performance liquid chromatographic determination of *cis*-dichlorodiamineplatinum(II) in plasma ultrafiltrate. *J. Pharm. Biomed. Anal.*, 13 (1995) 1173-1178.
- 891 Drozd, M.L., Lockwood, G., Miller, B. and Rosi, D.: A column-switching high-performance liquid chromatographic assay for a novel cytotoxic thioxanthone derivative (WIN 33377) in mouse plasma with toxicokinetic results from a mouse LD₁₀ study. *J. Chromatogr. B*, 670 (1995) 145-151.
- 892 Gurley, L.R., Umbarger, K.O., Kim, J.M., Bradbury, E.M. and Lehnert, B.E.: Development of a high-performance liquid chromatographic method for the analysis of staurosporine. *J. Chromatogr. B*, 670 (1995) 125-138.
- 893 Innocenti, F., Danesi, R., Di Paolo, A., Agen, C., Mardini, D., Bocci, G. and Del Tacca, M.: Plasma and tissue disposition of paclitaxel (taxol) after intraperitoneal administration in mice. *Drug Metab. Disp.*, 23 (1995) 713-717.
- 894 John, C.S., Vilner, B.J., Gulden, M.E., Efange, S.M.N., Langason, R.B., Moody, T.W. and Bowen, W.D.: Synthesis and pharmacological characterization of 4-[¹²⁵I]-N-(N-benzylpiperidin-4-yl)-4-iodobenzamide: a high affinity σ receptor ligand for potential imaging of breast cancer. *Cancer Res.*, 55 (1995) 3022-3027.
- 895 Lyon, A.W., Duran, G. and Raisys, V.A.: Determination of thalidomide by high-performance liquid chromatography: methodological strategy for clinical trials. *Clin. Biochem.*, 28 (1995) 467-470; C.A., 123 (1995) 159980z.
- 896 Mamber, S.W., Mikkilineni, A.B., Pack, E.J., Rosser, M.P., Wong, H., Ueda, Y. and Forenza, S.: Tubulin polymerization by paclitaxel (taxol) phosphate prodrugs after metabolic activation with alkaline phosphatase. *J. Pharmacol. Exp. Ther.*, 274 (1995) 877-883.
- 897 Oh, I.J., Lee, J.M., Lee, Y.B., Shin, S.C. and Choi, B.G.: Stability of antineoplaston A 10 in aqueous solution. *Arch. Pharmacol. Res.*, 18 (1995) 75-78; C.A., 123 (1995) 93088f.
- 898 Placidi, L., Cretton-Scott, E., de Sousa, G., Rahmani, R., Placidi, M. and Sommadossi, J.-P.: Disposition and metabolism of the angiogenic moderator O-(chloroacetyl-carbamoyl) fumagillo (TNP-470; AGM-1470) in human hepatocytes and tissue microsomes. *Cancer Res.*, 55 (1995) 3036-3042.
- 899 Poon, G.K., Raynaud, F.I., Mistry, P., Odell, D.E., Kelland, L.R., Harrap, K.R., Barnard, C.F.J. and Murrer, B.A.: Metabolic studies of an orally active platinum anticancer drug by liquid chromatography-electrospray ionization mass spectrometry. *J. Chromatogr. A*, 712 (1995) 61-66.
- 900 Qing, W.-G., Powell, K.L., Stoica, G., Szumlanski, C.L., Weinshilboum, R.M. and Macleod, M.C.: Toxicity and metabolism in mice of 2,6-dithiopurine, a potential chemopreventive agent. *Drug Metab. Disp.*, 23 (1995) 854-860.
- 901 Robin, H., Senan, S., Workman, P. and Graham, M.A.: Development and validation of a sensitive solid-phase-extraction and high-performance liquid chromatography assay for the bioreductive agent tirapazamine and its major metabolites in mouse and human plasma for pharmacokinetically guided dose escalation. *Cancer Chemother. Pharmacol.*, 36 (1995) 266-270; C.A., 123 (1995) 131853m.
- 902 Rodgers, A.H., Subramanian, S. and Morgan, L.R., Jr.: High-performance liquid chromatographic determination of 4,4'-dihydroxybenzophenone-2,4-dinitrophenylhydrazone in plasma. *J. Chromatogr. B*, 670 (1995) 365-368.
- 903 Siethoff, C., Nigge, W. and Linscheid, M.: The determination of ifosfamide in human blood serum using LC/MS. *Fresenius J. Anal. Chem.*, 352 (1995) 801-805.

See also 611, 667, 682, 710.

32g. Other drug categories

- 904 Acheampong, A.A., Shackleton, M. and Tang-Liu, D.D.-S.: Comparative ocular pharmacokinetics of brimonidine after a single dose application to the eyes of albino and pigmented rabbits. *Drug Metab. Disp.*, 23 (1995) 708-712.
- 905 Bourrinet, P., Feldman, H., Dencausse, A., Chambon, C. and Bonnemain, B.: High-performance liquid chromatographic determination of ibotritidol in plasma, urine and bile. *J. Chromatogr. B*, 670 (1995) 369-372.
- 906 Chen, I.-W., Vastag, K.J. and Lin, J.H.: High-performance liquid chromatographic determination of a potent and selective HIV protease inhibitor (L-735,524) in rat, dog and monkey plasma. *J. Chromatogr. B*, 672 (1995) 111-117.
- 907 Faulkner, A. and DeMontigny, P.: High-performance liquid chromatographic determination of denatonium benzoate in ethanol with 5% polyvinylpyrrolidone. *J. Chromatogr. A*, 715 (1995) 189-194.
- 908 Firdaous, I., Hassoun, A., Otte, J.B., Reding, R., Squifflet, J.P., Besse, T. and Wallemacq, P.E.: HPLC-microparticle enzyme immunoassay specific for tacrolimus in whole blood of hepatic and renal transplant patients. *Clin. Chem. (Washington)*, 41 (1995) 1292-1296.
- 909 Heitzmann, T., Neidlein, R., Freund, P. and Strein, K.: Pharmacokinetics of the new oral blood glucose-lowering agent (-)-2-(4-tert.-butylphenoxy)-7-(4-chlorophenyl)-heptanic acid sodium salt in mice, rats and dogs. *Arzneim.-Forsch.*, 67 (1995) 868-872.
- 910 Hvattum, E., Normann, P.T., Jamieson, G.C., Lai, J.-J. and Skotland, T.: Detection and quantitation of gadolinium chelates in human serum and urine by high-performance liquid chromatography and post-column derivatization of gadolinium with Arsenazo III. *J. Pharm. Biomed. Anal.*, 13 (1995) 927-932.
- 911 Josefsson, M., Zackrisson, A.-L. and Norlander, B.: Sensitive high-performance liquid chromatographic analysis of amlodipine in human plasma with amperometric detection and a single-step solid-phase sample preparation. *J. Chromatogr. B*, 672 (1995) 310-313.
- 912 Letellier, S., Maupas, B., Gramond, J.P., Guyon, F. and Gareil, P.: Determination of the formation constant for the inclusion complex between rutin and methyl-β-cyclodextrin. *Anal. Chim. Acta*, 315 (1995) 357-363.

- 913 Li, C., Chauret, N., Ducharme, Y., Trimble, L.A., Nicoll-Griffith, D.A. and Yerger, J.A.: Integrated application of capillary HPLC/continuous-flow liquid secondary ion mass spectrometry to discovery stage metabolism studies. *Anal. Chem.*, 67 (1995) 2931-2936.
- 914 Lin, S.: Direct liquid chromatographic separation of 2,3-dihydro-2-ethylbenzofuran-2-carboxylic acid enantiomers using a cellulose carbamate-based chiral stationary phase column. *J. Liq. Chromatogr.*, 18 (1995) 2611-2619.
- 915 Macher, M. and Wintersteiger, R.: Improved electrochemical detection of diuretics in high-performance liquid chromatographic analysis by postcolumn on-line photolysis. *J. Chromatogr. A*, 709 (1995) 257-264.
- 916 Nadeau, R.W., Satoh, H., Scheide, S., Crowl, R., Conroy, R., Garland, W.A. and Liberato, D.J.: A comparison of mass balance, pharmacokinetics and disposition of [$^{14}\text{C}(\text{U})$]- and [^{125}I]-recombinant human interleukin-2 in cynomolgus monkeys. *Drug Metab. Disp.*, 23 (1995) 904-909.
- 917 Núñez, M., Ferguson, J.E., Machacek, D., Jacob, G., Oda, R.P., Lawson, G.M. and Landers, J.P.: Detection of hypoglycemic drugs in human urine using micellar electrokinetic chromatography. *Anal. Chem.*, 67 (1995) 3668-3675.
- 918 Preechagoon, Y. and Charles, B.G.: Analysis of cisapride in neonatal plasma using high-performance liquid chromatography with a base-stable column and fluorescence detection. *J. Chromatogr. B*, 670 (1995) 139-143.
- 919 Rajagopalan, P., Gao, Z., Chu, C.K., Schinazi, R.F., McClure, H.M. and Boudinot, F.D.: High-performance liquid chromatographic determination of (-)- β -D-2,6-diaminopurine dioxolane and its metabolite, dioxolane guanosine, using ultraviolet and on-line radiochemical detection. *J. Chromatogr. B*, 672 (1995) 119-124.
- 920 Reith, M.K., Sproles, G.D. and Cheng, L.K.: Human metabolism of dolasetron mesylate, a 5-HT₃ receptor antagonist. *Drug Metab. Disp.*, 23 (1995) 806-812.
- 921 Roberts, W.L., Reynolds, K.M., Heimer, R., Jatlow, P.I. and Rainey, P.M.: Pyrimethamine analysis by enzyme inhibition and HPLC assays. *Am. J. Clin. Pathol.*, 104 (1994, Pub. 1995) 82-88; *C.A.*, 123 (1995) 159951r.
- 922 Salem, I.I., Mena, P., Gallardo, V. and Ruiz, M.A.: High-pressure liquid chromatographic determination of the stability of benzoyl peroxide in pharmaceuticals. *S.T.P. Pharma Prat.*, 5 (1995) 238-241; *C.A.*, 123 (1995) 122933k.
- 923 Scurei, D.: (New analytical system to characterize finished cosmetic products. I. Chromatographic identification of phenolic glycosides from poplar buds extracts in cosmetic lotions). *Rev. Chim. (Bucharest)*, 46 (1995) 379-383; *C.A.*, 123 (1995) 65534e.
- 924 Stavchansky, S., Demirbas, S., Reyderman, L. and Chai, C.-K.: Simultaneous determination of dextrophan and guaifenesin in human plasma by liquid chromatography with fluorescence detection. *J. Pharm. Biomed. Anal.*, 13 (1995) 919-925.
- 925 Stubbs, R.J. and Harker, A.J.: Automated high-performance liquid chromatographic method for the determination of a neuraminidase inhibitor (GG167) in human serum by pre-column fluorescence derivatisation using benzoin. *J. Chromatogr. B*, 670 (1995) 279-285.
- 926 Sumiyoshi, H., Fujiwara, Y., Ohune, T., Yamaoka, N., Tamura, K. and Yamakido, M.: High-performance liquid chromatographic determination of irinotecan (CPT-11) and its active metabolite (SN-38) in human plasma. *J. Chromatogr. B*, 670 (1995) 309-316.
- 927 Tegeler, A., Ruess, W. and Gmahl, E.: Determination of amphoteric surfactants in cosmetic cleansing products by high-performance liquid chromatography on a cation-exchange column. *J. Chromatogr. A*, 715 (1995) 195-198.
- 928 Torrado, S., Fraile, S., Torrado, J.J. and Selles, E.: Comparison of reversed-phase liquid chromatography with colorimetry for analysis of phenolphthalein preparations. *J. Pharm. Biomed. Anal.*, 13 (1995) 1167-1172.
- 929 Vucina, J.L.: Comparative investigation of some Tc-99m radiopharmaceuticals in gel chromatography analysis. *J. Serb. Chem. Soc.*, 60 (1990) 501-505; *C.A.*, 123 (1995) 65926j.
- 930 Wang, C.P., Lim, H.-K., Chan, K.W., Scatina, J. and Sisenwine, S.F.: High performance liquid chromatographic isolation and spectroscopic characterization of three major metabolites from the plasma of rats receiving rapamycin (Sirolimus) orally. *J. Liq. Chromatogr.*, 18 (1995) 2559-2568.
- 931 Weyland, J.W., Rooselaar, J. and Stern, A.G.: 5-Methoxysoralen and 8-methoxysoralen in sun cosmetics and fragrance. Analysis and market survey. *Mitt. Geb. Lebensmittelunters. Hyg.*, 85 (1994) 382-394; *C.A.*, 123 (1995) 122695j.
- 932 Wring, S.A., Hughes, S.C. and Shenoy, B.E.V.: Voltammetric study of GR127935 at a glassy carbon electrode, and optimized conditions for its determination by high-performance liquid chromatography with coulometric detection. *Anal. Proc.*, 32 (1995) 257-261; *C.A.*, 123 (1995) 101906p.
- 933 Wu, W.-N. and Mutter, M.S.: Biotransformation of linogliride, a hypoglycemic agent in laboratory animals and humans. *J. Pharm. Biomed. Anal.*, 13 (1995) 857-867.
- 934 Zhao, N., Lo, L., Berova, N., Nakanishi, K., Tywiak, A.A., Ludens, J.H. and Haupert, G.T., Jr.: Na,K-ATPase inhibitors from bovine hypothalamus and human plasma are different from ouabain: nanogram scale CD structural analysis. *Biochemistry*, 34 (1995) 9893-9896.
- For additional information see *C.A.*:
123 (1995) 179627t.
- See also 307, 658, 706, 709, 746, 1003.
- 32h. *Toxicological and forensic applications*
- 935 Aderjan, R., Hotfmann, S., Schmitt, G. and Skopp, G.: Morphine and morphine glucuronides in serum of heroin consumers and in heroin-related deaths determined by HPLC with native fluorescence detection. *J. Anal. Toxicol.*, 19 (1995) 163-168.
- 936 Bloemhof, H. and Greijdanus, Uges, D.R.A.: (Determination of drugs of abuse). *Ziekenhuisfarmacie*, 11 (1995) 88-92; *C.A.*, 123 (1995) 104456c.
- 937 Clark, C.R., DeRuiter, J., Valaer, A. and Noggle, F.T.: Gas chromatographic-mass spectrometric and liquid chromatographic analysis of designer butanamines related to MDMA. *J. Chromatogr. Sci.*, 33 (1995) 328-337.

938 Noggle, F.T., Clark, C.R. and DeRuiter, J.: GC-MS and LC of addition products formed from the reaction of allylbenzene and related arylpropenes with acetonitrile and sulfuric acid. *J. Chromatogr. Sci.*, 33 (1995) 256-262.

939 Sukbuntherng, J., Hutchaleelaha, A., Chow, H.-H. and Mayersohn, M.: Separation and quantitation of the enantiomers of methamphetamine and its metabolites in urine by HPLC: precolumn derivatization and fluorescence detection. *J. Anal. Toxicol.*, 19 (1995) 139-147.

940 Theodoridis, G., Papadoyannis, I., Tsoukali-Papadopoulou, H. and Vasilikiotis, G.: A comparative study of different solid phase extraction procedures for the analysis of alkaloids of forensic interest in biological fluids by RP-HPLC/diode array. *J. Liq. Chromatogr.*, 18 (1995) 1973-1995.

See also 328, 631, 636, 639, 732, 736, 948.

32i. Plant extracts

941 Fang, W., Wu, Y., Zhou, J., Chen, W. and Fang, Q.: Determination of 10-deacetyl baccatin III and baccatin III in *Taxus yunnanensis*. *J. Chin. Pharm. Sci.*, 4 (1995) 47-50; C.A., 123 (1995) 18022b.

942 Rickling, B., Hans, B., Kramarczyk, R., Krumbiegel, G. and Weyhenmeyer, R.: Two high-performance liquid chromatographic assays for the determination of free and total silibinin diastereomers in plasma using column switching with electrochemical detection and reversed-phase chromatography with ultraviolet detection. *J. Chromatogr. B*, 670 (1995) 267-277.

943 Sheu, S.-J. and Kuo, C.-H.: A comparative study of processed *Notopterygium rhizome*. *Chin. Pharm. J. (Taipei)*, 47 (1995) 13-21; C.A., 123 (1995) 65632k.

944 Vandenberghe, D.R., Vergauwe, A.N., van Montagu, M. and van den Eeckhout, E.G.: Simultaneous determination of artemisinin and its bioprecursors in *Artemisia annua*. *J. Nat. Prod.*, 58 (1995) 798-803; C.A., 123 (1995) 65938q.

For additional information see C.A.:

123 (1995) 93409m, 93412g, 153033w, 179607m.

See also 189, 192, 193, 194, 259, 314, 315, 316, 319, 322, 634, 653, 923.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

945 Luque de Castro, M.D., Fernández-Romero, J.M. and Quiles, R.: Determination of inorganic ions of clinical interest: state-of-the-art and trends. *J. Pharm. Biomed. Anal.*, 13 (1995) 797-808 - a review with 235 refs.

33b. Complex mixtures and profiling (single compounds by cross-reference only)

946 Shi, H., Ma, Y. and Ma, Y.: A simple and fast method to determine and quantify urinary creatinine. *Anal. Chim. Acta*, 312 (1995) 79-83.

See also 204, 245, 261, 265, 283, 302, 335, 337, 339, 346, 347, 357, 536, 567, 613, 615, 618, 621, 637, 641, 751.

34. FOOD ANALYSIS

34b. Complex mixtures (single compounds by cross-reference only)

947 Avalos, J., Fontan, G.P. and Rodriguez, E.: Simultaneous HPLC quantification of two dermatotoxins, 5-methoxysoralen and falcarinol, in healthy celery. *J. Liq. Chromatogr.*, 18 (1995) 2069-2076.

948 Kirilyuk, S.D. and Smirnov, L.P.: Effect of various environment factors on protein and polypeptide composition of different tissues of fish. I. Liquid chromatographic study. In: Sidorov, V.S. and Vysotskaya, R.U. (Editors), *Biokhim. Metody Ekol. Toksikol. Issled.*, Ross. Akad. Nauk, Karel, Nauch. Tsentr, Petrozavodsk, 1993, pp. 41-46; C.A., 123 (1995) 104551e.

949 Moersel, J.T., Huth, M. and Seifert, K.: (Food irradiation. Detection by coupled TLC-HPLC). *LaborPraxis*, 19 (1995) 24-26; C.A., 123 (1995) 110356m.

For additional information see C.A.:

123 (1995) 142004m.

See also 173, 174, 178, 179, 186, 195, 198, 215, 219, 260, 274, 282, 310, 343, 344, 355, 365, 417, 434, 499, 632, 673, 678, 700, 701, 950, 997, 998, 1001.

34c. Organoletically important compounds (flavors, odors, volatiles)

950 Piggott, J.R., Clyne, J., Paterson, A. and Conner, J.M.: Preparative HPLC as a tool for concentrating flavor compounds from distilled beverages. In: Cantagrel, R. (Editor), *Symp. Sci. Int. Cognac*, 1st 1992, Tec & Doc-Lavoisier, Paris, 1993, pp. 464-467; C.A., 123 (1995) 168174s.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

951 Krylov, A.I.: Chromatographic analysis in environmental expert examinations. *J. Anal. Chem.*, 50 (1995) 210-219; C.A., 123 (1995) 73798j - a review with 96 refs.

952 Oikawa, K., Imaizumi, N. and Nonomura, M.: (Ion chromatography). *Bunseki*, (1994) 11-118; C.A., 123 (1995) 180045h - a review with 100 refs.

For additional information see C.A.:

123 (1995) 187079f.

See also 153, 156, 165, 622, 788, 789.

35b. *Air pollution (complex mixtures; single compounds by cross-reference only)*

- 953 Camel, V. and Caude, M.: Trace enrichment methods for the determination of organic pollutants in ambient air. *J. Chromatogr. A.*, 710 (1995) 3-19 - a review with 74 refs.
 954 Frenzel, W., Rauterberg-Wulff, A. and Schepers, D.: Serial potentiometric and conductometric detection in fast non-suppressed ion chromatography applied to the analysis of filter collected airborne particulates. *Fresenius J. Anal. Chem.*, 353 (1995) 123-127.
 955 Xu, X.B., Chu, S.G. and Song, N.: Application of chromatographic studies of air pollution in China. *J. Chromatogr. A.*, 710 (1995) 21-37 - a review with 116 refs.

See also 156, 158, 162, 165, 205, 206, 654.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

- 956 Schröder, H.F.: Polar organic pollutants in the Elbe river. Liquid chromatographic-mass spectrometric and flow-injection analysis-mass spectrometric analyses demonstrating changes in quality and concentration during the unification process in Germany. *J. Chromatogr. A.*, 712 (1995) 123-140.

See also 127, 159, 176, 184, 185, 325, 329, 331, 655, 665, 666, 733, 734, 739, 740, 741, 989, 997, 999.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

- 957 Huettenhain, S.H., Wilhelm, C., Holley, C., Windrich, J., Arnold, J. and Kampe, M.: Separation of pyrene and hexachlorobenzene by middle pressure liquid extraction (MPLE) of soil. *Chemosphere*, 31 (1995) 2747-2753; C.A., 123 (1995) 110838v.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. Surfactants

- 958 Felber, H., Hegetschweiler, K., Mueller, M., Odermatt, R. and Wampfler, B.: (Quantitative determination of phosphonates in detergents). *Chimia*, 49 (1995) 179-181; C.A., 123 (1995) 86629e.
 959 Groska, L., Daradics, L. and Palinkas, J.: (Production control of betain-type surfactants by high-performance liquid chromatography (HPLC)). *Olaj. Szappan, Kosmet.*, 42 (1993) 19-21; C.A., 123 (1995) 147380r.
 960 Hui, F. and Rosset, R.: Isolation of polybutenylsuccinimide-type dispersant from monograde and multigrade lubricating oils by classical liquid-adsorption chromatography on a Florisil column. *Anal. Chim. Acta*, 314 (1995) 161-167.
 961 Lawrence, J.G.: Chromatographic analysis of surfactants. In: Cullum, D.C. (Editor), *Introd. Surfactant Anal.*, Blackie, Glasgow, 1994, pp. 207-233; C.A., 123 (1995) 147374x - a review with 52 refs.

- 962 Ye, M.Y., Walkup, R.G. and Hill, K.D.: Separation of octylphenol polyether alcohols surfactants by capillary column SFC and HPLC. *J. Liq. Chromatogr.*, 18 (1995) 2309-2322.

See also 927, 982.

36b. Antioxidants and preservatives

See 658, 680, 844.

36c. Complex mixtures, technical products and unidentified compounds

- 963 Godard, M., Bodinier, J.-L. and Vasseur, G.: Effect of mineralogical reactions on trace during redistribution in mantle rocks during percolation processes: A chromatographic approach. *Earth Planet. Sci. Lett.*, 133 (1995) 449-461; C.A., 123 (1995) 149213n.
 964 Nemeth, Z.: The peak tracking role of the PRIMA method in liquid chromatography. *ACH-Models Chem.*, 13 (1994) 835-845; C.A., 123 (1995) 101715a.
 965 Rovira, J., Aparicio, I. and Almarcha, M.: (Determination of xylene musk in aquatic organisms). *Aliment., Equipos Tecnol.*, 14 (1995) 79-81; C.A., 123 (1995) 137782j.
 966 Sokolov, O.M., Kochergina, G.G. and Maier, L.V.: (Chromatography of technical lignins on various gels). *Izv. Vyssh. Vchebn. Zaved., Lesn. Zh.*, (1994) 80-85; C.A., 123 (1995) 86286r.
 967 Voorhees, K.J.: Analysis of soot produced from the combustion of polymeric materials. *ACS Symp. Ser.*, 599 (1995) 393-406; C.A., 123 (1995) 113807v.
 968 Vuillard, L., Madern, D., Franzetti, B. and Rabilloud, T.: Halophilic protein stabilization by the mild solubility agents nonionic sulfobetaines. *Anal. Biochem.*, 230 (1995) 290-294.

See also 960, 977.

38. INORGANIC COMPOUNDS

38a. Cations

- 969 Bettinelli, M. and Spezia, S.: Determination of trace elements in sea water by ion chromatography-inductively coupled plasma mass spectrometry. *J. Chromatogr. A*, 709 (1995) 275-281.
 970 Bricker, T.M. and Houk, R.S.: Speciation of selenium in human serum by size exclusion chromatography and inductively coupled plasma mass spectrometry. *Spec. Publ. - R.Soc. Chem.*, 163(Frontiers in Analytical Spectroscopy) (1995) 109-116; C.A., 123 (1995) 164365g.
 971 Byrdy, F.A., Olson, L.K., Vela, N.P. and Caruso, J.A.: Chromium speciation by anion-exchange high-performance liquid chromatography with both inductively coupled plasma atomic emission spectroscopic and inductively coupled plasma mass spectrometric detection. *J. Chromatogr. A*, 712 (1995) 311-320.

- 972 Chuveleva, E.A., Firsova, L.A. and Kharitonov, O.V.: (Separation of rare earth elements by displacement complexing chromatography with diethylenetriaminepentaacetic acid solution at elevated pH on various lots of KU-2 cation exchanger). *Zh. Prikl. Khim. (S.-Peterburg)*, 67 (1994) 1787-1790; C.A., 123 (1995) 61926m.
- 973 Chuveleva, E.A., Peshkov, A.S., Kvaritonov, O.V. and Firsova, L.A.: (Effect of cadmium ions and their introduction into the system on the effectiveness of the chromatographic separation of yttrium and europium). *Zh. Fiz. Khim.*, 69 (1995) 1132-1134; C.A., 123 (1995) 131374f.
- 974 Doepke, T., Braun, N. and Wuensch, G.: Determination of halide traces in refractory metals by ion chromatography. In: Hirschfeld, D. (Editor), *Nichtmet. Met. '92*, DGM Informationsges, Oberursel, 1992, pp. 101-108; C.A., 123 (1995) 101519q.
- 975 Firsova, L.A. and Chuveleva, E.A.: (Study of the feasibility of using chelating resins for separating rare-earth elements). *Zh. Fiz. Chim.*, 69 (1995) 358-361; C.A., 123 (1995) 73542w.
- 976 Grudpan, K., Laiwraungrath, A. and Sooksamiti, P.: Flow injection spectrophotometric determination of uranium with in-valve ion-exchange column preconcentration and separation. *Analyst (Cambridge)*, 120 (1995) 2107-2110.
- 977 Hesse, D. and Giersberg, R.: (The use of ion chromatography for the determination of anions and cations in the leather manufacture). *Leder*, 46 (1995) 46-61; C.A., 123 (1995) 173350a.
- 978 Hu, W., Miyazaki, A., Tao, H., Itoh, A., Umemura, T. and Haraguchi, H.: Direct determination of inorganic ions at sub-ppb levels by ion chromatography using water as a mobile phase. *Anal. Chem.*, 67 (1995) 3713-3716.
- 979 Ilie, V.A. and Patroescu, C.: (Liquid chromatographic separation and determination of some transition metal and platinum-group metal cations as chelates with mercaptobenzothiazole). *Rev. Chim. (Bucharest)*, 46 (1995) 478-484; C.A., 123 (1995) 131372d.
- 980 Janvion, P., Motellier, S. and Pitsch, H.: Ion-exchange mechanisms of some transition metals on a mixed-bed resin with a complexing eluent. *J. Chromatogr. A*, 715 (1995) 105-115.
- 981 Jones, P. and Beere, H.G.: Ion chromatography determination of trace silver ions using hydrophylic resins with chemiluminescence detection. *Anal. Proc.*, 32 (1995) 169-171; C.A., 123 (1995) 73665p.
- 982 Klyuchnikova, N.V., Denisova, L.V., Balyatinskaya, L.N. and Akmetzhanov, I.S.: (Trace determination of heterocharged ions in nonionic surfactants by ion chromatography). *Zavod. Lab.*, 61 (1995) 13-15; C.A., 123 (1995) 173566a.
- 983 Manjarekar, S.V. and Argekar, A.P.: Cation exchange chromatographic separation of zinc in mixed solvents and non-aqueous hydrobromic acid media. *Anal. Lett.*, 28 (1995) 1711-1727.
- 984 Mansur, M.S., Mushtag, A. and Muhammad, A.: Separation of ^{111}Ag from neutron irradiated natural palladium. *Radiochim. Acta*, 68 (1995) 161-162; C.A., 123 (1995) 187069c.
- 985 Mohite, B.S. and Zambare, D.N.: Column chromatographic separation of gallium(III) from other elements using poly(dibenzo-18-crown-6). *Indian J. Chem., Sect. A: Inorg., Bio-inorg., Phys., Theor. Anal. Chem.*, 34A (1995) 323-325; C.A., 122 (1995) 280815g.
- 986 Na, C.-k., Nakano, T., Tazawa, K., Sakagawa, M. and Ito, T.: A systematic and practical method of liquid chromatography for the determination of Sr and Nd isotopic ratios and REE concentrations in geological samples. *Chem. Geol.*, 123 (1995) 225-237; C.A., 123 (1995) 187169k.
- 987 Okada, T.: (Chromatographic separation of ions. A review on separation mechanisms). *Bunseki Kagaku*, 44 (1995) 579-601; C.A., 123 (1995) 190043f - a review with 179 refs.
- 988 Padarauskas, A.V.: Ion-pair chromatography of barium, copper, and yttrium with the use of complex formation. *J. Anal. Chem.*, 50 (1995) 465-469; C.A., 123 (1995) 187068b.
- 989 Powell, M.J., Boomer, D.W. and Wiederin, D.R.: Determination of chromium species in environmental samples using high-pressure liquid chromatography direct injection nebulization and inductively coupled plasma mass spectrometry. *Anal. Chem.*, 67 (1995) 2474-2478.
- 990 Ramanujam, A., Achuthan, P.V., Dhami, P.S., Gopalakrishna, V., Kannan, R. and Mathur, J.N.: Extraction chromatographic separation of promethium from high active waste solutions of Purex origin. *Solvent Extr. Ion Exch.*, 13 (1995) 301-312; C.A., 123 (1995) 159549r.
- 991 Schmuckler, G. and Prior, A.: Separation of the platinum group metals by a chromatographic method. *Precious Met.*, 18th (1994) 299-305; C.A., 123 (1995) 175394s.
- 992 Sharma, R.K.: Column chromatographic preconcentration and separation of cobalt in a vitamin, an alloy and an ore using chelating resin containing acenaphthenequinone monoxime supported over naphthalene. *Analyst (Cambridge)*, 120 (1995) 2203-2205.
- 993 Stella, R., Ganzerli Valentini, M.T. and Maggi, L.: Separation of U(IV) and U(VI) by ion exchange chromatography on lanthanum ammonium oxalate. *Appl. Radiat. Isot.*, 46 (1995) 1-5; C.A., 123 (1995) 73541v.
- 994 Sutheimer, S.H. and Cabaniss, S.E.: Aqueous Al(III) speciation by high performance cation exchange chromatography with fluorescence detection of the aluminium-lumogallion complex. *Anal. Chem.*, 67 (1995) 2342-2349.
- 995 Takaya, M. and Sawatari, K.: Speciation of vanadium(IV) and vanadium(V) using ion-exchange chromatography and ICP-AES. *Ind. Health*, 32 (1994) 165-178; C.A., 123 (1995) 162826j.
- 996 Tian, S. and Schwedt, G.: Separation of chromate and tungstate by reversed-phase high-performance chromatography using rutin as chelating reagent. *J. Chromatogr. A*, 709 (1995) 381-386.
- For additional information see C.A.:
123 (1995) 60626b, 61968b, 175479y.
- See also 40, 74, 119, 127, 665, 667, 670, 945.
- 38b. Anions*
- 997 Blanco, D., Martinez, L., Mangas, J.J., Dapena, E. and Gutierrez, D.: Determination of nitrate and nitrite in tap water and vegetables by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 2445-2456.
- 998 Chadha, R.K., Lawrence, J.F. and Ratnayake, W.M.N.: Ion chromatographic determination of cyanide released from flaxseed under autohydrolysis conditions. *Food Addit. Contam.*, 12 (1995) 527-533; C.A., 123 (1995) 110334c.

- 999 Gamoh, K., Kiyomoto, W., Ayumi, K., Matsushima, A. and Sawamoto, H.: (Analytical methods for the determination of trace ion species in environmental water. III. Instrumental comparison in determination methods for sulphite ion by liquid chromatography). *Kochi Daigaku Gakujutsu Kenkyou Hokoku Shizen Kagaku*, 43 (1994) 91-99; C.A., 123 (1995) 122514z.
- 1000 Sakuragawa, A., Takano, Y., Suzuki, M. and Okutani, T.: (Ion chromatographic determination of micro amounts of sulphate ion using a chitosan loaded column). *Bunseki Kagaku*, 44 (1995) 477-481; C.A., 123 (1995) 187101g.

For additional information see C.A.:
123 (1995) 73775z, 110347j, 131430w.

See also 40, 45, 74, 147, 945, 958, 974, 977, 978, 982, 987.

38d. *Volatile inorganic compounds*

- 1001 Makaga-Kabinda-Massard, E. and Maujean, A.: (Simultaneous determination of sulfur dioxide and ascorbic acid in Champagne wines by HPLC with electrochemical detection. Study of the ascorbic acid-sulfur dioxide interaction). *Bull. O.I.V.*, 67 (1994) 753-767; C.A., 123 (1995) 110320v.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 1002 Trubetskoy, V.S. and Torchilin, V.P.: Fast and specific labeling of antibody fragments with multiple atoms of heavy metal radioisotopes. *Anal. Biochem.*, 229 (1995) 345-347.
- 1003 Vucina, J.L.: Comparative quality control of some bone-seeking technetium-99m radiopharmaceuticals by gel chromatography. *J. Radioanal. Nucl. Chem.*, 200 (1995) 291-298; C.A., 123 (1995) 93231x.
- 1004 Vucina, J.L.: Effect of the eluent composition in gel chromatography of $^{99m}\text{Tc}(\text{Sn})$ -methylenediphosphonate. *J. Radioanal. Nucl. Chem.*, 199 (1995) 347-352; C.A., 123 (1995) 123325a.

See also 929, 984, 990.