



ELSEVIER

**Bibliography Section****Liquid Column Chromatography****1. REVIEWS AND BOOKS**

- 2421 Blau, K. and Halket, J.M. (Editors): (*Handbook of Derivatives for Chromatography*). Maruzen, Tokyo, 1996, 374 p.; C.A., 124 (1996) 249250m.
- 2422 Brown, P.R., Grushka, E. (Editors): (*Advances in Chromatography*). Dekker, New York, 1996, 443 p.; C.A., 124 (1996) 249249t.
- 2423 Buchberger, W.: Electrochemical detectors - tailor made techniques for liquid chromatography and capillary electrophoresis? *Fresenius J. Anal. Chem.*, 354 (1996) 797-802.
- 2424 Freitag, R. and Horvath, C.: Chromatography in the downstream processing of biotechnological products. *Adv. Biochem. Eng/Biotechnol.*, 53(Downstream Processing Biosurfactants/Carotenoids) (1996) 17-59; C.A., 124 (1996) 169578j - a review with 383 refs.
- 2425 Ito, Y. and Conway, W.D. (Editors): (*High Speed Countercurrent Chromatography*). In: *Chem. Anal.*, Vol. 132, Wiley, New York, 1996, 454 p.; C.A., 124 (1996) 134422b.
- 2426 Stamm, C.: (Ion chromatographic analysis. Part 2.) *Schweiz. Lab.-Z.*, 53 (1996) 19-23; C.A., 124 (1996) 248840s - a review without refs.
- 2427 Stock, R. and Rice, C.B.F. (Editors): (*Chromatographic Methods*). Blackie, Glasgow, 1995, 424 p.; C.A., 124 (1996) 105031t.
- 2428 Stöckl, D., Franzini, C., Kratochvíla, J., Middle, J., Ricos, C., Siekmann, L. and Thienpont, L.M.: Analytical specification of reference methods compilation and critical discussion. *Eur. J. Clin. Chem. Clin. Biochem.*, 34 (1996) 319-337 - a review.
- 2429 Toft, J.: Evolutionary rank analysis applied to multidetectional chromatographic structures. *Chemom. Intell. Lab. Syst.*, 29 (1995) 189-212; C.A., 124 (1996) 10592p - a review with 63 refs.
- 2430 Van de Merbel, N.C., Lingeman, H. and Brinkman, U.A.T.: Sampling and analytical strategies in on-line bioprocess monitoring and control. *J. Chromatogr. A*, 725 (1996) 13-27 - a review with 37 refs.

See also 2449, 2452, 2456, 2494, 2495, 2500, 2504, 2508, 2532, 2533, 2540, 2551, 2554, 2560, 2561, 2565, 2598, 2602, 2603, 2620, 2634, 2649, 2653, 2654, 2655, 2659, 2660, 2668, 2671, 2744, 2750, 2754, 2830, 2834, 2865, 2890, 2914, 3010, 3019, 3025, 3259, 3306, 3347, 3467, 3471, 3478, 3482, 3483, 3490, 3599, 3677, 3723, 3740, 3748, 3766, 3772, 3796.

**2. FUNDAMENTALS, THEORY AND GENERAL****2a. General**

- 2431 Carr, P.W., Tan, L.C. and Park, J.H.: Revisionist look at solvophobic driving forces in reversed-phase liquid chromatography. III. Comparison of the behavior of nonpolar and polar solutes. *J. Chromatogr. A*, 724 (1996) 1-12.
- 2432 Claessens, H.A., van Straten, M.A. and Kirkland, J.J.: Effect of buffers on silica-based column stability in reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 728 (1996) 259-270.
- 2433 Danilov, A.V., Mustaeva, I.G., Vaginina, I.V. and Baru, M.B.: Size-exclusion chromatography on soft packing material under axial compression. *J. Chromatogr. A*, 732 (1996) 17-25.
- 2434 Dolan, J.W.: A second shot at some troubleshooting problems. *LC-GC*, 13 (1995) 940-943; C.A., 124 (1996) 134102x.
- 2435 Dolan, J.W.: Injection problems. *LC-GC*, 14 (1996) 20-24; C.A., 124 (1996) 163353k.
- 2436 Dolan, J.W.: Injection problems. *LC-GC Int.*, 9 (1996) 70-75.
- 2437 Dolan, J.W.: Using a gradient scouting run to get started. *LC-GC Int.*, 9 (1996) 130-136.
- 2438 Dolan, J.W., Kern, J.R. and Culley, T.: Water problems - a case study. *LC-GC Int.*, 9 (1996) 194-200.
- 2439 Ettre, S.: M.S. Tswett and the 1918 Nobel prize in chemistry. *Chromatographia*, 42 (1996) 343-351.
- 2440 Ford, J.C. and Ko, J.: Comparison of methods for extracting linear solvent strength gradient parameters from gradient chromatographic data. *J. Chromatogr. A*, 727 (1996) 1-11.
- 2441 Garrido Frenich, A., Martínez Galera, M., Martínez Vidal, J.L. and Gil García, M.D.: Partial least-squares and principal component regression of multi-analyte high-performance liquid chromatography with diode-array detection. *J. Chromatogr. A*, 727 (1996) 27-38.
- 2442 Guan, H. and Guiuchon, G.: Reproducibility, accuracy, and correction of isotherm data measured by chromatography. *J. Chromatogr. A*, 724 (1996) 39-54.
- 2443 Hayashi, Y. and Matsuda, R.: Noise and measurement precision in analytical instruments. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 35-44; C.A., 124 (1996) 75099b.
- 2444 Herten, S.: ACS award in chromatography. *Chem. Eng. News*, 74 (1996) 82; C.A., 124 (1996) 201131w.

- 2445 Kimata, K., Kobayashi, M., Hosoya, K., Araki, T. and Tanaka, N.: Chromatographic separation based on isotopic chirality. *J. Am. Chem. Soc.*, 118 (1996) 759-762; C.A., 124 (1996) 201910z.
- 2446 Li, X. and Fritz, J.S.: Novel additives for the separation of organic compounds by high-performance liquid chromatography. *J. Chromatogr. A*, 728 (1996) 235-247.
- 2447 Lough, W.J., Mills, M.J. and Maltas, J.: Analyte adsorption in liquid chromatography valve injectors for samples in non-eluting solvents. *J. Chromatogr. A*, 726 (1996) 67-75.
- 2448 Metting, H.J. and Coenegracht, P.M.J.: Neural networks in high-performance liquid chromatography optimization: response surface modeling. *J. Chromatogr. A*, 728 (1996) 47-53.
- 2449 Outinen, K., Lehtonen, P. and Vuorela, H.: (Optimization methods in liquid chromatography. The principles). *Kem.-Kemi*, 22 (1995) 592-596; C.A., 124 (1996) 192842z - a review with 54 refs.
- 2450 Pecanac, D., Karljikovic-Rajic, K. and Radulovic, D.: Copper(II)-tartrate system as a potential mobile phase additive for enantioseparation of some chiral pharmaceuticals. *Pharmazie*, 51 (1996) 124-125.
- 2451 Seaver, S., Przybytek, J. and Roelofs, N.: Solvent selection Part III-Solvent life and degradation. *LC-GC*, 13 (1995) 860-864; C.A., 124 (1996) 75092u.
- 2452 Stamm, C.: (Ion chromatographic analysis. Part 1). *Schweiz. Lab.-Z.*, 52 (1995) 393-397; C.A., 124 (1996) 192416p - a review without refs.
- For additional information see C.A.:  
124 (1996) 127885b.
- See also 2427, 2429, 2470, 2490, 2522, 2564, 2568, 2576, 2589, 2660, 2661, 2710, 3487, 3549, 3817.
- 2b. Thermodynamics and theoretical relationships*
- 2453 Azzaoui, K. and Morin-Allory, L.: Comparison and quantification of chromatographic retention mechanisms and three stationary phases using structure-retention relationships. *Chromatographia*, 42 (1996) 389-395.
- 2454 Bruzzoniti, M.C., Mentasti, E., Sacchero, G. and Sarzanini, C.: Retention model for singly and doubly charged analytes in ion-interaction chromatography. *J. Chromatogr. A*, 728 (1996) 55-65.
- 2455 Felinger, A. and Guiochon, G.: Multicomponent interferences in overloaded gradient elution chromatography. *J. Chromatogr. A*, 724 (1996) 27-37.
- 2456 Han, D., Ye, M. and Shi, L.: (Non-size exclusion effect in aqueous gel permeation chromatography). *Sepu*, 13 (1995) 432-436; C.A., 124 (1996) 163738q - a review with 69 refs.
- 2457 Heron, S. and Tchapla, A.: Retention mechanism in reversed-phase liquid chromatography: specific effects of propiophenyl and multifunctional (cyanopropyl-octadecyl) bonded silicas. *J. Chromatogr. A*, 725 (1996) 205-218.
- 2458 Hong, H., Wang, L., Han, S. and Zou, G.: Prediction [of] adsorption coefficients ( $K_{oc}$ ) for aromatic compounds by HPLC retention factors ( $K'$ ). *Chemosphere*, 32 (1996) 343-351; C.A., 124 (1996) 144564p.
- 2459 Kantoci, D.: A study of an empirical equation for the evaluation of separation efficiency in chromatography. *J. Liq. Chromatogr.*, 19 (1996) 853-866.
- 2460 Lavine, B.K. and Hendayana, S.: Band broadening in micellar liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 101-123.
- 2461 Li, J.: A simplified exponentially modified Gaussian function for modelling chromatographic peaks. *J. Chromatogr. Sci.*, 33 (1995) 568-572.
- 2462 Lim, B.-G., Ching, C.-B. and Tan, R.B.H.: Determination of competitive adsorption isotherms of enantiomers on a dual-site adsorbent. *Sep. Technol.*, 5 (1995) 213-228; C.A., 124 (1996) 98757d.
- 2463 Reese, C.E., Huang, L., Hsu, S.-H., Tripathi, S. and Lochmüller, C.H.: Universal retention indices and their prediction in reversed-phase liquid chromatography based on principal component analysis and target testing. *J. Chromatogr. Sci.*, 34 (1996) 101-110.
- 2464 Row, K.H. and Larin, A.V.: A chromatographic theory based on the concept of a layer of equilibrium adsorption. *Korean J. Chem. Eng.*, 12 (1995) 442-447; C.A., 124 (1996) 67760v.
- 2465 Row, K.H. and Larin, A.V.: Experimental study of influence of slight deviation from adsorption isotherm linearity on elution peak profiles. *J. Chem. Eng. Jpn.*, 28 (1995) 851-853; C.A., 124 (1996) 127821c.
- 2466 Sajonz, P., Zhong, G. and Guiochon, G.: Influence of the concentration dependence of the mass transfer properties on chromatographic band profiles. II. Accuracy of the determination of isotherm data by frontal analysis. *J. Chromatogr. A*, 731 (1996) 1-25.
- 2467 Vailaya, A. and Horvath, C.: Enthalpy-entropy compensation in hydrophobic interaction chromatography. *J. Phys. Chem.*, 100 (1996) 2447-2455; C.A., 124 (1996) 98750w.
- 2468 Verdu-Andres, J., Reig, F.B. and Campins-Falco, P.: H-Point standard additions method for resolution of overlapped chromatographic peaks with a conventional fluorescence detection. Determination of phenol and cresols in waters. *Chromatographia*, 42 (1996) 283-289.
- 2469 Wan, Q.-H., Davies, M.C., Shaw, P.N. and Barrett, D.A.: Retention behavior of ionizable isomers in reversed-phase liquid chromatography: a comparative study of porous graphitic carbon and octadecyl bonded silica. *Anal. Chem.*, 68 (1996) 437-446.
- 2470 Xu, Y. and Liapis, A.I.: Modelling and analysis of the elution stage of "perfusion chromatography". Effects of intraparticle convective velocity and microsphere size on system performance. *J. Chromatogr. A*, 724 (1996) 13-25.
- For additional information see C.A.:  
124 (1996) 169673m.
- See also 2442, 2479, 2524, 2528, 2544, 2546, 2553, 2570, 2587, 2592, 2617, 2646, 2665, 2694, 2875, 3281, 3417, 3417, 3425, 3460, 3632, 3690, 3804.
- 2c. Relationship between structure and chromatographic behaviour*
- For additional information see C.A.:  
124 (1996) 192801k.
- See also 2509, 2582, 2708, 2716, 3459, 3551, 3762, 3767.

## 2d. Measurement of physico-chemical and related values

- 2471 Albertsson, A.-C. and Karlsson, S.: Chromatographic fingerprinting as a means to predict degradation mechanisms. *J. Environ. Polym. Degrad.*, 4 (1996) 51-53; C.A., 124 (1996) 203279z.
- 2472 Berthod, A., Mallet, A.I. and Bully, M.: Measurement of partition coefficients in waterless biphasic liquid systems by counter-current chromatography. *Anal. Chem.*, 68 (1996) 431-436.
- 2473 Cheong, W.J., Chun, S.H. and Lee, G.Y.: Determination of solvent basicity scale,  $\beta$ , of mixed solvents for three chromatographic solvent systems: 2-propanol/hexane, ethyl acetate/hexane, and methanol/water. *J. Liq. Chromatogr.*, 19 (1996) 277-291.
- 2474 Denuziere, A., Yagoubi, N., Baillet, A. and Ferrier, D.: An improved statistical parameter allowing elaborate comparison of polymer molecular weight distribution by gel filtration chromatography: Application to chitosan. *S.T.P. Pharma Sci.*, 5 (1995) 481-485; C.A., 124 (1996) 211821k.
- 2475 Huang, N.-J. and Sundberg, D.C.: A gel permeation chromatography method to determine grafting efficiency during graft copolymerization. *Polymer*, 35 (1994) 5693-5698; C.A., 124 (1996) 56936f.
- 2476 Kilz, P.: (Quality control of polymers by gel permeation chromatography (GPC)). *GIT Fachz. Lab.*, 39 (1995) 944-949; C.A., 124 (1996) 57250q.
- 2477 Shibukawa, M., Aoki, A. and Oguma, K.: A new liquid chromatographic method for the determination of acid dissociation constants. *Anal. Sci.*, 11 (1995) 915-922; C.A., 124 (1996) 127811z.
- 2478 Silva, A.: Molecular weight distribution analysis of wood pulp cellulose by size exclusion chromatography. Avail. *Univ. Microfilms Int.*, Order No.: DA9604330, 1996, 1125 p.; C.A., 124 (1996) 205364x.
- 2479 Velayudhan, A. and Horvath, C.: Isotherm measurement by frontal chromatography in the presence of an adsorbing mobile phase modulator. *Ind. Eng. Chem. Res.*, 35 (1995) 1173-1179; C.A., 124 (1996) 205911y.
- 2480 Yamagami, C. and Masaki, Y.: Hydrophobicity parameters determined by reversed-phase liquid chromatography. X. Relationship between capacity factor and octanol-water partition coefficients of monosubstituted thiophenes. *Chem. Pharm. Bull.*, 43 (1995) 2238-2242.

For additional information see C.A.:

124 (1996) 67666u, 90884x.

See also 2491, 2534, 2543, 2548, 2593, 2698, 2700, 2960, 3451, 3462, 3463, 3464, 3706, 3764.

## 3. GENERAL TECHNIQUES

## 3a. Apparatus and accessories

- 2481 Baram, G.I.: Portable liquid chromatograph for mobile laboratories. I. Aims. *J. Chromatogr. A*, 728 (1996) 387-399.

- 2482 Barka, G.: Method and apparatus for preparation of solvent mixtures in gradient-elution high-pressure liquid chromatography. *Ger. Offen.* DE 4,412,703 (Cl. G01N30/36), 26 Oct. 1995, Appl. 4,412,703, Apr. 1994; 7 p.; C.A., 124 (1996) 75186c.
- 2483 Guenther, J., Krause, B., Mader, G. and Sitting, W.: Low pressure chromatography column with automatically movable cover for ions. *Ger. Offen.* DE 4,431,662 (Cl. B01D15/08), 7 Mar. 1996, Appl. 4,431,662, 6 Sep. 1994; 4 p.; C.A., 124 (1996) 235731g.
- 2484 Herraez-Hernandez, R., Campins-Falco, P. and Sevillano-Cabeza, A.: On-line derivatization into precolumns for the determination of drugs by liquid chromatography and column switching: determination of amphetamines in urine. *Anal. Chem.*, 68 (1996) 734-739.
- 2485 Huber, L.: A cost-effective HPLC system for easy validation. *Int. Lab.*, 26, January (1996) 14-17.
- 2486 Ladisch, M., Hamaker, K., Hendrickson, R. and Brewer, M.: Device for packing chromatographic stationary phases in columns. *PCT Int. Appl.* WO 95 34,361 (Cl. B01D15/08), 21 Dec. 1995, US Appl. 260,021, 15 Jun. 1994; 28 p.; C.A., 124 (1996) 120911y.
- 2487 Matsushima, Y., Kanazawa, H., Yamamoto, K., Takai, S., Sakurai, Y. and Okano, M.: (Chromatography and column filler for biological sample). *Jpn. Kokai Tokkyo Koho JP 07,318,551 [95,318,551]* (Cl. G01N30/48), 8 Dec. 1995, Appl. 94/108,643, 23 May 1994; 9 pp.; C.A., 124 (1996) 170005b.
- 2488 Schick, H.G.: Column for liquid chromatography. *U.S. US 5,482,628* (Cl. 210-198.2; B01D15/08), 9 Jan. 1996, Appl. 277,853, 15 Apr. 1994; 10 p.; C.A., 124 (1996) 179650r.
- 2489 Vissers, J., Chervet, J.-P. and Salzmann, J.-P.: A fully automated microautosampler for micro and capillary liquid chromatography. *Int. Lab.*, 26, January (1996) 12A-12H.

For additional information see C.A.:

124 (1996) 60085v, 71090u, 75045f, 81469p, 91158a, 105043y, 111723n, 149897j, 163656m, 163672p, 163715e, 179805v, 218894v, 219010r, 249080f.

See also 2539, 2577, 2654.

## 3b. Detectors and detection reagents

- 2490 Boque, R. and Rius, F.X.: Multivariate detection limits estimators. *Chemom. Intell. Lab. Syst.*, 32 (1996) 11-23; C.A., 124 (1996) 218721m.
- 2491 Busnel, J.-P., Degoulet, C., Nicolai, T., Woodley, W. and Patin, P.: Multangle light scattering and viscometric detector for size-exclusion chromatography. *J. Phys. III*, 5 (1995) 1501-1512; C.A., 124 (1996) 118640r.
- 2492 Dreux, M., Lafosse, M. and Morin-Allory, L.: The evaporative light scattering detector - a universal instrument for non-volatile solutes in LC and SFC. *LC-GC Int.*, 9 (1996) 148-156.
- 2493 Foster, M.D. and Synovec, R.E.: Liquid chromatographic sensing in water on a thin-clad optical fibre by mode-filtered light detection. *Anal. Chem.*, 68 (1996) 1456-1463.
- 2494 Goto, J.: (Detection-oriented derivatization for liquid chromatography). *Kuromatogurafu*, 16 (1995) 324-325; C.A., 124 (1996) 127591c - a review without refs.

- 2495 Hayakawa, K.: (Advances of indirect photometric detection in LC: inorganic ions and optically active compounds). *Kuromatogurafu*, 16 (1995) 314-317; C.A., 124 (1996) 127590b - a review with 11 refs.
- 2496 Li, G., Yu, J., Krull, I.S. and Cohen, S.: An improvement for the synthesis of a styrene-divinylbenzene copolymer based, 6-aminoquinoline carbamate reagent. Applications for derivatization of amino acids, peptides, and proteins. *J. Liq. Chromatogr.*, 18 (1995) 3889-3918.
- 2497 Lin, Z.-P. and Aue, W.A.: Quantitation of perfectly co-eluting analytes on dual-channel-subtraction chromatograms. *J. Chromatogr. A*, 727 (1996) 101-109.
- 2498 Lores, M., Garcia, C.M. and Cela, R.: Selectable-power photoreactor for flow-injection analysis systems and high-performance liquid chromatography post-column photochemical derivatization. *J. Chromatogr. A*, 724 (1996) 55-65.
- 2499 Mank, A.J.G., de Nijs, H., Lingeman, H., Brinkman, U.A.T., Velthorst, N.H. and Gooijer, C.: Diode laser-based absorption detector for conventional-size liquid chromatography. *Appl. Spectrosc.*, 50 (1996) 28-34; C.A., 124 (1996) 163772w.
- 2500 Mank, A.J.G., Lingeman, H. and Gooijer, C.: Diode laser-based detection in liquid chromatography and capillary electrophoresis. *TrAC*, 15 (1996) 1-11 - a review.
- 2501 Novak, T.J.: An investigation of the luminescence of 10-methyl-acridinium carboxylate salts and evaluation as a chemiluminescent derivatizing reagent for high performance liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No.: DA9533457, 1994, 141 p.; C.A., 124 (1996) 192578t.
- 2502 Papadopoulou-Mourkidou, E. and Patsias, J.: Development of a semi-automated high-performance liquid chromatographic-diode array detection system for screening pesticides at trace levels in aquatic systems of the Axios River basin. *J. Chromatogr. A*, 726 (1996) 99-113.
- 2503 Pravda, M., Petit, C., Michotte, Y., Kauffmann, J.-M. and Vytras, K.: Study of a new solid carbon paste tyrosinase-modified amperometric biosensor for the determination of catecholamines by high-performance liquid chromatography. *J. Chromatogr. A*, 727 (1996) 47-54.
- 2504 Scheper, T.H., Hilmer, J.M., Lammers, F., Müller, C. and Reinecke, M.: Biosensors in bioprocess monitoring. *J. Chromatogr. A*, 725 (1996) 3-12 - a review with 26 refs.
- 2505 Seidel, B.S., Dübel, O., Faubel, W. and Ache, H.J.: Trace analysis of pollutants in water using the photothermal interferometry as HPLC detector. *Fresenius J. Anal. Chem.*, 354 (1996) 900-902.
- 2506 Voegel, P.D. and Baldwin, R.P.: Electrochemical detection with copper electrodes in liquid chromatography and capillary electrophoresis. *Int. Lab.*, 26, March (1996) 16A-16N.
- 2507 Watanabe, H., Yokoyama, Y. and Sato, H.: High-sensitivity conductivity detection in non-suppressed ion chromatography using sulfoisophthalic acid as eluent. *J. Chromatogr. A*, 727 (1996) 311-316.
- 2508 Webster, G.: (Is the diode array detector (DAD) the future standard in the quality control laboratory?) *LaborPraxis*, 19 (1995) 68-77; C.A., 124 (1996) 75080p - a review with 5 refs.
- See also 2423, 2441, 2468, 2642, 2674, 2682, 2704, 2717, 2721, 2763, 2767, 2817, 2874, 2896, 2916, 2941, 2955, 3083, 3244, 3362, 3427, 3489, 3519, 3520, 3546, 3639, 3702, 3769, 3834, 3837, 3838.
- 3c. *Sorbents and columns, packing procedures*
- 2509 Barrett, D.A., Brown, V.A., Shaw, P.N., Davies, M.C., Ritchie, H. and Ross, P.: Characterization of a range of alkyl-bonded silica HPLC stationary phases: chromatographic behavior of neutral, acidic, and basic test solutes. *J. Chromatogr. Sci.*, 34 (1996) 146-156.
- 2510 Bede, Á., Rippel, G., Szepesy, L. and Claessens, H.A.: Characterization of wide-pore reversed-phase columns for biopolymer separations. I. Single-parametric evaluation. *J. Chromatogr. A*, 728 (1996) 179-188.
- 2511 Benmoussa, A., Lacout, J.L., Loiseau, P.R. and Mikou, M.: Calcium phosphate hydroxyapatite and lead phosphate hydroxyapatite for separation by liquid chromatography. *Chromatographia*, 42 (1996) 177-180.
- 2512 Brindle, R., Albert, K., Harris, S.J., Tröltzsch, C., Horne, E. and Glennon, J.D.: Silica-bonded calixarenes in chromatography. I. Synthesis and characterization by solid-state NMR spectroscopy. *J. Chromatogr. A*, 731 (1996) 41-46.
- 2513 Chyuch, S.-D. and Liu, C.-Y.: Functional crown ether derivatives as sorbents for ligand exchange chromatography. *Fresenius J. Anal. Chem.*, 354 (1996) 278-283.
- 2514 Ciucanu, I.: Selective immobilization on silica gel of permethylated  $\beta$ -cyclodextrin for liquid chromatography. *J. Chromatogr. A*, 727 (1996) 195-201.
- 2515 Davydov, V.Ya. and Filatova, G.N.: Silica with bonded fullerenes as adsorbents for chromatography. *Proc.-Electrochem. Soc.*, 95-10 (1995) 1603-1609; C.A., 124 (1996) 163452s.
- 2516 Dawidowicz, A.L., Wasilewska, D. and Radkiewicz, S.: Siliceous materials with the surface polymer layer composed of dextran-polyimine mixture as column packings for liquid chromatography. *Chromatographia*, 42 (1996) 49-55.
- 2517 Denizli, A., Salih, B. and Piskin, E.: Congo Red and Cu(II) carrying poly(ethylene glycol dimethacrylate-hydroxyethyl methacrylate) microbeads as specific sorbents. Albumin adsorption/desorption. *J. Chromatogr. A*, 731 (1996) 57-63.
- 2518 Dumont, P.J.: New methods and materials for solid phase extraction and high performance liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No.: DA9540891, 1995, 124 p.; C.A., 124 (1996) 163457x.
- 2519 Dunlap, C.J., Carr, P.W. and McCormick, A.V.: A chromatographic comparison of the pore structures of zirconia high performance liquid chromatographic materials made by the polymerization induced colloidal aggregation and the oil-emulsion methods. *Chromatographia*, 42 (1996) 273-282.
- 2520 Girot, P. and Boschetti, E.: Passivated porous supports for immobilization of chromatographic separation of biologicals. U.S. US 5,470,463 (Cl. 210-198.2; B01D15/08), 28 Nov. 1995, US Appl. 901,326, 19 Jun. 1992; 22 p.; C.A., 124 (1996) 149883b.
- 2521 Glennon, J.D., Horne, E., Hall, K., Cocker, D., Kuhn, A., Harris, S.J. and McKerney, M.A.: Silica-bonded calixarenes in chromatography. II. Chromatographic retention of metal ions and amino acid ester hydrochlorides. *J. Chromatogr. A*, 731 (1996) 47-55.

For additional information see C.A.:

124 (1996) 134319y, 219000n, 246036k.

- 2522 Gowanlock, D., Bailey, R. and Cantwell, F.F.: Intra-particle sorption rate and liquid chromatographic bandbroadening in porous polymer packings. I. Methodology and validation of the model. *J. Chromatogr. A*, 726 (1996) 1-23.
- 2523 Grieb, S.J., Matlin, S.A. and Belenguer, A.M.: Flash chiral chromatography with cellulose tris(3,5-dimethylphenylcarbamate)-coated phases. Improved resolution of basic analytes. *J. Chromatogr. A*, 728 (1996) 195-199.
- 2524 Guan, H. and Guiuchon, G.: Study of physico-chemical properties of some packing materials. I. Measurements of the external porosity of packed columns by inverse size-exclusion chromatography. *J. Chromatogr. A*, 731 (1996) 27-40.
- 2525 Guenu, S. and Hennion, M.-C.: Prediction from liquid chromatographic data of obligatory backflush desorption from solid-phase extraction cartridges packed with porous graphitic carbon. *J. Chromatogr. A*, 725 (1996) 57-66.
- 2526 Haky, J.E., Blauvelt, T.M. and Wieserman, L.F.: Perfluorooctyl and perfluorobutyl bonded alumina stationary phases for high performance liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 307-332.
- 2527 Hosoya, K., Yoshizako, K., Shirasu, Y., Kimata, K., Araki, T., Tanaka, N. and Haganaka, J.: Molecularly imprinted uniform-size polymer-based stationary phase for high-performance liquid chromatography. Structural contribution of cross-linked polymer network on specific molecular recognition. *J. Chromatogr. A*, 728 (1996) 139-147.
- 2528 Hsieh, S. and Jorgenson, J.W.: Preparation and evaluation of slurry-packed liquid chromatography microcolumns with inner diameters from 12 to 30 µm. *Anal. Chem.*, 68 (1996) 1212-1217.
- 2529 Hyun, M.H. and Lee, J.B.: A new chiral stationary phase derived from cyclohexylamide derivative of (S)-naproxen for the liquid chromatographic resolution of enantiomers. *Bull. Korean Chem. Soc.*, 16 (1995) 977-980; C.A., 124 (1996) 105139j.
- 2530 Ihara, H., Nakamura, N., Nagaoka, S. and Hitayama, C.: Immobilization of lipid membrane on silica gels for high performance liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 231-235; C.A., 124 (1996) 75109e.
- 2531 Inoue, Y., Katsumata, Y., Tani, K. and Suzuki, Y.: Determination of the specific interactions of disubstituted aromatic compounds on hydrous zirconium oxide-loaded porous polymer resin. *Chromatographia*, 42 (1996) 396-400.
- 2532 Itoyama, K.: (Applications of chromatographic separation with chitosan). *Gekkan Fudo Kemikaru*, 12 (1996) 19-23; C.A., 124 (1996) 200326b - a review with 13 refs.
- 2533 Jaroniec, M.: Energetic heterogeneity of porous inorganic oxides: adsorption and chromatographic studies. *Stud. Surf. Sci. Catal.*, 99 (1996) 411-433; C.A., 124 (1996) 243053c - a review with 189 refs.
- 2534 Jaroniec, M.: Studies of interfacial properties of chemically bonded phases by sorption and liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 211-219; C.A., 124 (1996) 67686a.
- 2535 Li, J. and Cantwell, F.F.: Intra-particle sorption rate and liquid chromatographic. III. Diffusion in the polymer matrix as the cause of slow sorption. *J. Chromatogr. A*, 726 (1996) 37-44.
- 2536 Li, J., Litwinson, L.M. and Cantwell, F.F.: Intra-particle sorption rate and liquid chromatographic bandbroadening in porous polymer packings. II. Slow sorption rate on a microparticle packing. *J. Chromatogr. A*, 726 (1996) 25-36.
- 2537 Liao, J.-L., Li, Y.M. and Hjertén, S.: Continuous beds for micro-chromatography: reversed-phase chromatography. *Anal. Biochem.*, 234 (1996) 27-30.
- 2538 Lu, H. and Rutan, S.C.: Solvatochromic studies on reversed-phase liquid chromatographic phases. 2. Characterization of stationary and mobile phases. *Anal. Chem.*, 68 (1996) 1387-1393.
- 2539 Majors, R.E.: New chromatography columns and accessories at the 1996 Pittsburgh Conference, Part I. *LC-GC Int.*, 9 (1996) 214-227.
- 2540 Majors, R.E.: The role or column diameter in chromatography and capillary electrophoresis. *LC-GC*, 14 (1996) 10-19; C.A., 124 (1996) 98435x - a review with 17 refs.
- 2541 Matsui, J., Miyoshi, Y., Matsui, R. and Takeuchi, T.: Rod-type affinity media for liquid chromatography prepared by *in-situ*-molecular imprinting. *Anal. Sci.*, 11 (1995) 1017-1019; C.A., 124 (1996) 163783a.
- 2542 Nakagama, T., Murata, K., Hanno, Y., Sugimoto, Y., Ihara, T. and Hobo, T.: Study on preparation method of chiral stationary phases in high-performance liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 719-723; C.A., 124 (1996) 105132b.
- 2543 Nielsen, M.W.F. and van de Ven, H.J.F.M.: Characterization of (methoxymethyl)melamine resin by combined chromatographic/mass spectrometric techniques. *Rapid Commun. Mass Spectrom.*, 10 (1996) 74-81; C.A., 124 (1996) 88812d.
- 2544 Ohkuma, T., Dobashi, A. and Hara, S.: Delayed adsorption on slowly accessible stationary phase surfaces in porous liquid chromatographic column packings. *J. Chromatogr. A*, 728 (1996) 97-112.
- 2545 Okamoto, M., Kakamu, H., Nobuhara, K. and Ishii, D.: The effect of silver modified silica on the chromatographic behavior in high-performance liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 259-265; C.A., 124 (1996) 75124f.
- 2546 Park, J.H., Choe, J.J. and Jang, M.D.: Characterization of RPLC stationary phases based on linear solvation energy relationships. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 537-557; C.A., 124 (1996) 105131a.
- 2547 Pettersson, C., Porrvik, I. and Eriksson, E.: A liquid chromatography column with monolithic porous matrix. *PCT Int. Appl. WO 95 34,359* (Cl. B01D15/08), 21 Dec. 1995, SE Appl. 94/2,091, 14 Jun. 1994; 11 p.; C.A., 124 (1996) 120912z.
- 2548 Retuert, J.: Size exclusion chromatography study on the tetraethyl orthosilicate sol-gel derived system. *Bol. Soc. Chil. Quim.*, 40 (1995) 415-419; C.A., 124 (1996) 88626w.
- 2549 Schwarz, A. and Wilchek, M.: Chromatography adsorbents for immunoglobulins utilizing mercapto heterocyclic ligands. *PCT Int. Appl. WO 95 31279* (Cl. B01320/22), 23 Nov. 1995, US Appl. 243,861, 16 May 1994; 35 p.; C.A., 124 (1996) 149887f.

- 2550 Seibert, D.S., Poole, C.F. and Abraham, M.H.: Retention properties of spacer-bonded propanediol sorbent for reversed-phase liquid chromatography and solid-phase extraction. *Analyst (Cambridge)*, 121 (1996) 511-520.
- 2551 Svec, F.: (Why is the efficiency and selectivity of polymeric stationary phases for HPLC different from those based on silica?) *Chem. Listy*, 90 (1996) 103-108; C.A., 124 (1996) 177330n - a review with 38 refs.
- 2552 Takeuchi, T., Miwa, T., Hum, R. and Chu, J.: Retention behavior on ion exchangers dynamically modified by ions with alkyl chains in microcolumn liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 501-511; C.A., 124 (1996) 75138p.
- 2553 Tani, K. and Suzuki, Y.: Influence of titania matrix on retention behavior in reversed-phase liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 301-306; C.A., 124 (1996) 75128k.
- 2554 Tarasevich, Yu.I.: (Natural mineral adsorbents and semisynthetic adsorption materials prepared from them). *Ross. Khim. Zh.*, 39 (1995) 52-61; C.A., 124 (1996) 243036z - a review with 70 refs.
- 2555 Turowski, M., Kalisz, R., Lüllmann, C., Genieser, H.G. and Jastorff, B.: New stationary phases for the high-performance liquid chromatographic separation of nucleosides and cyclic nucleotides. Synthesis and chemometric analysis of retention data. *J. Chromatogr. A*, 728 (1996) 201-211.
- 2556 Yamaki, S., Isobe, T., Okuyama, T. and Shinoda, T.: Reversed-phase liquid chromatography on a microspherical carbon column at high temperature. *J. Chromatogr. A*, 728 (1996) 189-194.
- 2557 Yamamoto, F.M. and Rokushika, S.: Investigation stationary phase in reversed phase chromatography by thermodynamic treatment. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr. 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 605-609; C.A., 124 (1996) 7508x.
- 2558 Yang, Y.-B., Harrison, K. and Kindsvater, J.: Characterization of a novel stationary phase derived from a hydrophilic polystyrene-based resin for protein cation-exchange high-performance liquid chromatography. *J. Chromatogr. A*, 723 (1996) 1-10.
- 2559 Zimina, T., Smith, R.M., Highfield, J.C., Myers, P. and King, B.W.: Study of the flow development during the slurry packing of microcolumns for liquid chromatography. *J. Chromatogr. A*, 728 (1996) 33-45.

For additional information see C.A.:

- 124 (1996) 58483e, 75118g, 91166b, 91610s, 119735n, 127886c, 134369q, 140385h, 140386j, 143284j, 147225q, 177527g, 179646u, 180210d, 192752v, 192802m, 192940e, 197365f, 197399v, 197400p, 205804r, 219009x, 235171z.

- See also 2457, 2469, 2486, 2488, 2604, 2605, 2609, 2612, 2629, 2632, 2633, 2635, 2636, 2639, 2641, 2671, 2683, 2691, 2900, 2924, 2929, 2943, 2970, 3021, 3248, 3280, 3292, 3297, 3317, 3328, 3432, 3467, 3597, 3731, 3816, 3820, 3831, 3833.

### 3d. Quantitative analysis

- 2560 Jenke, D.R.: Chromatographic method validation: a review of current practices and procedures. I. General concepts and guidelines. *J. Liq. Chromatogr.*, 19 (1996) 719-736 - a review with 49 refs.
- 2561 Jenke, D.R.: Chromatographic method validation: a review of current practices and procedures. II. Guidelines for primary validation parameters. *J. Liq. Chromatogr.*, 19 (1996) 737-757 - a review with 37 refs.
- See also 3382, 3732.
- ### 3e. Preparative scale chromatography
- 2562 Ambedkar, S.S. and Deshpande, B.S.: Expanded bed chromatography and radial flow column chromatography: the approaches for adsorption chromatography of biomolecules on industrial scale. *Hind. Antibiot. Bull.*, 36 (1995, Pub. 1994) 164-172; C.A., 124 (1996) 111351p.
- 2563 Asplund, S.E. and Edvinsson, R.K.: A method for the rapid simulation of preparative liquid chromatography. *Comput. Chem. Eng.*, 20 (1996) 507-516; C.A., 124 (1996) 205930d.
- 2564 Chambers, T.K. and Fritz, J.S.: High-performance liquid chromatography with non-aqueous solvents. Relative elution strength on a polystyrene-divinylbenzene column. *J. Chromatogr. A*, 728 (1996) 271-278.
- 2565 Combs, G.L. and Cook, R.L.: Process chromatography. In: McLennan, F. and Kowalski, B.R. (Editors), *Process Anal. Chem.*, Blackie, Glasgow, 1995, pp. 106-158; C.A., 124 (1996) 218942j - a review with 21 refs.
- 2566 Frey, D.D.: Local-equilibrium behavior of retained pH and ionic strength gradients in preparative chromatography. *Biotechnol. Prog.*, 12 (1996) 65-72; C.A., 124 (1996) 81060y.
- 2567 Gruene, H.: (Development of methods and scale up by perfusion chromatography). *GIT Spez. Chromatogr.*, 15 (1995) 120-121; C.A., 124 (1996) 134345d.
- 2568 Jandera, P. and Fischer, J.: Chromatographic behaviour in reversed-phase high-performance liquid chromatography with micellar and submicellar mobile phases. *J. Chromatogr. A*, 728 (1996) 279-298.
- 2569 Jones, M.C., Nassimbene, R.D. and Wolfe, J.D.: Mixing and dispersion measurements on packed bed flows using a fiber-optic probe array. *Chem. Eng. Sci.*, 51 (1996) 1009-1021; C.A., 124 (1996) 235996x.
- 2570 Sajonz, P., Zhong, G. and Guiochon, G.: Influence of the concentration dependence of the mass transfer properties on the chromatographic band profiles. I. Apparent axial dispersion coefficient in frontal analysis. *J. Chromatogr. A*, 728 (1996) 15-23.
- 2571 Sarker, M., Yun, T. and Guiochon, G.: Study of the operating conditions of radial compression columns in preparative chromatography. *J. Chromatogr. A*, 728 (1996) 3-13.

- See also 2455, 2572, 2587, 2589, 2590, 2614, 2621, 2650, 2651, 2652, 2957, 3017, 3300, 3656, 3748, 3771.

## 3f. Programmed temperature, pressure, vapors, gradients

- 2572 Gallant, S.R., Vunnum, S. and Cramer, S.M.: Optimization of preparative ion-exchange chromatography of proteins: linear gradient separations. *J. Chromatogr. A*, 725 (1996) 295-314.
- 2573 Strel, B., Ceccato, A., Chiap, P., Hubert, P. and Crommen, J.: Injection-generated solvent and pH gradients for sample enrichment on injection of large volumes in microcolumn liquid chromatography. *Biomed. Chromatogr.*, 9 (1995) 254-256; *C.A.*, 124 (1996) 134356h.
- 2574 Welsch, T., Schmid, M., Kutter, J. and Kálmán, A.: Temperature of the eluent: a neglected tool in high-performance liquid chromatography? *J. Chromatogr. A*, 728 (1996) 299-306.
- 2575 Yang, F.J., Rpiington, P., Park, D., Lai, W., Watanabe, C., Shiro, N. and Witayel, A.: (Gradient micro HPLC for routine analysis). *LaborPraxis*, 19 (1995) 52-59; *C.A.*, 124 (1996) 134327z.

See also 2482, 3453, 3463, 3616.

## 4. SPECIAL TECHNIQUES

## 4a. Automation

- 2576 Gansen, M.: (Automatic control of control devices saves costs). *LaborPraxis*, 19 (1995) 60-64; *C.A.*, 124 (1996) 163747s.
- 2577 Hawk, G.L. and Hensley, P.F.: Automated sample preparation for LC and GC analysis using a nonrobotic workstation. *Int. Lab.*, 26, May (1996) 12-14.
- 2578 McDowall, R.D.: Ten steps to heaven? *LC-GC Int.*, 9 (1996) 202-212.

See also 2483, 2489, 2502, 2567, 2872, 2913, 2932, 2952, 3018, 3072, 3362, 3410, 3506, 3579, 3622, 3672.

## 4b. Computerization and modelling

- 2579 Bergman, D.L.: Investigation of chromatographic mechanism through the digital simulation of chromatographic peak shapes (longitudinal diffusion). Avail. *Univ. Microfilms Int.*, Order No.: DA9533458, 1995, 172 p.; *C.A.*, 124 (1996) 75094w.
- 2580 Hayes, T.L., Kohne, J.W. and Miller, T.L.: The functional testing of chromatographic software as part of computer validation. *LC-GC*, 13 (1995) 960-968; *C.A.*, 124 (1996) 134065n.
- 2581 Klatte, S.J. and Beck, T.L.: Microscopic simulation of solute transfer in reversed phase liquid chromatography. *J. Phys. Chem.*, 100 (1996) 5931-5934; *C.A.*, 124 (1996) 243352z.
- 2582 Lachmüller, C.H., Hsu, S.-H. and Reese, C.: Prediction of the retention behavior of ionizable compounds in reversed-phase LC using factor-analytical modeling. *J. Chromatogr. Sci.*, 34 (1996) 77-84.
- 2583 Lukulay, P.H. and McGuffin, V.L.: Comparison of solvent modulation with premixed mobile phases for the separation of corticosteroids by liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 4039-4062.
- 2584 Lukulay, P.H.S.: Parametric modulation: a novel computer-assisted optimization method for liquid chromatography. Avail. *Univ. Microfilms Int.*, Order No. DA9605906, 1996, 236 p.; *C.A.*, 124 (1996) 249215d.

- 2585 Marengo, E., Gennaro, M.C. and Abrigo, C.: Investigation by experimental design and regression models of the effect of five experimental factors on ion-interaction high-performance liquid chromatographic retention. *Anal. Chim. Acta*, 321 (1996) 225-236.
- 2586 McDowall, R.D.: Chromatographic data-acquisition systems and LIMS: is integration possible? *LC-GC*, 14 (1996) 28-31; *C.A.*, 124 (1996) 163766x.
- 2587 Miron, M., Szanya, T., Hanak, L., Argyelan, J. and Marton, G.: Mathematical modelling of displacement chromatography. *Hung. J. Ind. Chem.*, 23 (1995) 293-298; *C.A.*, 124 (1996) 149512e.
- 2588 Roth, C.M., Unger, K. and Lenhoff, A.M.: Mechanistic model of retention in protein ion-exchange chromatography. *J. Chromatogr. A*, 726 (1996) 45-56.
- 2589 Row, K.H. and Larin, A.V.: Computational algorithm to predict peak profiles in preparative high-performance chromatography. *Korean J. Chem. Eng.*, 12 (1995) 512-515; *C.A.*, 124 (1996) 120964t.
- 2590 Truedinger, U. and Kupka, A.: (Simulation software for the preparative HPLC). *LaborPraxis*, 19 (1995) 28-30; *C.A.*, 124 (1996) 116335w.
- 2591 Ulrich, N. and Wuensch, G.: (Expert systems to assist analytical chemistry. Expert system NINA for method setup in ion chromatography). *J. Pract. Chem./Chem.-Ztg.*, 337 (1995) 628-635; *C.A.*, 124 (1996) 104860a.
- 2592 Watanabe, H. and Sato, H.: Retention behavior of the second system peak in ion-exchange chromatography with dibasic acid eluent systems. *Anal. Sci.*, 11 (1995) 923-927; *C.A.*, 124 (1996) 127812a.
- 2593 Xu, Y.: The modelling and analysis of adsorption and elution stages of perfusion chromatographic systems. Avail. *Univ. Microfilms Int.*, Order No.: DA9603415, 1996, 273 p.; *C.A.*, 124 (1996) 235800d.

For additional information see *C.A.*:  
124 (1996) 127888e, 163809p, 219011s.

See also 2440, 2441, 2448, 2454, 2522, 2555, 2563, 2577, 2624, 2661, 3114, 3281, 3412, 3762.

## 4c. Combination with other physico-chemical techniques (MS, IR etc.)

- 2594 Behnke, B., Schlotterbeck, G., Tallarek, U., Strohschein, S., Tseng, L.-H., Keller, T., Albert, K. and Bayer, E.: Capillary HPLC-NMR coupling: high resolution  $^1\text{H}$  NMR spectroscopy in the nanoliter scale. *Anal. Chem.*, 68 (1996) 1110-1115.
- 2595 Caimi, R.J., Jr.: The development and application of liquid chromatography-combustion-high precision isotope ratio mass spectrometry (lcc-irms) (carbon isotope). Avail. *Univ. Microfilms Int.*, Order No.: DA9542551, 1995, 174 p.; *C.A.*, 124 (1996) 134232q.
- 2596 Hiraoka, K., Fukusawa, H., Matsushita, F. and Aizawa, K.: High-flow liquid chromatography/mass spectrometry interface using a parallel ion spray. *Rapid Commun. Mass Spectrom.*, 9 (1995) 1349-1355; *C.A.*, 124 (1996) 105189a.

- 2597 Hutton, T. and Major, H.J.: Characterizing biomolecules by electrospray ionization-mass spectrometry [ESI-MS] coupled to liquid chromatography and capillary electrophoresis. *Biochem. Soc. Trans.*, 23 (1995) 924-927; C.A., 124 (1996) 197491u.
- 2598 Lindon, J.C., Nicholson, J.K. and Wilson, I.D.: The development and application of coupled HPLC-NMR spectroscopy. *Adv. Chromatogr.*, 36 (1996) 315-382; C.A., 124 (1996) 192852c - a review with 70 refs.
- 2599 McCormack, A.L., Eng, J.K., DeRoos, P.C., Rudensky, A.Y. and Yates, J.R.: Microcolumn liquid chromatography-electrospray ionization tandem mass spectrometry. Analysis of immunological samples. *ACS Symp. Ser.*, 619(Biochemical and Biotechnological Applications of Electrospray Ionization Mass Spectrometry) (1996) 207-225; C.A., 124 (1996) 142980x.
- 2600 Munavalli, S., Jakubowski, E.M. and Durst, H.D.: Liquid chromatography/thermospray mass spectrometry of mustard and its metabolites. *J. Mass Spectrom.*, 30 (1995) 1716-1722; C.A., 124 (1996) 78843a.
- 2601 Somsen, G.W., Hooijsscher, E.W.J., Gooijer, C., Brinkman, U.A.T., Velthorst, N.H. and Visser, T.: Coupling of reversed-phase liquid column chromatography and Fourier transform infrared spectrometry using postcolumn on-line extraction and solvent elimination. *Anal. Chem.*, 68 (1996) 746-752.
- 2602 Stults, J.T., Gillette-Castro, B.L., Henzel, W.J., Bourell, J.H., O'Connell, K.L. and Nuwaysir, L.M.: Packed capillary HPLC-electrospray ionization mass spectrometry. In: Hancock, W.S. (Editor), *New Methods Pept. Mapp. Charact. Proteins*, CRC, Boca Raton, 1996, pp. 119-141; C.A., 124 (1996) 111286w - a review with 108 refs.
- 2603 Termonia, M.: (Coupling techniques in chromatography). *Chim. Nouv.*, 13 (1995) 1491-1493; C.A., 124 (1996) 163737p - a review with 14 refs.

For additional information see C.A.:

124 (1996) 127589h, 218722n, 249148j.

- See also 2500, 2543, 2619, 2644, 2659, 2669, 2672, 2698, 2722, 2726, 2750, 2781, 2810, 2825, 2832, 2833, 2835, 2844, 2849, 2884, 2886, 2889, 2897, 2981, 2983, 2992, 2994, 2998, 3010, 3018, 3024, 3143, 3179, 3223, 3249, 3250, 3283, 3285, 3304, 3305, 3306, 3309, 3318, 3322, 3330, 3356, 3401, 3431, 3458, 3469, 3486, 3495, 3561, 3566, 3572, 3574, 3580, 3589, 3605, 3611, 3612, 3615, 3654, 3672, 3673, 3676, 3686, 3704, 3714, 3721, 3726, 3729, 3738, 3740, 3754, 3761, 3777, 3781, 3789, 3800, 3807, 3813, 3819, 3823.

#### 4d. Affinity chromatography (advances)

- 2604 Björklund, M. and Hearn, M.T.W.: Synthesis of silica-based heparin-affinity adsorbents. *J. Chromatogr. A*, 728 (1996) 149-169.
- 2605 Denizli, A., Piskin, E. and Salih, B.: Metal chelate affinity chromatography: Congo Red and Cd(II) attached poly(EGDMA-HEMA) microbeads for specific albumin adsorption. *Turk. J. Chem.*, 19 (1995) 296-304; C.A., 124 (1996) 212025r.

- 2606 Johnson, R.D., Todd, R.J. and Arnold, F.H.: Multipoint binding in metal-affinity chromatography. II. Effect of pH and imidazole on chromatographic retention of engineered histidine-containing cytochromes c. *J. Chromatogr. A*, 725 (1996) 225-235.
- 2607 Kazmierski, W.M. and McDermed, J.: Synthesis of the carbonic acid benzotriazol-1-yl-ester-(2-biotinylamino)-9H-fluoren-9-yl-methyl ester: a convenient transient-biotinylation reagent for use in affinity chromatography. *Tetrahedron Lett.*, 36 (1995) 9097-9100; C.A., 124 (1996) 203029t.
- 2608 Kiyohara, S., Sasaki, M., Saito, K., Sugita, K. and Sugo, T.: Amino acid addition to epoxy-group-containing polymer chain grafted onto a porous membrane. *J. Membr. Sci.*, 109 (1996) 87-92; C.A., 124 (1996) 169621t.
- 2609 Koyama, T. and Terauchi, K.-i.: Synthesis and application of boronic acid-immobilized porous polymer particles: a novel packing for high-performance liquid affinity chromatography. *J. Chromatogr. B*, 679 (1996) 31-40.
- 2610 Li, Q.X. and Casida, J.E.: Affinity probes for the GABA-gated chloride channel: 5e-tert.-butyl-2-[4-(substituted-ethynyl)phenyl]-1,3-dithianes with photoactivatable, fluorescent, biotin, agarose and protein substituents. *Bioorg. Med. Chem.*, 3 (1995) 1675-1684; C.A., 124 (1996) 169907r.
- 2611 Liapis, A.I. and Unger, K.K.: The chemistry and engineering of affinity chromatography. In: Street, G. (Editor), *Highly Sel. Sep. Biotechnol.*, Blackie, Glasgow, 1994, pp. 121-162; C.A., 124 (1996) 169607t.
- 2612 Narinesingh, D. and Ngo, T.T.: Activation of supports containing hydroxyl groups using bis(4-nitrophenyl)carbonate. *Anal. Lett.*, 29 (1996) 547-564.
- 2613 Petro, M., Svec, F. and Frechet, J.M.J.: Immobilization of trypsin onto "molded macroporous poly(glycidyl methacrylate-co-ethylene dimethacrylate) rods and use of the conjugates as bioreactors and for affinity chromatography. *Biotechnol. Bioeng.*, 49 (1996) 355-363; C.A., 124 (1996) 139731m.
- 2614 Price, M.R. and Beyzavi, K.: Preparative immunoaffinity techniques. In: Wisodom, G.B. (Editor), *Pept. Antigens*, IRL Press, Oxford, 1994, pp. 117-137; C.A., 124 (1996) 169590g.
- 2615 Vannum, S., Gallant, S. and Cramer, S.: Immobilized metal affinity chromatography: displacer characteristics of traditional mobile phase modifiers. *Biotechnol. Prog.*, 12 (1996) 84-91; C.A., 124 (1996) 81061z.

For additional information see C.A.:

124 (1996) 49058t.

- See also 2520, 2541, 2549, 2787, 2995, 3066, 3241, 3248, 3259, 3340, 3375.

#### 4f. Trace analysis and preseparation techniques

- 2616 Beketov, V.I., Parchinskii, V.Z. and Zorov, N.B.: Effects of high-frequency electromagnetic treatment on the solid-phase extraction of aqueous benzene, naphthalene and phenol. *J. Chromatogr. A*, 731 (1996) 65-73.
- 2617 Bergård, A., Colmsjö, Å. and Melin, J.: Assessing breakthrough times for denuder samplers with emphasis on volatile organic compounds. *J. Chromatogr. A*, 723 (1996) 301-311.
- 2618 Horka, M., Kahle, V., Krejci, M. and Slais, K.: Analyte preconcentration procedure based on parallel-current open-tubular liquid chromatography. *J. Chromatogr. A*, 728 (1996) 311-318.

- 2619 Jin, Q., Zhang, H., Liang, F. and Jin, Q.: Some observation on the online anion-exchange preconcentration for microwave plasma torch atomic emission spectrometry. *J. Anal. At. Spectrom.*, 10 (1995) 875-879; C.A., 124 (1996) 74901v.  
 2620 Welch, C.J.: The determination of trace amounts of organic compounds. *Environ. Anal. Chem.*, (1995) 386-407; C.A., 124 (1996) 192843a - a review with 18 refs.

For additional information see C.A.:  
 124 (1996) 127650w, 218888w.

See also 2646, 2657, 2681, 2919, 3306, 3397, 3400, 3420, 3424, 3726, 3790, 3833.

#### 4g. Enantiomers, separation

- 2621 Bousquet, O.R., Braun, J. and le Goffic, F.: Countercurrent chromatography as an enzymic reactor for organic synthesis. *Tetrahedron Lett.*, 36 (1995) 8195-8196; C.A., 124 (1996) 116683h.  
 2622 Bunke, A., Jira, T. and Beyrich, T.: Influence of different types of cyclodextrins on the racemization of scopolamine-N-butylbromide. *J. Chromatogr. A*, 728 (1996) 441-445.  
 2623 Büschges, R., Linde, H., Mutschler, E. and Spahn-Langguth, H.: Chloroformates and isothiocyanates derived from 2-arylpropionic acids as chiral reagents: synthetic routes and chromatographic behaviour of the derivatives. *J. Chromatogr. A*, 725 (1996) 323-334.  
 2624 Cabrera, K., Jung, M., Fluck, M. and Schuring, V.: Determination of enantiomerization barriers by computer simulation of experimental elution profiles obtained by high-performance liquid chromatography on a chiral stationary phase. *J. Chromatogr. A*, 731 (1996) 315-321.  
 2625 Fischer, C., Modler, A., Moinet, C. and Fiaud, J.-C.: Enantioselective resolution of axially chiral compounds and interconvertible enantiomers by high-performance liquid chromatography. *J. Chromatogr. A*, 728 (1996) 433-439.  
 2626 Gao, R.Y., Yang, G.S., Shen, H.X., Chen, R.Y., Dai, Q. and Wang, Q.S.: Direct chiral high performance liquid chromatography separation of O,O-diethyl-(*p*-methyl-benzenesulfonamido)-, aryl (alkyl)-methylphosphonate. *J. Liq. Chromatogr.*, 19 (1996) 171-178.  
 2627 Gasparrini, F., Misiti, D., Pierini, M. and Villani, C.: Enantioselective chromatography on brush-type chiral stationary phases containing totally synthetic selectors. Theoretical aspects and practical applications. *J. Chromatogr. A*, 724 (1996) 79-90.  
 2628 Haglund, P.: Enantioselective separation of polychlorinated biphenyl atropisomers using chiral high-performance liquid chromatography. *J. Chromatogr. A*, 724 (1996) 219-228.  
 2629 Kennedy, J.H.: Comparison of chiral separations on polysaccharide chiral stationary phases to an improved Pirkle phase. *J. Chromatogr. A*, 725 (1996) 219-224.  
 2630 Kunath, A., Theil, F. and Jähnisch, K.: Influence of the kind of the alcoholic modifier on chiral separation on a Chiralpak AD column. *J. Chromatogr. A*, 728 (1996) 249-257.  
 2631 Meyer, V.R.: Accuracy in the chromatographic determination of extreme enantiomeric ratios: a critical reflection. *Chirality*, 7 (1995) 567-571; C.A., 124 (1996) 197382j.  
 2632 Minguillón, C., Franco, P. and Oliveros, L.: Bonded cellulose-derived high-performance liquid chromatography chiral stationary phases. II. Influence of the porosity of the silica gel matrix on performance. *J. Chromatogr. A*, 728 (1996) 415-422.  
 2633 Minguillón, C., Franco, P., Oliveros, L. and López, P.: Bonded cellulose-derived high-performance liquid chromatography chiral stationary phases. I. Influence of the degree of fixation on selectivity. *J. Chromatogr. A*, 728 (1996) 407-414.  
 2634 Oi, S.: (Chiral molecular recognition with axially dissymmetric binaphthyl compounds via chromatography). *Kagaku to Kogyo (Tokyo)*, 49 (1996) 127-130; C.A., 124 (1996) 175008q - a review with 11 refs.  
 2635 Pirkle, W.H. and Welch, C.J.: Some thoughts on the coupling of dissimilar chiral columns or the mixing of chiral stationary phases for the separation of enantiomers. *J. Chromatogr. A*, 731 (1996) 322-326.  
 2636 Pirkle, W.H., Koscho, M.E. and Wu, Z.: High-performance liquid chromatographic separation of the enantiomers of N-aryloxa-zolinones, N-aryl thiazolinones and their sulfur derivatives on a synthetic chiral stationary phase. *J. Chromatogr. A*, 726 (1996) 91-97.  
 2637 Riering, H. and Sieber, M.: Covalently bonded permethylated cyclodextrins, new selectors for enantiomeric separations by liquid chromatography. *J. Chromatogr. A*, 728 (1996) 171-177.  
 2638 Sonnet, P.E., Oliver, J.E., Waters, R.M., King, G. and Panicker, S.: A potential chiral derivatizing agent for 1,2-diglycerides. *Chem. Phys. Lipids*, 78 (1995) 203-208; C.A., 124 (1996) 111623d.  
 2639 Stalcup, A.M. and Gahm, K.H.: A sulfated cyclodextrin chiral stationary phase for high-performance liquid chromatography. *Anal. Chem.*, 68 (1996) 1369-1374.  
 2640 Swartz, M.E.: Separation and detection of enantiomeric compounds by liquid chromatography and chiral micellar electrokinetic capillary chromatography. Avail. *Univ. Microfilms Int.*, Order No.: DA9601875, 1995, 231 p.; C.A., 124 (1996) 192902u.  
 2641 Szemán, J., Ganzler, K., Salgó, A. and Szejtli, J.: Effect of the degree of substitution of cyclodextrin derivatives on chiral separations by high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr. A*, 728 (1996) 423-431.  
 2642 Yamamoto, A., Matsunaga, A., Hayakawa, K., Miyazaki, M. and Nishimura, M.: Novel approach to the measurement of enantiomeric purity by high-performance liquid chromatography using a polarized photometric detector equipped with a split cell. *J. Chromatogr. A*, 727 (1996) 55-59.  
 2643 Yang, J. and Hage, D.S.: Role of binding capacity versus binding strength in the separation of chiral compounds on protein-based high-performance liquid chromatography columns. Interactions of D- and L-tryptophan with human serum albumin. *J. Chromatogr. A*, 725 (1996) 273-285.

For additional information see C.A.:  
 124 (1996) 149906m, 163714d.

See also 2445, 2450, 2462, 2514, 2529, 2542, 2828, 2862, 2908, 2923, 2924, 2929, 2931, 2936, 2943, 2949, 2954, 2959, 2962, 3276, 3287, 3293, 3296, 3297, 3300, 3380, 3473, 3478, 3484, 3488, 3492, 3497, 3503, 3510, 3511, 3512, 3518, 3526, 3533, 3536, 3541, 3551, 3553, 3572, 3583, 3588, 3593, 3594, 3597, 3602, 3607, 3620, 3644, 3679, 3690, 3691, 3696, 3705, 3716.

#### 4h. Other special techniques

- 2644 Balogh, M.P.: Adaptation of high-performance liquid chromatographic methods to multiple spectral analyses. *J. Chromatogr. A*, 728 (1996) 351-357.
- 2645 Buttler, T., Nilsson, C., Gorton, L., Marko-Varga, G. and Laurell, T.: Membrane characterisation and performance of microdialysis probes intended for use as bioprocess sampling units. *J. Chromatogr. A*, 725 (1996) 41-56.
- 2646 Fedotov, P.S., Maruytina, T.A., Grebneva, O.N. and Spivakov, B.Ya.: Influence of kinetic properties of extraction systems on the separation and preconcentration of some elements by countercurrent chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 575-585; *C.A.*, 124 (1996) 74896x.
- 2647 Freitag, R., Splitt, H. and Reif, O.-W.: Controlled mixed-mode interaction chromatography on membrane adsorbers. *J. Chromatogr. A*, 728 (1996) 129-137.
- 2648 García Sánchez, F., Navas Diaz, A. and García Pareja, A.: Micellar liquid chromatography of plant growth regulators detected by derivative fluorometry. *J. Chromatogr. A*, 723 (1996) 227-233.
- 2649 Gennaro, M.C.: Reversed-phase ion-interaction chromatography. Retention mechanisms and dependence on different factors development of analytical method. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 527-533; *C.A.*, 124 (1996) 74802p - a review with 11 refs.
- 2650 Hotier, G. and Nicoud, R.-M.: Separation by simulated moving bed chromatography with dead volume correction by desynchronization of periods. *Eur. Pat. Appl. EP 688, 589* (Cl. B01D15/02), 27 Dec. 1995, FR Appl. 94/7,763, 22 Jun. 1994; 21 p.; *C.A.*, 124 (1996) 91617z.
- 2651 Hotier, G., Couenne, N., Cohen, C. and Nicoud, R.-M.: Separation by simulated moving bed chromatography with correction for dead volume by length shrink. *Eur. Pat. Appl. EP 688,590* (Cl. B01D15/02), 27 Dec. 1995, FR Appl. 94/7,764, 22 Jun. 1994; 12 p.; *C.A.*, 124 (1996) 91618a.
- 2652 Hotier, G., Couenne, N., Cohen, C. and Toussaint, J.-M.: Separation by simulated moving bed chromatography with correction for dead volume. *Eur. Pat. Appl. EP 688,588* (Cl. B01D15/02), 27 Dec. 1995, FR Appl. 94/7,762, 22 Jun. 1994; 12 p.; *C.A.*, 124 (1996) 91616y.
- 2653 Ito, Y.: pH-Peak-focusing and pH-zone-refining countercurrent chromatography. *Chem. Anal.*, 132 (1996) 121-175; *C.A.*, 124 (1996) 201195v - a review with 27 refs.
- 2654 Ito, Y.: Principle apparatus and methodology of high-speed countercurrent chromatography. *Chem. Anal.*, 132 (1996) 3-44; *C.A.*, 124 (1996) 192826x - a review with 67 refs.
- 2655 Lee, Y.W.: Dual countercurrent chromatography. *Chem. Anal.*, 132 (1996) 93-105; *C.A.*, 124 (1996) 192829a - a review with 14 refs.
- 2656 Liu, S.: Development and applications of capillary-based analysis systems (electroosmotic flow, microbore column chromatography). Avail. *Univ. Microfilms Int.*, Order No.: DA9541756, 1995, 211 p.; *C.A.*, 124 (1996) 134106b.
- 2657 Maryutina, T.A., Fedotov, P.S. and Spivakov, B.Ya.: Application of various two-phase liquid systems for preconcentration and separation of trace elements by countercurrent chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 623-639; *C.A.*, 124 (1996) 74897y.
- 2658 Oka, H.: Foam countercurrent chromatography of bacitracin complex. *Chem. Anal.*, 132 (1996) 107-120; *C.A.*, 124 (1996) 197364e.
- 2659 Oka, H.: High-speed countercurrent chromatography/mass spectrometry. *Chem. Anal.*, 132 (1996) 73-91; *C.A.*, 124 (1996) 192828z - a review with 12 refs.
- 2660 Schaufelberger, D.E.: Analytical high-speed countercurrent chromatography. *Chem. Anal.*, 132 (1996) 45-70; *C.A.*, 124 (1996) 192827y - a review with 63 refs.
- 2661 Shibukawa, A. and Nakagawa, T.: Theoretical study of high-performance frontal analysis: a chromatographic method for determination of drug-protein interaction. *Anal. Chem.*, 68 (1996) 447-454.
- 2662 Shinomiya, K., Muto, M., Kabasawa, Y., Fales, H.M. and Ito, Y.: Protein separation by improved cross-axis coil planet centrifuge with eccentric coil assemblies. *J. Liq. Chromatogr.*, 19 (1996) 415-424.
- 2663 Thede, R., Haberland, D. and Below, E.: Autocatalytic reactions in a liquid chromatographic reactor. *J. Chromatogr. A*, 728 (1996) 401-406.
- 2664 Yang, S.: Characterization of chemical interactions in micellar liquid chromatography and micellar electrokinetic capillary chromatography (phenols, surfactants). Avail. *Univ. Microfilms Int.*, Order No.: DA9542250, 1995, 205 p.; *C.A.*, 124 (1996) 134384r.
- 2665 Zhong, G. and Guiochon, G.: Steady-state concentration boundary profile for the injection of a single-component feed in counter-current chromatography. *J. Chromatogr. A*, 728 (1996) 25-32.
- See also 2425, 2472, 2567, 2575, 2608, 2670, 2915, 2946, 3105, 3261, 3417, 3440, 3481, 3758, 3769, 3778, 3796, 3810.

## 5. HYDROCARBONS AND HALOGEN DERIVATIVES

### 5a. Aliphatic hydrocarbons

- 2666 Shi, B. and Davis, B.H.: High-performance liquid chromatographic separation of pairs of isotopic labeled (deuterium/protium) molecules. *J. Chromatogr. A*, 731 (1996) 351-354.

See also 2650, 2651, 2652.

## 5b. Cyclic hydrocarbons, fullerenes

- 2667 Beltrán, J.L., Ferrer, R. and Guiteras, J.: Determination of polycyclic aromatic hydrocarbons by HPLC with spectrofluorimetric detection and wavelength programming. *J. Liq. Chromatogr.*, 19 (1996) 477-488.
- 2668 Biggs, W.R. and Fetzer, J.C.: Analytical techniques for large polycyclic aromatic hydrocarbons: a review. *TrAC*, 15 (1996) 196-206.
- 2669 Bonfanti, L., Careri, M., Mangia, A., Manini, P. and Maspero, M.: Simultaneous identification of different classes of hydrocarbons and determination of nitro-polycyclic aromatic hydrocarbons by means of particle beam liquid chromatography-mass spectrometry. *J. Chromatogr. A*, 728 (1996) 359-369.
- 2670 Chee, K.K., Wong, M.K. and Lee, H.K.: Optimization of microwave-assisted solvent extraction of polycyclic aromatic hydrocarbons in marine sediments using a microwave extraction system with high-performance liquid chromatography-fluorescence detection and gas chromatography-mass spectrometry. *J. Chromatogr. A*, 723 (1996) 259-271.
- 2671 Davydov, V.Ya.: Liquid chromatography of fullerenes and study of adsorption properties of fullerenes crystals. *Stud. Surf. Sci. Catal.*, 99 (1996) 899-914; C.A., 124 (1996) 243067k - a review with 30 refs.
- 2672 Galceran, M.T. and Moyano, E.: Determination of hydroxy polycyclic aromatic hydrocarbons by liquid chromatography-mass spectrometry. Comparison of atmospheric pressure chemical ionization and electrospray. *J. Chromatogr. A*, 731 (1996) 75-84.
- 2673 Hackl, C., Lintelmann, J., Sauerbrey, R. and Kettrup, A.: Preliminary experiments for the analysis of tetrahydrotetrols of benz[a]anthracene and benzo[a]pyrene derived from their hemoglobin adducts using a coupled-column high-performance liquid chromatographic system. *J. Chromatogr. A*, 730 (1996) 153-159.
- 2674 Hatrik, S., Lehotay, J., Cakrt, M. and Brandstetr, R.: Computer-assisted method for the threshold optimization of fluorescence detection conditions in HPLC of polycyclic aromatic hydrocarbons. *J. Liq. Chromatogr.*, 18 (1995) 4149-4164.
- 2675 Hayatsu, H., Hayatsu, T., Arimoto, S. and Sakamoto, H.: A short-column technique for concentrating mutagens/carcinogens having polycyclic structures. *Anal. Biochem.*, 235 (1996) 185-190.
- 2676 Holden, K.M.L., Lewis, A.C., Bartle, K.D. and Mitchell, S.C.: Characterization of polycyclic aromatic compounds (PAC) in coal products and related materials by GC and coupled LC-GC techniques. *Coal Sci. Technol.*, 24 (1995) 1935-1938; C.A., 124 (1996) 180741c.
- 2677 Jinno, K. and Saito, Y.: Separation of fullerenes by liquid chromatography: molecular recognition mechanisms in liquid chromatographic separation. *Adv. Chromatogr.*, 36 (1996) 65-125; C.A., 124 (1996) 192596x.
- 2678 Kayali, M.N., Rubio-Barroso, S. and Polo Diez, L.M.: Separation study of PAHs by HPLC using a micellar SDS mobile phase and short chain columns. *J. Liq. Chromatogr.*, 19 (1996) 759-770.
- 2679 Kiss, G., Varga-Puchony, Z. and Hlavay, J.: Determination of polycyclic aromatic hydrocarbons in precipitation using solid-phase extraction and column liquid chromatography. *J. Chromatogr. A*, 725 (1996) 261-272.
- 2680 Koehl, W., Amin, S., Staretz, M.E., Ueng, Y., Yamazaki, H., Tateishi, T., Guengerich, F.P. and Hecht, S.S.: Metabolism of 5-methylchrysene and 6-methylchrysene by human hepatic and pulmonary cytochrome P450 enzymes. *Cancer Res.*, 56 (1996) 316-324.
- 2681 Michor, G., Carron, J., Bruce, S. and Cancilla, D.A.: Analysis of 23 polynuclear aromatic hydrocarbons from natural water at the sub-nmol level using solid-phase disk extraction and mass-selective detection. *J. Chromatogr. A*, 732 (1996) 85-99.
- 2682 Nirmaier, H.-P., Fischer, E., Meyer, A. and Henze, G.: Determination of polycyclic aromatic hydrocarbons in water samples using high-performance liquid chromatography with amperometric detection. *J. Chromatogr. A*, 730 (1996) 169-175.
- 2683 Ohta, H., Jinno, K., Saito, Y., Fetzer, J.C., Pesek, J.J., Matyska, M.T. and Chen, Y.-L.: Effect of temperature on the mechanism of retention of fullerenes in liquid chromatography using various alkyl bonded stationary phases. *Chromatographia*, 42 (1996) 56-62.
- 2684 Podosyeonova, N.G.: The regularities of the fullerene molecules adsorption chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 251-258; C.A., 124 (1996) 74894v.
- 2685 Pospisil, M., Pecka, K. and Vanura, J.: Determination of polycyclic aromatic hydrocarbon in petroleum oil fractions. *Pet. Coal*, 37 (1995) 34-39; C.A., 124 (1996) 95987t.
- 2686 Prade, K.: Chromatographic separation of tar-borne PAH in a soil. *Z. Pflanzenernaehr. Bodenkd.*, 158 (1995) 361-363; C.A., 124 (1996) 116157q.
- 2687 Rebscher, H. and Pyell, U.: Instrumental developments in capillary electrochromatography. *Chromatographia*, 42 (1996) 171-176.
- 2688 Rivera, L., Curto, M.J.C., Pais, P., Galceran, M.T. and Puignou, G.L.: Solid-phase extraction for the selective isolation of polycyclic aromatic hydrocarbons, azaarenes and heterocyclic aromatic amines in charcoal-grilled meat. *J. Chromatogr. A*, 731 (1996) 85-94.
- 2689 Rodríguez Delgado, M.A., Sánchez, M.J., González, V. and García Montelongo, F.: Influence of several parameters on efficiency and peak shape in the micellar liquid chromatography of polynuclear aromatic hydrocarbons. *J. Liq. Chromatogr.*, 19 (1996) 187-200.
- 2690 Rodriguez, M.A., Sanchez, M.J., Gonzalez, V. and Montelongo, F.G.: Determination of polynuclear aromatic hydrocarbons in environmental samples by micellar liquid chromatography. *J. High Resolut. Chromatogr.*, 19 (1996) 111-116.
- 2691 Saito, Y., Ohta, H., Nagashima, H., Itoh, K., Jinno, K. and Pesek, J.J.: Separation of fullerenes with liquid-crystal bonded silica phases in microcolumn high performance liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 243-250; C.A., 124 (1996) 74893u.
- 2692 Shou, M., Krausz, K.W., Gonzalez, F.J. and Gelboin, H.V.: Metabolic activation of the potent carcinogen dibenz[a,h]anthracene by cDNA-expressed human cytochromes P450. *Arch. Biochem. Biophys.*, 328 (1996) 201-207.

- 2693 Sun, Y.-P., Ma, B., Lawson, G.E., Bunker, C.E. and Rollins, H.W.: Effects of photochemical reactions of pyrene in alcohol and aqueous solvent systems on spectroscopic analyses. *Anal. Chim. Acta.*, 319 (1996) 379-386.
- 2694 Sun, Z.L., Song, L.J., Zhang, X.T. and Hu, Z.D.: Study on the relationship between retention behavior and molecular structure parameters of substituted benzene derivatives in RPLC. *Chromatographia*, 42 (1996) 43-48.
- 2695 Viola, A. and Cao, G.: Rapid direct analysis of *p*-xylene oxidation products by reversed-phase high-performance liquid chromatography. *J. Chromatogr. Sci.*, 34 (1996) 27-33.
- 2696 Yan, X., Wu, Y. and Sun, Y.: Direct determination of coronene in heavy oil fraction by high-performance liquid chromatography. *J. Chromatogr. A*, 724 (1996) 337-342.

For additional information see C.A.:  
124 (1996) 192586u.

See also 2446, 2481, 2552, 2564, 2616, 2625, 3755.

#### 5c. Halogen derivatives

- 2697 Van Bavel, B., Järemo, M., Karlsson, L. and Lindström, G.: Development of a solid phase carbon trap for simultaneous determination of PCDDs, PCDFs, PCBs, and pesticides in environmental samples using SFE-LC. *Anal. Chem.*, 68 (1996) 1279-1283.

See also 2628.

#### 5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

- 2698 Bartle, K.D., Herod, A.A., John, P., Johnson, B.R., Johnson, C.A.F., Kandiyoti, R., Parker, J.E. and Smith, G.P.: Molecular mass distributions of coal products relevant to pyrolysis, liquefaction and combustion. *Coal Sci. Technol.*, 24 (1995) 1435-1438; C.A., 124 (1996) 150482b.
- 2699 Daucik, P.: Determination of group composition of crude oil products by liquid chromatography. *Pet. Coal.*, 37 (1995) 68-69; C.A., 124 (1996) 92147h.
- 2700 Herod, A.A. and Kandiyoti, R.: Characterization of large molecular mass materials in coal tar pitch fractions separated by planar chromatography. *Coal Sci. Technol.*, 24 (1995) 949-952; C.A., 124 (1996) 150489j.
- 2701 Oba, K. and Hugener, M.: Characterization of polymer-modified bituminous roofing membranes using chromatography. *Mater. Struct.*, 28 (1995) 534-544; C.A., 124 (1996) 95575g.

For additional information see C.A.:  
124 (1996) 149170s, 150340d.

See also 2686, 3836.

#### 6. ALCOHOLS

- 2702 Blehaut, J., Charton, F. and Nicoud, R.-M.: Separation of fatty alcohol stereoisomers on a large-scale high-performance simulated moving bed. *LC-GC Int.*, 9 (1996) 228-238.

- 2703 Cataldi, T.R.I., Centonze, D. and Desimoni, E.: Sensitive determination of ethanol in low-alcohol samples by ion-exclusion chromatography with EC detection using a ruthenium-based inorganic film electrode. *Food Chem.*, 55 (1996) 17-21; C.A., 124 (1996) 85156g.
- 2704 Haj-Yehia, A.I. and Benet, L.Z.: Determination of alcohols by high-performance liquid chromatography with fluorimetric detection after precolumn derivatization with 2-(4-carboxyphenyl)-6-methoxybenzofuran. *J. Chromatogr. A*, 724 (1996) 107-115.
- 2705 Kakemoto, M., Murakami, K. and Ogawa, H.: (Simultaneous determination of ethanol and glucose by HPLC with an immobilized enzyme membrane electrode). *Nippon Kagaku Kaishi*, (1996) 141-145; C.A., 124 (1996) 173826f.
- 2706 Moldovan, Z., Delgado Luque, M.V., Otal Salaverri, E., Suárez, A., Andreozzi, R., Insola, A. and Lebrato Martínez, J.: Determination of polyethylene glycols in water by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 723 (1996) 243-249.
- 2707 Motte, J.C., Windey, R. and Delafortrie, A.: High-sensitivity fluorescence derivatization for the determination of hydroxy compounds in aqueous solution by high-performance liquid chromatography. *J. Chromatogr. A*, 728 (1996) 333-341.

See also 2568, 2695, 2895, 3751.

#### 7. PHENOLS

- 2708 Barroso, C.G., Rodríguez, M.C., Guillén, D.A. and Pérez-Bustamante, J.A.: Analysis of low molecular mass phenolic compounds, furfural and 5-hydroxymethylfurfural in Brandy de Jerez by high-performance liquid chromatography-diode array detection with direct injection. *J. Chromatogr. A*, 724 (1996) 125-129.
- 2709 Bartolomé, B., Hernández, T., Bengoechea, M.L., Quesada, C., Gómez-Cordovés, C. and Estrella, I.: Determination of some structural features of procyanidins and related compounds by photodiode-array detection. *J. Chromatogr. A*, 723 (1996) 19-26.
- 2710 Bosch Reig, F., Campins Falcó, P. and Verdú Andrés, J.: H-Point standard additions method for resolution of overlapping chromatographic peaks with diode array detection by using area measurements. Determination of phenol and cresols in waters. *J. Chromatogr. A*, 726 (1996) 57-66.
- 2711 Bru, E.R., Barroso, C.G., Cela, R. and Perez-Bustamante, J.A.: Development of rotary and continuous liquid-liquid extraction technique for phenolic compounds in wine. *Analyst (Cambridge)*, 121 (1996) 297-302.
- 2712 Celotti, E., Ferrarini, R., Zironi, R. and Conte, L.S.: Resveratrol content of some wines obtained from dried Valpolicella grapes: Recioto and Amarone. *J. Chromatogr. A*, 730 (1996) 47-52.
- 2713 García-Parrilla, C., Heredia, F.J. and Troncoso, A.M.: Phenols HPLC analysis by direct injection of sherry wine vinegar. *J. Liq. Chromatogr.*, 19 (1996) 247-258.
- 2714 Guillén, D.A., Barroso, C.G. and Pérez-Bustamante, J.A.: Automation of sample preparation as a preliminary stage in the high-performance liquid chromatographic determination of polyphenolic compounds in sherry wines. *J. Chromatogr. A*, 730 (1996) 39-46.

- 2715 Guillén, D.A., Barroso, C.G. and Pérez-Bustamante, J.A.: Selection of column and gradient for the separation of polyphenols in sherry wine by high-performance liquid chromatography incorporating internal standards. *J. Chromatogr. A*, 724 (1996) 117-124.
- 2716 Guolan, H., Weihua, Z. and Zhiren, Z.: Separation of positional isomers of chlorophenols by reverse phase HPLC. *J. Liq. Chromatogr.*, 19 (1996) 899-909.
- 2717 Knutsson, M., Mathiasson, L. and Jönsson, J.Å.: Supported liquid membrane work-up in combination with liquid chromatography and electrochemical detection for the determination of chlorinated phenols in natural water samples. *Chromatographia*, 42 (1996) 165-170.
- 2718 Korba, T., Popl, M. and Novotna, M.: Evaluation of methods for the determination of chlorophenols in surface water. *Fresenius J. Anal. Chem.*, 355 (1996) 91-93.
- 2719 Leira, E., Botana, A. and Cela, R.: Resolution of complex mixtures of non-flavonoid polyphenols by column-switching high-performance liquid chromatography using octadecylsilica and graphitized carbon columns. *J. Chromatogr. A*, 724 (1996) 67-78.
- 2720 Lukkari, I., Irgum, K., Lindgren, P. and Liden, J.: Online determination of pyrocatechol, protocatechic acid, pyrogallol and gallic acid in black liquor by sequential injection and high-performance liquid chromatography. *Process Control Qual.*, 7 (1995) 185-193; *C.A.*, 124 (1996) 59585b.
- 2721 Paterson, B., Cowie, C.E. and Jackson, P.E.: Determination of phenols in environmental waters using liquid chromatography with electrochemical detection. *J. Chromatogr. A*, 731 (1996) 95-102.
- 2722 Roeder, A., Lam, T.M.L. and Galensa, R.: (Thermospray-LCMS-)investigations of proanthocyanidins and other polyphenols in malt, beer, and hop. *Monatsschr. Brauwiss.*, 48 (1995) 390-396; *C.A.*, 124 (1996) 173774n.
- 2723 Suárez, B., Picinelli, A. and Mangas, J.J.: Solid-phase extraction and high-performance liquid chromatographic determination of polyphenols in apple musts and ciders. *J. Chromatogr. A*, 727 (1996) 203-209.
- 2724 Tassaneeyakul, W., Birkett, D.J., Edwards, J.W., Veronese, M.E., Tasaneeyakul, W., Tukey, R.H. and Miners, J.O.: Human cytochrome P450 isoform specificity in the regioselective metabolism of toluene and o-, m- and p-xylene. *J. Pharmacol. Exp. Ther.*, 276 (1996) 101-108.
- 2725 Thompson, M.J., Ballinger, L.N., Cross, S.E. and Roberts, M.S.: High-performance liquid chromatographic determination of phenol, 4-nitrophenol, β-naphthol and a number of their glucuronide and sulphate conjugates in organ perfusate. *J. Chromatogr. B*, 677 (1996) 117-122.

For additional information see *C.A.*:  
124 (1996) 218983y.

See also 2446, 2468, 2481, 2526, 2531, 2564, 2616, 2673, 3425.

## 8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

### 8a. Flavonoids

- 2726 Coward, L., Kirk, M., Albin, N. and Barnes, S.: Analysis of plasma isoflavones by reversed-phase HPLC-multiple reaction ion monitoring-mass spectrometry. *Clin. Chim. Acta*, 247 (1996) 121-142.
- 2727 Kijima, H., Ide, T., Otsuka, H. and Takeda, Y.: Alangiflavoside, a new flavonol glycoside from the leaves of *Alangium prennifolium*. *J. Natural Prod.*, 58 (1995) 1753-1755.
- 2728 Palomino, O.M., Gomez-Serranillos, P., Carretero, E. and Villar, A.: High-performance liquid chromatography of flavonoids from *Sideritis* species. *J. Chromatogr. A*, 731 (1996) 103-108.
- For additional information see *C.A.*:  
124 (1996) 97874c, 156217r.
- See also 2709, 3734.
- 8b. Aflatoxins and other mycotoxins
- 2729 Delgado, T., Gómez-Cordovés, C. and Scott, P.M.: Determination of alternariol and alternariol methyl ether in apple juice using solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. A*, 731 (1996) 109-114.
- 2730 Franco, C.M., Fente, C.A., Vazquez, B., Cepeda, A., Lallaoui, L., Prognon, P. and Mahuzier, G.: Simple and sensitive high-performance liquid chromatography-fluorescence method for the determination of citrinin. Application to the analysis of fungal cultures and cheese extracts. *J. Chromatogr. A*, 723 (1996) 69-75.
- 2731 Gökmen, V. and Acar, J.: Rapid reversed-phase liquid chromatographic determination of patulin in apple juice. *J. Chromatogr. A*, 730 (1996) 53-58.
- 2732 Roch, O.G., Blunden, G., Haig, D.J., Coker, R.D. and Gay, C.: Determination of aflatoxins in groundnut meal by high-performance liquid chromatography: a comparison of two methods of derivatization of aflatoxin B<sub>1</sub>. *Br. J. Biomed. Sci.*, 52 (1995) 312-316; *C.A.*, 124 (1996) 143876e.
- 2733 Solfrizzo, M. and Visconti, A.: Simultaneous high-performance liquid chromatographic determination of visoltricin, acuminatepyrone and chlamydosporols in *Fusarium* cultures on maize. *J. Chromatogr. A*, 730 (1996) 69-73.
- 2734 Vazquez, B.I., Fente, C., Franco, C., Cepeda, A., Prognon, P. and Mahuzier, G.: Simultaneous high-performance liquid chromatographic determination of ochratoxin A and citrinin in cheese by time-resolved luminescence using terbium. *J. Chromatogr. A*, 727 (1996) 185-193.
- 2735 Vesely, Z. and Bohacenko, I.: Immunoaffinity extraction of aflatoxins and their HPLC determination in raw materials, foods and feeds. *Potraviny. Vedy.*, 13 (1995) 451-460; *C.A.*, 124 (1996) 143909t.
- 2736 Zhou, Y., He, G., Peng, W., Yao, J., Ni, E. and Hu, P.: (Determination of aflatoxin B<sub>1</sub> in human liver tissues with high performance liquid chromatography). *Sepu*, 13 (1995) 455-457; *C.A.*, 124 (1996) 109089r.

## 8c. Other compounds with heterocyclic oxygen (incl. tannins)

- 2737 Ball, S.E., Thiel, V.E., Tio, C.O., Lim, H.-K., Kao, J., Sisenwine, S.F. and Scatina, J.: [ $^{14}\text{C}$ ]-Ethoxycoumarin metabolism by precision-cut rat hepatic slices. *Drug Metab. Disp.*, 24 (1996) 383-389.
- 2738 Cadahia, E., Conde, E., Garcia-Vallejo, M.C. and Fernandez de Simon, B.: Gel permeation chromatographic study of the molecular weight distribution of tannins in the wood, bark and leaves of *Eucalyptus* spp. *Chromatographia*, 42 (1996) 95-100.
- 2739 Du Preez, J.L., Lötter, A.P. and Guillot, J.K.: Kinetics and mechanism of degradation of the heterocyclic cage compound NPG1-01 in aqueous solution. *Pharmazie*, 51 (1996) 223-227.

For additional information see C.A.:  
124 (1996) 48179w.

## 9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

- 2740 Bastos, J.K., Burandt, C.L., Jr., Nanayakkara, N.P., Bryant, L. and McChesney, J.D.: Quantitation of aryltetralin lignans in plant parts and among different populations of *Podophyllum peltatum* by reversed-phase high-performance liquid chromatography. *J. Natural Prod.*, 59 (1996) 406-408.
- 2741 Bechmann, I.E.: Determination of formaldehyde in frozen fish with formaldehyde dehydrogenase using a flow injection system with an incorporated gel-filtration chromatography column. *Anal. Chim. Acta*, 320 (1996) 155-164.
- 2742 Cordeiro, C. and Freire, A.P.: Methylglyoxal assay in cells as 2-methylquinoxaline using 1,2-diaminobenzene as derivatizing reagent. *Anal. Biochem.*, 234 (1996) 221-224.
- 2743 Führer, U., Deissler, A. and Ballschmiter, K.: Determination of biogenic halogenated methyl-phenyl ethers (halogenated anisoles) in the picogram.m<sup>-3</sup> range in air. *Fresenius J. Anal. Chem.*, 354 (1996) 333-343.
- 2744 Karst, U.: (Determination of aldehydes and ketones using 2,4-dinitrophenylhydrazine as derivatizing agent). *GIT Fachz. Lab.*, 40 (1996) 49-50; C.A., 124 (1996) 249159p - a review with 11 refs.
- 2745 Londero, D. and Lo Greco, P.: Automated high-performance liquid chromatographic separation with spectrofluorometric detection of a malondialdehyde-thiobarbituric acid adduct in plasma. *J. Chromatogr. A*, 729 (1996) 207-210.
- 2746 Steinert, J., Khalaf, H. and Rimpler, M.: High-performance liquid chromatographic separation of some naturally occurring naphthoquinones and anthraquinones. *J. Chromatogr. A*, 723 (1996) 206-209.
- 2747 Xu, R., Huang, X., Morgan, T.D., Prakash, O., Kramer, K.J. and Hawley, M.D.: Characterization of products from the reactions of N-acetyldopamine quinone with N-acetylhistidine. *Arch. Biochem. Biophys.*, 329 (1996) 56-64.

See also 2497, 2498, 2695, 2708, 2810, 2853, 3110, 3452, 3751, 3765, 3769.

## 10. CARBOHYDRATES

## 10a. Mono and oligosaccharides. Structural studies

- 2748 Araujo, B.: Evaluation by chromatography of contamination in sugar mills and concentration of glucose and fructose ( $\alpha$ -D-glucopyranose and  $\beta$ -D-fructofuranose). *Int. Sugar J.*, 97 (1995) 642-645; C.A., 124 (1996) 149132f.
- 2749 Bernal, J.L., del Nozal, M.J., Toribio, L. and del Alamo, M.: HPLC analysis of carbohydrates in wines and instant coffees using anion-exchange chromatography coupled to pulsed amperometric detection. *J. Agric. Food Chem.*, 44 (1996) 507-511; C.A., 124 (1996) 115650h.
- 2750 Black, G.E. and Fox, A.: Liquid chromatography with electrospray ionization tandem mass spectrometry. Profiling carbohydrates in whole bacterial cell hydrolyzates. *ACS Symp. Ser.*, 619(Biochemical and Biotechnological Applications of Electrospray Ionization Mass Spectrometry) (1996) 81-105; C.A., 124 (1996) 197306n - a review with 34 refs.
- 2751 Dubois, M.-F., Rigal, L. and Gaset, A.: (Comparative determination of products produced by alkaline degradation of hexoses). *Can. J. Chem.*, 73 (1995) 1582-1590; C.A., 124 (1996) 105182t.
- 2752 Harsch, G. and Heimann, R.: (Training of analytical thinking as exemplified in detection of carbohydrates in foods). *Prax. Naturwiss., Chem.*, 44 (1995) 19-23; C.A., 124 (1996) 54954m.
- 2753 Kim, Y.-J., Grodzinsky, A.J. and Plaas, A.H.K.: Compression of cartilage results in differential effects on biosynthetic pathways for aggrecan, link protein and hyaluronan. *Arch. Biochem. Biophys.*, 328 (1996) 331-340.
- 2754 Koizumi, K.: (Separation and analysis of carbohydrates). *Oyo Toshitsu Kagaku*, 41 (1994) 465-471; C.A., 124 (1996) 87496s - a review with 10 refs.
- 2755 Kvasnicka, F., Ahmadová-Vavrousová R., Frias, J., Price, K.R. and Kadlec, P.: A rapid HPLC method for the determination of raffinose family of oligosaccharides in pea seeds. *J. Liq. Chromatogr.*, 19 (1996) 135-147.
- 2756 Lichon, M.J. and James, K.W.: Determination of dietary fiber and carbohydrate composition of foods by a modified version of the Englyst method. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 54-61.
- 2757 Miki, K., Butler, R., Moore, D. and Davidson, G.: Rapid and simultaneous quantification of rhamnose, mannitol, and lactulose in urine by HPLC for estimating intestinal permeability in pediatric practice. *Clin. Chem. (Washington)*, 42 (1996) 71-75.
- 2758 Nakagawa, H., Zheng, M., Hakomori, S.-i., Tsukamoto, Y., Kawamura, Y. and Takahashi, N.: Detailed oligosaccharide structures of human integrin  $\alpha 5\beta 1$  analyzed by a three-dimensional mapping technique. *Eur. J. Biochem.*, 237 (1996) 76-85.
- 2759 Rooyakers, D.R., van Eijk, H.M.H. and Deutz, N.E.P.: Simple and sensitive multi-sugar-probe gut permeability test by high-performance liquid chromatography with fluorescence labelling. *J. Chromatogr. A*, 730 (1996) 99-105.
- 2760 Saska, M. and Diack, M.: Separation of inositol isomers and derivatives from sugars and sugar alcohols. *U.S. US 5,482,631 (Cl. 210-635; B01D15/08) 9 Jan. 1996, Appl. 319,240, 6 Oct. 1994; 8 p.; C.A., 124 (1996) 235370p.*

- 2761 Takaha, T., Yanase, M., Takata, H., Okada, S. and Smith, S.M.: Potato D-enzyme catalyzes the cyclization of amylose to produce cycloamylose, a novel cyclic glucan. *J. Biol. Chem.*, 271 (1996) 2902-2908.
- 2762 Thurl, S., Müller-Werner, B. and Sawatzki, G.: Quantification of individual oligosaccharide compounds from human milk using high-pH anion-exchange chromatography. *Anal. Biochem.*, 235 (1996) 202-206.
- 2763 Torto, N., Marko-Varga, G., Gorton, L., Stålbrand, H. and Tjerneld, F.: On-line quantitation of enzymatic mannan hydrolysates in small-volume bioreactors by microdialysis sampling and column liquid chromatography-integrated pulsed electrochemical detection. *J. Chromatogr. A*, 725 (1996) 165-175.
- 2764 Trotter, A., Hofsommer, H.-J. and Rubach, K.: (Carbohydrate analysis with anion-exchange chromatography and pulsed amperometric detection with the Dionex Dx-300 chromatography module). *Fluess. Obst.*, 61 (1994) 581-589; C.A., 124 (1996) 200430f.
- 2765 Tyagarajan, K., Townsend, R.R. and Forte, J.G.: The  $\beta$ -subunit of the rabbit H,K-ATPase: a glycoprotein with all terminal lactosamine units capped with  $\alpha$ -linked galactose residues. *Biochemistry*, 35 (1996) 3238-3246.
- 2766 Vazquez-Blanco, M.E., Lopez-Hernandez, J., Vazquez-Oderiz, M.L., Simal-Lozano, J. and Gonzales-Castro, M.J.: Determination of pectins by HPLC. A comparison of two methods applied to green beans (*Phaseolus vulgaris*). *J. Chromatogr. Sci.*, 33 (1995) 551-553.
- 2767 Wilson, R., Cataldo, A. and Andersen, C.P.: Determination of total nonstructural carbohydrates in tree species by high-performance anion-exchange chromatography with pulsed amperometric detection. *Can. J. For. Res.*, 25 (1995) 2022-2028; C.A., 124 (1996) 197385n.
- 2768 Yagi, F. and Tadera, K.: Substrate specificity and transglucosylation catalyzed by cycad  $\beta$ -glucosidase. *Biochim. Biophys. Acta*, 1289 (1996) 315-321.
- 2769 Zhang, Z., Renen, Z. and Liu, G.: High-performance liquid chromatographic analysis of hexosamines, hexosaminolts, N-acetylhexosamines and N-acetylhexosaminolts by ultraviolet and fluorescence detection at picomole levels. *J. Chromatogr. A*, 730 (1996) 107-114.
- 2770 Zhang, Z., Zhang, R. and Liu, G.: Analysis of the 9-fluorenylmethoxycarbonyl hydrazide labelling of neutral and sialic acid-containing oligosaccharides by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 728 (1996) 343-350.
- 2771 Zook, C.M. and LaCourse, W.R.: Pulsed amperometric detection of carbohydrates in fruit juices following high-performance anion-exchange chromatography. *Curr. Sep.*, 14 (1995) 48-52; C.A., 124 (1996) 143892g.
- For additional information see C.A.:  
124 (1996) 53942u, 59725x, 192929h.
- See also 2492, 2506, 2705, 2786, 2798, 2805, 3161, 3185, 3231, 3444.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 2772 Albertini, R., Rindi, S., Passi, A., Bardoni, A., Salvini, R., Pallavicini, G. and de Luca, G.: The effect of cornea proteoglycans on liposome peroxidation. *Arch. Biochem. Biophys.*, 327 (1996) 209-214.
- 2773 Campbell, M.A., Winter, A.D., Ilic, M.Z. and Handley, C.J.: Catabolism and loss of proteoglycans from cultures of bovine collateral ligament. *Arch. Biochem. Biophys.*, 328 (1996) 64-72.
- 2774 Itano, N., Oguri, K., Nagayasu, Y., Kusano, Y., Nakanishi, H., David, G. and Okayama, M.: Phosphorylation of a membrane-intercalated proteoglycan, syndecan-2, expressed in a stroma-inducing clone from a mouse Lewis lung carcinoma. *Biochem. J.*, 315 (1996) 925-930.
- 2775 Reantragoon, S., Arrigo, L.M., Dweck, H.S. and Rosenfeld, L.: Suppression of endothelin-1 production in cultured human umbilical vein endothelial cells by heparin fractions separated by strong anion exchange chromatography. *Arch. Biochem. Biophys.*, 327 (1996) 234-238.
- 2776 Shreshinskaya, G.M., Kozlova, Yu.I., Evtushenko, L.I., Taran, V.V., Shashkov, A.S. and Naumova, I.B.: (Teichoic acid of *Nocardiopsis* ssp. *VKM Ac-1457* cell wall). *Biokhimiya (Moscow)*, 61 (1996) 378-383.
- 2777 Takagaki, K., Nakamura, T., Shibata, S., Higuchi, T. and Endo, M.: Characterization and biological significance of sialyloligosaccharides synthesized in cultured human skin fibroblasts. *J. Biochem. (Tokyo)*, 119 (1996) 697-702.
- 2778 Vijayagopal, P., Figueiroa, J.E., Guo, Q., Fontenot, J.D. and Tao, Z.: Marked alteration of proteoglycan metabolism in cholesterol-enriched human arterial smooth muscle cells. *Biochem. J.*, 315 (1996) 995-1000.
- 2779 Wang, H.X., Ng, T.B., Ooi, V.E.C., Liu, W.K. and Chang, S.T.: A polysaccharide-peptide complex from cultured mycelia of the mushroom *Tricholoma mongolicum* with immunoenhancing and antitumor activities. *Biochem. Cell Biol.*, 74 (1996) 95-100.
- 2780 Zvyagintseva, T.N., Isakov, V.V., Shevchenko, N.M. and Elyakova, L.A.: (Method of isolating branched 1 $\rightarrow$ 3;1 $\rightarrow$ 6- $\beta$ -D-glucooligo- and -polysaccharides). U.S.S.R. SU 1,329,153 (Cl. C07H1/08), 20 Mar. 1995, Appl. 3,909,862, 10 Jun. 1995; C.A., 124 (1996) 117884m.

For additional information see C.A.:

124 (1996) 49829v, 235379y.

See also 2474, 2478, 2766, 3055, 3113.

#### 10c. Glycoproteins and their constituents

- 2781 Apffel, A., Chakel, J.A., Hancock, W.S., Souders, C., M'Timkulu, T. and Pungor, E., Jr.: Application of high-performance liquid chromatography-electrospray ionization mass spectrometry and matrix-assisted laser-desorption ionization time-of-flight mass spectrometry in combination with selective enzymatic modifications in the characterization of glycosylation patterns in single-chain plasminogen activator. *J. Chromatogr. A*, 732 (1996) 27-42.
- 2782 Atmani, F., Mizon, J. and Khan, S.R.: Identification of uronic acid-rich protein as urinary bikunin, the light chain of inter- $\alpha$ -inhibitor. *Eur. J. Biochem.*, 236 (1996) 984-990.

- 2783 Carron, J.A., Bates, R.C., Smith, A.I., Tetozi, T., Arellano, A., Gordon, D.L. and Burns, G.F.: Factor H co-purifies with thrombospondin isolated from platelet secretate. *Biochim. Biophys. Acta*, 1289 (1996) 305-311.
- 2784 Chambers, S.J., Langdon, M.J., Williams, P.A. and White, R.J.: Combined differential light scattering with size-exclusion chromatography techniques for monitoring conformational changes in  $\kappa$ -carrageenan. In: Phillips, G.O., Williams, P.A. and Wedlock, D.J. (Editors), *Gums Stab. Food Ind. 7, [Proc. Int. Conf.]*, 7th 1993, IRL Press, Oxford, 1994, pp. 37-53; C.A., 124 (1996) 173777r.
- 2785 Cladman, W.M., Watt, M.-A.V., Dini, J.-P. and Mellors, A.: The *Pasteurella haemolytica* O-sialoglycoprotein endopeptidase is inhibited by zinc ions and does not cleave fetuin. *Biochim. Biophys. Res. Commun.*, 220 (1996) 141-146.
- 2786 Hanisch, F.-G., Stadie, T.R.E., Deutzmann, F. and Peter-Katalinic, J.: MUC1 glycoforms in breast cancer. Cell line T47D as a model for carcinoma-associated alterations of O-glycosylation. *Eur. J. Biochem.*, 236 (1996) 318-327.
- 2787 Hervé, F., Millot, M.-C., Eap, C.B., Duché, J.-C. and Tillement, J.-P.: Two-step chromatographic purification of human plasma  $\alpha_1$ -acid glycoprotein. Its application to the purification of rare phenotype samples of the protein and their study by chromatography on immobilized metal chelate affinity adsorbent. *J. Chromatogr. B*, 678 (1996) 1-14.
- 2788 Huang, Y., Toyoda, H., Koshishi, I., Toida, T. and Imanari, T.: Determination of a depolymerized holothurian glycosaminoglycan in plasma after intravenous administration by postcolumn HPLC. *Chem. Pharm. Bull.*, 43 (1995) 2182-2186.
- 2789 Jorge, P. and Abdul-Wajid, A.: Sialyl-Tn-KLH, glycoconjugate analysis and stability by high-pH anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD). *Glycobiology*, 5 (1995) 759-764; C.A., 124 (1996) 114775r.
- 2790 Kamemura, K., Furuichi, Y., Umekawa, H. and Takahashi, T.: Purification and characterization of a pod lectin from Great Northern bean, *Phaseolus vulgaris* L. *Biochim. Biophys. Acta*, 1289 (1996) 87-94.
- 2791 Lee, K.B. and Lee, Y.C.: Conformational studies of glycopeptides by energy transfer. Introduction of fluorophore at specific branches of biantennary glycopeptides. *J. Biol. Chem.*, 271 (1996) 1462-1469.
- 2792 Okino, N., Kawabata, S., Saito, T., Hirata, M., Takagi, T. and Iwanaga, S.: Purification, characterization, and cDNA cloning of a 27-kDa lectin (L10) from horseshoe crab hemocytes. *J. Biol. Chem.*, 270 (1995) 31008-31015.
- 2793 Plummer, T.H., Jr., Phelan, A.W. and Tarentino, A.L.: Porcine fibrinogen glycopeptides: substrates for detecting endo- $\beta$ -N-acetylglucosaminidases F<sub>2</sub> and F<sub>3</sub>. *Anal. Biochem.*, 235 (1996) 98-101.
- 2794 Song, K.-S., Cho, S.-M., Lee, J.H., Kim, H.-M., Han, S.-B., Ko, K.-S. and Yoo, I.-D.: B-Lymphocyte-stimulating polysaccharide from mushroom *Phellinus linteus*. *Chem. Pharm. Bull.*, 43 (1995) 2105-2108.
- 2795 Van Damme, E.J.M., Briké, F., Winter, H.C., van Leuven, F., Goldstein, I.J. and Peumans, W.J.: Molecular cloning of two different mannose-binding lectins from tulip bulbs. *Eur. J. Biochem.*, 236 (1996) 419-427.
- 2796 Van Willigen, G., Hers, I., Gorter, G. and Akkerman, J.-W.N.: Exposure of ligand-binding sites on platelet integrin  $\alpha_{IIb}\beta_3$  by phosphorylation of the  $\beta_3$  subunit. *Biochem. J.*, 314 (1996) 769-779.
- 2797 Wang, C. and Barofsky, D.F.: Temperature effect on the separation of glycopeptides by reverse phase HPLC. In: Hatano, H. and Hanai T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, Singapore, 1995, pp. 377-383; C.A., 124 (1996) 49815n.
- 2798 Youings, A., Chang, S.-C., Dwek, R.A. and Scragg, I.G.: Site-specific glycosylation of human immunoglobulin G is latered in four rheumatoid arthritis patients. *Biochem. J.*, 314 (1996) 621-630.
- For additional information see C.A.:  
124 (1996) 76674x.
- See also 3043, 3097, 3160.
- ## 11. ORGANIC ACIDS AND LIPIDS
- ### 11a. Organic acids and simple esters
- 2799 Arin, M.J., Diez, M.T. and Resines, J.A.: High-performance liquid chromatographic determination of phenolic acids isolated from plants. *J. Liq. Chromatogr.*, 18 (1995) 4183-4192.
- 2800 Bitman, R., Swords, B., Lilion, K. and Tygi, G.: Inhibitors of lipid phosphatidate receptors: N-palmitoyl-serine and N-palmitoyl-tyrosine phosphoric acids. *J. Lipid Res.*, 37 (1996) 391-398.
- 2801 Boucher, J.-L., Delaforge, M., Salaün, J.-P., Pinot, F., Durst, F., Pflieger, P. and Mioskowski, C.: *In vitro* hydroxylation and epoxidation of some isomeric lauric acid analogs by rat liver microsomes. Identification of metabolites and effects of clofibrate or phenobarbital pretreatment. *Drug Metab. Disp.*, 24 (1996) 462-468.
- 2802 Buratti, M., Fustinoni, S. and Colombi, A.: Fast liquid chromatographic determination of urinary *trans,trans*-muconic acid. *J. Chromatogr. B*, 677 (1996) 257-263.
- 2803 Chang, M.S., Boeglin, W.E., Guengerich, F.P. and Brash, A.R.: Cytochrome P450-dependent transformations of 15R- and 15S-hydroperoxyeicosatetraenoic acids: stereoselective formation of epoxy alcohol products. *Biochemistry*, 35 (1996) 464-471.
- 2804 DeMar, J.C., Jr., Wensel, T.G. and Anderson, R.E.: Biosynthesis of the unsaturated 14-carbon fatty acids found on the N termini of photoreceptor-specific proteins. *J. Biol. Chem.*, 271 (1996) 5007-5016.
- 2805 Doughty, E.: Separation of organic acids, carbohydrates, and fermentation products by HPLC. *Lebensm.-Biotechnol.*, 12 (1995) 100-101; C.A., 124 (1996) 53947z.
- 2806 Eisele, T.A.: Determination of D-malic acid in apple juice. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 50-54.
- 2807 Fauconnier, N., Bee, A., Massart, R. and Dardoize, F.: Direct determination of organic acids in a ferrofluid ( $\gamma$ -Fe<sub>2</sub>O<sub>3</sub>) by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 783-797.

- 2808 Gao, Z., Liu, P. and Fu, C.: (Determination of low molecular carboxylic acids in *Portulaca Oleracea* L. by ion exclusion chromatography). *Sepu.*, 14 (1996) 50-52; *C.A.*, 124 (1996) 169694u.
- 2809 Gatti, R., Bousquet, E., Bonazzi, D. and Cavrini, V.: Determination of carboxylic acid salts in pharmaceuticals by high-performance liquid chromatography after precolumn fluorogenic labeling. *Biomed. Chromatogr.*, 10 (1996) 19-24; *C.A.*, 124 (1996) 185767a.
- 2810 Gioacchini, A.M., Roda, A., Galletti, G.C., Bocchini, P., Manetta, A.C. and Baraldini, M.: High-performance liquid chromatographic-electrospray mass spectrometric analysis of phenolic acids and aldehydes. *J. Chromatogr. A*, 730 (1996) 31-37.
- 2811 Glowniak, K., Zgórka, G. and Kozyra, M.: Solid-phase extraction and reversed-phase high-performance liquid chromatography of free phenolic acids in some *Echinacea* species. *J. Chromatogr. A*, 730 (1996) 25-29.
- 2812 Goromaru, T., Sera, S. and Ikeda, H.: (Isotopic fractionation of benzoic acid and hippuric acid from their deuterated analogs by high performance liquid chromatography). *Radioisotopes*, 44 (1995) 693-700; *C.A.*, 124 (1996) 49812j.
- 2813 Hashimoto, A., Yamasaki, K., Kokusenya, Y., Miyamoto, T., Nakai, H. and Sato, T.: Study on "signal" constituents for the evaluation of animal crude drugs. II. Organic acids. *Chem. Pharm. Bull.*, 43 (1995) 2195-2199.
- 2814 Iwase, H., Takatori, T., Nagao, M., Iwadate, K., Takahashi, M., Nakajima, M., Takahashi, K. and Shimizu, T.: Superoxide anion reduces the ability of myeloperoxidase to damage lipids. *Biochem. Biophys. Res. Commun.*, 219 (1996) 625-632.
- 2815 Kunii, Y., Otsuka, M., Kashino, S., Takeuchi, H. and Ohmori, S.: 4-Hydroxypipeolic acid and pipeolic acid in *Acacia* species: their determination by HPLC, its application to leguminous plants, and configuration of 4-hydroxypipeolic acid. *J. Agric. Food Chem.*, 44 (1996) 483-487; *C.A.*, 124 (1996) 111355t.
- 2816 Lord, B.S. and Stringham, R.W.: Liquid chromatographic determination of organic acids used as pharmaceutical counterions. *Anal. Chem.*, 68 (1996) 1067-1070.
- 2817 Marcato, B. and Cecchin, G.: Analysis of mixtures containing free fatty acids and mono-, di- and triglycerides by high-performance liquid chromatography coupled with evaporative light-scattering detection. *J. Chromatogr. A*, 730 (1996) 83-90.
- 2818 Maslovská, J., Duda, J. and Witusik, A.: Use of 2,4-dinitrophenylhydrazone of glyoxalic acid for the determination of glyoxilic acid by the chromatographic-spectrophotometric method and by differential pulse polarography. *Fresenius J. Anal. Chem.*, 355 (1996) 154-156.
- 2819 Miwa, H., Yamamoto, M., Futata, T., Kan, K. and Asano, T.: Thin-layer chromatography and high-performance liquid chromatography for the assay of fatty acid compositions of individual phospholipids in platelets from non-insulin-dependent diabetes mellitus patients: effect of eicosapentaenoic acid ethyl ester administration. *J. Chromatogr. B*, 677 (1996) 217-223.
- 2820 Miwa, H., Yamamoto, M., Kan, K., Futata, T. and Asano, T.: High-performance liquid chromatographic measurements of urinary hydroxycarboxylic acids as an index of the metabolic control in non-insulin-dependent diabetic patients. *J. Chromatogr. B*, 679 (1996) 1-6.
- 2821 Peldszus, S., Huck, P.M. and Andrews, S.A.: Determination of short-chain aliphatic, oxo- and hydroxyacids in drinking water at low microgram per liter concentrations. *J. Chromatogr. A*, 723 (1996) 27-34.
- 2822 Pinkernell, U., Effkemann, S., Nitzsche, F. and Karst, U.: Rapid high-performance liquid chromatographic method for the determination of peroxyacetic acid. *J. Chromatogr. A*, 730 (1996) 203-208.
- 2823 Prucha, M., Wray, V. and Pieper, K.N.: Metabolism of 5-chloro-substituted muconolactones. *Eur. J. Biochem.*, 237 (1996) 357-366.
- 2824 Rosolowsky, M., Falck, J.R. and Campbell, W.B.: Metabolism of arachidonic acid by canine polymorphonuclear leukocytes synthesis of lipoxygenase and omega-oxidized metabolites. *Biochim. Biophys. Acta*, 1300 (1996) 143-150.
- 2825 Schneider, G. and Schmidt, J.: Liquid chromatography-electrospray ionization mass spectrometry for analysing plant hormone conjugates. *J. Chromatogr. A*, 728 (1996) 371-375.
- 2826 Shen, Y., Obuseng, V., Grönberg, L. and Jönsson, J.Å.: Liquid membrane enrichment for the ion chromatographic determination of carboxylic acids in soil samples. *J. Chromatogr. A*, 725 (1996) 189-197.
- 2827 Song, D.-J. and Hsu, K.-Y.: Determination of *p*-aminobenzoic acid and its metabolites in rabbit plasma by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. B*, 677 (1996) 69-75.
- 2828 Van Overbeke, A., Baeyens, W. and Dewaele, C.: Enantiomeric separation of six 2-arylpropionic acids after pre-column derivatization with various amines and alcohols on a cellulose-based chiral stationary phase. *Anal. Chim. Acta*, 321 (1996) 245-261.
- 2829 Waksmundzka-Hajnos, M., Bieganowska, M.L. and Petruczynik, A.: High-performance liquid chromatographic separation of some 2- and 4-hydroxy derivatives of 2'-benzoylbenzoic acid on Florisil and silica gel columns. *J. Chromatogr. A*, 730 (1996) 195-201.
- 2830 Weiss, J. and Reinherd, S.: (Aliphatic carboxylic acids. A difficult chromatographic problem). *GIT Spez. Chromatogr.*, 15 (1995) 105-111; *C.A.*, 124 (1996) 105084n - a review with 16 refs.
- 2831 Wu, T., Ikezoni, T., Angus, C.W. and Shelhamer, J.H.: Tumor necrosis factor- $\alpha$  induces the 85-kDa cytosolic phospholipase A<sub>2</sub> gene expression in human bronchial epithelial cells. *Biochim. Biophys. Acta*, 1310 (1996) 175-184.
- 2832 Yamane, M. and Abe, A.: High-performance liquid chromatography-thermospray mass spectrometry of 5,6-dihydroxyeicosatrienoate-1,5-lactone from tissue homogenates. *J. Chromatogr. B*, 678 (1996) 339-343.
- 2833 Yamane, M., Abe, A., Shimizu, S. and Nakajima, M.: Subcellular localization of docosahexaenoic acid and arachidonic acid  $\omega$ -hydroxylation activity in the brain, liver and colonic adenocarcinoma. *J. Chromatogr. A*, 730 (1996) 91-98.

For additional information see *C.A.*:

124 (1996) 66741c, 111366x, 197564v, 201642p.

See also 2426, 2531, 2648, 2695, 2720, 2751, 2862, 3486, 3751, 3828.

## 11b. Prostaglandins

- 2834 Kumlin, M.: Analytical methods for the measurement of leukotrienes and other eicosanoids in biological samples from asthmatic subjects. *J. Chromatogr. A*, 725 (1996) 29-40 - a review with 56 refs.
- 2835 Oda, Y., Mano, N. and Asakawa, N.: Simultaneous determination of thromboxane B<sub>2</sub>, prostaglandin E<sub>2</sub> and leukotriene B<sub>4</sub> in whole blood by liquid chromatography/mass spectrometry. *J. Mass Spectrom.*, 30 (1995) 1671-1678; C.A., 124 (1996) 45874h.
- 2836 Qiu, D.W., Hui, K.P., Lee, C.W., Lim, T.K. and Tan, W.C.: Simplified method for measuring urinary leukotriene E<sub>4</sub>. *J. Chromatogr. B*, 677 (1996) 152-155.

See also 2824.

## 11c. Lipids and their constituents

- 2837 Beckman, B.S., Mallia, C. and Clejan, S.: Molecular species of phospholipids in a murine stem-cell line responsive to erythropoietin. *Biochem. J.*, 314 (1996) 861-867.
- 2838 Bosque, F., Lacoste, F. and Mordret, F.: (Identification of rapeseed methyl esters incorporated in diesel fuel by liquid chromatography-gas chromatography). *Oil, Corps Gras, Lipides*, 2 (1995) 311-315; C.A., 124 (1996) 150170Y.
- 2839 Choudhari, K.B., J Jayanthi, S., Murty, R.B. and Matharu, R.P.: A high-performance liquid chromatographic method for the analysis of lipids from lyophilized formulations. *J. Chromatogr. A*, 724 (1996) 343-347.
- 2840 Demopoulos, C.A., Antonopoulou, S., Andrikopoulos, N.K. and Kapoulas, V.M.: Isolation and complete separation of lipids from natural sources. *J. Liq. Chromatogr.*, 19 (1996) 521-535.
- 2841 Demopoulos, C.A., Kyrili, M., Antonopoulou, S. and Andrikopoulos, N.K.: Separation of several main glycolipids into classes and partially into species by HPLC an UV-detection. *J. Liq. Chromatogr.*, 19 (1996) 771-781.
- 2842 Dupuis, F., Denizot, Y., Fixe, P., Dulery, C. and Praloran, V.: PAF and haematopoiesis. X. Macrophage colony-stimulating factor and granulocyte macrophage colony-stimulating factor enhance platelet-activating factor acetylhydrolase production by human blood-derived macrophages. *Biochim. Biophys. Acta*, 1311 (1996) 27-32.
- 2843 Kühn, G., Weidner, S., Just, U. and Hohner, G.: Characterization of technical waxes. Comparison of chromatographic techniques and matrix-assisted laser-desorption/ionization mass spectrometry. *J. Chromatogr. A*, 732 (1996) 111-117.
- 2844 Kusaka, T., Ishihara, S., Sakaida, M., Mifune, A., Nakano, Y., Tsuda, K., Ikeda, M. and Nakano, H.: Composition analysis of normal plant triacylglycerols and hydroperoxidized rac-1-stearoyl-2-oleoyl-3-linoleoyl-sn-glycerols by liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *J. Chromatogr. A*, 730 (1996) 1-7.
- 2845 Kwon, G., Bohrer, A., Han, X., Corbett, J.A., Ma, Z., Gross, R.W., McDaniel, M.L. and Turk, J.: Characterization of the sphingomyelin content of isolated pancreatic islets. Evaluation of the role of sphingomyelin hydrolysis in the action of interleukin-1 to induce islet overproduction of nitric oxide. *Biochim. Biophys. Acta*, 1300 (1996) 63-72.

- 2846 Matthaeus, B., Loesing, R. and Fiebig, H.-J.: Determination of inositol phosphates IP3-IP6 in rapeseed and rapeseed meal by an HPLC method, part 2: investigation of rapeseed and rapeseed meal and comparison with other methods. *Fett Wiss. Technol.*, 97 (1995) 372-374; C.A., 124 (1996) 173746e.
- 2847 Muethling, J.: (Separation of isomeric gangliosides by anion-exchange HPLC).  *GIT Fachz. Lab.*, 39 (1995) 897-898; C.A., 124 (1996) 140083h.
- 2848 Natowicz, M.R., Prence, E.M., Chaturvedi, P. and Newburg, D.S.: Urine sulfatides and the diagnosis of metachromatic leukodystrophy. *Clin. Chem. (Washington)*, 42 (1996) 232-238.
- 2849 Neff, W.E. and Byrdwell, W.C.: Triacylglycerol analysis by high performance liquid chromatography-atmospheric pressure chemical ionization mass spectrometry: *Crepis alpina* and *Vernonia galamensis* seed oils. *J. Liq. Chromatogr.*, 18 (1995) 4165-4181.
- 2850 Ni, Y.-C., Wong, T.-Y., Lloyd, R.V., Heinze, T.M., Shelton, S., Casciano, D., Kadlubar, F.F. and Fu, P.P.: Mouse liver microsomal metabolism of chloral hydrate, trichloroacetic acid, and trichloroethanol leading to induction of lipid peroxidation via a free radical mechanism. *Drug Metab. Disp.*, 24 (1996) 81-90.
- 2851 Ou, Z., Ogamo, A., Kawai, Y. and Nakagawa, Y.: Quantitation of choline glycerophospholipids that contain carboxylate residues by fluorometric high-performance liquid chromatography. *J. Chromatogr. A*, 724 (1996) 131-136.
- 2852 Rodriguez, P.E.A., Maggio, B. and Cumar, F.A.: Acid and enzymatic hydrolysis of the internal sialic acid residue in native and chemically modified ganglioside GM1. *J. Lipid Res.*, 37 (1996) 382-390.
- 2853 Saeki, A., Tsuchida, T., Yamada, R. and Miyazawa, T.: (Measurement of lipid hydroperoxides in human blood by using chemiluminescence). *Optronics*, 162 (1995) 142-146; C.A., 124 (1996) 49822n.
- 2854 Schlegelmilch, F., Siemanowski, W. and Wiebrock, L.: Development of an HPLC method for determining the free polyglycerol content of polyglycerol caprates. *Fett Wiss. Technol.*, 97 (1995) 408-411; C.A., 124 (1996) 185118q.
- 2855 Schnurr, K., Belkner, J., Ursini, F., Schewe, T. and Kühn, H.: The selenoenzyme phospholipid hydroperoxide glutathione peroxidase controls the activity of the 15-lipoxygenase with complex substrates and preserves the specificity of the oxygenation products. *J. Biol. Chem.*, 271 (1996) 4653-4658.
- 2856 Takayama, M.: Simplified quantitative determination of phospholipids in amniotic fluid by high-performance liquid chromatography. *Med. Sci. Res.*, 23 (1995) 717-718; C.A., 124 (1996) 49814m.
- 2857 Van Dijken, P., de Haas, J.-R., Craxton, A., Erneux, C., Shears, S.B. and van Haastert, P.J.M.: A novel, phospholipase C-independent pathway of inositol 1,4,5-trisphosphate formation in *Dictyostelium* and rat liver. *J. Biol. Chem.*, 270 (1995) 29724-29731.
- 2858 Wada, M., Nakashima, K., Kuroda, N., Akiyama, S. and Imai, K.: Sensitive flow-injection method with peroxyoxalate chemiluminescence detection combined with preparative high-performance liquid chromatography for determination of choline-containing phospholipids in human serum. *J. Chromatogr. B*, 678 (1996) 129-136.

For additional information see C.A.:  
124 (1996) 173803w.

See also 2545, 2638, 2817, 2862, 2964, 3300, 3438.

**11d. Lipoproteins and their constituents**

- 2859 Brown, A.J., Dean, R.T. and Jessup, W.: Free and esterified oxysterol: formation during copper-oxidation of low density lipoprotein and uptake by macrophages. *J. Lipid Res.*, 37 (1996) 320-335.
- 2860 Hoeg, J.M., Vaismann, B.L., Demoskey, S.J., Jr., Meyn, S.M., Talley, G.D., Hoyt, R.F., Jr., Feldman, S., Berard, A.M., Sakai, N., Wood, D. et al.: Lecithin:cholesterol acyltransferase overexpression generates hyperalpha-lipoproteinemia and a nonatherogenic lipoprotein pattern in transgenic rabbits. *J. Biol. Chem.*, 271 (1996) 4396-4402.
- 2861 Kitamura, T., Ito, S., Kato, Y., Sasamoto, K. and Okazaki, M.: (Analysis of serum lipoproteins by high performance gel filtration chromatography). *Bunseki Kagaku*, 45 (1996) 103-106; C.A., 124 (1996) 169670h.
- 2862 Lass, A., Belkner, J., Esterbauer, H. and Kühn, H.: Lipoxygenase treatment renders low-density lipoprotein susceptible to Cu<sup>2+</sup>-catalysed oxidation. *Biochem. J.*, 314 (1996) 577-585.
- 2863 Suwabe, A., Mason, R.J. and Voelker, D.R.: Calcium dependent association of surfactant protein A with pulmonary surfactant: application to simple surfactant protein A purification. *Arch. Biochem. Biophys.*, 327 (1996) 285-291.
- 2864 Winger, T.M., Ludovice, P.J. and Chaikof, E.L.: Purification of synthetic lipopeptide conjugates by liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 4117-4125.

**12. ORGANIC PEROXIDES**

See 2850, 2853, 2878.

**13. STEROIDS**

**13a. General techniques**

- 2865 Makin, H.L.J., Honour, J.W. and Shackleton, C.H.L.: General method of steroid analysis, Part 1. Extraction, purification and measurement of steroids by high-performance liquid chromatography, gas chromatography and mass spectrometry. In: Makin, H.L.J., Gower, D.B. and Kirk, D.N. (Editors), *Steroid Anal.*, Blackie, Glasgow, 1995, pp. 114-228; C.A., 124 (1996) 249151e - a review with many refs.
- 2866 Salo, Sirén, H., Volin, P., Wiedmer, S. and Vuorela, H.: Structure-retention relationships of steroid hormones in reversed-phase liquid chromatography and micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 728 (1996) 83-88.

For additional information see C.A.:  
124 (1996) 111720h.

See also 2875, 3054, 3472, 3646.

**13b. Pregnan and androstane derivatives**

- 2867 Bagnani, R., Ramazza, V., Zucchi, M., Simonella, A., Leone, F., Bellini, A. and Fanelli, R.: Analysis of dexamethasone and betadexamethasone in bovine urine by purification with an "online" immunoaffinity chromatography-high performance liquid chromatography system and determination by gas-chromatography-mass spectrometry. *Anal. Biochem.*, 235 (1996) 119-126.
- 2868 Eisen, C., Meyer, C., Dressendorfer, R., Strasburger, C., Decker, H. and Wehling, M.: Binding-labelled photoactivatable aldosterone and progesterone derivatives as ligands for affinity chromatography, fluorescence immunoassays and photoaffinity labelling. *Eur. J. Biochem.*, 237 (1996) 514-518.
- 2869 Kanda, T., Shirota, O. and Ohtsu, Y.: (Simultaneous determination of cortisol and cortisone in saliva by semimicrocolumn high-performance liquid chromatography). *Kuromatogurafi*, 16 (1995) 332-333; C.A., 124 (1996) 135886t.
- 2870 Leopold, C.S. and Friend, D.R.: *In vivo* pharmacokinetic study for the assessment of poly(L-aspartic acid) as a drug carrier for colon-specific drug delivery. *J. Pharmacokin. Biopharm.*, 23 (1995) 397-406.
- 2871 Mueller, H.W. and Eitel, J.: Quality control in the determination of cortisol in plasma/serum by using, on every sample, two different three-step separation methods including ultrafiltration, restricted-access high-performance liquid chromatography and reversed-phase high-performance liquid chromatography, and contrasting results to immunoassays. *J. Chromatogr. B*, 678 (1996) 137-150.
- 2872 Walker, V.R., Dombi, G.W., Gutai, J.P., Wade, D.D., Swartz, K.H., Liu, H. and Schroeder, R.R.: Semiautomated method for the quantitation of plasma or sera androstenedione, testosterone, and dihydrotestosterone in population studies. *Anal. Biochem.*, 234 (1996) 194-203.

For additional information see C.A.:  
124 (1996) 66740b.

See also 2583, 3495, 3619, 3681, 3712, 3726.

**13c. Estrogens**

- 2873 Goeroeg, S., Brlik, J., Csehi, A., Halmos, Z., Herenyi, B., Horvath, P., Fravetz, F. and Bor, D.: Analysis of steroids. Part 47. Estimation of impurity profiles of drugs and related materials, part 13: identification of impurities in estradiol. *Anal. Methods Instrum.*, 2 (1995) 154-157; C.A., 124 (1996) 97856y.
- 2874 Oosterkamp, A.J., Villaverde Herriaiz, M.T., Irth, H., Tjaden, U.R. and van der Greef, J.: Reversed-phase liquid chromatography coupled on-line to receptor affinity detection based on the human estrogen receptor. *Anal. Chem.*, 68 (1996) 1201-1206.
- 2875 Sadlej-Sosnowska, N.: Thermodynamic parameters of the formation of a complex between cyclodextrins and steroid hormones. *J. Chromatogr. A*, 728 (1996) 89-95.
- 2876 Spencer, B.J. and Purdy, W.C.: High-performance liquid chromatographic separation of equilin, estrone, and estrone derivatives with cyclodextrins as mobile phase additives. *J. Liq. Chromatogr.*, 18 (1995) 4063-4080.

See also 3675.

## 13d. Sterols

- 2877 Feingold, K.R., Spady, D.K., Pollock, A.S., Moser, A.H. and Grunfeld, C.: Endotoxin, TNF and IL-1 decrease cholesterol 7 $\alpha$ -hydroxylase mRNA levels and activity. *J. Lipid Res.*, 37 (1996) 223-228.
- 2878 Lercker, G., Bortolomeazzi, R., Pizzale, L. and Vichi, S.: Thermal degradation of single 7-cholesteryl acetate hydroperoxide. *Chromatographia*, 42 (1996) 29-33.
- 2879 Nakajima, M., Yamato, S., Wakabayashi, H. and Shimada, K.: High-performance liquid chromatographic determination of cholesterol and cholestanol in human serum by precolumn derivatization with 2-[2(isocyanate)ethyl]-3-methyl-1,4-naphthoquinone combined with platinum catalyst reduction and electrochemical detection. *Biol. Pharm. Bull.*, 18 (1995) 1762-1764; *C.A.*, 124 (1996) 140088p.
- 2880 Swami, S., Szaltryd, C. and Kraemer, F.B.: Effects of streptozotocin-induced diabetes on low density lipoprotein receptor expression in rat adipose tissue. *J. Lipid Res.*, 37 (1996) 229-236.

For additional information see *C.A.*:  
124 (1996) 185802h.

See also 3742.

## 13e. Bile acids and alcohols

For additional information see *C.A.*:  
124 (1996) 50156e.

## 13f. Ecdysones and other insect steroid hormones

- 2881 Gildengorn, V.D.: Reversed-phase affinity chromatography of ecdysteroids with boronic acid-containing eluents. *J. Chromatogr. A*, 730 (1996) 147-152.

## 14. STEROID GLYCOSIDES AND SAPONINS

- 2882 Kanamori, H., Terauchi, M., Fuse, J.-i., Nobusou, M., Sakamoto, I., Eto, S.-y. and Tanabe, K.: Saponin type of commercial *Panacis japonici* rhizoma. *Nat. Med.*, 49 (1995) 293-297; *C.A.*, 123 (1996) 97372n.
- 2883 Palagiano, E., Zollo, F., Minale, L., Iorizzi, M., Bryan, P., McClintock, J. and Hopkins, T.: Isolation of 20 glycosides from starfish *Henricia downeyae*, collected in the gulf of Mexico. *J. Natural Prod.*, 59 (1996) 348-354.
- 2884 Park, M.K., Park, J.H., Hwang, G.S., Lee, M.Y. and Park, I.J.: Analysis of ginsenosides by thermospray LC/MS. *Koryo Insam Hakhoechi*, 19 (1995) 134-137; *C.A.*, 124 (1996) 212191s.

See also 3734.

## 15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

## 15a. Terpenes

- 2885 Bazylak, G., Rosiak, A. and Shi, C.-Y.: Systematic analysis of glucoriridoids from *Penstemon serrulatus* Menz. by high-performance liquid chromatography with pre-column solid-phase extraction. *J. Chromatogr. A*, 725 (1996) 177-187.
- 2886 Bicchi, C. and Rubiolo, P.: High-performance liquid chromatographic-particle beam mass spectrometric analysis of sesquiterpene lactones with different carbon skeletons. *J. Chromatogr. A*, 727 (1996) 211-221.
- 2887 Lin, L.-J., Lee, K.R. and Shiao, M.-S.: Partition coefficients in octanol-water and liquid chromatographic behavior of eight oxygenated triterpenes which are paired stereo- and postional isomers. *J. Chin. Chem. Soc. (Taipei)*, 42 (1995) 981-985; *C.A.*, 124 (1996) 117651h.
- 2888 Nilsson, U., Bergh, M., Shao, L.P. and Karlberg, A.-T.: Analysis of contact allergenic compounds in oxidized  $\alpha$ -limonene. *Chromatographia*, 42 (1996) 199-205.

## 15b. Essential oils

- 2889 Buiarelli, F., Cartoni, G.P., Cocciali, F. and Leone, T.: Analysis of bitter essential oils from orange and grapefruit by high-performance liquid chromatography with microbore columns. *J. Chromatogr. A*, 730 (1996) 9-16.
- 2890 Calvarano, M.: (Use of HPLC methods for determining specific components of citrus juices and essential oils). *Essenze Deriv. Agrum.*, 64 (1994) 471-516; *C.A.*, 124 (1996) 155610b - a review with many refs.
- 2891 Kosman, V.M., Blinova, K.F. and Zenkevich, I.G.: (Identification and determination of eugenol in *Geum urbanum* L. roots). *Rastit. Resur.*, 31 (1995) 93-97; *C.A.*, 124 (1996) 127231k.

For additional information see *C.A.*:  
124 (1996) 242431n.

## 15c. Bitter substances

- 2892 Hughes, P.S.: Preparative regime for the purification of bitter acids derived from hops (*Humulus lupulus* L.). *J. Chromatogr. A*, 731 (1996) 327-330.

## 16. NITRO AND NITROSO COMPOUNDS

- 2893 Caton, J.E. and Griest, W.H.: Determination of explosives and some metabolites of TNT in biological and environmental samples by liquid chromatography on a mixed-mode C<sub>18</sub>-anion column. *J. Liq. Chromatogr.*, 19 (1996) 661-677.
- 2894 Lewin, U., Efer, J. and Engewald, W.: High-performance liquid chromatographic analysis with electrochemical detection for residues of explosives in water samples around a former ammunition plant. *J. Chromatogr. A*, 730 (1996) 161-167.
- 2895 McCarthy, R.M., Farmer, P. and Sheehan, D.: Binding of 2-hydroxy-5-nitrobenzyl alcohol to rat alpha class glutathione S-transferases; evidence for binding at tryptophan 21. *Biochim. Biophys. Acta*, 1293 (1996) 185-190.

- 2896 Murayama, M. and Dasgupta, P.K.: Liquid chromatographic determination of nitro-substituted polynuclear aromatic hydrocarbons by sequential electrochemical and fluorescence detection. *Anal. Chem.*, 68 (1996) 1226-1232.
- 2897 Volmer, D.A., Lay, J.O., Jr., N'Billedeau, S.M. and Vollmer, D.L.: Detection and confirmation of N-nitrosodialkylamines using liquid chromatography - electrospray ionization coupled on-line with a photolysis reactor. *Anal. Chem.*, 68 (1996) 546-552.

For additional information see C.A.:

124 (1996) 60909s, 150290n, 218996e.

See also 2669, 3637.

## 17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

### 17a. Amines and polyamines

- 2898 Carbonnelle, P., Boukortt, S., Lison, D. and Buchet, J.P.: Determination of toluenediamines in urine of workers occupationally exposed to isocyanates by high-performance liquid chromatography. *Analyst (Cambridge)*, 121 (1996) 663-669.
- 2899 Chiba, T., Akizawa, T., Mastukawa, M., Pan-Hou, H., Kawai, N. and Yoshioka, M.: Large-scale purification and further characterization of spidamine and joramine from venom of spider, *Nephila clavata*. *Chem. Pharm. Bull.*, 43 (1995) 2177-2181.
- 2900 Chriswanto, H. and Wallace, G.G.: Characterization of polyaniline using chromatographic studies. *Chromatographia*, 42 (1996) 191-198.
- 2901 Clark, C.R., DeRuiter, J. and Noggle, F.T.: Analysis of 1-(3-methoxy-4,5-methylenedioxophenyl)-2-propanamine (MDMA) derivatives synthesized from nutmeg oil and 3-methoxy-4,5-methylenedioxobenzaldehyde. *J. Chromatogr. Sci.*, 34 (1996) 34-42.
- 2902 DeRuiter, J., Clark, C.R. and Noggle, F.T.: LC and GC-MS analysis of 4-bromo-2,5-dimethoxyphenethylamine (Nexus) and 2-propanamine and 2-butanamine analogues. *J. Chromatogr. Sci.*, 33 (1995) 583-590.
- 2903 Desplanques, D. and Calves, N.: (Identification and determination of biogenic amines in meals of animal origin). *Ann. Falsif. Expert. Chim. Toxicol.*, 88 (1995) 201-211; C.A., 124 (1996) 173792s.
- 2904 Dobberpuhl, D.A., Hoekstra, J.C. and Johnson, D.C.: Pulsed electrochemical detection at gold electrodes applied to monoamines and diamines following their chromatographic separation. *Anal. Chim. Acta*, 322 (1996) 55-62.
- 2905 Malle, P., Valle, M. and Bouquelet, S.: Assay of biogenic amines involved in fish decomposition. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 43-49.
- 2906 Monser, L.I. and Greenway, G.M.: Liquid chromatographic determination of methylamines. Determination of trimethylamine in fish samples using a porous graphitic carbon stationary phase. *Anal. Chim. Acta*, 322 (1996) 63-68.
- 2907 Moret, S. and Conte, L.S.: High-performance liquid chromatographic evaluation of biogenic amines in foods. An analysis of different methods of sample preparation in relation to food characteristics. *J. Chromatogr. A*, 729 (1996) 363-369.

- 2908 Morris, N.M., Greally, B.R. and Cairns, P.M.: An investigation of factors influencing the resolution of chiral alkyl aromatic amines on a dinitrobenzoyl-(S)-leucine HPLC column. *J. Liq. Chromatogr.*, 19 (1996) 489-502.

- 2909 Pietsch, J., Hampel, S., Schmidt, W., Brauch, H.-J. and Worch, E.: Determination of aliphatic and alicyclic amines in water by gas and liquid chromatography after derivatization by chloroformates. *Fresenius J. Anal. Chem.*, 355 (1996) 164-173.
- 2910 Robert, A., Ducos, P. and Francin, J.M.: Determination of urinary 4,4'-methylenedianiline and its acetylated metabolites by solid-phase extraction and HPLC analysis with UV and electrochemical detection. *Int. Arch. Occup. Environ. Health*, 68 (1995) 44-51; C.A., 124 (1996) 196074y.

- 2911 Sanchez-Rodas, D., Hohaus, E. and Wenclawiak, B.: High-performance liquid chromatographic determination of primary amines in aqueous solutions after extraction and derivatization with 2,2-diphenyl-1-oxa-3-oxonia-2-boratanaphthalene (DOOB). *Fresenius J. Anal. Chem.*, 355 (1996) 187-189.

- 2912 Seiler, N., Delcros, J.-G., Cipolla, B., Chamaillard, L., Havouis, R., Quemener, V. and Moulinoux, J.-P.: Observations with an S-adenosylmethionine decarboxylase inhibitor in rats with prostatic adenocarcinoma. *Arzneim.-Forsch.*, 46 (1996) 311-315.

- 2913 Van Eijk, H.M.H., Rooyakers, D.R. and Deutz, N.E.P.: Automated determination of polyamines by high-performance liquid chromatography with simple sample preparation. *J. Chromatogr. A*, 730 (1996) 115-120.

See also 2688, 2827, 2923.

### 17b. Catecholamines and their metabolites

- 2914 Hibi, K. and Kudo, K.: (Simultaneous measurement of VMA, HVA, and creatinine. Using LC system with autosampler with microplate). *Jasco Rep.*, 37 (1995) 74-75; C.A., 124 (1996) 111317g - a review without refs.
- 2915 Ma, Y., Sokoloski, E. and Ito, Y.: pH-Zone refining counter-current chromatography of polar catecholamines using di(2-ethylhexyl)phosphoric acid as a ligand. *J. Chromatogr. A*, 724 (1996) 348-353.
- 2916 Róna, K., Ary, K., Gachályi, B. and Klebovich, I.: Determination of  $\alpha$ -methyldopa in human plasma by validated high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 730 (1996) 125-131.
- 2917 Swahn, C.-G., Halldin, C., Naagren, K. and Farde, L.: An HPLC-method for determination of ligand metabolism during PET studies with [ $^{11}\text{C}$ ]metaraminol and [ $^{11}\text{C}$ ]-norepinephrine. *Nud. Med. Biol.*, 22(8, New Radio-Tracers and Methods of Quality Assurance for Nuclear Medicine Applications) (1995) 1045-1048; C.A., 124 (1996) 169506j.
- 2918 Tsai, T.H., Chou, C.J., Tsai, W.J. and Chen, C.F.: Magnolol inhibits arachidonic acid-induced rabbit platelet serotonin release, measure by microbore HPLC with electrochemical detection. *J. Liq. Chromatogr.*, 19 (1996) 949-955.

See also 2503, 2747, 2946.

## 17c. Urea and guanidine derivatives

- 2919 Shintani, H.: Comparison of solid-phase extraction and dialysis on pretreatment efficiency of blood urea analysis. *J. Chromatogr. Sci.*, 34 (1996) 92-94.

## 17d. Other amine derivatives and amides (excl. peptides)

- 2920 Bellec, G., Cauvin, J.M., Salaun, M.C., le Calvé, K., Dréano, Y., Gouériou, H., Ménez, J.F. and Berthou, F.: Analysis of N-nitrosamines by high-performance liquid chromatography with post-column photohydrolysis and colorimetric detection. *J. Chromatogr. A*, 727 (1996) 83-92.
- 2921 Fang, J.Y. and Tsai, Y.H.: Stability of nonivamide and sodium nonivamide acetate in solution at various pH values. *Pharmazie*, 51 (1996) 191-192.
- 2922 Ghanayem, B.I. and Burka, L.T.: Excretion and identification of methacrylonitrile metabolites in the bile of male F344 rats. *Drug Metab. Disp.*, 24 (1996) 390-394.
- 2923 Kleiderigg, O.P., Posch, K. and Lindner, W.: Synthesis and application of a new isothiocyanate as a chiral derivatizing agent for the indirect resolution of chiral amino alcohols and amines. *J. Chromatogr. A*, 729 (1996) 33-42.
- 2924 Lin, C.-e., Lin, F.-k. and Lin, C.-h.: Enantioseparation of amino acids and aminoalcohols on chiral stationary phases derived from (S or R)-N-phenylalanyl-(S or R)-1-( $\alpha$ -naphthyl)ethylamine. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 701-711; C.A., 124 (1996) 75136m.
- 2925 Mengerink, Y., Mure, M., de Brabander, E.M.M. and van der Wal, Sj.: Exclusion chromatography of polypropylenamine dendrimers. *J. Chromatogr. A*, 730 (1996) 75-81.
- 2926 Tsai, T.-R., Cham, T.-M., Chen, K.-C., Chen, C.-F. and Tsai, T.-H.: Determination of acetylcholine by on-line microdialysis coupled with pre- and post-microbore column enzyme reactors with electrochemical detection. *J. Chromatogr. B*, 678 (1996) 151-155.

## 18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS

## 18a. Amino acids and their derivatives

- 2927 Andre, M., Domanig, R., Riemer, E., Moser, H. and Groepelin, A.: Identification of the thermal degradation products of G-triiodothyronine sodium (liothyronine sodium) by reversed-phase high-performance liquid chromatography with photodiode-array UV and mass spectrometric detection. *J. Chromatogr. A*, 725 (1996) 287-294.
- 2928 Beppu, T., Shirahata, A. and Samejima, K.: Determination of hypusine and deoxyhypusine covalently bound to protein using submilligram amounts of protein samples by HPLC. *Biol. Pharm. Bull.*, 19 (1996) 1-5; C.A., 124 (1996) 197381h.
- 2929 Berthod, A., Liu, Y., Bagwill, C. and Armstrong, D.W.: Facile liquid chromatographic enantioresolution of native amino acids and peptides using a teicoplanin chiral stationary phase. *J. Chromatogr. A*, 731 (1996) 123-137.
- 2930 Brady, J.D., Sadler, I.H. and Fry, S.C.: Di-isodityrosine, a novel tetrameric derivative of tyrosine in plant cell wall proteins: a new potential cross-link. *Biochem. J.*, 315 (1996) 323-327.

- 2931 Brückner, H. and Wachsmann, M.: Liquid chromatographic separation of amino acid enantiomers on a silica-bonded chiral s-triazine column. *J. Chromatogr. A*, 728 (1996) 447-454.
- 2932 Carducci, C., Birarelli, M., Leuzzi, V., Santagata, G., Serafini, P. and Antonozzi, I.: Automated method for the measurement of amino acids in urine by high-performance liquid chromatography. *J. Chromatogr. A*, 729 (1996) 173-180.
- 2933 Chen, X. and Sato, M.: HPLC determination of primary amino acids using electrochemiluminescence of ruthenium complex. *J. Flow Injection Anal.*, 12 (1995) 216-222; C.A., 124 (1996) 218990y.
- 2934 De Jonge, L.H. and Breuer, M.: Evaluation of systematic errors due to deproteinization, calibration and storage of plasma for amino acid assay by ion-exchange chromatography. *J. Chromatogr. B*, 677 (1996) 61-68.
- 2935 Dorresteijn, R.C., Berwald, L.G., Zomer, G., de Gooijer, C.D., Wieten, G. and Beuvery, E.C.: Determination of amino acids using o-phthalaldehyde-2-mercaptoethanol derivatization. Effect of reaction conditions. *J. Chromatogr. A*, 724 (1996) 159-167.
- 2936 El-Waziry, A.M., Tomita, Y., Ling, J.R. and Onodera, R.: Measurement of total and separate stereoisomers of diaminopimelic acid in rumen bacteria by high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 53-59.
- 2937 Galletti, G.C., Bocchini, P., Chiavari, G. and Fabbri, D.: High-performance liquid chromatography versus gas chromatography in the analysis of proteinaceous painting ligands. *Fresenius J. Anal. Chem.*, 354 (1996) 381-383.
- 2938 Gilon, N., Astruc, A., Astruc, M. and Potin-Gautier, M.: Selenoamino acid speciation using HPLC-ETAAS following an enzymic hydrolysis of selenoprotein. *Appl. Organomet. Chem.*, 9 (1995) 623-628; C.A., 124 (1996) 81066e.
- 2939 Hazen, S.L., Hsu, F.F. and Heinecke, J.W.: p-Hydroxyphenyl-acetaldehyde is the major product of L-tyrosine oxidation by activated human phagocytes. A chloride-dependent mechanism for the conversion of free amino acids into reactive aldehydes by myeloperoxidase. *J. Biol. Chem.*, 271 (1996) 1861-1867.
- 2940 Khalifa, M. and Molnár-Perl, I.: The analysis of phenylthiocarbamyl amino acids in the presence of sodium chloride. *LC-GC Int.*, 9 (1996) 143-146.
- 2941 Kiba, N., Oyama, Y., Kato, A. and Furusawa, M.: Postcolumn co-immobilized leucine dehydrogenase-NADH oxidase reactor for the determination of branched-chain amino acids by high-performance liquid chromatography with chemiluminescence detection. *J. Chromatogr. A*, 724 (1996) 354-357.
- 2942 Klein, B.H. and Dudenhausen, J.W.: Ion-exchange chromatography and ion-pair chromatography. Complementation of HPLC analysis of amino acids in body fluids by pre-column derivatization usig ortho-phthalodialdehyde. *J. Liq. Chromatogr.*, 18 (1995) 4007-4028.
- 2943 Kuromizu, K., Abe, O., Izumiya, N. and Kodera, Y.: Preparation of a chiral stationary phase containing a L-cysteine residue and its application to enantioselective separation of dansyl amino acids. *Fukuoka Shika Daigaku Gakkai Zasshi*, 22 (1995) 359-369; C.A., 124 (1996) 197372f.
- 2944 Li, G.-D., Krull, I.S. and Cohen, S.A.: Electrochemical activity of 6-aminoquinolyl urea derivatives of amino acids and peptides. Application to high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. A*, 724 (1996) 147-157.

- 2945 Lindner, K.-J., Hartvig, P., Åkesson, C., Tyrefors, N., Sundin, A. and Långström, B.: Analysis of L-[methyl-<sup>14</sup>C]methionine and metabolites in human plasma by an automated solid-phase extraction and a high-performance liquid chromatographic procedure. *J. Chromatogr. B*, 679 (1996) 13-19.
- 2946 Ma, Y. and Ito, Y.: Affinity countercurrent chromatography using a ligand to stationary phase. *Anal. Chem.*, 68 (1996) 1207-1211.
- 2947 Martins, A.R. and Padovan, A.P.: A practical approach to improve the resolution of dansyl-amino acids by high-performance liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 467-476.
- 2948 Maruyama, W., Hashizume, Y., Matsubara, K. and Naoi, M.: Identification of 3-nitro-L-tyrosine, a product of nitric oxide and superoxide, as an indicator of oxidative stress in the human brain. *J. Chromatogr. B*, 676 (1996) 153-158.
- 2949 Messina, A., Girelli, A.M., Flieger, M., Sinibaldi, M., Sedmera, P. and Cvak, L.: Direct resolution of optically active isomers on chiral packings containing ergoline skeletons. 5. Enantioseparation of amino acid derivatives. *Anal. Chem.*, 68 (1996) 1191-1196.
- 2950 Oguri, S., Uchida, C., Mishina, M., Miki, Y. and Kakehi, K.: Determination of amino acids by pre-column fluorescence derivatization with 1-methoxycarbonylindolizine-3,5-dicarbaldehyde. *J. Chromatogr. A*, 724 (1996) 169-177.
- 2951 Okamoto, M.: (Separation of 3,4-dihydroxyphenylserine isomer). *Jpn. Kokai Tokkyo Koho JP 07,278,075 [95,278,075]* (Cl. C07C29/36), 24 Oct. 1995, Appl. 94/70,495, 8 Apr. 1994; 5 pp.; C.A., 124 (1996) 117994x.
- 2952 Ou, K., Wilkins, M.R., Yan, J.X., Gooley, A.A., Fung, Y., Sheumack, D. and Williams, K.L.: Improved high-performance liquid chromatography of amino acids derivatised with 9-fluorenylmethyl chloroformate. *J. Chromatogr. A*, 723 (1996) 219-225.
- 2953 Parent, X., Boyeau, E., Hanser, A.M. and Faller, B.: (Interference observed in creatinine determination in peritoneal dialysis fluids containing amino acids). *Ann. Biol. Clin.*, 53 (1995) 357-358; C.A., 124 (1996) 169676q.
- 2954 Peter, A., Tóth, G., Täräk, G. and Tourwé, D.: Separation of enantiomeric β-methyl amino acids and of β-methyl amino acid containing peptides. *J. Chromatogr. A*, 728 (1996) 455-465.
- 2955 Rizzo, V., Anesi, A., Montalbetti, L., Bellantoni, G., Trotti, R. and Melzi d'Eril, G.V.: Reference values of neuroactive amino acids in the cerebrospinal fluid by high-performance liquid chromatography with electrochemical and fluorescence detection. *J. Chromatogr. A*, 729 (1996) 181-188.
- 2956 Shang, S. and Wang, H.: (Reversed-phase high-performance liquid chromatographic Pico-Tag analysis of amino acids in eggs). *Sepu*, 14 (1996) 47-49; C.A., 124 (1996) 143917u.
- 2957 Simon, G., Hanak, L., Grévillot, G., Szanya, T. and Marton, G.: Amino acid separation by preparative temperature-swing chromatography with flow reversal. *J. Chromatogr. A*, 732 (1996) 1-15.
- 2958 Suido, M., Sakakibara, Y., Nakajima, H., Sakida, H. and Liu, M.-C.: Enzymic sulphation of dopa and tyrosine isomers by HepG2 human hepatoma cells: stereoselectivity and stimulation by Mn<sup>2+</sup>. *Biochem. J.*, 314 (1996) 151-158.
- 2959 Tabrizi-Fard, M.A. and Fung, H.-L.: Reversed-phase high-performance liquid chromatography method for the analysis of nitro-arginine in rat plasma and urine. *J. Chromatogr. B*, 679 (1996) 7-12.
- 2960 Tao, Z., Yu, C. and Huai, Q.: Determination of adsorption equilibrium constant and HETP of amino acids on dextran-type ion-exchangers column - a study on isocratic elution chromatography and moment method. *Funct. Polym.*, 28 (1996) 221-226; C.A., 124 (1996) 243343x.
- 2961 Venkatakrishnan, A., Abel, M.J., Campbell, R.A., Donahue, E.P., Uselton, T.C. and Flakk, P.J.: Whole blood analysis of gluconeogenic amino acids for estimation of de novo gluconeogenesis using pre-column o-phthalaldehyde derivatization and high-performance liquid chromatography. *J. Chromatogr. B*, 676 (1996) 1-6.
- 2962 Wang, S.-H., Kao, H.-Y., Chen, S.-T. and Wang, K.T.: Direct measurement of enantiomeric excess of synthetic tert-leucine with a chiral-ligand-exchange chromatography. *J. Chin. Chem. Soc. (Taipei)*, 42 (1995) 953-956; C.A., 124 (1996) 134380m.
- 2963 Zhao, Q., Sanier, F., Garreau, I., Lecoeur, C. and Piot, J.M.: Reversed-phase high-performance liquid chromatography coupled with second-order derivative spectroscopy for the quantitation of aromatic amino acids in peptides: application to hemorphins. *J. Chromatogr. A*, 723 (1996) 35-41.

For additional information see C.A.:

124 (1996) 146850c, 192910v, 218964t.

See also 2496, 2513, 2521, 2552, 2556, 2594, 2643, 2815, 2827, 2924, 2967, 3100, 3313, 3444, 3697, 3711.

#### 18b. Peptides, peptidic and proteinous hormones, growth factors

- 2964 Arthur, J.S.C. and Crawford, C.: Investigation of the interaction of m-calpain with phospholipids: calpain-phospholipid interactions. *Biochim. Biophys. Acta*, 1293 (1996) 201-206.
- 2965 Augustijns, P.F. and Borchardt, R.T.: Transport and metabolism of delta sleep-inducing peptide in cultured human intestinal epithelial cell monolayers. *Drug Metab. Disp.*, 23 Z(1995) 1372-1378.
- 2966 Barbosa, J., Sanz-Nebot, V. and Toro, I.: Solvatochromic parameter values and pH in acetonitrile-water mixtures. Optimization of mobile phase for the separation of peptides by high-performance liquid chromatography. *J. Chromatogr. A*, 725 (1996) 249-260.
- 2967 Baumann, M., Golabek, A., Lalowski, M. and Wisniewski, T.: Micropreparative gel electrophoresis of low-molecular-weight peptides: purification of highly insoluble amyloid peptide fragments. *Anal. Biochem.*, 236 (1996) 191-198.
- 2968 Bouhallab, S., Henry, G. and Boschetti, E.: Separation of small cationic bioactive peptides by strong ion-exchange chromatography. *J. Chromatogr. A*, 724 (1996) 137-145.
- 2969 Corradini, D. and Cannarsa, G.: A comparative study on the effect of hydrochloric, phosphoric, and trifluoroacetic acid in the reversed phase chromatography of angiotensins and related peptides. *J. Liq. Chromatogr.*, 18 (1995) 3919-3931.
- 2970 Corradini, D., Kalghatgi, K. and Horváth, C.: Effect of mobile phase additives on peptide retention in reversed-phase chromatography with pellicular and totally porous sorbents. *J. Chromatogr. A*, 728 (1996) 225-233.
- 2971 Edling, O., Bao, G., Feilisch, M., Unger, T. and Gohlke, P.: Moexipril, a new angiotensin-converting enzyme (ACE) inhibitor: Pharmacological characterization and comparison with enalapril. *J. Pharmacol. Exp. Ther.*, 275 (1995) 854-863.

- 2972 Gibbs, B.F., Alli, I. and Mulligan, C.N.: Simple and rapid high-performance liquid chromatographic method for the determination of aspartame and its metabolites in foods. *J. Chromatogr. A.*, 725 (1996) 372-377.
- 2973 Gurevich, L.S., Zharkovich, I.K. and Nezdoiminogo, E.L.: (Study of *Amanita verna* (Bull.: Fr.)VITT. and *A. virosa* (Fr.) Bertillon toxins by high-performance liquid chromatography). *Mikol. Fitopatol.*, 29 (1995) 32-39; *C.A.*, 124 (1996) 196264k.
- 2974 Hansen, P., Andersson, L. and Lindeberg, G.: Purification of cysteine-containing synthetic peptides via selective binding of the  $\alpha$ -amino group to immobilised  $\text{Cu}^{2+}$  and  $\text{Ni}^{2+}$  ions. *J. Chromatogr. A.*, 723 (1996) 51-59.
- 2975 Hashimoto, E., Kobayashi, N., Kubota, N., Tanaka, Y. and Yamamura, H.: Phosphorylated sites of M<sub>r</sub> 25,000 protein, a putative protein phosphatase 2A modulator, and phosphorylation of the synthetic peptide containing these sites by protein kinase C. *J. Biochem. (Tokyo)*, 119 (1996) 626-632.
- 2976 Hashimoto, T., Ohki, K. and Sakura, N.: Hydrolytic cleavage of pyroglutamyl-peptide bond. I. The susceptibility of pyroglutamyl-peptide bond to dilute hydrochloric acid. *Chem. Pharm. Bull.*, 43 (1995) 2068-2074.
- 2977 Hoijer, M.A., Melief, M.-J., Keck, W. and Hazenberg, M.P.: Purification and characterization of N-acetylmuramyl-L-alanine amidase from human plasma using monoclonal antibodies. *Biochim. Biophys. Acta*, 1289 (1996) 57-64.
- 2978 Janáky, T., Szabó, E., Baláspiri, L., Adi, B. and Penke, B.: High-performance separation methods in the analysis of a new peptide family: the galanins. *J. Chromatogr. B*, 676 (1996) 7-12.
- 2979 Janusz, M.J., Durham, S.L., Hare, C.M., Geary, J.L., Mandagere, A.K., Poole, J.C., Thompson, T.N., Xu, D., Angelastro, M.R., Burkhardt, J.P., Chen, T.M., et al.: Pharmacological evaluation of selected, orally active, peptidyl inhibitors of human neutrophil elastase. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1233-1238.
- 2980 Jonasson, P., Nilsson, J., Samuesson, E., Moks, T., Ståhl, S. and Uhlén, M.: Single-step trypsin cleavage of a fusion protein to obtain human insulin and its C-peptide. *Eur. J. Biochem.*, 236 (1996) 656-661.
- 2981 Kálmann, A., Thunecke, F., Schmidt, R., Schiller, P.W. and Horváth, C.: Isolation and identification of peptide conformers by reversed-phase high-performance liquid chromatography and NMR at low temperature. *J. Chromatogr. A*, 729 (1996) 155-171.
- 2982 Kumon, A., Kodama, H., Kondo, M., Yokoi, F. and Hiraishi, H.:  $\text{N}^{\omega}$ -Phosphoarginine phosphatase (17 kDa) and alkaline phosphatase as protein arginine phosphatases. *J. Biochem. (Tokyo)*, 119 (1996) 719-724.
- 2983 Liang, L.: The development of a continuous flow MALDI interface with an ion trap/reflection TOF mass spectrometer for the online LC/MS detection of peptides and proteins. Avail. *Univ. Microfilms Int.*, Order No.: DA9544987, 1995, 160 pp.; *C.A.*, 124 (1996) 169622u.
- 2984 Lockey, T.D. and Ourth, D.D.: Formation of pores in *Escherichia coli* cell membranes by a cecropin isolated from hemolymph of *Heliothis virescens* larvae. *Eur. J. Biochem.*, 236 (1996) 263-271.
- 2985 Mazzi, G., Fioravanzo, F. and Berti, E.: New marker of bone resorption: hydroxyproline-containing peptide. High-performance liquid chromatographic assay without hydrolysis as an alternative to hydroxyproline determination: a preliminary report. *J. Chromatogr. B*, 678 (1996) 165-172.
- 2986 Mehra, R.K., Miclat, J., Kodati, V.R., Abdullah, R., Hunter, T.C. and Mulchandani, P.: Optical spectroscopic and reverse-phase HPLC analyses of Hg(II) binding to phytochelatins. *Biochem. J.*, 314 (1996) 73-82.
- 2987 Miyanoshita, A., Hara, S., Sugiyama, M., Asaoka, A., Taniai, K., Yukihiko, E. and Yamakawa, M.: Isolation and characterization of a new member of the insect defensin family from a beetle, *Allomyrina dichotoma*. *Biochem. Biophys. Res. Commun.*, 220 (1996) 526-531.
- 2988 Moriyama, H., Anegayama, M. and Kato, Y.: Rapid separation of peptides and proteins on 2- $\mu\text{m}$  porous microspherical reversed-phase silica material. *J. Chromatogr. A*, 729 (1996) 81-85.
- 2989 Müller-Newen, G., Kähne, C., Keul, R., Hemmann, U., Müller-Esterl, W., Wijdenes, J., Brakenhoff, J.P.J., Hart, M.H.L. and Heinrich, P.C.: Purification and characterization of the soluble interleukin-6 receptor from human plasma and identification of an isoform generated through alternative splicing. *Eur. J. Biochem.*, 236 (1996) 837-842.
- 2990 Nagai, S., Okimura, K., Kaizawa, N., Ohki, K. and Kanatomo, S.: Study on surfactin, a cyclic depsipeptide. II. Synthesis of surfactin B2 produced by *Bacillus natto* KMD 2311. *Chem. Pharm. Bull.*, 44 (1996) 5-10.
- 2991 Nyberg, G., Sanderson, K., Andrén, P., Thörnwall, M., Einarsson, M., Danielson, B. and Nyberg, F.: Isolation of haemorphin-related peptides from filter membranes collected in connection with haemofiltration of human subjects. *J. Chromatogr. A*, 723 (1996) 43-49.
- 2992 Péter, A., Devadher, S., Laus, G. and Tourwé, D.: Liquid chromatography studies on the enzymatic degradation of luteinizing hormone-releasing hormone analogues with off-line identification by mass spectrometry. *J. Chromatogr. A*, 729 (1996) 137-142.
- 2993 Prechel, M.M., Orawski, A.T., Maggiora, L.L. and Simmons, W.H.: Effect of a new aminopeptidase P inhibitor, apstatin, on bradykinin degradation in the rat lung. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1136-1142.
- 2994 Ramsey, E.D., Lawrence, S.D., Games, D.E., Hewlins, M.J.E. and McDowell, M.A.: Determination of the presence of a diastereomeric impurity within a synthetic peptide by combined liquid chromatography-mass spectrometry techniques. *Anal. Commun.*, 33 (1996) 79-83; *C.A.*, 124 (1996) 242464a.
- 2995 Rhee, S.K., Chung, B.H., Choi, E.S., Sohn, J.H., Youn, D.J., Kim, M.K. and Lee, H.D.: Hirudin purification by affinity chromatography. *Ger. Offen.* DE 19,532,495 (Cl. C07K14/815), 14 Mar. 1996, KR Appl. 9,422,499, 7 Sep. 1994; 4 p.; *C.A.*, 124 (1996) 242294v.
- 2996 Rissler, K. and Engelmann, P.: Labeling of insulin with non-radioactive  $^{127}\text{I}$  and application to incorporation of radioactive  $^{125}\text{I}$  for use in receptor-binding experiments by high-performance liquid chromatography. *J. Chromatogr. B*, 679 (1996) 21-29, 387.
- 2997 Rousselle, J.-C., Massot, O., Delepine, M., Zifa, E., Rousseau, B. and Fillion, G.: Isolation and characterization of an endogenous peptide from rat brain interacting specifically with the serotonergic 1B receptor subtypes. *J. Biol. Chem.*, 271 (1996) 726-735.

2998 Silvestro, L. and Savu, S.R.: High-performance liquid chromatography/tandem mass spectrometry identification of salmon calcitonin degradation products in aqueous solution preparations. *Rapid Commun. Mass Spectrom.*, 10 (1996) 151-156; C.A., 124 (1996) 156197j.

2999 Sugimoto, T., Hayashi, T., Okita, A. and Morino, A.: Pharmacokinetics of the new thyrotropin releasing hormone analogue montireline hydrate. 3rd Communication: Identification of metabolites in rat urine. *Arzneim.-Forsch.*, 46 (1996) 127-133.

3000 Timoshchenko, A.S. and Rakitin, L.Yu.: Determination of peptide molecular weight and the number of amino acid residues by means of ion-exchange chromatography of amino acids. *Russ. J. Plant Physiol. (Transl. of Fiziol. Rast. (Moscow))*, 42 (1995) 853-857; C.A., 124 (1996) 81074f.

3001 Uchida, T., Yagi, A., Oda, Y., Nakada, Y. and Goto, S.: Instability of bovine insulin in poly(lactide- $\epsilon$ -glycolide) (PLGA). *Chem. Pharm. Bull.*, 44 (1996) 235-236.

3002 Vutla, N.B., Betageri, G.V. and Banga, A.K.: Transdermal iontophoretic delivery of enkephalin formulated in liposomes. *J. Pharm. Sci.*, 85 (1996) 5-8.

3003 Yamaki, S., Isobe, T., Okuyama, T. and Shinoda, T.: (High-performance liquid chromatography of peptides on a microspherical carbon column. (V). Chromatography under the basic conditions and regeneration of carbon packings). *Kuromatogafuji*, 16 (1995) 348-349; C.A., 124 (1996) 140096q.

3004 Yamaki, S., Isobe, T., Okuyama, T. and Shinoda, T.: High-performance liquid chromatography of peptides on a microspherical carbon column. *J. Chromatogr. A*, 729 (1996) 143-153.

3005 Yano, Y., Yokoyama, K., Tamiya, E. and Karube, I.: Direct evaluation of meat spoilage and the progress of aging using biosensors. *Anal. Chim. Acta*, 320 (1996) 269-276.

3006 Yomogida, S., Nagaoka, I. and Yamashita, T.: Purification of the 11- and 5-kDa antibacterial polypeptides from guinea pig neutrophils. *Arch. Biochem. Biophys.*, 328 (1996) 219-226.

3007 Yoshida, T.: Determination of reduced and oxidized glutathione in erythrocytes by high-performance liquid chromatography with ultraviolet absorbance detection. *J. Chromatogr. B*, 678 (1996) 157-164.

See also 2483, 2496, 2526, 2556, 2779, 2927, 2929, 2944, 2946, 2953, 2954, 3017, 3191, 3699, 3749, 3753.

#### 18c. Elucidation of structure of proteins and enzymes

3008 Becker, S., Bergman, T., Hjelmqvist, L., Jeck, R., Jörnvall, H., Leibrock, H. and Woenckhaus, C.: Photoaffinity labelling of lactate dehydrogenase from pig heart with a bifunctional NAD<sup>+</sup>-analogue. *Biochim. Biophys. Acta*, 1293 (1996) 277-283.

3009 Galzie, Z., Qasim, M.A. and Salahuddin, A.: Isolation and characterization of domain I of ovoxininhibitor. *Biochim. Biophys. Acta*, 1293 (1996) 113-121.

3010 Guzzetta, A.W. and Hancock, W.S.: Analyzing reversed phase peptide maps of recombinant human glycoproteins using LC/ES/MS. In: Hancock, W.S. (Editor), *New Methods Pept. Mapp. Charact. Proteins*, CRC, Boca Raton, 1996, pp. 181-217; C.A., 124 (1996) 111288y - a review with 111 refs.

3011 Jensen, P.E.H., Arbelaez, L.F., Shanbhag, V.P. and Stigbrand, T.: Preparation and characterization of a C-terminal fragment of pregnancy zone protein corresponding to the receptor-binding peptide from human  $\alpha_2$ -macroglobulin. *Biochim. Biophys. Acta*, 1293 (1996) 254-258.

3012 Weegels, P.L., Hamer, R.J. and Schofield, J.D.: RP-HPLC and capillary electrophoresis of subunits from glutenin isolated by SDS and Osborne fractionation. *J. Cereal Sci.*, 22 (1995) 211-224; C.A., 124 (1996) 173758k.

See also 2988, 3004, 3018, 3019, 3035, 3056, 3063, 3065, 3082, 3093, 3096, 3099, 3142, 3181, 3201, 3206, 3208.

#### 19. PROTEINS

##### 19a. General techniques

3013 Fang, F., Aguilar, M.-I. and Hearn, M.T.W.: Influence of temperature on the retention behaviour of proteins in cation-exchange chromatography. *J. Chromatogr. A*, 729 (1996) 49-66.

3014 Fang, F., Aguilar, M.-I. and Hearn, M.T.W.: Temperature-induced changes in the bandwidth behaviour of proteins separated with cation-exchange adsorbents. *J. Chromatogr. A*, 729 (1996) 67-79.

3015 Feng, W., Zhu, X., Zhang, L. and Geng, X.: Retention behaviour of proteins under conditions of column overload in hydrophobic interaction chromatography. *J. Chromatogr. A*, 729 (1996) 43-47.

3016 Gallant, S.R.: Nonlinear ion-exchange chromatography of proteins. Avail. *Univ. Microfilms Int.*, Order No. DA9608441, 1996, 372 p.; C.A., 124 (1996) 235855a.

3017 Hellberg, U., Ivarsson, J.-P. and Johansson, B.-L.: Characteristics of Superdex prep grade media for gel filtration chromatography of proteins and peptides. *Process Biochem. (Oxford)*, 31 (1996) 163-172; C.A., 124 (1996) 111361s.

3018 Hsieh, Y.L.F., Wang, H., Elicone, C., Mark, J., Martin, S.A. and Regnier, F.: Automated analytical system for the examination of protein primary structure. *Anal. Chem.*, 68 (1996) 455-462.

3019 Inglis, A.S., Reid, G.E. and Simpson, R.J.: Chemical techniques employed for the primary structural analysis of proteins and peptides. *EXS*, 73 (1995) 141-171; C.A., 124 (1996) 169551v - a review with many refs.

3020 Jacob, L.R., Heinen, A. and Britsch, L.: High performance size exclusion chromatography of proteins on Fractogel® EMD BioSEC (S) columns. *Int. Lab.*, 26, May (1996) 10H-10L.

3021 Liao, J.-L., Wang, Y. and Hjerten, S.: A novel support with artificial created recognition for the selective removal of proteins and for affinity chromatography. *Chromatographia*, 42 (1996) 259-262.

3022 Lookhart, G.L. and Bietz, J.A.: Protein extraction and sample handling techniques. In: Kruger, J.E. and Bietz, J.A. (Editors), *High-Perform. Liq. Chromatogr. Cereal Legume Proteins*, American Association of Cereal Chemists, St. Paul, 1994, pp. 51-65; C.A., 124 (1996) 50024k.

- 3023 Malhotra, O.P., Prabhakar, P. and Kayastha, A.M.: Application of short column gel permeation in the study of protein-protein interactions. *J. Biochem. Biophys. Methods*, 31 (1996) 23-30; C.A., 124 (1996) 197374h.
- 3024 Nakayama, H., Uchida, K., Shinkai, F., Shinoda, T., Okuyama, T., Seta, K. and Isobe, T.: Capillary column high-performance liquid chromatographic-electrospray ionization triple-stage quadrupole mass spectrometric analysis of proteins separated by two-dimensional polyacrylamide gel electrophoresis. Application to cerebellar protein mapping. *J. Chromatogr. A*, 730 (1996) 279-287.
- 3025 Shibusawa, Y.: Separation of proteins by high-speed counter-current chromatography. *Chem. Anal.*, 132 (1996) 385-414; C.A., 124 (1996) 111297a - a review with 41 refs.

For additional information see C.A.:  
124 (1996) 81467m.

See also 2496, 2510, 2511, 2517, 2520, 2558, 2572, 2588, 2599, 2647, 2662, 2983, 2988, 3010, 3029, 3083.

#### 19b. Proteins of cells, viruses and subcellular particles

- 3026 Baron, A.T., Errabolu, R., Dinusson, J. and Salisbury, J.L.: Centrin-based contractile fibers: chromatographic purification of centrin. *Methods Cell Biol.*, 47(Cilia and Flagella) (1995) 341-351; C.A., 124 (1996) 49819s.
- 3027 Oori, K.E., Aoyama, T., Souri, M., Jiang, L.L., Orii, K.O., Hayashi, S., Yamaguchi, S., Kondo, N., Orii, T. and Hashimoto, T.: Formation of the enzyme complex in mitochondria is required for function of trifunctional  $\beta$ -oxidation protein. *Biochem. Biophys. Res. Commun.*, 219 (1996) 773-777.
- 3028 Sanchirico, M., Tzellas, A., Fox, T.D., Conrad-Webb, H., Perlman, P.S. and Mason, T.L.: Relocation of the unusual VAR1 gene from the mitochondria to the nucleus. *Biochem. Cell Biol.*, 73 (1995) 987-995.

See also 3041.

#### 19c. Proteins synthesized by genetic manipulation, monoclonal antibodies

- 3029 Amari, J.V. and Mazsaroff, I.: Analysis of recombinant human interleukin-11 fusion protein derived from *Escherichia coli* lysate by combined size-exclusion and reversed-phase liquid chromatography. *J. Chromatogr. A*, 729 (1996) 113-124.
- 3030 Greve, K.F., Hughes, D.E., Richberg, P., Kats, and Karger, B.L.: Liquid chromatographic and capillary electrophoretic examination of intact and degraded fusion protein CTLA4Ig and kinetics of conformational transition. *J. Chromatogr. A*, 723 (1996) 273-284.
- 3031 Hamaker, K.H., Liu, J., Seely, R.J., Ladisch, C.M. and Ladisch, M.R.: Chromatography for rapid buffer exchange and refolding of secretory leukocyte protease inhibitor. *Biotechnol. Prog.*, 12 (1996) 184-189; C.A., 124 (1996) 196903t.

See also 3099, 3101.

#### 19d. Microbial and plant proteins

- 3032 Baldan, B., Navazio, L., Friso, L., Mariani, P. and Meggio, F.: Plant calreticulin is specifically and efficiently phosphorylated by protein kinase CK2. *Biochem. Biophys. Res. Commun.*, 221 (1996) 498-502.
- 3033 Cutruzzola, F., Travagliini Allocatelli, C., Brancaccio, A. and Brunori, M.: *Aplysia limacina* myoglobin cDNA cloning: an alternative mechanism of oxygen stabilization as studied by active-site mutagenesis. *Biochem. J.*, 314 (1996) 83-90.
- 3034 Hildén, P., Savolainen, K., Tyynelä, J., Vuento, M. and Kuusela, P.: Purification and characterization of a plasmin-sensitive surface protein of *Staphylococcus aureus*. *Eur. J. Biochem.*, 236 (1996) 904-910.
- 3035 Koh, M., van Driessche, G., Samyn, B., Hoff, W.D., Meyer, T.E., Cusanovich, M.A. and van Beeumen, J.J.: Sequence evidence for strong conservation of the photoactive yellow proteins from the halophilic phototrophic bacteria *Chromatium salexigens* and *Rhodospirillum salexigens*. *Biochemistry*, 35 (1996) 2526-2534.
- 3036 Macherel, D., Bourguignon, J., Forest, E., Faure, M., Sohen-Addad, C. and Douce, R.: Expression, lipoylation and structure determination of recombinant pea H-protein in *Escherichia coli*. *Eur. J. Biochem.*, 236 (1996) 27-33.
- 3037 Mioshi, S.-i. and Tu, A.T.: Phospholipase A<sub>2</sub> from *Naja naja sputatrix* venom is a muscarinic acetylcholine receptor inhibitor. *Arch. Biochem. Biophys.*, 328 (1996) 17-25.
- 3038 Morita, S., Fukase, M., Hoshino, K., Fukuda, Y., Yamaguchi, M. and Morita, Y.: Partial purification and characterization of a novel soybean protease which is inhibited by Kunitz nad Bowman-Birk trypsin inhibitors. *J. Biochem. (Tokyo)*, 119 (1996) 711-718.
- 3039 Pham, T.-N., Hayashi, K., Takano, R., Itoh, M., Eguchi, M., Shibata, H., Tanaka, T. and Hara, S.: A new family of serine protease inhibitors (Bombyx family) as established from the unique topological relation between the positions of disulfide bridges and reactive site. *J. Biochem. (Tokyo)*, 119 (1996) 428-434.
- 3040 Roobol-Boza, M. and Andersson, B.: Isolation of hydrophobic membrane proteins by perfusion chromatography - purification of photosystem II reaction centers from spinach chloroplasts. *Anal. Biochem.*, 235 (1996) 127-133.
- 3041 Roobol-Boza, M., Shochat, S., Tjus, S.E., Hagman, A., Gast, P. and Andersson, B.: Perfusion chromatography - a new procedure for very rapid isolation of integral photosynthetic membrane proteins. *Photosynth. Res.*, 46 (1995) 339-345; C.A., 124 (1996) 197288h.
- 3042 Sheldon, P.S. and Venis, M.S.: Purification and characterization of cytosolic and microsomal cyclophilins from maize (*Zea mays*). *Biochem. J.*, 315 (1996) 965-969.
- 3043 Sugiyama, H. and Natori, S.: Transmethylation reaction is essential for *Sarcophaga* lectin gene activation. *J. Biochem. (Tokyo)*, 119 (1996) 354-359.
- 3044 Talay, S.R., Grammel, M.P. and Chhatwal, G.S.: Structure of a group C streptococcal protein that binds to fibrinogen, albumin and immunoglobulin G via overlapping modules. *Biochem. J.*, 315 (1996) 577-582.

- 3045 Terabe, M., Kojima, S., Taguchi, S., Mormose, H. and Miura, K.-i.: New subtilisin-trypsin inhibitors produced by *Streptomyces*: primary structures and their relationship to other proteinase inhibitors from *Streptomyces*. *Biochim. Biophys. Acta*, 1292 (1996) 233-240.
- 3046 Van Damme, E.J.M., Barbe, A., Rouge, P., van Leuven, F. and Peumans, W.J.: Characterization and the molecular cloning of *Sambucus nigra* agglutinin V (nigrin b), a GalNAc-specific type-2 ribosome-inactivating protein from the bark of elderberry (*Sambucus nigra*). *Eur. J. Biochem.*, 237 (1996) 505-513.

See also 3012.

#### 19e. Proteins of blood, serum and blood cells

- 3047 Ando, K., Kikugawa, K. and Beppu, M.: Binding of anti-band 3 autoantibody to sialylated poly-N-acetyllactosaminyl sugar chains of band 3 glycoprotein on polyvinylidene difluoride membrane and Sepharose gel: further evidence for anti-band 3 autoantibody binding to the sugar chains of oxidized and senescent erythrocytes. *J. Biochem. (Tokyo)*, 119 (1996) 639-647.
- 3048 Duncker, B.P., Gauthier, S.Y. and Davies, P.L.: Evidence for a proprotein intermediate during maturation of Type II antifreeze protein in sea raven, *Hemitripterus americanus*. *Biochim. Biophys. Acta*, 1292 (1996) 312-316.
- 3049 Heino, P.U.: (Method for isolation and purification of the human blood plasma proteins transthyretin (TTR) and retinol-binding protein (RBP)). *Finn. FI* 95,806 (Cl. C07K1/16), 15 Dec. 1995, Appl. 93/5,161, 22 Nov. 1993; 24 pp.; C.A., 124 (1996) 170006c.
- 3050 Kamura, T., Tsuda, H., Yae, Y., Hattori, S., Ohga, S., Sjibata, Y., Kawabata, S. and Hamasaki, N.: An abnormal fibrinogen Fukuoka II (Gly- $\beta$ 15 $\rightarrow$ Cys) characterized by defective fibrin lateral association and mixed disulfide formation. *J. Biol. Chem.*, 270 (1995) 29392-29399.
- 3051 Lapuk, V.A., Chernyak, V.Ya. and Magretova, N.N.: (Some peculiarities of fragmentation of monoclonal rheumatoid immunoglobulin M with 'hot' trypsin). *Biokhimiya (Moscow)*, 61 (1996) 85-88.
- 3052 Leibl, H., Tomasits, R., Wolf, H.M., Eibl, M.M. and Mannhalter, J.W.: Method for the isolation of biologically active monomeric immunoglobulin A from a plasma fraction. *J. Chromatogr. B*, 678 (1996) 173-180.
- 3053 Suzuki, Y., Takada, J., Isaka, K., Takayama, M. and Grudzinskas, J.G.: (Isolation and purification of pregnancy-associated plasma protein A). *Nippon Sanka Fujinka Gakkai Zasshi*, 48 (1996) 17-24; C.A., 124 (1996) 197396s.
- 3054 Watanabe, S. and Sato, T.: Effects of free fatty acids on the binding of bovine and human serum albumin with steroid hormones. *Biochim. Biophys. Acta*, 1289 (1996) 385-396.

For additional information see C.A.:

124 (1996) 53714w, 197568z, 199808b.

See also 2517, 2549, 2605, 2609, 2612, 2783, 3072, 3176.

#### 19f. Structural and muscle proteins

- 3055 Bittner, K., Liszio, C., Blumberg, P., Schönher, E. and Kresse, H.: Modulation of collagen gel contraction by decorin. *Biochem. J.*, 314 (1996) 159-166.
- 3056 Diab, M., Wu, J.-J. and Eyre, D.R.: Collagen type IX from human cartilage: a structural profile of intermolecular cross-linking sites. *Biochem. J.*, 314 (1996) 327-332.
- 3057 Duval, M., Ma, X., Valet, J.-P. and Vincent, M.: Purification of developmentally regulated avian 400-kDa intermediate filament association protein. Molecular interactions with intermediate filament proteins and other cytoskeleton components. *Biochem. Cell Biol.*, 73 (1995) 671-657.
- 3058 Gong, Z., Ewart, K.V., Hu, Z., Fletcher, G.L. and Hew, C.L.: Skin antifreeze protein genes of the winter flounder, *Pleuronectes americanus*, encode distinct and active polypeptides without the secretory signal and prosequences. *J. Biol. Chem.*, 271 (1996) 4106-4112.
- 3059 Gopal, D. and Burke, M.: Myosin subfragment 1 hydrophobicity changes associated with different nucleotide-induced conformations. *Biochemistry*, 35 (1996) 506-512.
- 3060 Onodera, S., Kuboki, Y. and Tashiro, S.: Purification of a non-collagenous high-molecular-weight protein from bovine articular cartilage. *Biol. Pharm. Bull.*, 19 (1996) 11-17; C.A., 124 (1996) 197670b.
- 3061 Yamauchi, M., Chandler, G., Tanzawa, H. and Katz, E.P.: Cross-linking and the molecular packing of corneal collagen. *Biochim. Biophys. Res. Commun.*, 219 (1996) 311-315.
- 19g. Protamines, histones and other nuclear proteins
- 3062 Zardo, G., Santoro, R., D'Erme, M., Reale, A., Guidobaldi, L., Caiafa, P. and Strom, R.: Specific inhibitory effect of H1e histone somatic variant on *in vitro* DNA-methylation process. *Biochim. Biophys. Res. Commun.*, 220 (1996) 102-107.
- See also 3165.
- 19h. Chromoproteins and metalloproteins
- 3063 Bucci, E., Razynska, A., Kwnsa, H., Gryszyński, Z., Collins, J.H. and Franticelli, C.: Positive and negative cooperativities at subsequent steps of oxygenation regulate the allosteric behavior of multistate sebacylhemoglobin. *Biochemistry*, 35 (1996) 3418-3425.
- 3064 Cartel, N.J.: Purification of human brain metallothionein by organic and reversed-phase high-performance liquid chromatography under acidic conditions. *J. Chromatogr. B*, 676 (1996) 159-164.
- 3065 Fushitani, K., Higashiyama, K., Asao, M. and Hosokawa, K.: Characterization of the constituent polypeptides of the extracellular hemoglobin from *Lumbricus terrestris*: heterogeneity and discovery of a new linker chain L4. *Biochim. Biophys. Acta*, 1292 (1996) 273-280.
- 3066 Goetghebeur, M., Kermasha, S., Kensley, J. and Metche, M.: Purification and characterization of copper-m metallothionein from *Aspergillus niger* by affinity chromatography. *Biotechnol. Appl. Biochem.*, 22 (1995) 315-325; C.A., 124 (1996) 81077j.

- 3067 Iwahara, S.-i., Abe, Y. and Okazaki, T.: Identification of five embryonic hemoglobins of rat and ontogeny of their constituent globins during fetal development. *J. Biochem. (Tokyo)*, 119 (1996) 360-366.
- 3068 Jones, R.T., Shih, D.T., Fujita, T.S., Song, Y., Xiao, H., Head, C. and Kluger, R.: A doubly cross-linked human hemoglobin. Effects of cross-links between different subunits. *J. Biol. Chem.*, 271 (1996) 675-680.
- 3069 Matsunaga, T., Iwawaki, Y., Watanabe, K., Yamamoto, I., Kageyama, T. and Yoshimura, H.: Microsomal aldehyde oxygenase (MALDO): purification and characterization of a cytochrome P450 isozyme responsible for oxidation of 9-anthraldehyde to 9-anthracenecarboxylic acid in monkey liver. *J. Biochem. (Tokyo)*, 119 (1996) 617-625.
- 3070 Myers, S.R., Spinnato, J.A. and Pinorini, M.T.: Chromatographic characterization of hemoglobin benzo[a]pyrene-7,8-diol-9,10-epoxide adducts. *Fundam. Appl. Toxicol.*, 29 (1996) 94-101; *C.A.*, 124 (1996) 109499z.
- 3071 Nakayama, K., Okabe, M., Aoyagi, K., Yamanoshita, O., Okui, T., Ohyama, T. and Kasai, N.: Visualization of yellowish-orange luminescence from cuprous metallothioneins in liver of Long-Evans Cinnamon rat. *Biochim. Biophys. Acta*, 1289 (1996) 150-158.
- 3072 Papadea, C. and Cate, J.C., IV: Identification and quantification of hemoglobins A, F, S, and C by automated chromatography. *Clin. Chem. (Washington)*, 42 (1996) 57-63.
- 3073 Rohrer, J.S. and Avdalovic, N.: Separation of human serum transferrin isoforms by high-performance pellicular anion-exchange chromatography. *Protein Expression Purif.*, 7 (1996) 39-44; *C.A.*, 124 (1996) 197397t.
- 3074 Saito, S. and Hunziker, P.E.: Differential sensitivity of metallothionein-1 and -2 in liver of zinc-injected rat toward proteolysis. *Biochim. Biophys. Acta*, 1289 (1996) 65-70.
- 3075 San-Marina, S. and Nicholls, D.M.E.: Metal binding and ferritin immunoreactivity in a high molecular weight fraction from rat brain. *Biochim. Biophys. Acta*, 1310 (1996) 277-283.
- 3076 Tanaka, S. and Clark, L.A.: An assay for hemoglobin A<sub>1c</sub> testing. *Am. Clin. Lab.*, 14 (1995) 8-10; *C.A.*, 124 (1996) 169679t.
- 3077 Vassilyev, V.B., Kachurin, A.M., Rocco, G.-P., Beltramini, M., Salvato, B. and Gaitskhoki, V.S.: (A spectral study of the active center of ceruloplasmin during its copper depletion/repletion). *Biochimiya (Moscow)*, 61 (1996) 296-307.
- See also 2606.
- 19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*
- 3078 Anguita, G., Martin, R., Garcia, T., Morales, P., Haza, A.I., Gonzalez, I., Sanz, B. and Hernandez, P.E.: Immunological characterization of bovine casein fractionated by fast protein liquid chromatography (FPLC). *Milchwissenschaft*, 51 (1996) 21-25; *C.A.*, 124 (1996) 143898p.
- 3079 Chang, L.-A., Wu, P.-f. and Chang, C.C.: cDNA sequence analysis and expression of the A chain of  $\beta$ -bungarotoxin from *Bungarus multicinctus* (Taiwan banded krait). *Biochem. Biophys. Res. Commun.*, 221 (1996) 328-332.
- 3080 Cochet, S., Pesliakas, H., Kroviarski, Y., Cartron, J.P. and Bertrand, O.: Chromatography of bovine lactoserum on the immobilized metallized dye Drimarene Rubine R/K 5BL. Modulation of selectivity by metal exchange and application to purification of milk immunoglobulins. *J. Chromatogr. A*, 725 (1996) 237-247.
- 3081 Ensslin, M., Calvete, J.J., Thole, H.H., Sierralta, W.D., Adermann, K., Sanz, L. and Toepfer-Petersen, E.: Identification by affinity chromatography of boar sperm membrane-associated proteins bound to immobilized porcine zona pellucida. Mapping of the phosphorylethanolamine-binding region of spermadhesin AWN. *Biol. Chem. Hoppe-Seyler*, 376 (1995) 733-738; *C.A.*, 124 (1996) 138933s.
- 3082 Kratzschmar, J., Haendler, B., Eberspaecher, U., Roosterman, D., Donner, P. and Schleuning, W.-D.: The human cysteine-rich secretory protein (CRISP) family. Primary structure and tissue distribution of CRISP-1, CRISP-2 and CRISP-3. *Eur. J. Biochem.*, 236 (1996) 827-836.
- 3083 Kundu, S.K., Robey, W.G., Nabors, P., Lopez, M.R. and Buko, A.: Purification of commercial Coomassie brilliant blue R-250 and characterization of the chromogenic fractions. *Anal. Biochem.*, 235 (1996) 134-140.
- 3084 Sautin, Yu.Yu., Povorotny, V.Yu. and Mikosha, A.S.: (Prolactin-induced phosphorylation of proteins in adrenal cortex by protein kinase C). *Biochimiya (Moscow)*, 61 (1996) 34-41.
- 3085 Splitt, H., Mackenstedt, I. and Freitag, R.: Preparative membrane adsorber chromatography for the isolation of cow milk components. *J. Chromatogr. A*, 729 (1996) 87-97.
- 3086 Torre, M., Cohen, M.E., Corzo, N., Rodriguez, M.A. and Diez-Masa, J.C.: Perfusion liquid chromatography of whey proteins. *J. Chromatogr. A*, 729 (1996) 99-111.
- 3087 Young, M.K., Tseng, H.C., Fang, H., Liang, W. and Rothman, S.S.: Comparison of stored and secreted rat pancreatic digestive enzymes by mass spectrometry:  $\alpha$ -amylase. *Biochim. Biophys. Acta*, 1293 (1996) 63-71.
- For additional information see *C.A.*:  
124 (1996) 111721j, 111722k.
- See also 3780.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 3088 Heath, M.M., Rixon, K.C. and Harding, J.J.: Glycation-induced inactivation of malate dehydrogenase protection by aspirin and a lens molecular chaperone,  $\alpha$ -crystallin. *Biochim. Biophys. Acta*, 1315 (1996) 176-184.
- 3089 Lee, K., Rosales, J.L., Tang, D. and Wang, J.H.: Interaction of cyclin-dependent kinase 5 (CDK5) and neuronal Cdk5 activator in bovine brain. *J. Biol. Chem.*, 271 (1996) 1538-1543.
- 3090 Maruyama, E.: Biochemical characterization of mouse brain neocrin. *Biochem. J.*, 314 (1996) 895-901.
- 3091 Pant, H.C. and Veeranna: Neurofilament phosphorylation. *Biochem. Cell Biol.*, 73 (1995) 575-592.
- 3092 Pergami, P., Jaffe, H. and Safar, J.: Semipreparative chromatographic method to purify the normal cellular isoform of the prion protein in nondenatured form. *Anal. Biochem.*, 236 (1996) 63-73.
- 3093 Roquemore, E.P., Chevrier, M.R., Cotter, R.J. and Hart, G.W.: Dynamic O-GlcNAcylation of the small heat shock protein  $\alpha$ B-crystallin. *Biochemistry*, 35 (1996) 3578-3586.

See also 3024.

**19k. Proteins of neoplastic tissue and transformed cells**

See 2786.

**19l. Specific binding and receptor proteins**

- 3094 Arafa, H.M.M., Hamada, F.M.A., Elmazar, M.M.A. and Nau, H.: Fully automated determination of selective retinoic acid receptor ligands in mouse plasma and tissue by reversed-phase liquid chromatography coupled on-line with solid-phase extraction. *J. Chromatogr. A*, 729 (1996) 125-136.
- 3095 Ball, K.L. and Lane, D.P.: Human and plant proliferating cell nuclear antigens have a highly conserved binding site for the p53-inducible gene product p21<sup>WAF1</sup>. *Eur. J. Biochem.*, 237 (1996) 854-861.
- 3096 Beck, C.A., Zhang, Y., Weigel, N.L. and Edwards, D.P.: Two types of anti-progestins have distinct effects on site-specific phosphorylation of human progesterone receptor. *J. Biol. Chem.*, 271 (1996) 1209-1217.
- 3097 Castagna, L., Zarzur, J., Filipetti, M. and Landa, C.: Isolation and partial characterization of N-acetyl-D-galactosamine-binding lectins from *Epiphragmophora trenquelleonis* snail. *J. Biochem. (Tokyo)*, 119 (1996) 372-377.
- 3098 Gangopadhyay, A., Lazure, D.A., Kelly, T.M. and Thomas, P.: Purification and analysis of an 80-kDa carcinoembryonic antigen-binding protein from Kupffer cells. *Arch. Biochem. Biophys.*, 328 (1996) 151-157.

- 3099 Hashino, K., Uemori, Y., Kimizuka, F., Kato, I. and Titani, K.: A 31-kDa recombinant fibronectin cell-binding domain fragment: its binding to receptor, cell adhesive activity, and fusion proteins. *J. Biochem. (Tokyo)*, 119 (1996) 604-609.
- 3100 Mariller, C., Allain, F., Kouach, M. and Spik, G.: Evidence that human milk isolated cyclophilin B corresponds to a truncated form. *Biochim. Biophys. Acta*, 1293 (1996) 31-38.
- 3101 Schaaap, F.G., Specht, B., van der Vusse, G., Börchers, T. and Glatz, J.F.C.: One-step purification of rat heart-type fatty acid-binding protein expressed in *Escherichia coli*. *J. Chromatogr. B*, 679 (1996) 61-67.
- 3102 Schleicher, C.H., Santomé, J.A.: Purification, characterization, and partial amino acid sequencing of an amphibian liver fatty acid binding protein. *Biochem. Cell Biol.*, 74 (1996) 109-115.
- 3103 Velu, N.K., Karande, A.A. and Adiga, P.R.: Refolding of riboflavin carrier protein as probed by biochemical and immunological parameters. *Biochim. Biophys. Acta*, 1293 (1996) 231-237.

See also 2786, 3049, 3189.

**19m. Urinary proteins**

- 3104 Hayakawa, K., Terentyeva, E.A., Tanae, A., de Felice, C., Tanaka, T., Yoshikawa, K. and Yamauchi, K.: Urinary protein and albumin determinations by high-performance gel-permeation chromatography. *J. Liq. Chromatogr.*, 18 (1995) 3955-3968.

**19n. Other proteins (incl. proteinous inhibitors of enzymic activity)**

- 3105 Shinomiya, K., Inokuchi, N., Gnabre, J.N., Muto, M., Kabasawa, Y., Fales, H.M. and Ito, Y.: Countercurrent chromatographic analysis of ovalbumin obtained from various sources using the cross-axis coil planet centrifuge. *J. Chromatogr. A*, 724 (1996) 179-184.

See also 3009, 3048, 3109, 3176.

**20. ENZYMES AND ENZYME ACTIVITY ESTIMATION**

**20a. Oxidoreductases**

- 3106 Bock, A.-K., Kunow, J., Glasemacher, J. and Schönheit, P.: Catalytic properties, molecular composition and sequence alignments of pyruvate-ferredoxin oxidoreductase from the methanogenic archaeon *Methanosarcina barkeri* (strain Fusaro). *Eur. J. Biochem.*, 237 (1996) 35-44.
- 3107 Bülow, H.E., Mäbius, K., Bähr, V. and Bernhardt, R.: Molecular cloning and functional expression of the cytochrome P450 11B-hydrolases of the guinea pig. *Biochem. Biophys. Res. Commun.*, 221 (1996) 304-312.
- 3108 Chang, Y.K. and Chase, H.A.: Ion exchange purification of G6PDH from unclarified yeast cell homogenates using expanded bed adsorption. *Biotechnol. Bioeng.*, 49 (1996) 204-216; C.A., 124 (1996) 143630v.
- 3109 Chiba, M., Hensleigh, M., Nishime, J.A., Balani, S.K. and Lin, J.H.: Role of cytochrome P450 3A4 in human metabolism of MK-639, a potent human immunodeficiency virus protease inhibitor. *Drug Metab. Disp.*, 24 (1996) 307-314.
- 3110 Cohn, J.A., Tsai, L., Friguet, B. and Szweda, L.I.: Chemical characterization of a protein-4-hydroxy-2-nonenal cross-link: immunochemical detection in mitochondria exposed to oxidative stress. *Arch. Biochem. Biophys.*, 328 (1996) 158-164.
- 3111 Del Castillo-Olivares, A., Medina, M.A., Nunez de Castro, I. and Marquez, J.: Purification and characterization of a plasma membrane ferricyanide-utilizing NADH dehydrogenase from Ehrlich tumour cells. *Biochem. J.*, 314 (1996) 587-593.
- 3112 Dickinson, F.M.: The purification and some properties of the Mg<sup>2+</sup>-activated cytosolic aldehyde dehydrogenase of *Saccharomyces cerevisiae*. *Biochem. J.*, 315 (1996) 393-399.
- 3113 Fernandes, A.I. and Gregoridis, G.: Synthesis, characterization and properties of sialylated catalase. *Biochim. Biophys. Acta*, 1293 (1996) 90-96.
- 3114 Gilfoyle, D.J., Rodriguez-Lopez, J.N. and Smith, A.T.: Probing the aromatic-donor-binding site of horseradish peroxidase using site-directed mutagenesis and the suicide substrate phenylhydrazine. *Eur. J. Biochem.*, 236 (1996) 714-722.
- 3115 Hjelmqvist, L., Shafqat, J., Siddiqi, A.R. and Jörnvall, H.: Linking of isozyme and class variability patterns in the emergence of novel alcohol dehydrogenase functions. Characterization of isozymes in *Uromastix hardwickei*. *Eur. J. Biochem.*, 236 (1996) 563-570.

- 3116 Horio, T., Kubo, T. and Natori, S.: Purification and cDNA cloning of the alcohol dehydrogenase of the flesh fly *Sarcophaga peregrina*. A structural relationship between alcohol dehydrogenase and a 25 kDa protein. *Eur. J. Biochem.*, 237 (1996) 698-703.
- 3117 Iwakura, M. and Honda, S.: Stability and reversibility of thermal denaturation are greatly improved by limiting terminal flexibility of *Escherichia coli* dihydrofolate reductase. *J. Biochem. (Tokyo)*, 119 (1996) 414-420.
- 3118 List, B.M., Klatt, P., Werner, E.R., Schmid, K. and Mayer, B.: Overexpression of neuronal nitric oxide synthase in insect cells reveals requirement of haem for tetrahydrobiopterin binding. *Biochem. J.*, 315 (1996) 57-63.
- 3119 Lowe, P.N., Smith, D., Stammers, D.K., Riveros-Moreno, V., Moncada, S., Charles, I. and Boyhan, A.: Identification of the domains of neuronal nitric acid oxide synthase by limited proteolysis. *Biochem. J.*, 314 (1996) 55-62.
- 3120 Martinez, M.J., Ruiz-Duenas, F.J., Guillen, F. and Martinez, A.T.: Purified catalytic properties of two manganese peroxidase isoenzymes from *Pleurotus eryngii*. *Eur. J. Biochem.*, 237 (1996) 424-432.
- 3121 Matsuura, K., Deyashiki, Y., Bunai, Y., Ohya, I. and Hara, A.: Aldose reductase is a major reductase for isocaproaldehyde, a product of side-chain cleavage of cholesterol, in human and animal adrenal glands. *Arch. Biochem. Biophys.*, 328 (1996) 265-271.
- 3122 Mazumdar, A., Chatterjee, R., Adak, S., Ghosh, A., Mondal, C. and Banerjee, R.K.: Characterization of sheep lacrimal-gland peroxidase and its major physiological electron donor. *Biochem. J.*, 314 (1996) 413-419.
- 3123 Naud, I., Mayer, C., David, L., Breton, J., Gaillard, J. and Jouanneau, Y.: Identification of residues of *Rhodobacter capsulatus* ferredoxin I important for its interaction with nitrogenase. *Eur. J. Biochem.*, 237 (1996) 399-405.
- 3124 Ohgiya, N., Yokota, H., Takahashi, M., Komoro, S. and Yuasa, A.: Purification and properties of a new  $\beta$ -naphthoflavone inducible cytochrome P-450, aryl hydrocarbon hydroxylase from rat kidney. *Biochim. Biophys. Acta*, 1289 (1996) 122-130.
- 3125 Pannamperuma, K. and Croteau, R.: Purification and characterization of an NADPH-cytochrome P450 (cytochrome c) reductase from spearmint (*Mentha spicata*) glandular trichomes. *Arch. Biochem. Biophys.*, 329 (1996) 9-16.
- 3126 Rawat, U.B. and Rao, M.B.: Purification, kinetic characterization and involvement of tryptophan residue at the NADPH binding site of xylose reductase from *Neurospora crassa*. *Biochim. Biophys. Acta*, 1293 (1996) 222-230.
- 3127 Scheller, U., Zimmer, T., Kärgel, E. and Schunck, W.-H.: Characterization of the *n*-alkane and fatty acid hydroxylating cytochrome P450 forms 52A3 and 52A4. *Arch. Biochem. Biophys.*, 328 (1996) 245-254.
- 3128 Stephan, I., Tshisuka, B., Fetzner, S. and Lingens, F.: Quinaldine 4-oxidase from *Arthrobacter* sp. Rü61a, a versatile prokaryotic molybdenum-containing hydroxylase active towards N-containing heterocyclic compounds and aromatic aldehydes. *Eur. J. Biochem.*, 236 (1996) 155-162.
- 3129 Wang, J. and Letham, D.S.: Cytokinin oxidase - purification by affinity chromatography and activation by caffeic acid. *Plant Sci. (Shannon)*, 112 (1995) 161-166; *C.A.*, 124 (1996) 110528q.
- 3130 Yu, Y., Yamasaki, H., Kita, K. and Takamiya, S.: Purification and molecular characterization of a novel b5-type cytochrome of the parasitic nematode *Ascaris suum*. *Arch. Biochem. Biophys.*, 328 (1996) 165-172.
- See also 3008, 3069.
- 20b. Transferases (excl. E.C. 2.7.-.)
- 3131 Bohets, H.H., Nouwen, E.J., de Broe, M.E. and Dierickx, P.J.: The cytosolic glutathione S-transferase isoenzymes in the dog kidney cortex as compared with the corresponding MDCK renal cell line. *Biochim. Biophys. Acta*, 1311 (1996) 93-101.
- 3132 Danjoh, I. and Fujiyama, A.: Enzymic characterization of fission yeast farnesyl transferase. Recognition of the -CAAL motif at the C-terminus. *Eur. J. Biochem.*, 236 (1996) 847-851.
- 3133 Honke, K., Yamane, M., Ishii, A., Kobayashi, T. and Makita, A.: Purification and characterization of 3'-phosphoadenosine-5'-phosphosulfate:GalCer sulfotransferase from human renal cancer cells. *J. Biochem. (Tokyo)*, 119 (1996) 421-427.
- 3134 Horsch, M., Mayer, C. and Rast, D.M.: Stereochemical requirements of chitin synthase for ligand binding at the allosteric site for N-acetylglucosamine. *Eur. J. Biochem.*, 237 (1996) 476-482.
- 3135 Kura, T., Takahashi, Y., Takayama, T., Ban, N., Saito, T., Kuga, T. and Niitsu, Y.: Glutathione S-transferase- $\pi$  is secreted as a monomer into human plasma by platelets and tumor cells. *Biochim. Biophys. Acta*, 1292 (1996) 317-323.
- 3136 Lee, J.-F., Liaw, Y.-C. and Shaw, P.-C.: Overproduction, purification and characterization of M.EcoHK31, a bacterial methyltransferase with the polypeptides. *Biochem. J.*, 314 (1996) 321-326.
- 3137 Lenz, R. and Zenk, M.H.: Acetyl coenzyme A: salutaridinol-7-O-acetyltransferase from *Papaver somniferum* plant cell cultures. The enzyme catalyzing the formation of the baine in morphine biosynthesis. *J. Biol. Chem.*, 270 (1995) 31091-31096.
- 3138 Magnuson, B.A., Raju, R.V.S. and Sharma, R.K.: Distribution of myristoyl-CoA:protein N-myristoyl transferase activity in rabbit intestine. *Biochim. Biophys. Acta*, 1300 (1996) 119-124.
- 3139 Mainwaring, G.W., Nash, J., Davidson, M. and Green, T.: Isolation of a mouse Theta glutathione S-transferase active with methylene chloride. *Biochem. J.*, 314 (1996) 445-448.
- 3140 Paul, P., Kamisaka, Y., Marks, D.L. and Pagano, R.E.: Purification and characterization of UDP-glucose:ceramide glucosyltransferase from rat liver Golgi membranes. *J. Biol. Chem.*, 271 (1996) 2287-2293.
- 3141 Tan, K.-L. and Board, P.G.: Purification and characterization of a recombinant human Theta-class glutathione transferase (GSTT2-2). *Biochem. J.*, 315 (1996) 727-732.
- 3142 Tsai, S.-P., Wang, L.-Y., Yeh, H.-I. and Tam, M.F.: The 30 kDa protein co-purified with chick liver glutathione S-transferases is a carbonyl reductase. *Biochim. Biophys. Acta*, 1292 (1996) 289-292.
- 3143 Yeh, H.-I., Lee, J.-Y., Tsai, S.-P., Hsieh, C.-H. and Tam, M.F.: Rat kidney glutathione S-transferase 1 subunits have C-terminal truncations. *Biochem. J.*, 314 (1996) 1017-1025.
- 3144 Yuziuk, J.A., Nakagawa, H., Hasegawa, A., Kiso, M., Li, S.-C. and Li, Y.-T.: Two different sialidases, KDN-sialidase and regular sialidase in the starfish *Asterina pectinifera*. *Biochem. J.*, 315 (1996) 1041-1048.

- 20c. Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)
- 3145 Ando, S., Toku, T., Yano, T. and Inagaki, M.: Keratin 8 phosphorylation *in vitro* by cAMP-dependent protein kinase occurs within the amino- and carbonyl-terminal end domains. *Biochem. Biophys. Res. Commun.*, 221 (1996) 67-71.
- 3146 Bensen, E.S., Umphress, J.L., Traugh, J.A., Pinna, L.A. and Tuzon, P.T.: A membrane-bound protein kinase from rabbit reticulocytes is an active form of multipotential S6 kinase. *Biochim. Biophys. Acta*, 1292 (1996) 249-258.
- 3147 Chan, D.W., Mody, C.H., Ting, N.S.Y. and Less-Miller, S.P.: Purification and characterization of the double-stranded DNA-activated protein kinase, DNA-PK, from human placenta. *Biochem. Cell Biol.*, 74 (1996) 67-73.
- 3148 Clement-Chomienne, O. and Walsh, M.P.: Identification of protein kinase C isoenzymes in smooth muscle: partial purification and characterization of chicken gizzard PKC $\zeta$ . *Biochem. Cell Biol.*, 74 (1996) 51-65.
- 3149 Dutra, M.B., Silva, J.T., Mattos, D.C. and Panek, A.D.: Regulation of UDPG-pyrophosphorylase isoforms in *Saccharomyces cerevisiae* and their roles in trehalose metabolism. *Biochim. Biophys. Acta*, 1289 (1996) 261-269.
- 3150 Emanuelli, M., Raffaelli, N., Amici, A., Fanelli, M., Ruggieri, S. and Magni, G.: Three-minute high-performance liquid chromatographic assay for NMN adenylyltransferase using a 20-mm-long reversed-phase column. *J. Chromatogr. B*, 676 (1996) 13-18.
- 3151 Gehrmann, T., Vereb, G., Schmidt, M., Klix, D., Meyer, H.E., Varsányi, M. and Heilmeyer, L.M.G., Jr.: Identification of a 200 kDa polypeptide as type 3 phosphatidylinositol 4-kinase from bovine brain by partial protein and cDNA sequencing. *Biochim. Biophys. Acta*, 1311 (1996) 53-63.
- 3152 He, Q., Skog, S., Wu, C., Johansson, A. and Tribukait, B.: Existence of phosphorylated and dephosphorylated forms of cytosolic thymidine kinase (TK1). *Biochim. Biophys. Acta*, 1289 (1996) 25-30.
- 3153 Li, Z., Gillis, K.A., Hegg, L.A., Zhang, J. and Thurlow, D.L.: Effects of nucleotide substitutions within T-loop of precursor tRNAs on interaction with ATC/CTP: tRNA nucleotidyltransferase from *Escherichia coli* and yeast. *Biochem. J.*, 314 (1996) 49-53.
- 3154 Meier, R., Rouse, J., Cuenda, A., Nebreda, A.R. and Cohen, P.: Cellular stresses and cytokines activate multiple mitogen-activated-protein kinase kinase homologues in PC12 and KB cells. *Eur. J. Biochem.*, 236 (1996) 796-805.
- 3155 Szyszka, R., Bou, G. and Ballesta, J.P.G.: RAP kinase, a new enzyme phosphorylating the acidic P proteins from *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta*, 1293 (1996) 213-221.
- 3156 Walter, J., Schnölzer, M., Pyerin, W., Kinzel, V. and Kübler, D.: Induced release of cell surface protein kinase yields CK1- and CK2-like enzymes in tandem. *J. Biol. Chem.*, 271 (1996) 111-119.
- 20d. Hydrolases, acting on ester bonds (E.C. 3.1.-.-)
- 3157 Altieri, F., Matas, B., Ferraro, A. and Turano, C.: Purification of a 60-kDa protein from chicken liver associated with the internal matrix and closely related to carboxylesterases. *Eur. J. Biochem.*, 236 (1996) 806-813.
- 3158 Atkins Latham, K., Rajendran, S., Carmical, J.R., Lee, J.C. and Lloyd, R.S.: T4 endonuclease V exists in solution as a monomer and binds to target sites as a monomer. *Biochim. Biophys. Acta*, 1292 (1996) 324-334.
- 3159 Bernal, C. and Pidgeon, C.: Affinity purification of phospholipase A<sub>2</sub> on immobilized artificial membranes containing and lacking the glycerol backbone. *J. Chromatogr. A*, 731 (1996) 139-151.
- 3160 Bhanumathy, C.D. and Balasubramanian, A.S.: Evidence for a Zn<sup>2+</sup>-binding site in human serum butyrylcholinesterase. *Biochem. J.*, 315 (1996) 127-131.
- 3161 Endo, T., Fujiwara, T., Ikehara, Y. and Kobata, A.: Comparative study of the sugar chains of alkaline phosphatases purified from rat liver and rat AH-130 hepatoma cells. Occurrence of fucosylated high-mannose-type and hybrid-type sugar chains. *Eur. J. Biochem.*, 236 (1996) 579-590.
- 3162 Flores-Flores, C., Martinez-Martinez, A., Munoz-Delgado, E. and Vidal, C.J.: Conversion of acetylcholinesterase hydrophilic trimers into amphiphilic dimers and monomers. *Biochim. Biophys. Res. Commun.*, 219 (1996) 53-58.
- 3163 Haas, R., Jackson, B.C., Reinhold, B., Foster, J.D. and Rosenberry, R.L.: Glycoinositol phospholipid anchor and protein C-terminus of bovine erythrocyte acetylcholinesterase: analysis by mass spectrometry and by protein and DNA sequencing. *Biochem. J.*, 314 (1996) 817-825.
- 3164 Lee, S.B. and Rhee, S.G.: Molecular cloning, splice variants, expression, and purification of phospholipase C-84. *J. Biol. Chem.*, 271 (1996) 25-31.
- 3165 Li, W., Chen, H.Y. and Davie, J.R.: Properties of chicken erythrocyte histone deacetylase associated with the nuclear matrix. *Biochem. J.*, 314 (1996) 631-637.
- 3166 Ota, Y., Itabashi, Y. and Hasuo, M.: Measurement of positional specificity index of microbial lipases by chiral phase high-pressure liquid chromatography. *Biosci., Biotechnol., Biochem.*, 60 (1996) 145-146; C.A., 124 (1996) 139554f.
- 3167 Reardon, D.B., Wood, S.L., Brautigan, S.L., Bell, G.I., Dent, P. and Sturgill, T.W.: Activation of a protein tyrosine phosphatase and inactivation of Raf-1 by somatostatin. *Biochem. J.*, 314 (1996) 401-404.
- 3168 Somoskeöy, S., Rao, M. and Slobin, L.I.: Purification and characterization of a 5' to 3' exoribonuclease from rabbit reticulocytes that degrades capped and uncapped RNAs. *Eur. J. Biochem.*, 237 (1996) 171-179.
- 3169 Steinebach, V., Benen, J.A.E., Bader, R., Postma, P., de Vries, S. and Duine, J.A.: Cloning of the maoA gene that encodes aromatic amine oxidase of *Escherichia coli* W3350 and characterization of the overexpressed enzyme. *Eur. J. Biochem.*, 237 (1996) 584-591.
- 3170 Timmermans, M.Y.J., Reekmans, G., Teuchy, J.H.J. and Kupers, L.P.M.: Inhibition studies on calf pregastrin esterase: the enzyme has no functional thiol group. *Biochem. J.*, 314 (1996) 931-936.
- 3171 Woscholski, R., Waterfield, M.D. and Parker, P.J.: Purification and biochemical characterization of a mammalian phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase. *J. Biol. Chem.*, 270 (1995) 31001-31007.
- 3172 Wu, W., Liu, Y., Riedel, B., Wissing, J.B., Fischl, A.S. and Carman, G.M.: Purification and characterization of diacylglycerol pyrophosphate phosphatase from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 271 (1996) 1868-1876.

See also 3037, 3079.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.)*

- 3173 Berhow, Dora Smolensky, M.A.: Developmental and substrate specificity of hesperetin-7-O-glucosyltransferase activity in Citrus limon tissues using high-performance liquid chromatographic analysis. *Plant Sci. (Shanon)*, 112 (1995) 139-147; C.A., 124 (1996) 110494a.
- 3174 Cueva, R., Muñoz, M.D., Andaluz, E., Basco, R.D. and Larriba, G.: Preferential transfer of truncated oligosaccharides to the first sequon of yeast exoglycanase in *Saccharomyces cerevisiae* alg<sup>3</sup> cells. *Biochim. Biophys. Acta*, 1289 (1996) 336-342.
- 3175 Guadalupi, R., Bernard, M., Orlacchio, A., Foglietti, M.J. and Emiliani, C.: Purification and properties of human urinary  $\beta$ -D-mannosidase. *Biochim. Biophys. Acta*, 1293 (1996) 9-16.
- 3176 Lockey, T.D. and Ourth, D.D.: Purification and characterization of lysozyme from hemolymph of *Heliothis virescens* larvae. *Biochim. Biophys. Res. Commun.*, 220 (1996) 502-508.
- 3177 Srisomsap, C., Svasti, J., Surarit, R., Champattanachai, V., Sawangareetrakul, P., Boonpuan, K., Subhasitanont, P. and Chokchaichamnankit, D.: Isolation and characterization of an enzyme with  $\beta$ -glucosidase and  $\beta$ -fucosidase activities from *Dalbergia cochinchinensis* Pierre. *J. Biochem. (Tokyo)*, 119 (1996) 585-590.
- 3178 Trachuk, L.A., Shemyakina, T.M., Chestukhina, G.G. and Stepanov, V.M.: (Chitinases of *Bacillus cereus*: isolation and properties). *Biokhimiya (Moscow)*, 61 (1996) 357-368.
- 3179 Tull, D., Burgoyne, D.L., Chow, D.T., Withers, S.G. and Aebersold, R.: A mass spectrometry-based approach for probing enzyme active sites: identification of Glu 127 in *Cellulomonas fimi* exoglycanase as the residue modified by N-bromoacetyl cellobiosylamine. *Anal. Biochem.*, 234 (1996) 119-125.

For additional information see C.A.:

124 (1996) 49789j.

See also 2768.

20f. *Other hydrolases*

- 3180 Azarkan, M., Wintjens, R.T., Smolders, N., Nijs, M. and Looze, Y.: S-Pegylthiopapain, a versatile intermediate for the preparation of the fully active form of the cysteine proteinase archetype. *J. Chromatogr. A*, 724 (1996) 185-192.
- 3181 Castano, J.G., Mahillo, E., Arizti, P. and Arribas, J.: Phosphorylation of C8 and C9 subunits of the multicatalytic proteinase by casein kinase II and identification of the C8 phosphorylation sites by direct mutagenesis. *Biochemistry*, 35 (1996) 3782-3789.
- 3182 Chen, Y., Ma, Y. and Rando, R.R.: Solubilization, partial purification, and affinity labeling of the membrane-bound isoprenylated protein endoprotease. *Biochemistry*, 35 (1996) 3227-3237.
- 3183 Delaria, K. and Müller, D.: High-performance liquid chromatographic assay for trypase based on the hydrolysis of dansyl-vasoactive intestinal peptide. *Anal. Biochem.*, 236 (1996) 74-81.

- 3184 Fukusen, N. and Aoki, Y.: Purification and characterization of novel trypsin-like serine proteases from mouse spleen. *J. Biochem. (Tokyo)*, 119 (1996) 633-638.
- 3185 Geyer, H., Jacobi, I., Linder, D., Stirm, S., Bialojan, S., Strube, K.-H. and Geyer, R.: Glycosylation of recombinant ancrad from *Agrostostroma rhodostoma* after expression in mouse epithelial cells. *Eur. J. Biochem.*, 237 (1996) 113-127.
- 3186 Hashida, R., Kuwada, M., Chiba, K.-i., Horizoe, T., Shiota, H. and Nagai, Y.: A factor derived from polymorphonuclear leukocytes enhances interleukin-1-induced synovial cell collagenase and prostaglandin E<sub>2</sub> production in rats. *Eur. J. Biochem.*, 236 (1996) 517-522.
- 3187 Inoue, H., Hayashi, T., Huang, X.-P., Lu, J.-F., Athauda, S.B.P., Kong, K.-H., Yamagata, H., Ueda, S. and Takahashi, K.: Heterologous expression and site-directed mutagenesis studies on the activation mechanism and the roles of the basic residues in the prosegment of aspergillopepsinogen I. *Eur. J. Biochem.*, 237 (1996) 719-725.
- 3188 Itoh, M., Masuda, K., Ito, Y., Akizawa, T., Yoshioka, M., Imai, K., Okada, Y., Sato, H. and Seiki, M.: Purification and refolding of recombinant human proMMP-7 (pro-matrilysin) expressed in *Escherichia coli* and its characterization. *J. Biochem. (Tokyo)*, 119 (1996) 667-673.
- 3189 Kania, M.A., Demartino, G.N., Baumeister, W. and Goldberg, A.L.: The proteasome subunit C2, contains an important site for binding of the PA28 (11S) activator. *Eur. J. Biochem.*, 236 (1996) 510-516.
- 3190 Kuehn, L. and Dahlmann, B.: Proteasome activator PA28 and its interaction with 20 S proteasomes. *Arch. Biochem. Biophys.*, 329 (1996) 87-96.
- 3191 Lamango, N.S., Sajid, M. and Isaac, R.E.: The endopeptidase activity and the activation by Cl<sup>-</sup> of angiotensin-converting enzyme from the housefly, *Musca domestica*. *Biochem. J.*, 314 (1996) 639-646.
- 3192 Larcher, G., Cimon, B., Symoens, F., Tronchin, G., Chabasse, D. and Bouchara, J.-P.: A 33 kDa serine protease from *Scedosporium apiospermum*. *Biochem. J.*, 315 (1996) 119-126.
- 3193 Matsuki, H., Fujimoto, N., Iwata, K., Knäuper, V., Okada, Y. and Hayakawa, T.: A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 8 (neutrophil collagenase) using monoclonal antibodies. *Clin. Chim. Acta*, 244 (1996) 129-143.
- 3194 Meyer, S.L., Bozyczko-Coyne, D., Mallya, S.K., Spais, C.M., Bihovsky, R., Kawooya, J.K., Lang, D.M., Scott, R.W. and Siman, R.: Biologically active monomeric and heterodimeric recombinant human calpain I produced using the baculovirus expression system. *Biochem. J.*, 314 (1996) 511-519.
- 3195 Ohuchi, E., Azumano, I., Yoshida, S., Iwata, K. and Okada, Y.: A one-step sandwich immunoassay for human matrix metalloproteinase 7 (matrilysin) using monoclonal antibodies. *Clin. Chim. Acta*, 244 (1996) 181-198.
- 3196 Orriss, G.L., Runswick, M.J., Collinson, I.R., Miroux, B., Fearnley, I.M., Skehel, J.M. and Walker, J.E.: The  $\delta$ - and  $\epsilon$ -subunits of bovine F1-ATPase interact to form a heterodimeric subcomplex. *Biochem. J.*, 314 (1996) 695-700.
- 3197 Richards, G.P., Zintz, C., Chao, J. and Chao, L.: Purification and characterization of salivary kallikrein from an insectivore (*Scalopus aquaticus*): substrate specificities, immunoreactivity, and kinetic analyses. *Arch. Biochem. Biophys.*, 329 (1996) 104-112.

- 3198 Tanaka, T. and Yada, R.Y.: Expression of soluble cloned porcine pepsinogen 1 in *Escherichia coli*. *Biochem. J.*, 315 (1996) 443-446.
- 3199 Tang, C. and Capaldi, R.A.: Characterization of the interface between  $\gamma$  and  $\epsilon$  subunits of *Escherichia coli* F<sub>1</sub>-ATPase. *J. Biol. Chem.*, 271 (1996) 3018-3024.
- 3200 Tran, C.M. and Farley, R.A.: Photoaffinity labeling of the active site of the Na<sup>+</sup>/K<sup>+</sup>-ATPase with 4-azido-2-nitrophenyl phosphate. *Biochemistry*, 35 (1996) 47-55.
- 3201 Ueshima, S., Okada, K. and Matsuo, O.: Stabilization of plasmin by lysine derivatives. *Clin. Chim. Acta*, 245 (1996) 7-18.
- 3202 Wijffels, G., Fitzgerald, C., Gough, J., Riding, G., Elvin, C., Kemp, D. and Willadsen, P.: Cloning and characterization of angiotensin-converting enzyme from the dipteran species, *Haematobia irritans exigua*, and its expression in the maturing male reproductive system. *Eur. J. Biochem.*, 237 (1996) 414-423.
- 3203 Yoshikawa, K., Hayakawa, K., Katsumata, N., Tanaka, T., Kimura, T. and Yamauchi, K.: High-performance liquid chromatographic determination of lipoamidase (lipoyl-X hydrolase) activity with a novel substrate, lipoyl-6-aminoquinoline. *J. Chromatogr. B*, 679 (1996) 41-47.

See also 2977, 3204, 3221.

#### 20g. Lyases

- 3204 Aceto, A., Dragani, B., Allocati, N., Masulli, M., Petruzzelli, R. and Di Ilio, C.: Purification and characterization of a heterodimeric 23/20-kDa proteolytic fragment of bacterial glutathione transferase B1-1. *Arch. Biochem. Biophys.*, 328 (1996) 302-308.
- 3205 Ashmarina, L.I., Robert, M.-F., Elsliger, M.-A. and Mitchell, G.A.: Characterization of the hydroxymethylglutaryl-CoA-lyase precursor, a protein targeted to peroxisomes and mitochondria. *Biochem. J.*, 315 (1996) 71-75.
- 3206 Okuda, H., Nagata, S. and Misono, H.: A novel phenylserine dehydratase from *Pseudomonas pickettii* PS22: purification, characterization, and sequence of its phosphopyridoxyl peptide. *J. Biochem. (Tokyo)*, 119 (1996) 690-696.
- 3207 Rolland, N., Ruffet, M.-L., Job, D., Douce, R. and Droux, M.: Spinach chloroplast O-acetylserine (thiol)-lyase exhibits two catalytically non-equivalent pyridoxal-5'-phosphate-containing active sites. *Eur. J. Biochem.*, 236 (1996) 272-282.
- 3208 Sasisekharan, R., Venkataraman, G., Godavarti, R., Ernst, S., Cooney, C.L. and Langer, R.: Heparinase I from *Flavobacterium heparinum*. Mapping and characterization of the heparin binding domain. *J. Biol. Chem.*, 271 (1996) 3124-3131.
- 3209 Schmid, M., Berg, M., Hilbi, H. and Dimroth, P.: Malonate decarboxylase of *Klebsiella pneumoniae* catalyses the turnover of acetyl and malonyl thioester residues on a coenzyme-A-like prosthetic group. *Eur. J. Biochem.*, 237 (1996) 221-228.
- 3210 Stoltz, M. and Dördemann, D.: Purification, metal cofactor, N-terminal sequence and subunit composition of 5-aminolevulinic acid dehydratase from the unicellular green alga *Scenedesmus obliquus*, mutant C-2A'. *Eur. J. Biochem.*, 236 (1996) 600-609.
- 3211 Tsujimura, M., Odaka, M., Nagashima, S., Yohda, M. and Endo, I.: Photoreactive nitrile hydratase: the photoreaction site is located on the  $\alpha$  subunit. *J. Biochem. (Tokyo)*, 119 (1996) 407-413.

#### 20h. Isomerases

- 3212 Giacomelli, S., Leone, M.-G., Grima, J., Silvestrini, B. and Cheng, C.Y.: Astrocytes synthesize and secrete prostaglandin D synthetase *in vitro*. *Biochim. Biophys. Acta*, 1310 (1996) 269-276.
- 3213 Kristo, P., Saarelainen, R., Fagerström, R., Aho, S. and Korhola, M.: Protein purification, and cloning and characterization of the cDNA and gene for xylose isomerase of barley. *Eur. J. Biochem.*, 237 (1996) 240-246.
- 3214 Prucha, M., Peterseim, A., Timmis, K.N. and Pieper, D.H.: Muconolactone isomerase of the 3-oxoadipate pathway catalyzes dechlorination of 5-chloro-substituted muconolactones. *Eur. J. Biochem.*, 237 (1996) 350-356.

#### 20i. Ligases

- 3215 Purcarete, C., Simon, V., Prieur, D. and Herve, G.: Purification and characterization of carbamoyl-phosphate synthetase from the deep-sea hyperthermophilic archaeabacterium *Pyrococcus abyssi*. *Eur. J. Biochem.*, 236 (1996) 189-199.
- 3216 Uchida, Y., Kondo, N., Orii, T. and Hashimoto, T.: Purification and properties of rat liver peroxisomal very-long-chain acyl-CoA synthetase. *J. Biochem. (Tokyo)*, 119 (1996) 565-571.

#### 20j. Complex mixtures and incompletely identified enzymes

- 3217 Carlsson, H., Ljung, S. and Bülow, L.: Physical and kinetic effects on introduction of various linker regions in  $\beta$ -galactosidase/galactose dehydrogenase fusion enzymes. *Biochim. Biophys. Acta*, 1293 (1996) 154-160.
- 3218 Hermida-Matsumoto, M.L., Chock, P.B., Curran, T. and Yang, D.C.H.: Ubiquitylation of transcription factors c-Jun and c-Fos using reconstituted ubiquitylating enzymes. *J. Biol. Chem.*, 271 (1996) 4930-4936.
- 3219 Yamada, D., Sekiya, F. and Morita, T.: Isolation and characterization of carinactivase, a novel prothrombin activator in *Echis carinatus* venom with a unique catalytic mechanism. *J. Biol. Chem.*, 271 (1996) 5200-5207.

#### 21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS

##### 21a. Purines, pyrimidines, nucleosides, nucleotides

- 3220 Bofill, M., Fairbanks, L.D., Ruckemann, K., Lipman, M. and Simmonds, H.A.: T-Lymphocytes from AIDS patients are unable to synthesize ribonucleotides *de novo* in response to mitogenic stimulation. Impaired pyrimidine responses are already evident at early stages of HIV-1 infection. *J. Biol. Chem.*, 270 (1995) 29690-29697.
- 3221 Borovsky, D., Janssen, I., Vanden Broeck, J., Huybrechts, R., Verhaert, P., de Bondt, H.L., Bylemans, D. and de Loof, A.: Molecular sequencing and modeling of *Neobellieria bullata* trypsin. Evidence for translation control by *Neobellieria* trypsin-modulating oostatic factor. *Eur. J. Biochem.*, 237 (1996) 279-287.
- 3222 Bory, C., Chantin, C. and Boulieu, R.: Comparison of capillary electrophoretic and liquid chromatographic determination of hypoxanthine and xanthine for the diagnosis of xanthinuria. *J. Chromatogr. A*, 730 (1996) 329-331.

- 3223 Bothner, B., Chatman, K., Sarkisian, M. and Siuzdak, G.: Liquid chromatography mass spectrometry of antisense oligonucleotides. *Bioorg. Med. Chem. Lett.*, 5 (1995) 2863-2868; C.A., 124 (1996) 202879h.
- 3224 Boulieu, R.: Sample preparation for the determination of purine nucleotide analogues in tissues. *J. Chromatogr. A*, 729 (1996) 197-200.
- 3225 Chantin, C., Bonin, B., Boulieu, R. and Bory, C.: Liquid-chromatographic study of purine metabolism abnormalities in purine nucleoside phosphorylase deficiency. *Clin. Chem. (Washington)*, 42 (1996) 326-328.
- 3226 Cherwinski, H.M., Cohn, R.G., Cheung, P., Webster, D.J., Xu, Y.-Z., Caufield, J.P., Young, J.M., Nakano, G. and Ransom, J.T.: The immunosuppressant leflunomide inhibits lymphocyte proliferation by inhibiting pyrimidine biosynthesis. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1043-1049.
- 3227 Child, K.F., Ning, X.-H. and Bolling, S.F.: Simultaneous detection of nucleotides, nucleosides and oxidative metabolites in myocardial biopsies. *J. Chromatogr. B*, 678 (1996) 181-186.
- 3228 Cho, S.-C.: Facile analysis of ribonucleotides in D,L- $\alpha$ -difluoromethylornithine-treated mouse lymphoid cells by high-performance anion-exchange column chromatography. *J. Chromatogr. B*, 678 (1996) 344-348.
- 3229 Edgecombe, M., McLennan, A.G. and Fisher, M.J.: Characterization of the binding of diadenosine 5',5''-P<sub>1</sub>P<sub>4</sub>-tetraphosphate (Ap<sub>4</sub>A) to rat liver cell membranes. *Biochem. J.*, 314 (1996) 687-693.
- 3230 Eisenberg, E.J. and Cundy, K.C.: High-performance liquid chromatographic determination of cytosine-containing compounds by precolumn fluorescence derivatization with phenacyl bromide: application to antiviral nucleosides and nucleotides. *J. Chromatogr. B*, 679 (1996) 119-127.
- 3231 Fritsch, M., Geilen, C.C. and Reutter, W.: Determination of cytidine 5'-monophospho-N-acetylneurameric acid pool size in cell culture scale using high-performance anion-exchange chromatography with pulsed amperometric detection. *J. Chromatogr. A*, 727 (1996) 223-230.
- 3232 Gerstner, J.A.: Economics of displacement chromatography - a case study: Purification of oligonucleotides. *BioPharm (Eugene)*, 9 (1996) 30-35; C.A., 124 (1996) 242275q.
- 3233 Green, A.P., Burzynski, J., Helveston, N.M., Prior, G.M., Wunner, W.H. and Thompson, J.A.: HPLC purification of synthetic oligodeoxyribonucleotides containing base- and backbone-modified sequences. *BioTechniques*, 19 (1995) 836-841; C.A., 124 (1996) 81062a.
- 3234 Gualix, J., Abal, M., Pintor, J., Garcia-Carmona, F. and Miras-Portugal, M.T.: Nucleotide vesicular transporter of bovine chromaffin granules. Evidence for a mnemonic regulation. *J. Biol. Chem.*, 271 (1996) 1957-1965.
- 3235 Huszar, E., Barat, E. and Kollai, M.: Isocratic high-performance liquid chromatographic determination of plasma adenosine. *Chromatographia*, 42 (1996) 318-322.
- 3236 Kazoka, H.: (High-performance liquid chromatography of flurafur, 5-fluorouracil and uracil on silica gel). *Khim.-Farm. Zh.*, 29 (1995) 49-51; C.A., 124 (1996) 212232f.
- 3237 Keuzenkamp-Jansen, C.W., Leegwater, P.A.J., de Abreu, R.A., Lambooy, M.A.H., Bokkerink, J.P.M. and Trijbels, J.M.F.: Thiopurine methyltransferase: a review and a clinical pilot study. *J. Chromatogr. B*, 678 (1996) 15-22.
- 3238 Leeds, J.M., Graham, M.J., Truong, L.A. and Cummins, L.L.: Quantitation of phosphorothioate oligonucleotides in human plasma. *Anal. Biochem.*, 235 (1996) 36-43.
- 3239 Liu, S.Y. and Rosazza, J.P.N.: Ion-pair reversed-phase HPLC identification of sugar nucleotides in cell free extracts of *Streptomyces griseus*. *J. Liq. Chromatogr.*, 18 (1995) 4081-4095.
- 3240 Nath, R.G., Ocano, J.E. and Chung, F.: Detection of 1,N<sup>2</sup>-propanodeoxyguanosine adducte as potential endogenous DNA lesions in rodent and human tissues. *Cancer Res.*, 56 (1996) 452-456.
- 3241 Reck, F.: Synthesis of uridine-5-propylamine derivatives and their use in affinity chromatography of N-acetylglucosaminyl-transferase I and II. *Carbohydr. Res.*, 276 (1995) 321-335; C.A., 124 (1996) 146695f.
- 3242 Roy, S.K., Gupta, E. and Dolan, M.E.: Pharmacokinetics of O<sup>6</sup>-benzylguanine in rats and its metabolism by rat liver microsomes. *Drug Metab. Disp.*, 23 (1995) 1294-1299.
- 3243 Schmidt, J.G., Nielsen, P.E. and Orgel, L.E.: Separation of "uncharged" oligodeoxynucleotide analogs by anion-exchange chromatography at high pH. *Anal. Biochem.*, 235 (1996) 239-241.
- 3244 Seki, T., Yamaji, K., Orita, Y., Moriguchi, S. and Shinoda, A.: Simultaneous determination of uric acid and creatinine in biological fluids by column-switching liquid chromatography with ultraviolet detection. *J. Chromatogr. A*, 730 (1996) 139-145.
- 3245 Villanueva, G.R., Monte, M.J., Barbero, E.R., Serrano, M.A. and Marin, J.J.G.: Evidence for dual effect of bile acids on thymidine anabolism and catabolism by the regenerating rat liver. *Biochim. Biophys. Acta*, 1289 (1996) 136-144.
- 3246 Villegas, S. and Brunton, L.L.: Separation of cyclic GMP and cyclic AMP. *Anal. Biochem.*, 235 (1996) 102-103.
- 3247 Vu, C.Q., Lu, P., Chen, C. and Jacobson, M.K.: 2'-Phospho-cyclic ADP-ribose, a calcium-mobilizing agent derived from NADP. *J. Biol. Chem.*, 271 (1996) 4747-4754.
- 3248 Wheatley, J.B., Lytle, M.H., Hocker, M.D. and Schmidt, D.E., Jr.: Salt-induced immobilizations of DNA oligonucleotides on an epoxide-activated high-performance liquid chromatographic affinity support. *J. Chromatogr. A*, 726 (1996) 77-90.
- 3249 Witters, E., Roef, L., Newton, R.P., van Dongen, W., Esmans, E.L. and van Onckelen, H.A.: Quantitation of cyclic nucleotides in biological samples by negative electrospray tandem mass spectrometry coupled to ion suppression liquid chromatography. *Rapid Commun. Mass Spectrom.*, 10 (1996) 225-231; C.A., 124 (1996) 197494x.
- For additional information see C.A.:  
124 (1996) 56573k, 66753h, 140099t.
- See also 2541, 2555, 2812, 3150, 3480, 3508, 3643, 3645, 3649, 3650, 3668, 3686.
- 21b. *Nucleic acids, RNA*
- 3250 Hayashi, N., Suzuki, M., Titani, K. and Taniguchi, H.: Structural studies of tRNAs by capillary high performance liquid chromatography/electrospray mass spectrometry. *Nucleic Acids Symp. Ser.*, 34(Twentysecond Symposium on Nucleic Acids Chemistry, 1995) (1995) 153-154; C.A., 124 (1996) 197677j.

3251 Imai, T., Sumi, Y., Hatakeyama, M., Fujimoto, K., Kawaguchi, H., Hayashida, N., Shiozaki, K., Terada, K., Yajima, H., et al.: Selective isolation of DNA or RNA using single-stranded DNA affinity latex particles. *J. Colloid Interface Sci.*, 177 (1996) 245-249; C.A., 124 (1996) 111372w.

3252 Yamazaki, Y., Tomizawa, H., Miyanaga, A., Ishikawa, O., Nakatani, S. and Moriyama, H.: Separation of nucleic acids by high-performance ion-exchange chromatography. *Nucleic Acids Symp. Ser.*, 34(Twentysecond Symposium on Nucleic Acids Chemistry, 1995) (1995) 149-150; C.A., 124 (1996) 197386p.

#### 21c. Nucleic acids, DNA

3253 Chaplen, F.W.R., Fahl, W.E. and Cameron, D.C.: Detection of methylglyoxal as a degradation product of DNA and nucleic acid components treated with strong acid. *Anal. Biochem.*, 236 (1996) 262-269.

3254 Hnatowich, D.J., Mardirossian, G., Fogarasi, M., Sano, T., Smith, C.L., Cantor, C.R., Ruszkowski, M. and Winnard, P., Jr.: Comparative properties of a technetium-99m-labeled single-stranded natural DNA and a phosphorothioate derivative *in vitro* and in mice. *J. Pharmacol. Exp. Ther.*, 276 (1996) 326-334.

3255 Jemilohun, P.F., Clark, C.W. and Archibald, E.R.: Characterization of bacterial cell membrane attachment sites of plasmid R6K. *Biochem. Biophys. Res. Commun.*, 221 (1996) 186-192.

3256 Mauthé, R.J., Marsch, G.A. and Turteltaub, K.W.: Improved high-performance liquid chromatography analysis of  $^{32}\text{P}$ -postlabeled 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine-DNA adducts using in-like precolumn purification. *J. Chromatogr. B*, 679 (1996) 91-101.

3257 Namsaraev, E.A., Lanzov, V.A. and Akhmedov, A.T.: Partial purification and characterization of two types of homologous DNA pairing activity from rat testis nuclei. *Biochim. Biophys. Acta*, 1305 (1996) 172-180.

3258 Pham, T.T., Chilapagari, S. and Suarez, A.R.: Preparation of pure plasmid or cosmid DNA using single-strand affinity matrix and gel-filtration spin columns. *BioTechniques*, 20 (1996) 492-497; C.A., 124 (1996) 195180t.

3259 Wittmeyer, J. and Formosa, T.: Identifying DNA replication complex components using protein affinity chromatography. *Methods Enzymol.*, 262(DNA Replication) (1995) 415-430; C.A., 124 (1996) 77501a - a review with 37 refs.

For additional information see C.A.:  
124 (1996) 50192p.

See also 3251, 3252.

#### 21d. Structural studies on RNA and RNA mapping

See 3250.

#### 21e. Structural studies on DNA and DNA mapping

For additional information see C.A.:  
124 (1996) 197401q.

#### 22. ALKALOIDS

3260 Chuang, W.-C., Chu, C.-Y. and Sheu, S.-J.: Determination of the alkaloids in *Evodiae Fructus* by high-performance liquid chromatography. *J. Chromatogr. A*, 727 (1996) 317-323.

3261 Cooper, R.A., Bowers, R.J., Beckham, C.I. and Huxtable, R.J.: Preparative separation of pyrrolizidine alkaloids by high-speed counter-current chromatography. *J. Chromatogr. A*, 732 (1996) 43-50.

3262 Lee, D.-U., Eibler, E., Mayer, K.K. and Wiegreb, W.: Stereochemical elucidation of the reaction products of  $\alpha$ -narcotine with ethyl chloroformate. *Chem. Pharm. Bull.*, 43 (1995) 1995-1997.

3263 Pedersen, S.E. and Papineni, R.V.L.: Interaction of d-tubocurarine analogs with the torpedo nicotinic acetylcholine receptor. Methylation and stereoisomerization affect site-selective competitive binding and binding to the noncompetitive site. *J. Biol. Chem.*, 270 (1995) 31141-31150.

3264 Rasmussen, B.B. and Brøsen, K.: Determination of theophylline and its metabolites in human urine and plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 676 (1996) 169-174.

3265 Schreiber-Deturmeny, E. and Bruguerolle, B.: Simultaneous high-performance liquid chromatographic determination of caffeine and theophylline for routine drug monitoring in human plasma. *J. Chromatogr. B*, 677 (1996) 305-312.

3266 Suzuki, H., Harada, M., Satake, M. and Mursito, B.: Determination of arecoline and arecaidine in areca by reversed-phase ion-pair high-performance liquid chromatography. *Nat. Med.*, 49 (1995) 303-307; C.A., 124 (1996) 66721w.

3267 Velcheva, M.P., Werner, C. and Hesse, M.: Reversed-phase high-performance liquid chromatographic separation of epimeric alkaloid N-oxides from *Thalictrum simplex* L. *J. Chromatogr. A*, 730 (1996) 63-67.

3268 Verma, R.K., Uniyal, G.C., Singh, S.P., Sharma, J.R. and Gupta, M.M.: Reverse-phase high-performance liquid chromatography of sennosides in *Cassia angustifolia*. *Phytochem. Anal.*, 7 (1996) 73-75; C.A., 124 (1996) 242447x.

3269 Wickremesinhe, E.R.M. and Arteca, R.N.: HPLC separation of cephalotaxine, harringtonine and homoharringtonine from callus and root cultures of *Cephalotaxus harringtonia*. *J. Liq. Chromatogr.*, 19 (1996) 889-897.

3270 Wu, S.-C., Chen, F.-S. and Li, J.-H.: Stability-indicating HPLC method for the determination of morphine and atropine in morphine-atropine injections. *Chin. Pharm. J. (Taipei)*, 47 (1995) 457-468; C.A., 124 (1996) 185807p.

For additional information see C.A.:  
124 (1996) 66711t, 66723y, 185731j.

See also 2622, 3480, 3560, 3729, 3734, 3737.

## 23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. *Porphyrins and other pyrroles*

- 3271 Johnson, C.G., Caron, S. and Blough, N.V.: Combined liquid chromatography/mass spectrometry of the radical adducts of a fluorescamine-derivatized nitroxide. *Anal. Chem.*, 68 (1996) 867-872.
- 3272 Miksik, I., Holan, V. and Deyl, Z.: Avian eggshell pigments and their variability. *Comp. Biochem. Physiol.*, 113B (1996) 607-612.

23c. *Indole derivatives and plant hormones (gibberellins)*

- 3273 Kele, M. and Ohmacht, R.: Determination of serotonin released from coffee wax by liquid chromatography. *J. Chromatogr. A*, 730 (1996) 59-62.
- 3274 Markova, E. and Fojtakova, A.: High-performance liquid chromatographic profiling of indolylacryloylglycine and its possible precursors in body fluids. *J. Chromatogr. A*, 730 (1996) 133-137.
- 3275 Pei, X.-F., Yu, Q.-s., Lu, B., Greig, N.H. and Brossi, A.: Practical total synthesis of physostigmines and of phenserines: a synopsis. *Heterocycles*, 42 (1996) 229-236; *C.A.*, 124 (1996) 202704x.
- 3276 Smith, D.J., Appleton, M.L., Carlson, J.R. and Yost, G.S.: Identification of  $\beta$ -glucuronidase-resistant diastereomeric glucuronides of 3-hydroxy-3-methoxindole formed during 3-methylindole metabolism in goats. *Drug Metab. Disp.*, 24 (1996) 119-125.
- 3277 Thornton-Manning, J., Appleton, M.L., Gonzalez, F.J. and Yost, G.S.: Metabolism of 3-methylindole by vaccinia-expressed P450 enzymes: correlation of 3-methyleneindolenine formation and protein-binding. *J. Pharmacol. Exp. Ther.*, 276 (1996) 21-29.
- 3278 Tsai, T.-H. and Chen, C.-F.: In vivo microdialysis determination of collagen-induced serotonin release in rat blood. *J. Chromatogr. A*, 730 (1996) 121-123.
- 3279 Tuomi, T. and Rosenqvist, H.: Detection of abscisic, gibberellic and indole-3-acetic acid from plants and microbes. *Plant Physiol. Biochem. (Paris)*, 33 (1995) 725-734; *C.A.*, 124 (1996) 111636k.

See also 2648, 3283, 3285, 3704.

23d. *Pyridine derivatives*

- 3280 Czajkowska, T., Jaroniec, M. and Buszewski, B.: Retention of pyridinecarboxylic acids on monomeric and polymeric alkylamide phases. *J. Chromatogr. A*, 728 (1996) 213-224.
- 3281 Lee, D.W. and Kim, H.S.: Retention behavior of oxadiazoles, carboxamides and quinolones in reversed-phase liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 467-473; *C.A.*, 124 (1996) 75131f.
- 3282 Skinner, T.L., Meurer, K.A. and Yokel, R.A.: HPLC quantitation of a very hydrophilic 3-hydroxypyridin-4-one chelator using a simple separation procedure and the baseline file subtraction method. *J. Chromatogr. Sci.*, 34 (1996) 52-57.

See also 3285, 3324, 3537, 3568.

23e. *Other N-heterocyclic compounds*

- 3283 Galceran, M.T., Moyano, E., Puignou, L. and Pais, P.: Determination of heterocyclic amines by pneumatically assisted electrospray liquid chromatography-mass spectrometry. *J. Chromatogr. A*, 730 (1996) 185-194.
- 3284 Imai, J., Yoshida, I., Murayama, K., Sakai, Y., Shimizu, H., Sumi, T., Iguchi, T., Kawai, M. and Yamaguchi, S.: Determination of the 8-methyl ether of xanthurenic acid in human serum by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. B*, 679 (1996) 204-207.
- 3285 Richling, E., Herderich, M. and Schreier, P.: High performance liquid chromatography-electrospray tandem mass spectrometry (HPLC-ESI-MS-MS) for the analysis of heterocyclic aromatic amines (HAA). *Chromatographia*, 42 (1996) 7-11.
- 3286 Sorensen, L.K. and Petersen, A.K.: Determination of benzimidazoles in milk by HPLC. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 306-307; *C.A.*, 124 (1996) 143872a.

See also 2629, 2636, 2688, 2914, 2953, 3244, 3291, 3484, 3585, 3622, 3670, 3708.

## 24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

- 3287 Allenmark, S., Andersson, C. and Widell, P.: Chromatographic resolution of some cyclic sulphoximides derived from prochiral and chiral sulphoxides. *Chirality*, 7 (1995) 541-546; *C.A.*, 124 (1996) 97359p.
- 3288 Andersson, J.T. and Sielex, K.: Dimethylbenzothiophenes and methylidibenzothiophenes in crude oils from different sources. *J. High Resolut. Chromatogr.*, 19 (1996) 49-53.
- 3289 Annable, P.L.: Determination of alkyl sulfonic acids in pharmaceuticals by ion chromatography. *J. Chromatogr. A*, 724 (1996) 199-206.
- 3290 Bona, M., Christen, M.-O. and Burgot, J.-L.: On the lipophilicity of the 1,2-dithiole-3-thione nucleus. *Chem. Pharm. Bull.*, 43 (1995) 1894-1896.
- 3291 Cox, C. and Hee, S.S.Q.: Synthesis of 2-thiothiazolidine-4-carboxylic acid and its chromatography in rat and human urine. *J. Chromatogr. B*, 679 (1996) 85-90.
- 3292 Elfakir, C. and Dreux, M.: Simultaneous analysis of intact and desulfated glucosinolates with a porous graphitized carbon column. *J. Chromatogr. A*, 727 (1996) 71-82.
- 3293 Fischer, M.B., Lawton, M.P., Atta-Adjei, E., Philpot, R.M. and Rettie, A.E.: Selectivity of flavin-containing monooxygenase 5 for the (S)-sulfoxidation of short-chain aralkyl sulfides. *Drug Metab. Disp.*, 23 (1995) 1431-1433.
- 3294 Full, G. and Schreier, P.: Covalent chromatography as efficient "clean-up" technique in the analysis of 8-mercapto-p-menthan-3-one. *Colloq.-Int. Natl. Rech. Agron.*, 75(Bioflavour 95) (1995) 113-116; *C.A.*, 124 (1996) 143865a.
- 3295 Okada, T.: Design of chromatographic and electrophoretic separation based on ion-pair formation of aromatic disulfonates with polyammonium ions. *Anal. Chem.*, 68 (1996) 1158-1163.

- 3296 Roussel, C., Popescu, C. and Shibata, T.: A factorial design approach for studying the HPLC chiral separation of N-arylthiazoline-2-(thio)one atropisomers on CHIRALCEL OJ. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 77-89; C.A., 124 (1996) 75122d.
- 3297 Tanaka, M., Yamazaki, H. and Hakusui, H.: Direct HPLC separation of enantiomers of pantoprazole and other benzimidazole sulphoxides using cellulose-based chiral stationary phases in reversed-phase mode. *Chirality*, 7 (1995) 612-615; C.A., 124 (1996) 212226g.

For additional information see C.A.:  
124 (1996) 111441t.

See also 2480, 2636, 2816, 3319.

## 25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

- 3298 Brearley, C.A. and Hanke, D.E.: Inositol phosphates in the duckweed *Spirodella polyrhiza* L. *Biochem. J.*, 314 (1996) 215-225.
- 3299 Brearley, C.A. and Hanke, D.E.: Metabolic evidence for the order of addition of individual phosphate esters to the myo-inositol moiety of inositol hexakisphosphate in the duckweed *Spirodella polyrhiza* L. *Biochem. J.*, 314 (1996) 225-233.
- 3300 Bringezu, F., Dobner, B., Stritzel, R., Elsner, B. and Nuhn, P.: Preparative separation of the diastereomers of methyl branched-chain phosphatidylcholines. *J. Chromatogr. A*, 724 (1996) 367-372.
- 3301 Matthaeus, B., Loesing, R. and Fiebig, H.J.: Determination of inositol phosphates IP3-IP6 in rapeseed and rapeseed meal by a HPLC method. Part 1. Method. *Fett Wiss. Technol.*, 97 (1995) 289-291; C.A., 124 (1996) 53938x.
- 3302 Schieffer, G.W.: Preliminary examination of a new post-column photolysis-molybdate reaction detection system for the determination of organophosphorus compounds by high performance liquid chromatography. *Instrum. Sci. Technol.*, 23 (1995) 255-263; C.A., 124 (1996) 105105v.
- 3303 Van Dijken, P., Bergsma, J.C.T., Hiemstra, H.S., de Vries, B., van der Kaay, J. and van Haastert, P.J.M.: *Dictyostelium discoideum* contains three inositol monophosphatase activities with different substrate specificities and sensitivities to lithium. *Biochem. J.*, 314 (1996) 491-495.

See also 2626, 2776, 2800, 2842, 2846, 2856, 2858, 3150, 3326.

## 26. ORGANOMETALLIC AND RELATED COMPOUNDS

### 26a. Organometallic compounds

- 3304 Falter, R. and Schöler, H.F.: A new pyrrolidinedithiocarbamate screening method for the determination of methyl-mercury relation in hair samples by HPLC-UV-PCO-CVAAS. *Fresenius J. Anal. Chem.*, 354 (1996) 492-493.

- 3305 Gailer, J. and Irgolic, K.J.: Retention behavior of arsenobetaine, arsenocholine, trimethylarsine oxide and tetramethylarsonium iodide on a styrene-divinylbenzene column with benzenesulfonates as ion-pairing reagents. *J. Chromatogr. A*, 730 (1996) 219-229.

- 3306 Ni, Z.: (Coupled chromatography-atomic spectrometry for the determination of trace organometallic compounds in environmental and biological samples). *Guangpuxue Yu Guangpu Fenxi*, 15 (1995) 87-106; C.A., 124 (1996) 163746r - a review with 88 refs.

- 3307 Pan, T.M., Wu, S.J., Huang, K.C., Wang, R.T., Wang, C.T. and Pan, T.C.: Determination of methyl mercury in fish and shellfish. *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 364-366; C.A., 124 (1996) 53934t.

- 3308 Park, M.K., Park, J.H., Han, S.B. and Park, I.H.: Analysis of organic germanium, Ge-132. *Anal. Sci. Technol.*, 8 (1995) 371-374; C.A., 124 (1996) 127272z.

See also 2938, 2986.

### 26c. Coordination compounds

- 3309 Careri, M., Mangia, A., Manini, P., Predieri, G., Licandro, E. and Papagni, A.: Characterization of chromium-carbene complexes by high-performance liquid chromatography-mass spectrometry with particle beam interface. *J. Chromatogr. A*, 730 (1996) 231-240.

- 3310 Gracia Bagur, M., Gázquez Evangelista, D. and Sanchez-Viñas, M.: Simultaneous determination of vanadium and molybdenum as N-benzoyl-N-phenylhydroxamate complexes by combining solvent extraction and liquid chromatography. *J. Chromatogr. A*, 730 (1996) 241-246.

- 3311 Liu, Q., Zhao, T.J. and Cheng, J.: Determination of platinum metals complexes with 2-(6-methyl-2-benzothiazolylazo)-5-diethylaminophenol by reversed-phase high performance liquid chromatography. *Microchim. Acta*, 122 (1996) 27-33; C.A., 124 (1996) 249055b.

- 3312 Lucena, J.J., Barak, P. and Hernández-Apaolaza, L.: Isocratic ion-pair high-performance liquid chromatographic method for the determination of various iron(III) chelates. *J. Chromatogr. A*, 727 (1996) 253-264.

- 3313 Nowack, B., Kari, F.G., Hilger, S.U. and Sigg, L.: Determination of dissolved and adsorbed EDTA species in waters and sediments by HPLC. *Anal. Chem.*, 68 (1996) 561-566.

For additional information see C.A.:  
124 (1996) 192571k, 218833z, 248652g.

See also 3695, 3805.

## 27. VITAMINS AND VARIOUS ANIMAL GROWTH FACTORS (NON-PEPTIDIC)

- 3314 Abidi, S.L. and Mounts, T.L.: Normal phase high-performance liquid chromatography of tocopherols on polar phases. *J. Liq. Chromatogr.*, 19 (1996) 509-520.

- 3315 Badwe, N., Garg, A., Dutta, P.K. and Sethi, P.D.: Estimation of vitamin E in pharmaceutical dosage forms using HPLC. *East Pharm.*, 38 (1995) 133-134; C.A., 124 (1996) 66715x.

- 3316 Balz, M., Schulte, E. and Thier, H.-P.: Simultaneous determination of retinol esters and tocopherol in foods using nitro-column HPLC. *Fett Wiss. Technol.*, 97 (1995) 445-448; C.A., 124 (1996) 200425h.
- 3317 Benmoussa, A., Delaurent, C., Lacout, J.-L., Loiseau, P.R. and Mikou, M.: Determination of cholecalciferol and related substances by calcium phosphate hydroxyapatite and calcium phosphate fluoroapatite high-performance liquid chromatography. *J. Chromatogr. A*, 731 (1996) 153-160.
- 3318 Careri, M., Mangia, A., Manini, P. and Taboni, N.: Determination of phylloquinone (vitamin K<sub>1</sub>) by high-performance liquid chromatography with UV detection and with particle beam - mass spectrometry. *Fresenius J. Anal. Chem.*, 355 (1996) 48-56.
- 3319 Han, D., Handelman, G.J. and Packer, L.: Analysis of reduced and oxidized lipoic acid in biological samples by high-performance liquid chromatography. *Methods Enzymol.*, 251(Biothiols, Part A) (1995) 315-325; C.A., 124 (1996) 49817q.
- 3320 Haruyama, M., Kosugi, E. and Okaya, Y.: (Determination of pantothenyl alcohol, pantothenyl ethyl ether in cosmetics by high performance liquid chromatography). *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 458-462; C.A., 124 (1996) 185123n.
- 3321 Hodam, J.R. and Creek, K.E.: Uptake and metabolism of [<sup>3</sup>H]retinoic acid delivered to human foreskin keratinocytes either bound to serum albumin or added directly to the culture medium. *Biochim. Biophys. Acta*, 1311 (1996) 102-110.
- 3322 Hu, Y., Mizoguchi, T., Kurimoto, Y. and Koyama, Y.: Isolation by high-pressure liquid chromatography of cis-trans isomers of β-apo-12'-carotenal and determination of their configurations by <sup>1</sup>H NMR spectroscopy. *Spectrochim. Acta, Part 1*, 51A (1995) 1949-1958; C.A., 124 (1996) 192880k.
- 3323 Jezequel-Cuer, M., le Moel, G., Mounie, J., Peynet, J., le Bizec, C., Vernet, M.H., Artur, Y., Laschi-Loquerie, A. and Troupel, S.: (Determination of α-tocopherol in serum or plasma by high performance liquid chromatography: optimization of the extraction procedure). *Ann. Biol. Clin.*, 53 (1995) 343-352; C.A., 124 (1996) 169676p.
- 3324 Kirsanov, N.B., Sizova, I.A., Tiaglov, B.V. and Yanenko, A.S.: (Quantitative analysis of nicotinamide and nicotinic acid in the reaction mixtures). *Biotehnologiya*, (1995) 41-43; C.A., 124 (1996) 169685s.
- 3325 Nyambaka, H. and Ryley, J.: An isocratic reversed-phase HPLC separation of the stereoisomers of the provitamin A carotenoids (α- and β-carotene) in dark green vegetables. *Food Chem.*, 55 (1995) 63-72; C.A., 124 (1996) 53990h.
- 3326 Ozone, K., Ueno, S. and Ishizaki, M.: (High-performance liquid chromatographic determination of riboflavin sodium phosphate based on its conversion to riboflavin with acid phosphase). *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 358-363; C.A., 124 (1996) 127209j.
- 3327 Raisig, A., Bartley, G., Scolnik, P. and Sandmann, G.: Purification in an active state and properties of the 3-step phytoene desaturase from *Rhodobacter capsulatus* overexpressed in *Escherichia coli*. *J. Biochem. (Tokyo)*, 119 (1996) 559-564.
- 3328 Sharpless, K.E., Brown Thomas, J., Sander, L.C. and Wise, S.A.: Liquid chromatographic determination of carotenoids in human serum using an engineered C<sub>30</sub> and a C<sub>18</sub> stationary phase. *J. Chromatogr. B*, 678 (1996) 187-195.
- 3329 Shimada, K., Saito, Y. and Hirose, M.: Retention behavior of vitamin D N-acetylglucosaminides during high-performance liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 837-842.
- 3330 Shirley, M.A., Bennani, Y.L., Boehm, M.F., Breau, A.P., Pathirana, C. and Ulm, E.H.: Oxidative and reductive metabolism of 9-cis-retinoic acid in the rat. Identification of 13,14-dihydro-9-cis-retinoic acid and its taurine conjugate. *Drug Metab. Disp.*, 24 (1996) 293-302.
- 3331 Sliva, M.G. and Sanders, J.K.: Vitamin D in infant formula and enteral products by liquid chromatography: collaborative study. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 73-80.
- 3332 Staprans, I., Rapp, J.H., Pan, X.-M. and Feingold, K.R.: Oxidized lipids in the diet are incorporated by the liver into very low density lipoprotein in rats. *J. Lipid Res.*, 37 (1996) 420-430.
- 3333 Teissier, E., Walters-Laporte, E., Duhem, C., Luc, G., Fruchart, J.-C. and Duriez, P.: Rapid quantification of α-tocopherol in plasma and low- and high-density lipoproteins. *Clin. Chem. (Washington)*, 42 (1996) 430-435.
- 3334 Vahteristo, L.T., Ollilainen, V., Koivistoinen, P.E. and Varo, P.: Improvements in the analysis of reduced folate monoglutamates and folic acid in food by high-performance liquid chromatography. *J. Agric. Food Chem.*, 44 (1996) 477-482; C.A., 124 (1996) 115649q.
- 3335 Yamanaka, K., Matsuoka, M. and Banno, K.: Determination of thiamine in dried yeast by high-performance liquid chromatography using a clean-up column of CM-cellulose. *J. Chromatogr. A*, 726 (1996) 237-240.

See also 2511, 2760, 3442.

## 28. ANTIBIOTICS

- 3336 Anderson, K.L., Moats, W.A., Rushing, J.E., Wesen, D.P. and Papich, M.G.: Ampicillin and amoxicillin residue detection in milk, using microbial receptor assay (Charm II) and liquid chromatography methods, after extra-label administration of the drugs to lactating cows. *Am. J. Vet. Res.*, 57 (1996) 73-78; C.A., 124 (1996) 173804x.
- 3337 Baker, G.H., Blanchflower, S.E., Dorgan, E.J.J., Everett, J.R., Manger, B.R., Reading, C.R., Readshaw, S.A. and Shelley, P.: Further novel milbemycin antibiotics from *Streptomyces* sp. E225. Formation, isolation and structure elucidation. *J. Antibiot.*, 49 (1996) 272-280.
- 3338 Bu, Y., Zhao, T., Hou, Y., Zhu, T., Liu, H. and Sun, Y.: (Determination of oxytetracycline residue in shrimps by solid-phase extraction and high-performance liquid chromatography). *Beijing Daxue Xuebao, Ziran Kexueban*, 31 (1995) 415-421; C.A., 124 (1996) 53952x.
- 3339 Bygrave, J., Rose, M. and Tarbin, J.: A comparison of screening and quantitative test methods for the determination of penicillin in milk. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 266-268; C.A., 124 (1996) 85174m.
- 3340 Carson, M.C. and Breslyn, W.: Simultaneous determination of multiple tetracycline residues in milk by metal chelate affinity chromatography: collaborative study. *J. Assoc. Off. Anal. Chem.*, 79 (1996) 29-42.

- 3341 Castaneda Penalvo, G., Julien, E. and Fabre, H.: Cross validation of capillary electrophoresis and high-performance liquid chromatography for cefotaxime and related impurities. *Chromatographia*, 42 (1996) 159-164.
- 3342 Croubels, S. and van Peteghem, C.: Residues of tetracyclines in milk: evaluation of a commercial enzyme-immunological screening test by high-performance liquid chromatography. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 288-292; C.A., 124 (1996) 85176p.
- 3343 Del Nozal, M.J., Bernal, J.L., Pampliega, A., Marinero, P., López, M.I. and Coco, R.: High-performance liquid chromatographic determination of vancomycin in rabbit serum, vitreous and aqueous humour after intravitreal injection of the drug. *J. Chromatogr. A*, 727 (1996) 231-238.
- 3344 Dudrikova, E., Sokol, J., Nagy, J., Cabadaj, R. and Burdova, O.: Detection of oxytetracycline residues in bovine milk by HPLC. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 293-296; C.A., 124 (1996) 85177q.
- 3345 Fedeniuk, R.W., Ramamurthy, S. and McCurdy, A.R.: Application of reversed-phase liquid chromatography and prepacked C<sub>18</sub> cartridges for the analysis of oxytetracycline and related compounds. *J. Chromatogr. B*, 677 (1996) 291-297.
- 3346 Frigerio, E., Magi, N., Benecchi, A., Pianezzola, E. and Strolin Benedetti, M.: Determination of FCE 26644, a new polysulphonated derivative of distamycin A, in monkey plasma by reversed-phase ion-pair high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. A*, 729 (1996) 237-242.
- 3347 Harada, K.-I.: Separation of complicated antibiotics by high-speed countercurrent chromatography. *Chem. Anal.*, 132 (1996) 301-319; C.A., 124 (1996) 111296z - a review with 40 refs.
- 3348 Hassan, E.E., Darwish, I.A., El-Khordagui, L.K. and Khalil, S.A.: Direct HPLC assay of amoxycillin in urine: application to bioequivalence testing. *Alexandria J. Pharm. Sci.*, 9 (1995, Pub. 1995) 231-236; C.A., 124 (1996) 105432z.
- 3349 Hsu, M.-C., Chung, H.-C. and Lin, Y.-S.: Liquid chromatographic determination of cephalexin preparations: interlaboratory validation. *J. Chromatogr. A*, 727 (1996) 239-244.
- 3350 Jelinska, A. and Zajac, M.: Effect of amino acids and amines on the stability of cefoperazone. *Pharmazie*, 51 (1996) 162-164.
- 3351 Kajimura, Y. and Kaneda, M.: Fusaricidin A, a new depsipeptide antibiotic produced by *Bacillus polymyxa* KT-8. Taxonomy, fermentation, isolation, structural elucidation and biological activity. *J. Antibiot.*, 49 (1996) 129-135.
- 3352 Kamano, Y., Zhang, H.-P., Hino, A., Yoshida, M., Pettit, G.R., Herald, C.L. and Itokawa, H.: An improved source of bryostatin 10, *Bugula maritima* from the gulf of Aomori, Japan. *J. Natural Prod.*, 58 (1995) 1868-1875.
- 3353 Kamigiri, K., Suzuki, Y., Shibasaki, M., Morioka, M., Suzuki, K.-i., Tokunaga, T., Setiawan, B. and Rantiamodjo, R.M.: Kalimantanacin A, B and C, novel antibiotics from Alcaligenes sp. YL-026325. I. Taxonomy, fermentation, isolation and biological properties. *J. Antibiot.*, 49 (1996) 136-139.
- 3354 Kees, F., Wollenhofer, M., Bröhl, K. and Grobecker, H.: Bioavailability of cefpodoxime proxetil with co-administered acetyl cysteine. *Arzneim.-Forsch.*, 46 (1996) 435-438.
- 3355 Kim, W.-G., Kim, J.-P., Kim, C.-J., Lee, K.-H. and Yoo, I.-D.: Benzastatins A, B, C, and D: new free radical scavengers from *Streptomyces nitrosporeus* 30643. I. Taxonomy, fermentation, isolation, physico-chemical properties and biological activities. *J. Antibiot.*, 49 (1996) 20-25.
- 3356 Kobayashi, K., Sato, K., Mizuno, Y. and Katsumata, Y.: Capillary high-performance liquid chromatography-fast atom bombardment mass spectrometry of 24 cephal antibiotics. *J. Chromatogr. B*, 677 (1996) 275-290.
- 3357 Kramer, R.A., Herben, P.J., van Rens, P. and Aerts, M.M.L.: Confirmation of residues in milk of the  $\beta$ -lactam Inmec using coupled SPE-HPLC-microbiological assay. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 299; C.A., 124 (1996) 173676n.
- 3358 Krauwinkel, W.J.J. and Volkers-Kamermans, N.J.: Determination of penicillin-V in human plasma by high-performance liquid chromatography and solid-phase extraction. *J. Chromatogr. B*, 679 (1996) 129-135.
- 3359 Kwon, K., Il, Kim, M.H., Park, J.W. and Lee, C.H.: Pharmacokinetics of two cyclosporine formulations using FPIA and HPLC assay in volunteers. *Arch. Pharmacal Res.*, 18 (1995) 385-390; C.A., 124 (1996) 134727e.
- 3360 Lauser, G. and Bergner-Lang, G.: (Determination of aminoglycoside antibiotics by HPLC based on an automated precolumn derivatization with 9-fluorenylmethylchloroformate). *Dtsch. Lebenschsm.-Rundsch.*, 91 (1995) 390-396; C.A., 124 (1996) 173677h.
- 3361 Lee, H.S., Shim, H.O. and Yu, S.R.: High-performance liquid chromatographic determination of meropenem in rat plasma using column switching. *Chromatographia*, 42 (1996) 405-408.
- 3362 Lihl, S., Rehorek, A. and Petz, M.: High-performance liquid chromatographic determination of penicillins by means of automated solid-phase extraction and photochemical degradation with electrochemical detection. *J. Chromatogr. A*, 729 (1996) 229-235.
- 3363 Muratake, H., Abe, I. and Natsume, M.: Preparation of alkyl-substituted indoles in the benzene portion. Part 14. Synthesis of (+/-)-duocarmycin SA, natural (+)-duocarmycin SA and non-natural (-)-duocarmycin SA. *Chem. Pharm. Bull.*, 44 (1996) 67-79.
- 3364 Muth, P., Metz, R., Beck, H., Bolten, W.W. and Vergin, H.: Improved high-performance liquid chromatographic determination of amoxicillin in human plasma by means of column switching. *J. Chromatogr. A*, 729 (1996) 259-266.
- 3365 Negi, S., Yamanaka, M., Komatsu, Y., Tsuruoka, A., Tsukada, I. and Minami, N.: Studies on the stability of  $\Delta^2$  and  $\Delta^3$  cepham esters. I. Marked difference in stability between  $\Delta^2$  and  $\Delta^3$  cepham prodrug esters and application to the preparation of key intermediates for oral cepham synthesis. *Chem. Pharm. Bull.*, 43 (1995) 1998-2000.
- 3366 Negi, S., Yamanaka, M., Komatsu, Y., Tsuruoka, A., Tsukada, I. and Minami, N.: Studies on the stability of  $\Delta^2$  and  $\Delta^3$  cepham esters. II. Comparative stability studies of  $\Delta^2$  and  $\Delta^3$  cepahms at various pHs and the degradation process of  $\Delta^2$  cepham esters. *Chem. Pharm. Bull.*, 43 (1995) 1866-1871.

- 3367 Nogusa, H., Yano, T., Okuno, S., Hamana, H. and Inoue, K.: Synthesis of carboxymethylpullulan-peptide-doxorubicin conjugates and their properties. *Chem. Pharm. Bull.*, 43 (1995) 1931-1936.
- 3368 Ogawa, T., Ando, K., Tanaka, T., Uosaki, Y. and Matsuda, Y.: RES-1149-1 and -2, novel non-peptidic endothelin type B receptor antagonists produced by *Aspergillus* sp. I. Taxonomy of producing strain, fermentation, isolation, and physico-chemical and biological properties. *J. Antibiot.*, 49 (1996) 1-5.
- 3369 Ritieni, A., Fogliano, J., Nanno, D., Randazzo, G., Altomare, C., Perrone, G., Bottalico, A., Maddau, L. and Marras, F.: Paracelsin E, a new peltaibol from *Trichoderma saturnisporum*. *J. Natural Prod.*, 58 (1995) 1745-1748.
- 3370 Roggo, B.E., Petersen, F., Sills, M., Roesel, J.L., Moerker, T. and Peter, H.H.: Novel spirodihydrobenzofuranlactams as antagonists of endothelin and as inhibitor of HIV-1 protease produced by *Stachylobotrys* sp. I. Fermentation, isolation and biological activity. *J. Antibiot.*, 49 (1996) 13-19.
- 3371 Salom, D. and Abad, C.: Chromatographic purification and characterization of synthetic tryptophan-substituted gramicidin A analogues. *J. Chromatogr. A*, 725 (1996) 315-322.
- 3372 Sokol, J., Dudrikova, E., Cabadaj, R. and Matisova, E.: Solid-phase extraction and liquid-chromatography determination of oxytetracycline antibiotics in milk. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 308-309; *C.A.*, 124 (1996) 143873b.
- 3373 Song, D. and Au, J.L.-S.: Direct injection isocratic high-performance liquid chromatographic analysis of mitomycin C in plasma. *J. Chromatogr. B*, 676 (1996) 165-168.
- 3374 Sorensen, L.K. and Hansen, P.: Determination of spiramycin and tylosin in milk by HPLC. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 302-303; *C.A.*, 124 (1996) 143870y.
- 3375 Stubbings, G., Tarbin, J.A. and Shearer, G.: On-line metal chelate affinity chromatography clean-up for the high-performance liquid chromatographic determination of tetracycline antibiotics in animal tissues. *J. Chromatogr. B*, 679 (1996) 137-145.
- 3376 Takatsu, T., Takahashi, M., Kawase, Y., Enokita, R., Okazaki, T., Matsukawa, H., Ogawa, K., Sakaida, Y., Kagaasaki, T., Kinoshita, T., Nakajima, M. et al.: A-72363 A-1, A-2 and C, novel heparanase inhibitors from *Streptomyces nobilis* SANK 60192. I. Taxonomy of producing organism, fermentation, isolation, and structure elucidation. *J. Antibiot.*, 49 (1996) 54-60.
- 3377 Tolhurst, T.A., Negrusz, A., Libelt, B., Woods, E.F. and Levine, B.S.: Determination of ampicillin in New Zealand white rabbit plasma using column switching technique and HPLC. *Chromatographia*, 42 (1996) 223-226.
- 3378 Tomoda, H., Tabata, N., Yang, D.-J., Namatame, I., Tanaka, H. and Omura, S.: Pyripyropenes, novel ACAT inhibitors produced by *Aspergillus fumigatus* IV. Structure elucidation of pyripyropenes M to R. *J. Antibiot.*, 49 (1996) 292-298.
- 3379 Tsuda, M., Muraoka, Y., Nagai, M. and Takeuchi, T.: Poststatin, a new inhibitor of prolyl endopeptidase. IV. The chemical synthesis of poststatin. *J. Antibiot.*, 49 (1996) 287-291.
- 3380 Tsuda, M., Muraoka, Y., Nagai, M., Aoyagi, T. and Takeuchi, T.: Poststatin, a new inhibitor of prolyl endopeptidase. III. Optical resolution of 3-amino-2-hydroxyvaleric acid and absolute configuration of poststatin. *J. Antibiot.*, 49 (1996) 281-286.
- 3381 Tsuruoka, T., Fukuyasu, H., Ishii, M., Usui, T., Shibahara, S. and Inouye, S.: Inhibition of mouse tumor metastasis with nojirimycin-related compounds. *J. Antibiot.*, 49 (1996) 155-161.
- 3382 Tyaglov, B.V., Zvenigorodsky, V.I., Sizova, I.A., Polanuer, B.M. and Zhdanov, V.G.: Quantitative chromatographic determination of virginiamycin antibiotics. *J. Planar Chromatogr.-Mod. TLC*, 8 (1995) 374-377; *C.A.*, 124 (1996) 127211d.
- 3383 Ubukata, M., Morita, T.-i., Urumoto, M. and Osada, H.: Sparomycins A1 and A2, new inducers of the flat reversion of NRK cells transformed by temperature sensitive Rous sarcoma virus. II. Isolation, physico-chemical properties and structure elucidation. *J. Antibiot.*, 49 (1996) 65-70.
- 3384 Ueno, R. and Aoki, T.: A simplified determination of oxytetracycline in serum and muscle of cultured eel by high-performance liquid chromatography. *Gyobyo Kenkyu*, 30 (1995) 239-240; *C.A.*, 124 (1996) 85132w.
- 3385 Unglaub, W. and Maertlbauer, E.: (Detection of tetracycline in urine, serum, and tissue fluid from slaughter animals by enzyme immunoassay). *Fleischwirtschaft*, 75 (1995) 1231-1233; *C.A.*, 124 (1996) 85143a.
- 3386 Van Rens, P. and Aerts, M.M.L.: Comparison of HPLC and microbiological assay for the determination of bacitracin in milk. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 297-298; *C.A.*, 124 (1996) 173766m.
- 3387 Villani, P., Regazzi, M.B., Maserati, R., Viale, P., Alberici, F. and Giacchino, R.: Clinical and pharmacokinetic evaluation of a new lipid-based delivery system of amphotericin B in AIDS patients. *Arzneim.-Forsch.*, 46 (1996) 445-449.
- 3388 Warr, G.A., Veitch, J.A., Walsh, A.W., Hesler, G.A., Pirnik, D.M., Leet, J.E., Lin, P.-F.M., Medina, I.A., McBrien, K.D., Foreza, S., Clark, J.M. and Lam, K.S.: BMS-182123, A fungal metabolite that inhibits the production of TNF- $\alpha$  by macrophages and monocytes. *J. Antibiot.*, 49 (1996) 234-240.
- 3389 Wood, M.J., Lund, R. and Beavan, M.: Stability of vancomycin in plastic syringes measured by high-performance liquid chromatography. *J. Clin. Pharm. Ther.*, 20 (1995) 319-325; *C.A.*, 124 (1996) 211823n.
- 3390 Zajac, M., Muszalska, I., Nogowska, M. and Siwek, J.: Kinetics of hydrolysis of ceftriaxone in aqueous solution. *Pharmazie*, 51 (1996) 227-231.
- 3391 Zarzuelo, A., López, F.G., Santos, M. and Lanao, J.M.: Determination of piperacillin in biological samples by HPLC. *J. Liq. Chromatogr.*, 19 (1996) 601-610.

See also 2658, 3620, 3714.

## 29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

## 29a. General techniques

- 3392 Bicchi, C., Balbo, C., Binello, A. and D'Amato, A.: HPLC-UV determination of pesticide residues at 0.01 ppm in apple and pear pulp used for baby food. *J. High Resolut. Chromatogr.*, 19 (1996) 105-110.
- 3393 Fucci, G., Ciaravolo, S. and Mazza, G.: (Determination of carbendazim and thiabendazole residues in fruits and vegetables by high-performance liquid chromatography). *Ind. Aliment. (Pinerolo)*, 34 (1995) 1013-1015; C.A., 124 (1996) 200434k.
- 3394 Hernandez, F., Serrano, R., Miralles, M.C. and Font, N.: Gas and liquid chromatography and enzyme linked immuno sorbent assay in pesticide monitoring of surface water from the western mediterranean (comunidad Valencia, Spain). *Chromatographia*, 42 (1996) 151-158.
- 3395 Jaeger, C., Uveira, M.N. and Marty, P.: (Determination method of pesticides in wine: application to the determination of iprodione, folpel, vinclozoline, and procymidone). *Rev. Fr. Oenol.*, 155(Cah. Sci.) (1995) 5-8; C.A., 124 (1996) 173820z.
- 3396 Jimena Garcia, J.A., Giménez Plaza, J. and Cano Pavón, J.M.: Determination of active components in insecticide formulations by liquid chromatography and resolution of overlapped peaks by multivariate analysis and derivative spectrophotometry. *Anal. Chim. Acta*, 321 (1996) 273-278.
- 3397 Pichon, V., Chen, L., Durand, N., le Goffic, F. and Hennion, M.-C.: Selective trace enrichment on immunosorbents for the multiresidue analysis of phenylurea and triazine pesticides. *J. Chromatogr. A*, 725 (1996) 107-119.
- 3398 Pichon, V., Guenu, S., Dupas, S. and Hennion, E.M.-C.: (Multi-residue pesticide analysis in waters by liquid-solid extraction and liquid chromatography). *Spectra Anal.*, 24 (1995) 25-31; C.A., 124 (1996) 96912b.
- 3399 Scaroni, I., Bovolenta, A. and Previati, M.P.: (Evaluation of a multiresidue method for the determination of pesticide residues and some of their metabolites in wine. Note 2). *Ind. Bevande*, 24 (1995) 593-598; C.A., 124 (1996) 200433j.
- 3400 Vera-Avila, L.E., Padilla, P.C., Hernandez, M.G. and Meraz, J.L.L.: On-line preconcentration, cleanup and high-performance liquid chromatographic determination of chlorophenoxy acid herbicides in water. *J. Chromatogr. A*, 731 (1996) 115-122.
- 3401 Volmer, D.A., Vollmer, D.L. and Wilkes, J.G.: Multiresidue analysis of pesticides by electrospray LC-MS and LC-MS-MS. *LC-GC*, 14 (1996) 216-224; C.A., 124 (1996) 196401c.

For additional information see C.A.:  
124 (1996) 47785d.

See also 2481, 2502, 3735, 3740.

## 29b. Chlorinated insecticides

- 3402 Grainger, J., McClure, P.C., Liu, Z., Botero, B., Sirimanne, S., Patterson, D.G., Jr., Sewer, M., Gillyard, C., Kimata, K., et al.: Isomer identification of chlorinated dibenz-p-dioxins by orthogonal spectroscopic and chromatographic techniques. *Chemosphere*, 32 (1996) 13-23; C.A., 124 (1996) 163838x.

- 3403 Pillai, U.A., Ziegler, T.L., Wang, D.X., Kattnig, M.J., McClure, T., Liebler, D.C., Mayersohn, M. and Sipes, I.G.: 3,3',4,4'-Tetrachloroazobenzene adsorption, disposition, and metabolism in male Fischer 344 rats. *Drug Metab. Disp.*, 24 (1996) 238-244.

- 3404 Polese, L., Minelli, E.V. and Jardim, E.F.G.: Small-scale method for the determination of selected organochlorine pesticides in soil. *Fresenius J. Anal. Chem.*, 354 (1996) 474-476.
- 3405 Schurig, V., Glausch, A. and Fluck, M.: On the enantio-merization barrier of atropisomeric 2,2',3,3',4,6-hexachlorobiphenyl (PCB 132). *Tetrahedron: Asymmetry*, 6 (1995) 2161-2164; C.A., 124 (1996) 55187a.

See also 2697.

## 29c. Phosphorus insecticides

- 3406 Chang, M.J.W., Lin, C.Y., Lo, L.W. and Lin, R.S.: Biological monitoring of exposure to chlorpyrifos by high performance liquid chromatography. *Bull. Environ. Contam. Toxicol.*, 56 (1996) 367-374; C.A., 124 (1996) 78856g.
- 3407 Lacorte, S. and Barceló, D.: Improvements in the determination of organophosphorus pesticides in ground- and wastewater samples from interlaboratory studies by automated on-line liquid-solid extraction followed by liquid chromatography-diode array detection. *J. Chromatogr. A*, 725 (1996) 85-92.
- 3408 Tauler, R., Lacorte, S. and Barceló, D.: Application of multivariate self-modeling curve resolution to the quantitation of trace levels of organophosphorus pesticides in natural waters from interlaboratory studies. *J. Chromatogr. A*, 730 (1996) 177-183.
- 3409 Tolosa, I., Readman, J.W. and Mee, L.D.: Comparison of the performance of solid-phase extraction techniques in recovering organophosphorus and organochlorine compounds from water. *J. Chromatogr. A*, 725 (1996) 93-106.

See also 2511.

## 29d. Carbamates

- 3410 Chiron, S., Valverde, A., Fernandez-Alba, A. and Barceló, D.: Automated sample preparation for monitoring groundwater pollution by carbamate insecticides and their transformation products. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1346-1352.
- 3411 Izquierdo, A., Tena, M.T., Luque de Castro, M.D. and Valcárcel, M.: Supercritical fluid extraction of carbamate pesticides from soils and cereals. *Chromatographia*, 42 (1996) 206-212.
- 3412 Rouberty, F. and Fournier, J.: Optimization of HPLC separation of carbamate insecticides (carbofuran, hydroxycarbofuran and aldicarb) by experimental design methodology. *J. Liq. Chromatogr.*, 19 (1996) 37-55.

## 29e. Herbicides

- 3413 Balinova, A.: Ion-pairing mechanism in the solid-phase extraction and reversed-phase high-performance liquid chromatographic determination of acidic herbicides in water. *J. Chromatogr. A*, 728 (1996) 319-324.

- 3414 Del Valle, P.L., Muldoon, M.T., Karns, J.S., Nelson, J.O. and Mulbry, W.W.: Development of improved immunoassay and HPLC methods for the analysis of chlorodiamino-s-triazine in environmental samples. *J. Agric. Food Chem.*, 44 (1996) 945-952; C.A., 124 (1996) 168136b.
- 3415 Gennaro, M.C. and Giacosa, D.: Separation of triazine herbicides by ion-interaction HPLC and application to surface waters. *J. Liq. Chromatogr.*, 19 (1996) 149-160.
- 3416 Gennaro, M.C., Giacosa, D., Baglietto, C., Gennari, M. and Negré, M.: Simultaneous separation of phenylurea-, triazine- and phenoxyacid herbicides by reverse phase ion-interaction HPLC. Application to soil analysis. *J. Liq. Chromatogr.*, 19 (1996) 911-924.
- 3417 Gorth, S.M., Hogendoorn, E.A., Dijkman, E., van Zoonen, P. and Hoogerbrugge, R.: The optimization of step-gradient elution conditions in liquid chromatography: application to herbicide residue analysis using coupled-column reversed-phase LC. *Chromatographia*, 42 (1996) 17-24.
- 3418 Graham, K.N., Sarna, L.P., Webster, G.R.B., Gaynor, J.D. and Ng, H.Y.F.: Solid-phase microextraction of the herbicide metolachlor in runoff and tile-drainage water samples. *J. Chromatogr. A*, 725 (1996) 129-136.
- 3419 Ibañez, M., Picó, Y. and Mañes, J.: Influence of organic matter and surfactants on solid-phase extraction of diquat, paraquat and difenoquat from waters. *J. Chromatogr. A*, 727 (1996) 245-252.
- 3420 Ibañez, M., Picó, Y. and Mañes, J.: On-line liquid chromatographic trace enrichment and high-performance liquid chromatographic determination of diquat, paraquat and difenoquat in water. *J. Chromatogr. A*, 728 (1996) 325-331.
- 3421 Kirksey, K.B. and Mueller, T.C.: Effect of extraction systems on clomazone recovery from aged soil samples. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1519-1522.
- 3422 Kuan, J.W. and Pepperman, A.B.: High performance liquid chromatographic determination of alachlor in alginic-based formulations. *J. Liq. Chromatogr.*, 19 (1996) 645-659.
- 3423 Lian, H.-Z., Zhang, W.B., Jiang, Q. and Miao, J.: Simultaneous determination of metsulfuron-methyl, chlorsulfuron and ben-sulfuron-methyl in various formulations of sulfonylurea herbicides by HPLC-UV detection. *J. Liq. Chromatogr.*, 19 (1996) 207-216.
- 3424 Matuszczyk, G., Knopp, D. and Niessner, R.: (Enrichment of environmental pollutants. Preparation and characterization of an immunoabsorbent for 2,4-dichlorophenoxyacetic acid (2,4-D)). *Vom Wasser*, 85 (1995) 81-94; C.A., 124 (1996) 184869y.
- 3425 Nomura, N., Morishita, F., Kurata, S., Asamura, H., Imada, Y. and Arakida, Y.: An approximate calculation of the capacity factors of alkylphenols in RP-HPLC. In: Hatano, H. and Hanai, Y. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 611-619; C.A., 124 (1996) 75134j.
- 3426 Trocewicz, J.: Determination of herbicides in surface water by means of a supported liquid membrane technique and high-performance liquid chromatography. *J. Chromatogr. A*, 725 (1996) 121-127.

See also 3397.

#### 29f. Fungicides

- 3427 Arenas, R.V., Rahman, H. and Johnson, N.A.: Determination of thiabendazol residues in white and sweet potatoes by liquid chromatography with fluorescence detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1455-1458.
- 3428 Motohashi, N., Nagashima, H. and Meyer, R.: HPLC determination of the fungicides bitertanol, imazalil, and thiabendazole in banana. *Heterocycl. Commun.*, 1 (1995) 431-444; C.A., 124 (1996) 53995p.
- 3429 Nakashima, H. and Miyano, N.: (Studies on safety evaluation of antimicrobial and deodorant agents. Determination of fungicides in miticidal finished products and household paints). *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 433-439; C.A., 124 (1996) 168134z.
- 3430 Nakazato, M., Tadano, K., Ogawa, H., Ushiyama, H., Kawai, Y., Kobayashi, T., Tateishi, Y. and Tamura, Y.: (Determination of imazalil, diphenyl, thiabendazole, and o-phenylphenol in citrus fruits, and of imazalil and thiabendazole in banana, by high performance liquid chromatography). *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 392-397; C.A., 124 (1996) 115644j.
- 3431 Thomas, D.H., Lopez-Avila, V., Betowski, L.D. and van Emon, J.: Determination of carbendazim in water by high-performance immunoaffinity chromatography on-line with high-performance liquid chromatography with diode-array or mass spectrometric detection. *J. Chromatogr. A*, 724 (1996) 207-217.

See also 3632.

#### 29g. Other types of pesticides and various agrochemicals

- 3432 El-Shahawi, M.S. and Aldhaheri, S.M.: Preconcentration and separation of acaricides by polyether based polyurethane foam. *Anal. Chim. Acta*, 320 (1996) 277-287.
- 3433 Harbin, D.N.: Quantitation of cyfluthrin in liquid and solid formulations by reversed-phase liquid chromatography: collaborative study. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1335-1338.
- 3434 Pang, G.-F., Chao, Y.-Z., Liu, X.-S. and Fan, C.-L.: Multiresidue liquid chromatographic method for simultaneous determination of pyrethroid insecticides in fruits and vegetables. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1474-1480.

See also 3728.

### 30. SYNTHETIC AND NATURAL DYES

#### 30a. Synthetic dyes

- 3435 Gagliardi, L., de Orsi, D. and Cozzoli, O.: (The problem of analysing dyes in cosmetic products). *Cosmet., Toiletries, (Ed. Ital.)*, 16 (1995) 31-37; C.A., 124 (1996) 155619n.
- 3436 Henricsson, S. and Westerholm, R.: Liquid chromatographic method for analysing the colour marker Solvent Yellow 124, N-ethyl-N-[2-(1-isobutoxyethoxy)ethyl](4-phenylazophenyl)amine, in diesel fuels. *J. Chromatogr. A*, 723 (1996) 395-398.
- 3437 Zghal, H., Gueitin, C., Haloui, E. and Czchok, M.: Determination of Sunset Yellow FCF and Ponceau 4R in candies by high-performance liquid chromatography. *Sci. Aliments*, 15 (1995) 491-496; C.A., 124 (1996) 200441k.

For additional information see C.A.:  
124 (1996) 143867c.

*30b. Chloroplast and other natural pigments*

- 3438 Borel, P., Grolier, P., Armand, M., Partier, A., Lafont, H., Lairon, D. and Zazis-Braesco, V.: Carotenoids in biological emulsions: solubility surface-to-core distribution, and release from lipid droplets. *J. Lipid Res.*, 37 (1996) 250-261.
- 3439 Bridle, P. and Garcia-Viguera, C.: A simple technique for the detection of red wine adulteration with elderberry pigments. *Food Chem.*, 55 (1996) 111-113; C.A., 124 (1996) 200462t.
- 3440 Elbesgali, A.K., Breneton, R.G. and Rahmani, A.: Influence of the method of calculation of noise threshold on wavelength selection in window factor analysis of diode array high-performance liquid chromatography. *Analyst (Cambridge)*, 121 (1996) 585-590.
- 3441 Fukushima, M., Tanaka, S., Nakamura, H., Ito, S., Haraguchi, K. and Ogata, T.: Copper(II) binding abilities of molecular weight fractionated humic acids and their mixtures. *Anal. Chim. Acta*, 322 (1996) 173-185.
- 3442 Hagen, J.J., Washco, K.A. and Monnig, C.A.: Determination of retinoids by reversed-phase capillary liquid chromatography with amperometric electrochemical detection. *J. Chromatogr. B*, 677 (1996) 225-231.
- 3443 Harris, P.G. and Maxwell, J.R.: A novel method for the rapid determination of concentrations at high stratigraphic resolution in marine sediments. *Org. Geochem.*, 23 (1995) 853-856; C.A., 124 (1996) 96907d.
- 3444 Huyghues-Despointes, A. and Yaylayan, V.A.: Retro-aldol and redox reactions of Amadori compounds: mechanistic studies with variously labeled D-[<sup>13</sup>C]-glucose. *J. Agric. Food Chem.*, 44 (1996) 672-681; C.A., 124 (1996) 200485c.
- 3445 Ito, H., Ohtsuka, T. and Tanaka, A.: Conversion of chlorophyll b to chlorophyll a via 7-hydroxymethyl chlorophyll. *J. Biol. Chem.*, 271 (1996) 1475-1479.
- 3446 Minguez-Mosquera, M.I., Gandul-Rojas, B. and Garrido-Fernández, J.: Preparation of Cu(II) complexes of oxidized chlorophylls and their determination by thin-layer and high-performance liquid chromatography. *J. Chromatogr. A*, 731 (1996) 261-271.
- 3447 Nagao, A., During, A., Hoshino, C., Terao, J. and Olson, J.A.: Stoichiometric conversion of *all-trans*-β-carotene to retinal by pig intestinal extract. *Arch. Biochem. Biophys.*, 328 (1996) 57-63.
- 3448 Porcar, M., Bel, Y., Socha, R., Nemec, V. and Ferre, J.: Identification of pteridines in the firebug, *Pyrrhocoris apterus* (L.) (Heteroptera, Pyrrhocoridae) by high-performance liquid chromatography. *J. Chromatogr. A*, 724 (1996) 193-197.
- 3449 Trubetskoy, O.A., Trubetskaya, O.E., Afanas'eva, G.V. and Reznikova, O.I.: (Combination of gel chromatography and PAAG electrophoresis for preparative accumulation of humic acid fractions). *Izv. Akad. Nauk, Ser. Biol.*, (1995) 481-486; C.A., 124 (1996) 54643j.
- 3450 Wouters, J.: Analysis of natural organic dyes by high performance liquid chromatography. *Nouv. Sci. Technol.*, 13 (1995) 453-455; C.A., 124 (1996) 144770c.

See also 3316, 3322, 3325, 3327, 3328.

**31. PLASTICS AND THEIR INTERMEDIATES**

- 3451 Boyd, R.H., Chance, R.R. and Ver Strate, G.: Effective dimensions of oligomers in size exclusion chromatography. A molecular dynamic simulation study. *Macromolecules*, 29 (1996) 1182-1190; C.A., 124 (1996) 88331w.
- 3452 Egestad, B., Green, G., Sjöberg, P., Klasson-Wehler, E. and Gustafsson, J.: Chromatographic fractionation and analysis by mass spectrometry of conjugated metabolites of bis(2-ethylhexyl)phthalate in urine. *J. Chromatogr. B*, 677 (1996) 99-109.
- 3453 Glöckner, G., Engelhardt, H., Wolf, D. and Schultz, R.: Sample size and retention values in gradient liquid chromatography of synthetic polymers. *Chromatographia*, 42 (1996) 185-190.
- 3454 Haken, J.K. and Iddamalgoda, P.-I.: Validation of micro-fusion chromatographic analysis of polyester systems. *Prog. Org. Coat.*, 26 (1995) 101-111; C.A., 124 (1996) 88402v.
- 3455 Jackson, C., Larsen, B. and McEwen, C.: Comparison of most probable peak values as measured for polymer distributions by MALDI mass spectrometry and by size exclusion chromatography. *Anal. Chem.*, 68 (1996) 1303-1308.
- 3456 Jacques, B., Devaux, J., Legras, R. and Nield, E.: Study of an appropriate procedure for the chromatographic analysis of poly(ethylene terephthalate), after melt processing with triphenyl phosphite. *J. Chromatogr. A*, 725 (1996) 367-371.
- 3457 La, S.B., Okano, T. and Kataoka, K.: Preparation and characterization of the micelle-forming polymeric drug indomethacin-incorporated poly(ethylene oxide)-poly(β-benzyl L-aspartate) block copolymer micelles. *J. Pharm. Sci.*, 85 (1996) 85-90.
- 3458 Lehrle, R.S. and Sarson, D.S.: Degradation and selective desorption of polymer can cause uncertainty in MALDI (matrix assisted laser desorption ionization) measurements. *Polym. Degrad. Stab.*, 51 (1996) 197-204; C.A., 124 (1996) 233966p.
- 3459 Lochmüller, C.H. and Jiang, C.: Retention behavior of poly(methyl methacrylate) and poly(ethylene glycol) in reversed-phase liquid chromatography. *J. Chromatogr. Sci.*, 33 (1995) 561-567.
- 3460 Lochmüller, C.H., Moebus, M.A., Liu, Q. and Jiang, C.: Temperature effect on retention and separation of poly(ethylene glycol)s in reversed-phase liquid chromatography. *J. Chromatogr. Sci.*, 34 (1996) 69-76.
- 3461 Machtalère, G., Xie, J., Hui, F., Kolodziejczyk, H. and Rosset, R.: Characterization of engine motor oils dispersants of the polybutenylsuccinimide type by saponification, methylation and size exclusion chromatography. *Anal. Chim. Acta*, 322 (1996) 31-41.
- 3462 Meehan, E., Warner, F.P., Reid, S.P., Patterson, M. and Dawkins, J.V.: Characterization of poly(vinyl alcohol) by liquid chromatographic techniques. *Int. J. Polym. Anal. Charact.*, 2 (1995) 71-82; C.A., 124 (1996) 203506w.
- 3463 Mori, S.: Characterization of styrene-acrylonitrile copolymers by size-exclusion chromatography/stepwise gradient elution-liquid precipitation chromatography. *Int. J. Polym. Anal. Charact.*, 2 (1996) 185-192; C.A., 124 (1996) 203551g.
- 3464 Nakamura, M., Ofusa, Y. and Mori, S.: Observation of absence of mechanical degradation of high molecular weight polystyrenes during elution through a column in size exclusion chromatography. *J. Liq. Chromatogr.*, 19 (1996) 339-352.

- 3465 Philipsen, H.J.A., Klumperman, B., van Herk, A.M. and German, A.L.: Critical retention behaviour of polymers. A study on the influence of some practical parameters. *J. Chromatogr. A*, 727 (1996) 13-25.
- 3466 Scheiman, D.A., Alston, W.B., Sutter, J.K. and Turk, M.J.: Characterization of PMR monomer solutions and prepgs using HPLC analysis. *Nasa Conf. Publ.*, 1911 (1993) 18/1-18/10; C.A., 124 (1996) 57597h.
- 3467 Svec, F.: (Porous crosslinked polymer materials for biotechnologies). *Plasty Kauc.*, 32 (1995) 356-360; C.A., 124 (1996) 147922q - a review with 14 refs.

For additional information see C.A.:  
124 (1996) 57700m, 119810h.

See also 2471, 2475, 2476, 2491, 2701, 2706, 3743.

## 32. DRUG ANALYSIS

### 32a. Drug analysis, general techniques

- 3468 Bilke, H.W., Molnar, I. and Gernet, C.: High-performance liquid chromatographic separation of the impurities in a pharmaceutical raw material with the aid of computer simulation. *J. Chromatogr. A*, 729 (1996) 189-195.
- 3469 Burinsky, D.J., Armstrong, B.L., Oyler, A.R. and Dunphy, R.: Characterization of tepezolin and its related compounds by high-performance liquid chromatography/mass spectrometry. *J. Pharm. Sci.*, 85 (1996) 159-164.
- 3470 Carducci, C.N., Lucangioli, S.E., Rodriguez, V.G. and Fernández Otero, G.C.: Application of extraction disks in dissolution tests of clenbuterol and levothyroxine tablets by capillary electrophoresis. *J. Chromatogr. A*, 730 (1996) 313-319.
- 3471 Engel, G., Hofmann, U. and Kroemer, H.K.: Prediction of CYP2D6-mediated polymorphic drug metabolism (sparteine type) based on *in vitro* investigations. *J. Chromatogr. B*, 678 (1996) 93-103 - a review with 73 refs.
- 3472 Forgács, E. and Cserháti, T.: Relationship between the retention characteristics and physicochemical parameters of some steroid drugs in reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 728 (1996) 75-81.
- 3473 Garcia, K.E., Sandra, P., Baeyens, W.R.G., Ferraz, V., van der Weken, G. and van Overbeke, A.: Chiral separation of drugs by micro-LC with cellulose as a stationary phase. *Biomed. Chromatogr.*, 9 (1995) 285-286; C.A., 124 (1996) 66712u.
- 3474 Hannak, D., Scharbert, F. and Kattermann, R.: Stepwise binary gradient high-performance liquid chromatographic system for routine drug monitoring. *J. Chromatogr. A*, 728 (1996) 307-310.
- 3475 Ivanovic, D., Medenica, N., Nivaud-Guernet, E. and Guernet, M.: Study of the reversed-phase high-performance liquid chromatographic separation of some pharmaceutical ingredients. *J. High Resolut. Chromatogr.*, 19 (1996) 124-126.
- 3476 Law, B. and Appleby, J.R.G.: Re-evaluation of strong cation-exchange high-performance liquid chromatography for the analysis of basic drugs. *J. Chromatogr. A*, 725 (1996) 335-341.
- 3477 Lowrey, A.H. and Famini, G.R.: Using theoretical descriptors in quantitative structure-activity relationships: HPLC capacity factors for energetic materials. *Struct. Chem.*, 6 (1995) 357-365; C.A., 124 (1996) 47813m.
- 3478 Marzo, A. and Balant, L.P.: Investigation of xenobiotic metabolism by CYP2D6 and CYP2C19: importance of enantioselective analytical methods. *J. Chromatogr. B*, 678 (1996) 73-92 - a review with 105 refs.
- 3479 Nicolas, A.: Analytical strategies for the screening of veterinary drugs and their residues in edible products (Review), by Aerts et al.: an addendum. *J. Chromatogr. B*, 679 (1996) 210-211.
- 3480 Norta, M. and Schöpke, T.: An *in vitro* perfusion model for the determination of absorption properties of drugs in isolated rat small intestine. *Pharmazie*, 51 (1996) 101-103.
- 3481 Ong, S., Liu, H. and Pidgeon, C.: Immobilized-artificial-membrane chromatography: measurements of membrane partition coefficient and predicting drug membrane permeability. *J. Chromatogr. A*, 728 (1996) 113-128.
- 3482 Oravcová, J., Böhls, B. and Lindner, W.: Drug-protein binding studies. New trends in analytical and experimental methodology. *J. Chromatogr. B*, 677 (1996) 1-28 - a review with 241 refs.
- 3483 Prankerd, R.J.: Practical aspects of the stability testing of formulated drug products. *Aust. J. Hosp. Pharm.*, 25 (1995) 405-408; C.A., 124 (1996) 66303m - a review with 8 refs.
- 3484 Quaglia, M.G., Bossù, E., Porretta, G.C., Biava, M., Fioravanti, R., Romanelli, L. and Leonardi, A.: Chromatographic resolution and pharmacological investigation of some imidazole derivatives. *J. Chromatogr. A*, 729 (1996) 1-4.
- 3485 Shibukawa, A. and Nakagawa, T.: (Theoretical study of high-performance frontal analysis, a chromatographic method for analysis of drug-protein interaction). *Kuromatogurafi*, 16 (1995) 290-291; C.A., 124 (1996) 164150d.
- 3486 Sidelmann, U.G., Nicholls, A.W., Meadows, P.E., Gilbert, J.W., Lindon, J.C., Wilson, I.D. and Nicholson, J.K.: High-performance liquid chromatography directly coupled to <sup>19</sup>F and <sup>1</sup>H NMR for the analysis of mixtures of isomeric ester glucuronide conjugates of trifluoromethylbenzoic acids. *J. Chromatogr. A*, 728 (1996) 377-385.
- 3487 Soltes, L., Büschges, R., Spahn-Langguth, H., Mutschler, E. and Sebillle, B.: An alternative high-performance liquid chromatographic arrangement for drug quality control. *Pharmazie*, 51 (1996) 93-96.
- 3488 Stelzer, A., Jira, T., Meusel, M., Morgenstern, O. and Plümer, F.: Liquid chromatographic separation of chiral triazolophthalazines. *J. Chromatogr. A*, 729 (1996) 5-11.
- 3489 Swartz, M.E. and Brown, P.R.: Use of mathematically enhanced spectral analysis and spectral contrast techniques for the liquid chromatographic and capillary electrophoretic detection and identification of pharmaceutical compounds. *Chirality*, 8 (1996) 67-76; C.A., 124 (1996) 242479j.
- 3490 Vecerkova, J.: Systematic analytic procedure for identification of unknown medicaments in biological material. *Sb. Lek.*, 95 (1994) 357-361; C.A., 124 (1996) 163953f - a review with 3 refs.
- 3491 Ye, L. and Stewart, J.T.: A comparison study of various octadecylsilane (C 18) solid-phase extraction materials in HPLC analysis of selected drugs and their metabolites in water and serum samples. *Anal. Lett.*, 29 (1996) 395-407.

For additional information see C.A.:

124 (1996) 104848c, 212174p, 218931e.

See also 2494, 2523, 2561, 2568, 2623, 2630, 2632, 2633, 2637, 2641, 2781, 2809, 2873, 2994, 3289, 3740, 3747.

### 32b. Antirheumatics and antiinflammatory drugs

- 3492 Bischer, A., Zia-Amirhosseini, P., Iwaki, M., McDonagh, A.F. and Benet, L.Z.: Stereoselective binding properties of naproxen glucuronide diastereomers to proteins. *J. Pharmacokin. Biopharm.*, 23 (1995) 379-395.
- 3493 Cerretani, D., Micheli, L., Fiaschi, A.I. and Giorgi, G.: High-performance liquid chromatography of flufenamic acid in rat plasma. *J. Chromatogr. B*, 678 (1996) 365-368.
- 3494 Colgan, S.T., Hammen, P.D., Knutson, K.L. and Bordner, J.: Investigation of cyclam-containing mobile phases for the liquid chromatographic analysis of tenidap. *J. Chromatogr. Sci.*, 34 (1996) 111-114.
- 3495 Creaser, C.S., Feely, S.J., Houghton, E., Seymour, M. and Teale, P.: Online immunoaffinity chromatography-high-performance liquid chromatography-mass spectroscopy for the determination of dexamethasone. *Anal. Commun.*, 33 (1996) 5-8; C.A., 124 (1996) 135897x.
- 3496 Golich, T.G., Thomsen, J.R., Green, D.L. and Jersild, J.: Determination of minor impurities of (R)-1-acetoxymethyl-3-(7-[1-N,N-dimethylcarbamoyl]-6-(4-fluorophenyl)-indol-3-oyl)-1H,3H-pyrrolo[1,2-c]thiazol-3-yl)pyridinium chloride by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 4107-4116.
- 3497 Grubb, N.G., Rudy, D.W. and Hall, S.D.: Stereoselective high-performance liquid chromatographic analysis of ketoprofen and its acyl glucuronides in chronic renal insufficiency. *J. Chromatogr. B*, 678 (1996) 237-244.
- 3498 Hanses, A., Spahn-Langguth, H., Meiss, F. and Mutschler, E.: Pharmacokinetics and drug input characteristics for a diclofenac-codeine phosphate combination following oral and rectal administration. *Arzneim.-Forsch.*, 46 (1996) 57-63.
- 3499 Kees, F., Jehnrich, D. and Grobecker, H.: Simultaneous determination of acetylsalicylic acid and salicylic acid in human plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 172-177.
- 3500 Lovlin, R., Vakily, M. and Jamali, F.: Rapid, sensitive and direct chiral high-performance liquid chromatographic method for ketoprofen enantiomers. *J. Chromatogr. B*, 679 (1996) 196-198.
- 3501 Motohashi, N. and Saito, Y.: Rate constants for reaction of hydroxyl radicals with sulfapyridine and aminosalicylic acids. *Chem. Pharm. Bull.*, 44 (1996) 163-166.
- 3502 Neto, L.M.R., Andraus, M.H. and Salvadori, M.C.: Determination of phenylbutazone and oxyphenbutazone in plasma and urine samples of horses by high-performance liquid chromatography and gas chromatography-mass spectrometry. *J. Chromatogr. B*, 678 (1996) 211-218.
- 3503 Péhourcq, F., Lagrange, F., Labat, L. and Bannwarth, B.: Simultaneous measurement of flurbiprofen, ibuprofen, and ketoprofen enantiomer concentrations in plasma using L-leucinamide as the chiral coupling component. *J. Liq. Chromatogr.*, 18 (1995) 3969-3979.

- 3504 Ramakrishna, S., Fadnavis, N.W. and Diwan, P.V.: Comparative pharmacokinetic evaluation of compressed suppositories of diclofenac sodium in humans. *Arzneim.-Forsch.*, 46 (1996) 175-177.
- 3505 Slater, J.G., Feenstra, K.L., Hauer, M.J., Kloosterman, D.A., Parton, A.H., Sanders, P.E., Scott, G. and Speed, W.: Metabolism of the bisphosphate ester U-91502 in rats. *Drug Metab. Disp.*, 24 (1996) 65-73.
- 3506 Solà, J., Prunosa, J., Colom, H., Peraire, C. and Obach, R.: Determination of ketorolac in human plasma by high performance liquid chromatography after automated on-line solid phase extraction. *J. Liq. Chromatogr.*, 19 (1996) 89-99.
- 3507 Thomas, S.B. and Albaiz, S.J.: Simultaneous determination of the 5-lipoxygenase inhibitor "Zileuton" and its N-dehydroxylated metabolite in untreated rat urine by micellar liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 977-991.
- 3508 Van Os, E.C., McKinney, J.A., Zins, B.J., Mays, D.C., Schriver, Z.H., Sandborn, W.J. and Lipsky, J.J.: Simultaneous determination of azathioprine and 6-mercaptopurine by high-performance liquid chromatography. *J. Chromatogr. B*, 679 (1996) 147-154.
- 3509 Yang, M. and Qian, Y.: (A HPLC method for simultaneous determination of propafenone and hydroquinidine in human serum). *Zhongguo Yiyan Yaoxue Zazhi*, 15 (1995) 437-439; C.A., 124 (1996) 75418e.

For additional information see C.A.:

124 (1996) 97901j, 185772y.

See also 2644, 2835, 3457, 3475, 3487, 3613.

### 32c. Autonomic and cardiovascular drugs

- 3510 Aboul-Enein, H.Y. and Abou-Basha, L.I.: HPLC separation of nadolol and enantiomers on Chiralcel OD column. *J. Liq. Chromatogr.*, 19 (1996) 383-392.
- 3511 Aboul-Enein, H.Y. and Serignese, V.: Direct HPLC separation of indenolol enantiomers using a cellulose chiral stationary phase. *J. Liq. Chromatogr.*, 19 (1996) 933-938.
- 3512 Aboul-Enein, H.Y., Abou-Basha, L.I. and Bakr, S.A.: Direct enantioselective separation of some propranolol analogs by HPLC on normal and reversed cellulose chiral stationary phases. *Chirality*, 8 (1996) 153-156; C.A., 124 (1996) 242486j.
- 3513 Appanna, G., Tang, B.K., Müller, R. and Kalow, W.: A sensitive method for determination of cytochrome P4502D6 activity *in vitro* using bupranolol as substrate. *Drug Metab. Disp.*, 24 (1996) 303-306.
- 3514 Aromdee, C., Fawcett, J.P. and Ledger, R.: Sensitive high-performance liquid chromatographic assay for pilocarpine in biological fluids using fluorescence derivatisation. *J. Chromatogr. B*, 677 (1996) 313-318.
- 3515 Begoña Barroso, M., Alonso, R.M. and Jiménez, R.M.: Quantitative analysis of the loop diuretic torasemide in tablets and human urine by HPLC-EC. *J. Liq. Chromatogr.*, 19 (1996) 179-186.
- 3516 Begoña Barroso, M., Alonso, R.M. and Jiménez, R.M.: Simultaneous determination of the diuretics triamterene and furosemide in pharmaceutical formulations and urine by HPLC-EC. *J. Liq. Chromatogr.*, 19 (1996) 231-246.

- 3517 Bichara, N., Ching, M.S., Blake, C.L., Ghabrial, H. and Smallwood, R.A.: Propranolol hydroxylation and N-desisopropylation of cytochrome P4502D6. Studies using the yeast-expressed enzyme and NADPH<sub>2</sub> and cumene hydroperoxide-supported reactions. *Drug Metab. Disp.*, 24 (1996) 112-118.
- 3518 Butter, J.J., van den Berg, B.T.J., Portier, E.J.G., Kaiser, G. and van Boxtel, C.J.: Determination by HPLC with electrochemical detection of formoterol RR and SS enantiomers in urine. *J. Liq. Chromatogr.*, 19 (1996) 993-1005.
- 3519 Cieri, U.R.: Determination of reserpine and chlorothiazide in commercial tablets by liquid chromatography with fluorescence and UV absorbance detectors in series. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1384-1387.
- 3520 Elrod, L., Jr., Schmit, J.L. and Morley, J.A.: Determination of isoproterenol sulfate on surfaces using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. A*, 723 (1996) 235-241.
- 3521 Fischer, U., Wacke, R., Stange, J., Nitschke, F.P., Adam, U. and Drewelow, B.: Rapid HPLC assay for varapamil and its metabolites: use for application to *in vitro* studies. *Pharmazie*, 51 (1996) 220-223.
- 3522 Frye, R.F. and Branch, R.A.: Improved high-performance liquid chromatographic determination of debrisoquine and 4-hydroxy-debrisoquine in human urine following direct injection. *J. Chromatogr. B*, 677 (1996) 178-182.
- 3523 Gigosos, P.G., Fernández, T.F., Maríz, O.C., Sampayo, C.A.F., Abuín, C.F. and Sáez, A.C.: Rapid and simple determination of clenbuterol residues in retina by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 677 (1996) 167-171.
- 3524 Hendrickx, J., Bockx, M., Zwijnen, C., Borgmans, C., Mannens, G., Meuldermans, W. and Heykants, J.: Location of the hydroxyl functions in hydroxylated metabolites of nebivolol in different animal species and human subjects as determined by on-line high-performance liquid chromatography-diode-array detection. *J. Chromatogr. A*, 729 (1996) 341-354.
- 3525 Herber, R., Hercelin, B., van Cantfort, J., de Graeve, J., Fourel-Gigleux, S., Taguchi, T. and Magdalou, J.: Reduction and glucuronidation of naftazone by human and rat liver microsomes. *Drug Metab. Disp.*, 23 (1995) 1305-1314.
- 3526 Herráez-Hernández, R., Campins-Falcó, P. and Sevillano-Cabeza, A.: Application of column switching in high performance liquid chromatographic analysis of chlorthalidone enantiomers in untreated urine. *J. Liq. Chromatogr.*, 19 (1996) 403-414.
- 3527 Herrmann, R.: Bioäquivalenz einer Nifedipin-Retardformulierung im Vergleich zu einer Standardzubereitung in Steady-State. *Arzneim.-Forsch.*, 46 (1996) 28-34.
- 3528 Iyer, E.K. and Tipnis, H.P.: Determination of timolol in plasma by high performance liquid chromatography. *Indian Drugs*, 32 (1995) 287-288; C.A., 124 (1996) 105435c.
- 3529 Kadenatsi, I.B., Levchuk, S.N., Agapitova, I.V., Glezer, M.G., Dombrovskii, V.S., Mukumov, M.R. and Firsov, A.A.: (HPLC determination of gemfibrozil in pharmacokinetic studies). *Khim.-Farm. Zh.*, 29 (1995) 61-64; C.A., 124 (1996) 105446g.
- 3530 Kountourellis, J.E. and Ebete, K.O.: Determination of pimephixene in plasma and urine by high performance liquid chromatography. *Anal. Lett.*, 29 (1996) 775-784.
- 3531 Kubrak, Z.V., Popova, V.I. and Komaritsa, I.D.: (Application of gel-chromatography for determination of ethacizine in biological materials). *Farm. Zh. (Kiev)*, (1995) 75-77; C.A., 124 (1996) 185716h.
- 3532 Kugler, A.R., Olson, S.C. and Smith, D.E.: Disposition of quinapril and quinaprilat in the isolated perfused rat kidney. *J. Pharmacokin. Biopharm.*, 23 (1995) 287-305.
- 3533 Kusu, F., Matsumoto, K., Arai, K. and Takamura, K.: Determination of synephrine enantiomers in food and conjugated synephrine in urine by high-performance liquid chromatography with electrochemical detection. *Anal. Biochem.*, 235 (1996) 191-194.
- 3534 Lee, H., Shim, H.O. and Lee, H.S.: Simultaneous determination of losartan and active metabolites EXP3174 in rat plasma by HPLC with column switching. *Chromatographia*, 42 (1996) 39-42.
- 3535 Malucelli, A., Rizzo, A. and Meyer, H.H.D.: Purification of clenbuterol, salbutamol, and terbutaline by immunoaffinity chromatography (IAC). *Arch. Lebensmittelhyg.*, 46 (1995) 101-103; C.A., 124 (1996) 105418z.
- 3536 Matchett, M.W., Branch, S.K. and Jefferies, T.M.: Polar organic chiral separation of propranolol and analogs using a  $\beta$ -cyclodextrin bonded stationary phase. *Chirality*, 8 (1996) 126-130; C.A., 124 (1996) 242483f.
- 3537 Mawatari, K.-i., Nakamura, Y., Shimizu, R., Sato, S., Inuma, F. and Watanabe, M.: Fluorimetric determination of nicorandil in human plasma by a high-performance liquid chromatographic-postcolumn ultraviolet detection system equipped with on-line back-pressure tubing. *J. Chromatogr. B*, 679 (1996) 155-159.
- 3538 McGrath, G.J., O'Kane, E., Smyth, W.F. and Tagliaro, F.: Investigation of the electrochemical oxidation of clenbuterol at a porous carbon electrode, and its application to the determination of this  $\beta$ -agonist in bovine hair by liquid chromatography with coulometric detection. *Anal. Chim. Acta*, 322 (1996) 159-166.
- 3539 Minina, S.A., Zhurkovich, I.K., Sergeeva, E.M., Puchkova, E.M., Pozharitskaya, O.N. and Korbelainen, E.S.: (HPLC analysis of quaternary derivatives of ajmaline). *Khim.-Pharm. Zh.*, 29 (1995) 52-54; C.A., 124 (1996) 242430m.
- 3540 Miyabayashi, T., Okuda, T., Motohashi, M., Izawa, K. and Yashiki, T.: Quantitation of a new potent angiotensin II receptor antagonist, TCV-116, and its metabolites in human serum and urine. *J. Chromatogr. B*, 677 (1996) 123-132.
- 3541 Pham-Huy, C., Sahui-Gnassi, A., Massicot, F., Galons, H., Dutertre-Catella, H., Radenen, B., Warnet, J.-M., Jr., Duc, H.T. and Claude, J.R., Jr.: Analysis of (S)- and (R)-propranolol in human plasma and urine by a specific immunoenzymic assay versus HPLC. *Arch. Toxicol. Suppl.*, 17 (Toxicology in Transition) (1995) 522-527; C.A., 124 (1996) 134626w.
- 3542 Pietiläinen, H. and Saesmaa, T.: HPLC determination of pindolol benzoate and pindolol 2-methoxyphenylacetate. *J. Liq. Chromatogr.*, 19 (1996) 583-591.
- 3543 Scully, P., Meehan, E. and Kelly, J.G.: High-performance liquid chromatographic assay for diltiazem in small-volume blood specimens and application to pharmacokinetic studies in rats. *J. Chromatogr. A*, 729 (1996) 297-300.
- 3544 Sun, E.L., Feenstra, K.L., Bell, F.B., Sanders, P.E., Slatter, J.G. and Ulrich, R.G.: Biotransformation of lifibrol (U-83860) to mixed glyceride metabolites by rat and human hepatocytes in primary culture. *Drug Metab. Disp.*, 24 (1996) 221-234.

- 3545 Sypniewski, S. and Bald, E.: Determination of captopril and its disulphides in whole human blood and urine by high-performance liquid chromatography with ultraviolet detection and precolumn derivatization. *J. Chromatogr. A*, 729 (1996) 335-340.
- 3546 Turberg, M.P., Macy, T.D., Lewis, J.J. and Coleman, M.R.: Determination of ractopamine hydrochloride in swine and turkey tissues by liquid chromatography with coulometric detection. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1394-1402.
- 3547 Van Waarde, A., Visser, T.J., Posthumus, H., Elsinga, P., Anthoon, R.L., van Loenen-Weemaes, A.-M.A., Visser, G.M., Beaufort-Krol, G.C.M., Paans, A.M.J. and Vaalburg, W.: Quantification of the  $\beta$ -adrenoceptor ligand S-1'--[ $^{18}\text{F}$ ]fluorocarazolol in plasma of humans, rats and sheep. *J. Chromatogr. B*, 678 (1996) 253-260.
- 3548 Vurna, V. and Kanfer, I.: High-performance liquid chromatographic determination of phenylephrine in human serum with coulometric detection. *J. Chromatogr. B*, 678 (1996) 245-252.
- 3549 Wieling, J., Hendriks, G., Tamminga, W.J., Hempenius, J., Mensink, C.K., Oosterhuis, B. and Jonkman, J.H.G.: Rational experimental design for bioanalytical methods validation. Illustration using an assay method for total captopril in plasma. *J. Chromatogr. A*, 730 (1996) 381-394.
- See also 2739, 3470, 3480, 3713.
- 32d. Central nervous system drugs**
- 3550 Aboul-Enein, H.Y. and Aboul-Basha, L.I.: Determination of tranylcypromine in urine and pharmaceutical formulation by HPLC using Symmetry column. *J. Liq. Chromatogr.*, 19 (1996) 925-932.
- 3551 Aboul-Enein, H.Y., Serignese, V., Abou-Basha, L.I. and Bojarski, J.: Direct chiral HPLC separation of several barbiturates on Chiracel® OJ column: Substituent effects on the enantioselectivity. *Pharmazie*, 51 (1996) 159-162.
- 3552 Argekar, A.P. and Kunjir, S.S.: High performance liquid chromatographic determination of roxatidine acetate in bulk drug and pharmaceutical preparations. *Indian Drugs*, 32 (1995) 333-335; *C.A.*, 124 (1996) 97895k.
- 3553 Ascalone, V., Ripamonti, M. and Malavasi, B.: Stereospecific determination of amisulpride, a new benzamide derivative, in human plasma and urine by automated solid-phase extraction and liquid chromatography on a chiral column. Application to pharmacokinetics. *J. Chromatogr. B*, 676 (1996) 95-105.
- 3554 Bagli, M., Rao, M.L., Höflich, G., Kasper, S., Langer, M., Barlage, U., Beneke, M., Süverkrüp, R. and Möller, H.-J.: Pharmacokinetics of chlorprothixene after single intravenous and oral administration of three galenic preparations. *Arzneim.-Forsch.*, 46 (1996) 247-250.
- 3555 Bansal, R. and Aranda, J.V.: High performance liquid chromatography microassay for the simultaneous determination of fentanyl and its major metabolites in biological samples. *J. Liq. Chromatogr.*, 19 (1996) 353-364.
- 3556 Berezentseva, O.M., Klyachko, Yu.A., Nikulichev, D.B., Nikulicheva, S.I. and Padalkina, V.S.: (Combined use of chromatographic methods of analysis in detection of some narcotic analgesics in biological fluids). *Sud.-Med. Ekspert.*, 38 (1995) 30-32; *C.A.*, 124 (1996) 164160g.
- 3557 Cai, J. and Henion, J.: Elucidation of LSD *in vitro* metabolism by liquid chromatography and capillary electrophoresis coupled with tandem mass spectrometry. *J. Anal. Toxicol.*, 20 (1996) 27-37; *C.A.*, 124 (1996) 134718c.
- 3558 Chandrashekhar, T.G., Patel, P., Vyas, S.K. and Dutt, C.: High-performance liquid chromatographic determination of loratadine in tablets. *East. Pharm.*, 38 (1995) 141-143; *C.A.*, 124 (1996) 97899q.
- 3559 Chism, J.P. and Rickert, D.E.: The pharmacokinetics and extrahepatic clearance of remifentanil, a short acting opioid antagonist, in male beagle dogs during constant rate infusion. *Drug Metab. Disp.*, 24 (1996) 34-40.
- 3560 Coffman, B.L., Rios, G.R. and Tephly, T.R.: Purification and properties of two rat liver phenobarbital-inducible UDP-glucuronosyltransferases that catalyze the glucuronidation of opioids. *Drug Metab. Disp.*, 24 (1996) 329-333.
- 3561 Di Pietra, A.M., Gatti, R., Andrisano, V. and Cavrini, V.: Application of high-performance liquid chromatography with diode-array detection and on-line post-column photochemical derivatization to the determination of analgesics. *J. Chromatogr. A*, 729 (1996) 355-361.
- 3562 Dowrie, R.H., Ebbling, W.F., Mandema, J.W. and Stanski, D.R.: High-performance liquid chromatographic assay of propofol in human and rat plasma and fourteen rat tissues using electrochemical detection. *J. Chromatogr. B*, 678 (1996) 279-288.
- 3563 Ducharme, J., Abdullah, S. and Wainer, I.W.: Dextromethorphan as an *in vivo* probe for the simultaneous determination of CYP2D6 and CYP3A activity. *J. Chromatogr. B*, 678 (1996) 113-128.
- 3564 El Walily, A.M., El Gindy, A. and Wahbi, A.M.: Determination of astemizole using iodine charge-transfer complexation and high-performance liquid chromatography. *S.T.P. Pharma Prat.*, 5 (1995) 403-408; *C.A.*, 124 (1996) 127237s.
- 3565 El-Sayed, Y.M., Khidr, S.H. and Niazy, E.M.: High-performance liquid chromatographic assay for the determination of haloperidol in plasma. *J. Liq. Chromatogr.*, 19 (1996) 125-134.
- 3566 Eto, S., Tanaka, N., Noda, H. and Noda, A.: Chiral separation of 10,11-dihydro-10,11-trans-dihydroxycarbamazepine, a metabolite of carbamazepine with two asymmetric carbons, in human serum. *J. Chromatogr. B*, 677 (1996) 325-330.
- 3567 Gupta, R.N.: Simultaneous determination of zopiclone and its two major metabolites (N-oxide and N-desmethyl) in human biological fluids by column liquid chromatography after solid-phase extraction. *J. Liq. Chromatogr.*, 19 (1996) 699-709.
- 3568 Gupta, R.N. and Hansebout, R.R.: Optimization of the determination of 4-aminopyridine in human serum and urine by column liquid chromatography. *J. Chromatogr. B*, 677 (1996) 183-189.
- 3569 Herráez-Hernández, R., Campins-Falcó, P. and Sevillano-Cabeza, A.: Determination of amphetamine and related compounds in urine using on-line derivatization in octadecyl silica columns with 9-fluorenylmethyl chloroformate and liquid chromatography. *J. Chromatogr. B*, 679 (1996) 69-78.
- 3570 Ho, S.-T., Wang, J.-J., Hu, O.Y.-P., Chiang, P.-S. and Lee, S.-C.: Determination of nalbuphine by high-performance liquid chromatography with ultraviolet detection: application to human and rabbit pharmacokinetic studies. *J. Chromatogr. B*, 678 (1996) 289-296.

- 3571 Jones, D.R., Gorski, J.C., Hamman, M.A. and Hall, S.D.: Quantification of dextromethorphan and metabolites: a dual phenotypic marker for cytochrome P450 3A4/5 and 2D6 activity. *J. Chromatogr. B*, 678 (1996) 105-111.
- 3572 Katagi, M., Nishioka, H., Nakajima, K., Tsuchihashi, H., Fujima, H., Wada, H., Nakamura, K. and Makino, K.: Direct high-performance liquid chromatographic and high-performance liquid chromatographic-thermospray-mass spectrometric determination of enantiomers of methamphetamine and its main metabolites amphetamine and *p*-hydroxymethamphetamine in human urine. *J. Chromatogr. B*, 676 (1996) 35-43.
- 3573 Kircher, V. and Parlar, H.: Determination of  $\Delta^9$ -tetrahydrocannabinol from human saliva by tandem immunoaffinity chromatography-high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 245-255.
- 3574 Kleinschmitz, M., Herderich, M. and Schreier, P.: Determination of 1,4-benzodiazepines by high-performance liquid chromatography-electrospray tandem mass spectrometry. *J. Chromatogr. B*, 676 (1996) 61-67.
- 3575 Kondo, T., Yoshida, K., Yamamoto, M. and Tanayama, S.: Enantiomeric toxicokinetics of the new isoindoline anxiolytic pazinaclone in rats and dogs. *Arzneim.-Forsch.*, 46 (1996) 11-14.
- 3576 Kopeleent-Frank, H. and Mittlböck, M.: Stability-indicating high-performance liquid chromatographic assay for the simultaneous determination of dixyrazine and chlorprothixene in intravenous admixtures. *J. Chromatogr. A*, 729 (1996) 201-206.
- 3577 Koyama, K., Kikuno, T., Ichikizaki, K., Murakoshi, K. and Ueki, H.: (Determination of serum drug levels by HPLC in 3 cases of acute intoxication with major-tranquillizer or antidepressant drug). *Iryo*, 49 (1995) 782-787; *C.A.*, 124 (1996) 44572w.
- 3578 Liu, Z.C. and Utrecht, J.P.: Clozapine is oxidized by activated human neutrophils to a reactive nitrenium ion that irreversibly binds to the cells. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1476-1483.
- 3579 Lloyd, T.L., Gupta, S.K., Gooding, A.E. and Alianti, J.R.: Determination of alosetron in human plasma or serum by high-performance liquid chromatography with robotic sample preparation. *J. Chromatogr. B*, 678 (1996) 261-267.
- 3580 Madden, S., Maggs, J.L. and Park, B.K.: Bioactivation of carbamazepine in the rat *in vivo*. Evidence for the formation of reactive arene oxide(s). *Drug Metab. Disp.*, 24 (1996) 469-479.
- 3581 Maggs, J.L., Pirmohamed, M. and Park, B.K.: The metabolic formation of reactive intermediates from clozapine, a drug associated with agranulocytosis in man. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1463-1475.
- 3582 Malavasi, B., Locatelli, M., Ripamonti, M. and Ascalone, V.: Determination of amisulpride, a new benzamide derivative, in human plasma and urine by liquid-liquid extraction or solid-phase extraction in combination with high-performance liquid chromatography and fluorescence detection. Application to pharmacokinetics. *J. Chromatogr. B*, 676 (1996) 107-115.
- 3583 Malavasi, B., Ripamonti, M., Rouchouse, A. and Ascalone, V.: Stereoselective determination of unchanged and glucuron conjugated eliprodil, a new anti-ischaemic drug, in human plasma and urine by precolumn derivatization and column-switching high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 729 (1996) 323-333.
- 3584 Malhotra, B.K., Lemaire, M., Brouillard, J.F. and Sawchuk, R.J.: High-performance liquid chromatographic analysis of (S)- $\alpha$ -amino-5-phosphonomethyl[1,1'-biphenyl]-3-propanoic acid (EAB 515) in brain and blood microdialysate (on-line) and in plasma ultrafiltrate of freely moving rats. *J. Chromatogr. B*, 679 (1996) 167-176.
- 3585 Marathe, P.H., Larkin, K.D., Duncan, G.F., Greene, D.S. and Barbhaya, R.H.: Determination of a novel indolylpiperazine anti-migraine agent in rat, monkey, mouse and rabbit plasma by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 678 (1996) 269-277.
- 3586 Mariot, R. and Zangiroli, L.: Quantitative determination of imidazenil, a novel imidazobenzodiazepine carboxamide derivative, by normal-phase high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 190-193.
- 3587 Martinez de Munoz, D., Arenas, R. and Chávez González, O.: Liquid chromatographic assay in plasma of one of the members of a new series of anticonvulsants: D,L-3-hydroxy-3-ethyl-3-phenylpropionamide. *J. Chromatogr. B*, 678 (1996) 377-383.
- 3588 Mawa, R., Mis, D., Gagnieu, M.C., Grancher, D., Petit-Ramel, M., Bressolle, F. and Vallon, J.-J.: Simple high-performance liquid chromatographic separation of oxazepam and its diastereoisomeric glucuronides in serum. Applications in a pharmacokinetic study in sheep. *J. Chromatogr. B*, 677 (1996) 331-338.
- 3589 McLoughlin, D.A., Olah, T.V., Ellis, J.D., Gilbert, J.D. and Halpin, R.A.: Quantitation of the 5HT<sub>1D</sub> agonists MK-462 and sumatriptan in plasma by liquid chromatography-atmospheric pressure chemical ionization mass spectrometry. *J. Chromatogr. A*, 726 (1996) 115-124.
- 3590 Milleroux, L., Brault, M., Gualano, V. and Mignot, A.: High-performance liquid chromatographic determination of baclofen in human plasma. *J. Chromatogr. A*, 729 (1996) 309-314.
- 3591 Nyoni, E.C., Sitaram, B.R. and Taylor, D.A.: Determination of  $\Delta^9$ -tetrahydrocannabinol levels in brain tissue using high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 679 (1996) 79-84.
- 3592 O'Neal, C. and Poklis, A.: Sensitive HPLC for simultaneous quantification of lidocaine and its metabolites monoethylglycinexylidide and glycinexylidide in serum. *Clin. Chem. (Washington)*, 42 (1996) 330-331.
- 3593 Piperaki, S. and Parissi-Poulou, M.: Enantiomeric separation of zopiclone, its metabolites and products of degradation on a  $\beta$ -cyclodextrin bonded phase. *J. Chromatogr. A*, 729 (1996) 19-28.
- 3594 Reepmeyer, J.C.: Separation of R- and S-thalidomide by reversed-phase HPLC with  $\beta$ -cyclodextrin in the mobile phase. *Chirality*, 8 (1996) 11-17; *C.A.*, 124 (1996) 212251m.
- 3595 Ring, B.J., Binkley, S.N., Roskos, L. and Wrighton, S.A.: Effect of fluoxetine, norfluoxetine, sertraline and desmethyl sertraline on human CYP3A catalyzed 1'-hydroxy midazolam formation *in vitro*. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1131-1135.
- 3596 Ring, B.J., Catlow, J., Lindsay, T.J., Gillespie, T., Roskos, L.K., Cerimele, B.J., Swanson, S.P., Hamman, M.A. and Wrighton, S.A.: Identification of the human cytochromes P450 responsible for the *in vitro* formation of the major oxidative metabolites of the antipsychotic agent olanzapine. *J. Pharmacol. Exp. Ther.*, 276 (1996) 658-666.

- 3597 Risley, D.S. and Sharp, V.S.: Evaluation of a new pepsin enzyme chiral stationary phase for the optimized separation of sepronetine (S-norfluoxetine) from R-norfluoxetine. *J. Liq. Chromatogr.*, 19 (1996) 449-465.
- 3598 Róna, K., Ary, K., Gachályi, B., Klebovich, I. and Tomori, É.: Simultaneous determination of nerisopam, a novel anxiolytic agent showing polymorphic metabolism, and its N-acetyl metabolite from human plasma by a validated high-performance liquid chromatographic method. *J. Chromatogr. B*, 678 (1996) 63-72.
- 3599 Rudenko, B.A., Savchuk, S.A. and Brodskii, E.S.: Chromatographic determination of narcotic anesthetics. *J. Anal. Chem.*, 51 (1996) 167-185; *C.A.*, 124 (1996) 212171k - a review with 62 refs.
- 3600 Schmid, J., Greenblatt, D.J., von Moltke, L.L., Harmatz, J.S. and Shader, R.I.: N-Demethylation of amitriptyline *in vitro*: Role of cytochrome P-450 3A (CYP3A) isoforms and effect of metabolic inhibitors. *J. Pharmacol. Exp. Ther.*, 275 (1995) 592-597.
- 3601 St-Pierre, M.V. and Pang, K.S.: Concentration-dependent metabolism of diazepam in mouse liver. *J. Pharmacokin. Biopharm.*, 23 (1995) 243-266.
- 3602 Svensson, J.-O. and Gustafsson, L.L.: Determination of ketamine and norketamine enantiomers in plasma by solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. B*, 678 (1996) 373-376.
- 3603 Takayasu, T., Kakubari, I., Fukamachi, A., Mafune, E., Takasugi, N., Takayama, K. and Nagai, T.: Determination of timiperone in rat plasma by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. B*, 679 (1996) 161-165.
- 3604 Tanaka, M., Kato, K., Sudo, K.-i., Hakusui, H., Ohkubo, T., Sugawara, K., Kaneko, S., Otani, K., Chiba, K. and Ishizaki, T.: Simple and selective assay of 4-hydroxymephenytoin in human urine using solid-phase extraction and high-performance liquid chromatography with electrochemical detection and its preliminary application to phenotyping test. *J. Chromatogr. B*, 676 (1996) 87-94.
- 3605 Tang, W., Borel, A.G. and Abbott, F.S.: Conjugation of glutathione with a toxic metabolite of valproic acid, (E)-2-propyl-2,4-pentadienoic acid, catalyzed by rat hepatic glutathione-S-transferases. *Drug Metab. Disp.*, 24 (1996) 436-446.
- 3606 Tsina, I., Chu, F., Kalooostian, M., Pettibone, M. and Wu, A.: HPLC method for the determination of ketorolac in human plasma. *J. Liq. Chromatogr.*, 19 (1996) 957-967.
- 3607 Vletter, A.A., Olieman, W., Burm, A.G.L., Groen, K. and van Kleef, J.W.: High-performance liquid chromatographic assay of mepivacaine enantiomers in human plasma in the nanogram per milliliter range. *J. Chromatogr. B*, 678 (1996) 369-372.
- 3608 Wong, Y.N., Quon, C.Y., Holm, K.A., Burcham, D.L., Frey, N.L., Huang, S.-M. and Lam, G.N.: Pharmacokinetics and metabolism of EXP921, a novel cognitive enhancer, in rats. *Drug Metab. Disp.*, 24 (1996) 172-179.
- 3609 Yamasaki, H., Kitamura, T. and Moriyama, H.: (Simultaneous determination of antiepileptic drugs using column switching HPLC). *Kuromatografi*, 16 (1995) 288-289; *C.A.*, 124 (1996) 185766z.
- 3610 Ye, L. and Stewart, J.T.: HPLC determination of an ondansetron and diphenhydramine mixture in 0.9% sodium chloride injection. *J. Liq. Chromatogr.*, 19 (1996) 711-718.
- 3611 Yu, Z. and Westerlund, D.: Direct injection of large volumes of plasma in a column-switching system for the analysis of local anaesthetics. I. Optimization of semi-permeable surface precolumns in the system and characterization of some interference peaks. *J. Chromatogr. A*, 725 (1996) 137-147.
- 3612 Yu, Z. and Westerlund, D.: Direct injection of large volumes of plasma in a column-switching system for the analysis of local anaesthetics. II. Determination of bupivacaine in human plasma with an alkyldiol silica precolumn. *J. Chromatogr. A*, 725 (1996) 149-155.
- 3613 Zarapkar, S.S. and Dhanvate, A.A.: Simultaneous determination of chlorzoxazone and paracetamol from its pharmaceutical preparations using high performance liquid chromatography. *Indian Drugs* 1995, 32 (1995) 465-468; *C.A.*, 124 (1996) 97878g.
- 3614 Zhang, D., Evans, F.E., Freeman, J.P., Duhart, B., Jr. and Cerniglia, C.E.: Biotransformation of amitriptyline by *Cunninghamella elegans*. *Drug Metab. Disp.*, 23 (1995) 1417-1425.
- 3615 Zhao, Z., Koepfinger, K.A., Bundy, G.L., Banitt, L.S., Padbury, G.E., Hauer, M.J. and Sanders, P.E.: *In vitro* and *in vivo* biotransformation of 6,7-dimethyl-2,4-di-1-pyrrolidinyl-7H-pyrrolo[2,3-D]pyrimidine (U-89843) in the rat. *Drug Metab. Disp.*, 34 (1996) 187-198.

For additional information see *C.A.*:

124 (1996) 44620k, 75229u, 135871j, 169671j.

See also 2484, 2573, 2902, 3475, 3493, 3498, 3500, 3670, 3708, 3729, 3737.

### 32e. Chemotherapeutics (exc. cytostatics and antibiotics)

- 3616 All-Deeb, O.A., Abdel-Moety, E.M., Abounassif, M.A. and Alzaben, S.R.: Stability-indicating high-performance liquid chromatographic method for determination of norfloxacin in bulk form and tablets. *Boll. Chim. Farm.* 1995, 134 (1995) 497-502; *C.A.*, 124 (1996) 185757x.
- 3617 Batty, K.T., Davis, T.M.E., Thu, L.T., Binh, T.Q., Anh, T.K. and Ilett, K.F.: Selective high-performance liquid chromatographic determination of artesunate and  $\alpha$ - and  $\beta$ -dihydroartemisinin in patients with falciparum malaria. *J. Chromatogr. B*, 677 (1996) 345-350.
- 3618 Breda, M., Marrari, P., Pianezzola, E. and Strolin Benedetti, M.: Determination of ethambutol in human plasma and urine by high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 729 (1996) 301-307.
- 3619 Chiba, M., Nishime, J.A. and Lin, J.H.: Potent and selective inactivation of human liver microsomal cytochrome P-450 isoforms by L-754,394, an investigational human immune deficiency virus protease inhibitor. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1527-1534.
- 3620 Easmon, J., Heinisch, G., Holzer, W. and Matusczak, B.: On the synthesis of pyridazomycin congeners. Part 2: L- $\alpha$ -Amino acids bearing a terminal azinium or diazinium system. *Pharmazie*, 51 (1996) 76-83.
- 3621 Foda, N.H.: High performance liquid chromatographic determination of nalidixic acid in tablets. *J. Liq. Chromatogr.*, 18 (1995) 4135-4147.

- 3622 Guyonnet, J., Pacaud, M., Richard, M., Doisi, A., Spavone, F. and Hellings, P.: Routine determination of flumequine in kidney tissue of pig using automated liquid chromatography. *J. Chromatogr. B*, 679 (1996) 177-184.
- 3623 Hormazabal, V., Steffenak, I. and Yndestad, M.: Simultaneous extraction and determination of florfenicol and the metabolite florfenicol amine in sediment by high-performance liquid chromatography. *J. Chromatogr. A*, 724 (1996) 364-366.
- 3624 Jessa, M.J., Barrett, D.A., Shaw, P.N. and Spiller, R.C.: Rapid and selective high-performance liquid chromatographic method for the determination of metronidazole and its active metabolite in human plasma, saliva and gastric juice. *J. Chromatogr. B*, 677 (1996) 374-379.
- 3625 Lau, Y.Y., Hanson, G.D. and Carel, B.J.: Determination of rifampin in human plasma by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 676 (1996) 147-152.
- 3626 Lin, C.-L., Hong, C.-C. and Kondo, F.: Simultaneous determination of residual sulfonamides in the presence and absence of *p*-aminobenzoic acid by high-performance liquid chromatography. *Microbios*, 83 (1995) 175-183; C.A., 124 (1996) 105444c.
- 3627 Malan, C.E.P., de Villiers, M.M., Handford, J.S. and Swanepoel, E.: HPLC determination of closantel in multi-component anthelmintic suspensions and tablets. *Pharmazie*, 51 (1996) 250.
- 3628 Mberu, E.K., Muhiia, D.K., Minyiri, G.O., Njonde, E.W. and Watkins, W.M.: Measurement of physiological concentrations of dapsone and its monoacetyl metabolite: a miniaturised assay for liquid or filter paper-absorbed samples. *J. Chromatogr. B*, 677 (1996) 385-387.
- 3629 Metz, R., Muth, P., Ferger, M., Bolten, W.W. and Vergin, H.: Improved determination of sulfadiazine in human plasma and urine by high-performance liquid chromatography. *J. Chromatogr. A*, 729 (1996) 243-249.
- 3630 Muth, P., Metz, R., Siems, B., Bolten, W.W. and Vergin, H.: Sensitive determination of nitrofurantoin in human plasma and urine by high-performance liquid chromatography. *J. Chromatogr. A*, 729 (1996) 251-258.
- 3631 Pal, R., Dwivedi, A.K., Kulshreshtra, D.K. and Singh, S.: HPLC determination method for antifilarial cardenolides in active fraction from *Streblus asper*. *Indian Drugs*, 32 (1995) 531-533; C.A., 124 (1996) 242491g.
- 3632 Plakas, S.M., El Said, K.R., Stehly, G.R. and Roybal, J.E.: Optimization of a liquid chromatographic method for determination of malachite green and its metabolites. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1388-1394.
- 3633 Pruñónosa, J., Solà, J., Peraire, C., Pla, F., Lavergne, O. and Obach, R.: Determination of the anti-platelet-activating factor BN-50727 and metabolites in human urine by high-performance liquid chromatography using solid-phase extraction. *J. Chromatogr. B*, 677 (1996) 388-392.
- 3634 Ricci, M.C. and Cross, R.F.: High performance liquid chromatographic analyses of sulphonamides and dihydrofolate reductase inhibitors. I. Separations in methanol-modified solutions. *J. Liq. Chromatogr.*, 19 (1996) 365-381.
- 3635 Ricci, M.C. and Cross, R.F.: High performance liquid chromatographic analyses of sulphonamides and dihydrofolate reductase inhibitors. II. Separations in acetonitrile modified solutions, ternary gradient studies & flow programming. *J. Liq. Chromatogr.*, 19 (1996) 547-564.
- 3636 Scalia, S., Guarneri, M. and Menegatti, E.: Assay of triclosan in deodorant sticks and soaps by supercritical fluid extraction and HPLC. *J. Soc. Cosmet. Chem.*, 45 (1994) 35-42; C.A., 124 (1996) 97230q.
- 3637 Shinde, V.M., Tendolkar, N.M. and Desai, B.S.: Simultaneous determination of nitrofurantoin and trimethoprim in capsules by HPLC. *Indian Drugs*, 32 (1995) 283-286; C.A., 124 (1996) 97885g.
- 3638 Sorensen, L.K. and Petersen, A.K.: Determination of thiabendazole and levamisole in milk by HPLC. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 304-305; C.A., 124 (1996) 143871z.
- 3639 Viñas, P., López Erroz, C., Campillo, N. and Hernández-Córdoba, M.: Determination of sulphonamides in foods by liquid chromatography with postcolumn fluorescence derivatization. *J. Chromatogr. A*, 726 (1996) 125-131.
- 3640 Wanwimolruk, S. and Pratt, E.L.: A simple HPLC assay for proguanil and its active metabolite cycloguanil: application to oxidation phenotyping. *J. Liq. Chromatogr.*, 18 (1995) 4097-4105.
- 3641 Wanwimolruk, S., Wong, S.M., Zhang, H. and Coville, P.F.: Simultaneous determination of quinine and a major metabolite 3-hydroxyquinine in biological fluids by HPLC without extraction. *J. Liq. Chromatogr.*, 19 (1996) 293-305.
- 3642 Zerilli, A., Lucas, D., Berthou, F., Bardou, L.G. and Ménez, J.-F.: Determination of cytochrome P 450 2E1 activity in microsomes by thin-layer chromatography using [2-<sup>14</sup>C]chlorzoxazone. *J. Chromatogr. B*, 677 (1996) 156-160.
- For additional information see C.A.:  
124 (1996) 66739h, 66743e.
- See also 3337, 3365, 3366, 3370, 3383, 3427, 3508.
- 32f. Cytostatics
- 3643 Braess, J., Pförtner, J., Kaufmann, C.C., Ramsauer, B., Unterhalt, M., Hiddemann, W. and Schleyer, E.: Detection and determination of the major metabolites of [<sup>3</sup>H]cytosine arabinoside by high-performance liquid chromatography. *J. Chromatogr. B*, 676 (1996) 131-140.
- 3644 Chimonczyk, Z., Kscinska, H. and Cibulsky, J.: Direct determination of the enantiomeric purity of (R)-(+)-aminoglutethimide and its main metabolite. *Arch. Pharm. (Weinheim)*, 328 (1995) 745-749; C.A., 124 (1996) 127242q.
- 3645 Compagnon, P., Thiberville, L., Moore, N., Thuiliez, C. and Lacroix, C.: Simple high-performance liquid chromatographic method for the quantitation of 5-fluorouracil in human plasma. *J. Chromatogr. B*, 677 (1996) 380-383.
- 3646 Cserháti, T. and Forgács, E.: Use of principal component analysis for the study of the retention behaviour of anticancer drugs on a  $\beta$ -cyclodextrin polymer-coated silica column. *J. Chromatogr. A*, 728 (1996) 67-73.
- 3647 Fuchs, W.S., Leary, W.P., van der Meer, M.J., Gay, S., Witschital, K. and von Nieciecki, A.: Pharmacokinetics and bioavailability of tamoxifen in postmenopausal healthy women. *Arzneim.-Forsch.*, 46 (1996) 418-422.

- 3648 Hsu, L.C.: High performance reverse phase liquid chromatographic method for topotecan trace analysis. *J. Liq. Chromatogr.*, 19 (1996) 611-624.
- 3649 Jarugula, V.R. and Boudinot, F.D.: High-performance liquid chromatographic determination of 5-fluorouracil and its pro-drugs, tegafur and 4-deoxy-5-fluorouracil, in rat plasma. *J. Chromatogr. B*, 677 (1996) 199-203.
- 3650 Keuzenkamp-Jansen, C.W., van Baal, J.M., de Abreu, R.A., de Jong, J.G.N., Zuidenr, R. and Trijbels, J.M.F.: Detection and identification of 6-methylmercapto-8-hydroxypurine, a major metabolite of 6-mercaptopurine, in plasma during intravenous administration. *Clin. Chem. (Washington)*, 42 (1996) 380-386.
- 3651 Loos, W.J., Stoter, G., Verweij, J. and Schellens, J.H.M.: Sensitive high-performance liquid chromatographic fluorescence assay for the quantitation of topotecan (SKF 104864-A) and its lactone ring-opened product (hydroxy acid) in human plasma and urine. *J. Chromatogr. B*, 678 (1996) 309-315.
- 3652 MacCallum, J., Cummings, J., Dixon, J.M. and Miller, W.R.: Solid-phase extraction and high-performance liquid chromatographic determination of tamoxifen and its major metabolites in plasma. *J. Chromatogr. B*, 678 (1996) 317-323.
- 3653 Marrari, P., Pianezzola, E. and Benedetti, M.S.: Determination of tallimustine in human plasma by high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 133-139.
- 3654 Martinsen, A. and Gynther, J.: Liquid chromatography-thermospray mass spectrometry of toremifene and its derivatives. *J. Chromatogr. A*, 724 (1996) 358-363.
- 3655 Pérez-Urizar, J., Picazo, Y.F., Navarro-González, B., Flores-Murieta, F.J. and Castaneda-Hernández, G.: A new rapid and economical high performance liquid chromatographic assay with electrochemical detection for the determination of etoposide (VP-16) in human plasma samples. *J. Liq. Chromatogr.*, 19 (1996) 939-947.
- 3656 Rao, K.V. and Bhakuni, R.S.: A large scale process for paclitaxel and other taxanes from the needles of *Taxus media* Hickssii and *Taxus floridana* using reverse phase column chromatography. *J. Liq. Chromatogr.*, 19 (1996) 427-447.
- 3657 Rentsch, K.M., Schwendener, R.A. and Hänseler, E.: Determination of mitoxantrene in mouse whole blood and different tissues by high-performance liquid chromatography. *J. Chromatogr. B*, 679 (1996) 185-192.
- 3658 Royer, I., Monserrat, B., Sonnier, M., Wright, M. and Cresteil, T.: Metabolism of docetaxel by human cytochromes P450: interactions with paclitaxel and other antineoplastic drugs. *Cancer Res.*, 56 (1996) 58-65.
- 3659 Song, D., Hsu, L.-F. and Au, J.L.-S.: Binding of taxol to plastic and glass containers and protein under *in vitro* conditions. *J. Pharm. Sci.*, 85 (1996) 29-31.
- 3660 Sonnichsen, D.S., Liu, Q., Schuetz, J.D., Pappo, A. and Relling, M.V.: Variability in human cytochrome P450 paclitaxel metabolism. *J. Pharmacol. Exp. Ther.*, 275 (1995) 566-575.
- 3661 Studenberg, S.D., DeAngelis, D.V. and Woolley, J.L.: High performance liquid chromatographic assay with fluorescence detection for the analysis of 1954U89, 1,3-diamino-7-(1-ethyl-propyl)-8-methyl-7H-pyrrolo-(3,2-f)-quinazoline, in plasma. *J. Liq. Chromatogr.*, 19 (1996) 823-836.
- 3662 Supko, J.G., Phillips, L.R. and Malspeis, L.: Specific high-performance liquid chromatographic assay with ultraviolet detection for the determination of 1-(2-chloroethyl)-3-sarcosinamide-1-nitrosourea in plasma. *J. Chromatogr. B*, 677 (1996) 351-362.
- 3663 Volkov, S.K. and Grodnitskaya, E.I.: (Methods for increasing the effectiveness of the chromatographic purification of vinblastine). *Khim.-Farm. Zh.*, 29 (1995) 58-59; *C.A.*, 124 (1996) 185511n.
- 3664 Wyss, R., Bucheli, F. and Hess, B.: Determination of the arotinoid mofarotene in human, rat and dog plasma by high-performance liquid chromatography with automated column switching and ultraviolet detection. *J. Chromatogr. A*, 729 (1996) 315-322.
- 3665 Yamazoe, K., Horiuchi, T., Sugiyama, T. and Katagiri, Y.: Simultaneous high-performance liquid chromatographic determination of carboplatin, epirubicin hydrochloride and mitomycin C in a Lipiodol emulsion. *J. Chromatogr. A*, 726 (1996) 241-245.

For additional information see *C.A.*:

124 (1996) 44618r, 105448j, 156238y.

See also 3236, 3269, 3308, 3363, 3367, 3731.

### 32g. Other drug categories

- 3666 Abdel-Moety, E.M., Al-Deeb, O.A. and Khattab, N.A.: Determination of dextromethorphan hydrobromide in bulk form and dosage formulations by high performance liquid chromatography. *J. Liq. Chromatogr.*, 18 (1995) 4127-4134.
- 3667 Al-Dhawailie, A.A., Abdulaziz, M.A., Tekle, A. and Matar, K.M.: A simple, specific, and rapid high performance liquid chromatographic assay for glibenclamide in plasma. *J. Liq. Chromatogr.*, 18 (1995) 3981-3990.
- 3668 Albertoni, F., Pettersson, B., Ohlman, S. and Peterson, C.: Analysis of azathioprine and 6-mercaptopurine in plasma in renal transplant recipients after administration with oral azathioprine. *J. Liq. Chromatogr.*, 18 (1995) 3991-4005.
- 3669 Ashton, D.S., Beddell, C.R., Cockerill, G.S., Gohil, K., Gowrie, C., Robinson, J.E., Slater, M.J. and Valko, K.: Binding measurements of indolocarbazole derivatives to immobilised human serum albumin by high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 194-198.
- 3670 Barbato, F., La Rotunda, M.I. and Quaglia, F.: Retention behavior of anti-emetic serotonin antagonists in reversed-phase high-performance liquid chromatography. *Farmaco*, 50 (1995) 875-880; *C.A.*, 124 (1996) 127271y.
- 3671 Bergstrom, K.A., Halldin, C., Kuikka, J.T., Swahn, C.-G., Tihonen, J., Hiltunen, J., Lansmies, E. and Farde, L.: The metabolite pattern of [ $^{123}\text{I}$ ]β-CIT determined with a gradient HPLC method. *Nucl. Med. Biol.*, 22(8, New Radio-Tracers and Methods of Quality Assurance for Nuclear Medicine Applications) (1995) 971-976; *C.A.*, 124 (1996) 169412a.
- 3672 Boppana, V.K., Miller-Stein, C. and Schaefer, W.H.: Direct plasma liquid chromatographic-tandem mass spectrometric analysis of granisetron and its 7-hydroxy metabolite utilizing internal surface reversed-phase guard columns and automated column switching devices. *J. Chromatogr. B*, 678 (1996) 227-236.

- 3673 Budzikiewicz, H. and Spahl, W.: (Quality control of dental resin composites by gas chromatography and liquid chromatography-mass spectrometry). *GIT Fachz. Lab.*, 39 (1995) 1043-1044; C.A., 124 (1996) 127050a.
- 3674 Castelnovo, P.: Determination of the calcium antagonist SIM6080 and its N- and O-demethylated metabolites in plasma, urine and tissues by high-performance liquid chromatography. *J. Chromatogr. B*, 676 (1996) 77-85.
- 3675 Chandrasekaran, A., Osman, M., Adelman, S.J., Warsheski, J., Scatina, J.A. and Sisenwine, S.F.: Determination of 17 $\alpha$ -dihydroequilenin in rat, rabbit and monkey plasma by high-performance liquid chromatography with fluorimetric detection. *J. Chromatogr. B*, 676 (1996) 69-75.
- 3676 Chou, H.J., Yates, R.L., Haverty, D.C. and Wenninger, J.A.: Determination of 2-ethylhexyl-4-(N-methyl-N-nitrosamino)benzoate in commercial sunscreen and cosmetic products. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1378-1383.
- 3677 Dasgupta, A. and Timmerman, T.G.: Therapeutic monitoring of Tacrolimus (FK 506) using liquid chromatography. A review. *J. Liq. Chromatogr.*, 18 (1995) 4029-4038 - a review with 32 refs.
- 3678 Ebete, K.O. and Koundourellis, J.E.: Determination of diphenylpyraline in plasma and urine by high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 319-323.
- 3679 Ferretti, R., Gallinella, B., la Torre, F. and Lusi, A.: Direct high-performance liquid chromatography resolution on a chiral column of dextfenfluramine and its impurities, in bulk raw drug and pharmaceutical formulations. *J. Chromatogr. A*, 731 (1996) 340-345.
- 3680 Franklin, M. and Odontiadis, J.: Determination of naloxone in human plasma by high-performance liquid chromatography with coulometric detection. *J. Chromatogr. B*, 679 (1996) 199-203.
- 3681 Ghanbarpour, A. and Amini, M.: Analysis of dexamethasone sodium phosphate injection and ophthalmic solution by HPLC, kinetic interpretation and determination of shelf-life. *J. Sci. Islamic Repub. Iran.*, 6 (1995) 226-230; C.A., 124 (1996) 211816n.
- 3682 Granneman, G.R., Braeckman, R.A. and Erdman, K.A.: Determination of a new 5-lipoxygenase inhibitor, zileuton, and its inactive N-dehydroxylated metabolite in plasma by high performance liquid chromatography. *Clin. Pharmacokinet.*, 29(Suppl. 2) (1995) 1-8; C.A., 124 (1996) 164154h.
- 3683 Grubb, M.F., Robinson, C.A., White, S.J., Krauthauer, C.L. and Shum, L.: Determination of a HIV protease inhibitor (DMP 450) in animal and human plasma by solid-phase extraction and high-performance liquid chromatography. *J. Chromatogr. B*, 678 (1996) 303-308.
- 3684 Hannan, S.L., Ridout, G.A. and Jones, A.E.: Determination of the potent antiprotozoal compound atovaquone in plasma using liquid-liquid extraction followed by reversed-phase high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 678 (1996) 297-302.
- 3685 King, L.E. and Vieth, R.: Extraction and measurement of pamidronate from bone samples using automated pre-column derivatization, high-performance liquid chromatography and fluorescence detection. *J. Chromatogr. B*, 678 (1996) 325-330.
- 3686 Kourany-Lefoll, E. and Cyr, T.D.: Determination of acyclovir (Zovirax) and quanine by microbore high-performance liquid chromatography with confirmation by atmospheric-pressure chemical ionization mass spectrometry. *Can. J. Appl. Spectrosc.*, 40 (1995) 155-159; C.A., 124 (1996) 185779f.
- 3687 Lampen, A., Christians, U., Guengerich, F.P., Watkins, P.B., Kolars, J.C., Bader, A., Gonschior, A.-K., Dralle, H., Hackbart, I. and Sewing, K.-F.: Metabolism of the immunosuppressant Tacrolimus in the small intestine: cytochrome P450, drug interactions, and interindividual variability. *Drug Metab. Disp.*, 23 (1995) 1315-1324.
- 3688 Lau, Y.Y., Hanson, G.D. and Carel, B.J.: Determination of rifabutin in human plasma by high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 676 (1996) 125-130.
- 3689 Lau-Cam, C., Theofanopoulos, V. and Spireas, S.S.: Simplified HPLC method with spectrophotometric detection for the assay of clofibric acid in rat plasma. *J. Liq. Chromatogr.*, 18 (1995) 3945-3954.
- 3690 Loun, B. and Hage, D.S.: Chiral separation mechanisms in protein-based HPLC columns. 2. Kinetic studies of (R)- and (S)-warfarin binding to immobilized human serum albumin. *Anal. Chem.*, 68 (1996) 1218-1225.
- 3691 Matsuoka, M., Banno, K. and Sato, T.: Analytical chiral separation of a new quinolone compound in biological fluids by high-performance liquid chromatography. *J. Chromatogr. B*, 676 (1996) 117-124.
- 3692 Merdink, J., Decosta, K., Mathews, J.M., Jones, C.B., Okita, J.R. and Okita, R.T.: Hydroxylation of lauramide diethanolamine by liver microsomes. *Drug Metab. Disp.*, 24 (1996) 180-186.
- 3693 Mikami, E., Itoh, Y., Ohno, T. and Hayakawa, J.: (Rapid determination of drugs in pharmaceutical preparations by liquid chromatography VII. Determination of eprazinone hydrochloride, pirenzepine hydrochloride, valethamate bromide, mepenzolate bromide, traniast and ketotifen fumarate in pharmaceutical preparations). *Iyakuhin Kenkyu*, 26 (1995) 695-700; C.A., 124 (1996) 97870y.
- 3694 Oka, H., Ikai, Y., Hayakawa, J., Harada, K.-i., Suzuki, M., Shimizu, A., Hayashi, T., Takeba, K., Nakazawa, H. and Ito, Y.: Separation of ivermectin components by high-speed counter-current chromatography. *J. Chromatogr. A*, 723 (1996) 61-68.
- 3695 Okazaki, O., Kurata, T., Yoshioka, N. and Hakusui, H.: Pharmacokinetics and stability of caldiamide sidum in rats. *Arzneim.-Forsch.*, 46 (1996) 79-83.
- 3696 Oliveros, L., Minguillón, C., Serkiz, B., Meunier, F., Volland, J.-P. and Cordi, A.A.: Preparative isolation of the eutomer of cyclothiazide by high-performance liquid chromatography on a cellulose-derived chiral stationary phase with toluene-acetone as the mobile phase. *J. Chromatogr. A*, 729 (1996) 29-32.
- 3697 Ono, I., Matsuda, K. and Kanno, S.: Determination of N-(trans-4-isopropylcyclohexylcarbonyl)-D-phenylalanine in human plasma by solid-phase extraction and column-switching high-performance liquid chromatography with ultraviolet detection. *J. Chromatogr. B*, 678 (1996) 384-387.
- 3698 Ono, Y., Noda, A., Zaima, Y., Jitsufuchi, N., Eto, S. and Noda, H.: Determination of isonicotinic acid in the presence of isoniazid and acetylisoniazid. Studies on isonicotinic acid formation from isoniazid in isolated rat hepatocytes. *J. Chromatogr. B*, 677 (1996) 339-343.

- 3699 Oyler, A.R., Armstrong, B.L., Cha, J.Y., Zhou, M.X., Yang, Q., Robinson, R.I., Dunphy, R. and Burinsky, D.J.: Hydrophilic interaction chromatography on amino-silica phases complements reversed-phase high-performance liquid chromatography and capillary electrophoresis for peptide analysis. *J. Chromatogr. A*, 724 (1996) 378-383.
- 3700 Palwal, J.K. and Gupta, R.C.: Tissue distribution and pharmacokinetics of centchroman. A new nonsteroidal postcoital contraceptive agent and its 7-desmethyl metabolite in female rats after a singler oral dose. *Drug Metab. Disp.*, 24 (1996) 148-155.
- 3701 Prabhu, S.V., Cooper, S.E., Mathre, D.J., Natishan, T.K., Ellison, D.K. and Twy, P.C.: High-performance, purity-indicating liquid chromatographic method for the oxytocin antagonist, L-368,899. *J. Chromatogr. A*, 731 (1996) 331-335.
- 3702 Richter, K., Oertel, R. and Kirch, W.: New sensitive method for the determination of hydrochlorothiazide in human serum by high-performance liquid chromatography with electrochemical detection. *J. Chromatogr. A*, 729 (1996) 293-296.
- 3703 Rossi, D.T., Hallak, H. and Bradford, L.: Liquid chromatographic assay for a butenolide endothelin antagonist (PD 156707) in plasma. *J. Chromatogr. B*, 677 (1996) 299-304.
- 3704 Sanwald, P., David, M. and Dow, J.: Use of electrospray ionization liquid chromatography-mass spectrometry to study the role of CYP2D6 in the *in vitro* metabolism of 5-hydroxytryptamine receptor antagonists. *J. Chromatogr. B*, 678 (1996) 53-61.
- 3705 Schall, R., Müller, F.O., Hundt, H.K.L., Duursema, L., Groenewoud, G. and Middle, M.V.: Study of the effect of miglitol on the pharmacokinetics and pharmacodynamics of warfarin in healthy males. *Arzneim.-Forsch.*, 46 (1996) 41-46.
- 3706 Shishoo, C.J., Jain, K.S., Rathod, I.S., Thakkar, B.J., Brambhatt, S.B., Gandhi, T.P., Bangaru, R. and Goyal, R.K.: Synthesis and quantitative structure-activity relationships of antihyperlipaemic 2-substituted thieno[3,2-d]pyrimidin-4(3H)-ones. *Arzneim.-Forsch.*, 46 (1996) 273-276.
- 3707 Somekh, V. and Stolk, L.M.L.: (Determination of laxatives). *Ziekenhuisfarmacie*, 11 (1995) 193-195; *C.A.*, 124 (1996) 193136j.
- 3708 Srinivas, J.S., Marwah, A.K. and Rao, B.E.: Paired-ion HPLC determination of ondansetron in pure and dosage form. *East Pharm.* 1995, 38 (1995) 149-150; *C.A.*, 124 (1996) 66754j.
- 3709 Stancato, F.A., Srinivas, N.R. and Knupp, C.A.: Effect of temperature on the high performance liquid chromatographic separation of the anti-HIV agents, didanosine and stavudine. *Biomed. Chromatogr.*, 10 (1996) 29-31; *C.A.*, 124 (1996) 185768b.
- 3710 Sun, S.F., Hssiao, C.L. and Bolton, S.: HPLC study of the binding of sulfapyrazone to serum albumin. *Chromatographia*, 42 (1996) 401-404.
- 3711 Swanepoel, E., de Villiers, M.M. and du Preez, J.L.: Fluorimetric method of analysis for D-norpseudoephedrine hydrochloride, glycine and L-glutamic acid by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A*, 729 (1996) 287-291.
- 3712 Takano, T. and Hata, S.: High-performance liquid chromatographic determination of finasteride in human plasma using direct injection with column switching. *J. Chromatogr. B*, 676 (1996) 141-146.
- 3713 Tanaka, M., Ishikawa, F. and Hakusui, H.: Isolation and identification of seven metabolites of a water-soluble platelet aggregation inhibitor in rat urine. *Xenobiotica*, 25 (1995) 1247-1257; *C.A.*, 124 (1996) 44698s.
- 3714 Taylor, P.J., Jones, A., Balderson, G.A., Lynch, S.V., Norris, R.L.G. and Pond, S.M.: Sensitive, specific quantitative analysis of tacrolimus (FK506) in blood by liquid chromatography-electrospray tandem mass spectrometry. *Clin. Chem. (Washington)*, 42 (1996) 279-285.
- 3715 Tleugabulova, D. and González Pérez, I.: Reversed-phase high-performance liquid chromatographic study of thimerosal stability in Cuban recombinant hepatitis B vaccine. *J. Chromatogr. A*, 729 (1996) 219-227.
- 3716 Toreson, H. and Eriksson, B.-M.: Determination of fluvastatin enantiomers and the racemate in human blood plasma by liquid chromatography and fluorometric detection. *J. Chromatogr. A*, 729 (1996) 13-18.
- 3717 Van Tonder, E.C., de Villiers, M.M., Handford, J.S., Malan, C.E.P. and du Preez, J.L.: Simple, robust and accurate high-performance liquid chromatography method for the analysis of several anthelmintics in veterinary formulations. *J. Chromatogr. A*, 729 (1996) 267-272.
- 3718 Woolf, E.J. and Matuszewski, B.: Simultaneous determination of L-702,007, a HIV-1 reverse transcriptase inhibitor, and its 6-hydroxy metabolite in human plasma by column-switching high-performance liquid chromatography with fluorescence detection. *J. Chromatogr. A*, 729 (1996) 211-217.
- 3719 Yamashita, K., Murakami, H., Okuda, T. and Motohashi, M.: High-performance liquid chromatographic determination of pioglitazone and its metabolites in human serum and urine. *J. Chromatogr. B*, 677 (1996) 141-146.
- 3720 Yamato, S., Nakajima, M. and Shimada, K.: Simultaneous determination of chlorpheniramine and maleate by high-performance liquid chromatography using tetra-n-butylammonium phosphate as an ion-pair reagent. *J. Chromatogr. A*, 731 (1996) 346-350.

For additional information see *C.A.*:

124 (1996) 193161p, 197366g.

See also 2813, 2854, 2921, 2999, 3264, 3282, 3320, 3359, 3435, 3619, 3636, 3736, 3753, 3770.

### 32h. Toxicological and forensic applications

- 3721 Achilli, G., Cellerino, G.P., Melzi d'Erl, G.V. and Tagliaro, F.: Determination of illicit drugs and related substances by high-performance liquid chromatography with an electrochemical coulometric-array detector. *J. Chromatogr. A*, 729 (1996) 273-277.
- 3722 Akasaka, K., Ohrui, H., Meguro, H and Yasumoto, T.: Fluorimetric determination of diarrhetic shellfish toxins in scallops and mussels by high-performance liquid chromatography. *J. Chromatogr. A*, 729 (1996) 381-386.
- 3723 Binder, S.R.: Analysis of drugs of abuse in biological fluids by liquid chromatography. *Adv. Chromatogr.*, 36 (1996) 201-271; *C.A.*, 124 (1996) 167568g - a review with 413 refs.

- 3724 Carmody, E.P., James, K.J. and Kelly, S.S.: Diarrhetic shellfish poisoning: evaluation of enzyme-linked immunosorbent assay methods for determination of dinophysistoxin-2. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1403-1408.
- 3725 Hummert, C., Shen, J.L. and Luckas, B.: Automatic high-performance liquid chromatographic method for the determination of diarrhetic shellfish poison. *J. Chromatogr. A*, 729 (1996) 387-392.
- 3726 Huopalaiti, R.P. and Henion, J.D.: Application of supercritical fluid extraction and high performance liquid chromatography/mass spectrometry for the determination of some anabolic agents directly from bovine tissue samples. *J. Liq. Chromatogr.*, 19 (1996) 69-87.
- 3727 Jung, J.H., Sim, C.J. and Lee, C.-O.: Cytotoxic compounds from a two-sponge association. *J. Natural Prod.*, 58 (1995) 1722-1726.
- 3728 Kuijpers, E.A.P., den Hartigh, J., Savelkoul, J.F. and de Wolff, F.A.: A method for the simultaneous identification and quantitation of five superwarfarin rodenticides in human serum. *J. Anal. Toxicol.*, 19 (1995) 557-562.
- 3729 Tyrefors, N., Hyllbrant, B., Ekman, L., Johansson, M. and Långström, B.: Determination of morphine, morphine-3-glucuronide and morphine-6-glucuronide in human serum by solid-phase extraction and liquid chromatography-mass spectrometry with electrospray ionisation. *J. Chromatogr. A*, 729 (1996) 279-285.
- 3730 Van Boeckelaer, J., Meyer, E., Clauwaert, K., Lambert, W., Piette, M. and de Leenheer, A.: Analysis of zopiclone (Imovane) in postmortem specimens by GC-MS and HPLC with diode-array detection. *J. Anal. Toxicol.*, 20 (1996) 52-54; *C.A.*, 124 (1996) 109144e.

For additional information see *C.A.*:

124 (1996) 163952e.

See also 2484, 2600, 2673, 2736, 2902, 2973, 3070, 3557, 3577, 3798, 3825.

### 32i. Plant extracts

- 3731 Aboul-Enein, H.Y. and Serignese, V.: Liquid chromatographic determination of taxol and related derivatives using a new poly-fluorinated reversed-phase column. *Anal. Chim. Acta*, 319 (1996) 187-190.
- 3732 Bastos, J.K., Burandt, C.L., Jr., Nanayakkara, N.P.D., Bryant, L. and McChesney, J.D.: Quantitation of aryl tetralin lignans in plant parts and among different populations of *Podophyllum peltatum* by reversed-phase high-performance liquid chromatography. *J. Nat. Prod.*, 59 (1996) 406-408; *C.A.*, 124 (1996) 242433q.
- 3733 Constant, H.L., Cordell, G.A., West, D.P. and Johnson, J.H.: Separation and quantification of capsaicinoids using complexation chromatography. *J. Natural Prod.*, 58 (1995) 1925-1928.
- 3734 Lin, S.-J., Tseng, H.-H., Wen, K.-C. and Suen, T.-T.: Determination of gentiopicroside, mangiferin, palmatine, berberine, baicalin, wogonin and glycyrrhizin in the traditional Chinese medicinal preparation Sann-Joong-Kuey-Jian-Tang by high-performance liquid chromatography. *J. Chromatogr. A*, 730 (1996) 17-23.

- 3735 Martínez Galera, M., Martínez Vidal, J.L., Garrido Frenich, A. and Gil García, M.D.: Determination of cypermethrin, fenvalerate and *cis*- and *trans*-permethrin in soil and groundwater by high-performance liquid chromatography using partial least-squares regression. *J. Chromatogr. A*, 727 (1996) 39-46.
- 3736 Micali, G., Lanuzza, F. and Currò, P.: High-performance liquid chromatographic determination of the biologically active principle hypericin in phytotherapeutic vegetable extracts and alcoholic beverages. *J. Chromatogr. A*, 731 (1996) 336-339.
- 3737 Pal, S.C., Kasture, S.B., Kasture, V.S. and Chatterjee, B.P.: Separation and estimation of ephedrine by HPLC. *East Pharm.*, 38 (1995) 157-158; *C.A.*, 124 (1996) 97900h.
- 3738 Wolfender, J.-L. and Hostettmann, K.: Applications of liquid chromatography-mass spectrometry to the investigation of medicinal plants. *Recent Adv. Phytochem.*, 29 (1995) 189-215; *C.A.*, 124 (1996) 197639y.

For additional information see *C.A.*:

124 (1996) 49830p, 111378c, 126771z, 127225m, 212188w.

See also 2811, 2885, 2886, 3268, 3269, 3656.

### 33. CLINICO-CHEMICAL APPLICATIONS

#### 33a. General papers and reviews

See 2834.

#### 33b. Complex mixtures and profiling (single compounds by cross-reference only)

- 3739 Frohm, M., Gunne, H., Bergman, A.-C., Agerberth, B., Bergman, T., Boman, A., Lidén, S., Jörnvall, H. and Boman, H.G.: Biochemical and antibacterial analysis of human wound and blister fluid. *Eur. J. Biochem.*, 237 (1996) 86-92.

See also 2726, 2757, 2802, 2820, 2848, 2872, 2879, 2932, 2942, 2945, 2953, 2959, 2961, 3027, 3053, 3072, 3076, 3104, 3222, 3225, 3238, 3244, 3323, 3823.

### 34. FOOD ANALYSIS

#### 34a. General papers and reviews

- 3740 Careri, M., Mangia, A. and Musci, M.: Applications of liquid chromatography-mass spectrometry interfacing systems in food analysis: pesticide, drug and toxic substance residues. *J. Chromatogr. A*, 727 (1996) 153-184 - a review with 162 refs.
- 3741 Romero, C., Perez-Andujar, O., Olmedo, A. and Jimenez, S.: Detection of cow's milk in ewe's or goat's milk by HPLC. *Chromatographia*, 42 (1996) 181-184.

See also 3316, 3392.

## 34b. Complex mixtures (single compounds by cross-reference only)

- 3742 Biedermann, M., Grob, K. and Bronz, M.: Determination of stigmastadiene in edible oils by HPLC-HPLC-UV. *Riv. Ital. Sostanze Grasse*, 72 (1995) 397-401; C.A., 124 (1996) 143884f.
- 3743 Brauer, B., Funke, T. and Schulenberg-Schell, H.: (Detection and determination of stabilizers in rigid PVC). *Dtsch. Lebensm.-Rundsch.*, 91 (1995) 381-385; C.A., 124 (1996) 200428m.
- 3744 Chen, B.H. and Fu, S.C.: Comparison of extraction methods and column types for the determination of additives by liquid chromatography. *J. Liq. Chromatogr.*, 19 (1996) 625-643.
- 3745 Ehlers, D., Pfister, M., Bork, W.-R. and Toffel-Nadolny, P.: HPLC analysis of tonka bean extracts. *Z. Lebensm.-Unters. Forsch.*, 201 (1995) 278-282; C.A., 124 (1996) 53939y.
- 3746 Osterneyer, U.: (Determination of sweeteners and preservatives in fish products by HPLC). *Dtsch. Lebensm.-Rundsch.*, 91 (1995) 307-311; C.A., 124 (1996) 143856y.
- 3747 Robb, E.J., Kausche, F.M. and Belschner, A.P.: Methods for detection of residues in milk after parenteral treatment with ceftiofur sodium - an evaluation using common milk residue screening tests and a quantitative HPLC method following either a single injection (Study I) or multiple injections (Study II) of 2.2 mg of ceftiofur free acid equivalents per kg of body weight. *Int. Dairy Fed. Spec. Issue*, 9505(Symposium on Residues of Antimicrobial Drugs and Other Inhibitors in Milk, 1995) (1995) 195-200; C.A., 124 (1996) 115679z.
- 3748 Sato, K.: (Application of pseudo-moving bed chromatographic separation apparatus to food manufacture). *Kagaku Sochi*, 38 (1996) 56-60; C.A., 124 (1996) 200342d - a review with 7 refs.

See also 2532, 2688, 2703, 2705, 2707, 2708, 2711, 2712, 2714, 2722, 2723, 2729, 2730, 2731, 2732, 2734, 2735, 2748, 2749, 2752, 2756, 2764, 2771, 2805, 2806, 2810, 2846, 2889, 2890, 2903, 2907, 2920, 2956, 2972, 3078, 3085, 3086, 3273, 3286, 3301, 3307, 3325, 3334, 3336, 3338, 3339, 3342, 3344, 3357, 3360, 3372, 3374, 3384, 3385, 3386, 3393, 3395, 3399, 3427, 3428, 3430, 3437, 3439, 3444, 3479, 3535, 3632, 3638, 3639, 3722, 3725, 3751, 3752, 3755, 3785, 3786, 3824, 3835.

## 34c. Organoleptically important compounds (flavors, odors, volatiles)

- 3749 Galletti, G.C. and Bocchini, P.: High-performance liquid chromatography with electrochemical detection of aspartame with a post-column photochemical reactor. *J. Chromatogr. A*, 729 (1996) 393-398.
- 3750 Langourieux, S. and Crouzet, J.: Determination of aroma compounds.  $\beta$ -Cyclodextrin complex formation constants by dynamic coupled column liquid chromatography. *Analisis*, 23 (1995) 333-336; C.A., 124 (1996) 163817q.
- 3751 Lübbe, M., le Quéré, J.-L. and Barron, D.: Prefractionation of aroma extracts from fat-containing food by high-performance size-exclusion chromatography. *J. Chromatogr. A*, 729 (1996) 371-379.
- 3752 Primus, T.M., Johnston, J.J. and Griffin, D.L.: Solid phase extraction/high performance liquid chromatography method for the determination of methyl anthranilate residues in blueberries. *J. Liq. Chromatogr.*, 19 (1996) 393-402.

- 3753 Tachibana, M., Aoyama, M., Taniguchi, R., Anabuki, K. and Kamata, K.: (Determination of cyclamate in foods by high-performance liquid chromatography with conductivity detection). *Tokyo-to Suginami-ku Eisei Shikensho Nenpo*, 13 (1995) 70-73; C.A., 124 (1996) 53948a.

For additional information see C.A.:  
124 (1996) 53949b.

## 35. ENVIRONMENTAL ANALYSIS

## 35a. General papers and reviews

- 3754 Slobodnik, J., Hogenboom, A.C., Louter, A.J.H. and Brinkman, U.A.T.: Integrated system for on-line gas and liquid chromatography with a single mass spectrometric detector for the automated analysis of environmental samples. *J. Chromatogr. A*, 730 (1996) 353-371.

See also 2697, 3402, 3424, 3822, 3841.

## 35b. Air pollution (complex mixtures; single compounds by cross-reference only)

See 2715, 2743, 2896, 3821.

## 35c. Water pollution (complex mixtures; single compounds by cross-reference only)

- 3755 Moret, S., Amici, S., Bortolomeazzi, R. and Lercker, G.: Determination of polycyclic aromatic hydrocarbons in water and water-based alcoholic beverages. *Z. Lebensm.-Unters. Forsch.*, 201 (1995) 322-326; C.A., 124 (1996) 174178h.

- 3756 Oleksy-Frenzel, J. and Jekel, M.: On-line multicomponent determination of organic compounds in water following gel permeation chromatographic separation. *Anal. Chim. Acta*, 319 (1996) 165-175.

- 3757 Oleksy-Frenzel, J., Wischnack, S. and Jekel, M.: (Determination of organic group parameters AOCl, AOBr and AOI in municipal waste water). *Vom Wasser*, 85 (1995) 59-67; C.A., 124 (1996) 184490t.

- 3758 Sundd, S., Prasad, B.B.: Evaluation of metal fractions in river sediments and waters: application of chelating chromatography-differential pulse anodic stripping voltammetry. *Talanta*, 42 (1995) 1395-1409; C.A., 124 (1996) 65859k.

For additional information see C.A.:  
124 (1996) 211310t.

- See also 2468, 2493, 2502, 2505, 2525, 2675, 2679, 2681, 2682, 2706, 2710, 2716, 2717, 2718, 2721, 2821, 2894, 2909, 3313, 3394, 3397, 3398, 3400, 3407, 3408, 3409, 3410, 3413, 3415, 3418, 3419, 3420, 3426, 3432, 3443, 3735, 3759, 3790, 3803, 3806, 3808, 3813, 3814, 3818, 3829, 3832.

35d. *Soil pollution (complex mixtures; single compounds by cross-reference only)*

See 2826, 3313, 3404, 3411, 3416, 3421, 3735, 3829.

### 36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

See 2888.

#### 36a. Surfactants

3759 Chee, K.K., Wong, M.K. and Lee, H.K.: Optimization of sample preparation techniques for the determination of 4-nonylphenol in water and sediment. *J. Liq. Chromatogr.*, 19 (1996) 259-275.

3760 Desbène, P.L., Portet, F.I. and Goussot, G.J.: Quantitative trace analysis of surfactant mixtures by reversed-phase high-performance liquid chromatography with refractometric detection. *J. Chromatogr. A*, 730 (1996) 209-218.

3761 Fernandez, P., Alder, A.C., Suter, M.J.-F. and Giger, W.: Determination of the quaternary ammonium surfactant ditallowdimethylammonium in digested sludges and marine sediments by supercritical fluid extraction and liquid chromatography with postcolumn ion-pair formation. *Anal. Chem.*, 68 (1996) 921-929.

3762 Forgacs, E. and Cserhati, T.: Use of alumina support for the separation of ethoxylated oligomer surfactants according to the length of the ethyleneoxide chain. *Anal. Lett.*, 29 (1996) 321-340.

3763 Ibrahim, N.M.A. and Wheals, B.B.: Oligomeric separation of alkylphenol ethoxylate surfactants on silica using aqueous acetonitrile eluents. *J. Chromatogr. A*, 731 (1996) 171-177.

3764 Marosi, T., Dekany, I. and de Keizer, A.: (Modified liquid chromatography and flow microcalorimetry for determination of surfactant adsorption). *Magy. Kem. Foly.*, 102 (1996) 78-82; C.A., 124 (1996) 243403s.

3765 McPherson, J.L., Nichols, J.H., Barditch-Crovo, P. and Hamzeh, F.M.: Determination of the spermicide nonoxynol-9 in vaginal lavage by high-performance liquid chromatography. *J. Chromatogr. B*, 677 (1996) 204-208.

3766 Miszkiewicz, W. and Szymanowski, J.: (HPLC analysis of non-ionic surfactants). *Wiad. Chem.*, 49 (1995) 415-428; C.A., 124 (1996) 120728g - a review with 54 refs.

3767 Németh-Kiss, V.: Separation of ethoxylated tributylphenol oligomers on porous graphitic carbon column. *J. Liq. Chromatogr.*, 19 (1996) 217-229.

3768 Sun, C., Baird, M., Anderson, H.A. and Brydon, D.L.: Separation of broadly distributed nonylphenol ethoxylates and determination of ethylene oxide oligomers in textile lubricants and emulsions by high-performance liquid chromatography. *J. Chromatogr. A*, 731 (1996) 161-169.

3769 Trathnigg, B. and Kollroser, M.: Mapping of polyethers using two-dimensional liquid chromatography with coupled density an RI detection. *Int. J. Polym. Anal. Charact.*, 1 (1995) 301-313; C.A., 124 (1996) 235572f.

See also 3777.

#### 36b. Antioxidants and preservatives

3770 Haruyama, M. and Okaya, Y.: (Determination of cationic preservatives in cosmetics by high performance liquid chromatography). *Jpn. J. Toxicol. Environ. Health*, 41 (1995) 367-374; C.A., 124 (1996) 126828y.

See also 3744.

#### 36c. Complex mixtures, technical products and unidentified compounds

3771 Botaro, V.R., Curvelo, A.A.S. and de Groote, R.A.M.C.: Fractionation of acetosolv. sugar cane bagasse lignin by preparative gel permeation chromatography. In: Veloso, D.P. and Ruggiero, R. (Editors), *Proc. Braz. Symp. Chem. Lignins Other Wood Comp.*, 3rd, Universidade Federal de Minas Gerais, Departamento de Quimica, Belo Horizonte, 1993, pp. 195-199; C.A., 124 (1996) 120461b.

3772 Fregeau, N.L. and Linehart, K.L.: Isolation of marine natural products by high-speed countercurrent chromatography. *Chem. Anal.*, 132 (1996) 265-300; C.A., 124 (1996) 126947m - a review with 75 refs.

3773 Herold, A.A., Johnson, B.R., Bartle, K.D. and Kandiyoti, R.: Large molecules in coal. Where do they hide? *Coal Sci. Technol.*, 24 (1995) 315-318; C.A., 124 (1996) 180704t.

3774 Huang, C.B.: Contaminant determination in cold rolling oil using HPSEC with RI and PDA detection. *J. Liq. Chromatogr.*, 19 (1996) 867-888.

3775 Lim, C.K.: Analysis of aryltetrahydrophthalene lignans and their glucoside conjugates in podophyllin resin by high-performance liquid chromatography. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 487-496; C.A., 124 (1996) 75173w.

3776 Ralph, J.P. and Catcheside, D.E.A.: Size-exclusion chromatography of solubilised low-rank coal. *J. Chromatogr. A*, 724 (1996) 97-105.

3777 Rovellini, P., Cortesi, N. and Fideli, E.: (LAS surfactant biodegradation: monitoring of metabolic intermediates by LC-MS). *Riv. Ital. Sostanze Grasse*, 72 (1995) 381-389; C.A., 124 (1996) 184877z.

3778 Thode, K., Kresse, M., Pfefferer, D., Semmler, W. and Mueller, R.H.: Mini-HIC for the determination of the surface hydrophobicity of magnetite dispersions. In: *World Meet. Pharm., Biopharm. Pharm. Technol.*, 1st, APGI, Chatenay Malabry, 1995, pp. 509-510; C.A., 124 (1996) 242506r.

For additional information see C.A.:

124 (1996) 149010q, 192800j, 218959v, 234715t.

See also 2676, 2696, 2838, 2893, 3795.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

- 3779 Huyghe, B.G., Liu, X., Sutjipto, S., Sugarman, B.J., Horn, M.T., Shepard, H.M., Scandella, C.J. and Shabram, P.: Purification of a type 5 recombinant adenovirus encoding human p53 by column chromatography. *Hum. Gene Ther.*, 6 (1995) 1403-1416; C.A., 124 (1996) 111371v.
- 3780 Jana, N.R., Halder, S. and Bhattacharya, S.: Thyroid hormone induces 52 kDa soluble protein in goat testis Leydig cell which stimulates androgen release. *Biochim. Biophys. Acta*, 1292 (1996) 209-214.

See also 2852.

38. INORGANIC COMPOUNDS

38a. Cations

- 3781 Akiyama, R., Takenaka, A. and Sugizaki, M.: Determination of antimony(III) and antimony(V) in glasses by ion chromatography/inductively coupled plasma atomic emission spectrometry. *Asahi Garasu Kenkyu Hokoku* 1994, 44 (1995) 13-18; C.A., 124 (1996) 248916w.
- 3782 Buzzoniti, M.C., Mentasti, E., Sarzanini, C., Braglia, M., Cocito, G. and Kraus, J.: Determination of rare earth elements by ion chromatography. Separation procedure optimization. *Anal. Chim. Acta*, 322 (1996) 49-54.
- 3783 Burnett, W.C. and Yeh, C.C.: Separation of protactinium from geochemical materials via extraction chromatography. *Radioact. Radiochem.*, 6 (1995) 22-32; C.A., 124 (1996) 192595w.
- 3784 Butt, S.B., Riaz, M. and Ul-Haq, E.: Simultaneous assay of copper, iron and uranium phosphate media by ion-pair liquid chromatography. *Chromatographia*, 42 (1996) 34-38.
- 3785 Cobb, J., Warwick, P., Carpenter, R.C. and Morrison, R.T.: Determination of strontium-90 in milk samples using a controlled precipitation clean-up step prior to ion-chromatography. *Sci. Total Environ.*, 173/174 (1995) 179-186; C.A., 124 (1996) 80757u.
- 3786 Coni, E., Alimonti, A., Bocca, A., la Torre, F., Menghetti E., Miraglia, E. and Caroli, S.: Speciation of trace elements in human milk by high performance liquid chromatography combined with inductively coupled plasma-mass spectrometry. *Trace Elem. Electrolytes*, 13 (1996) 26-32; C.A., 124 (1996) 200443n.
- 3787 Ehrling, C., Schmidt, U. and Liebscher, H.: Analysis of chromium(III)-fluoride-complexes by ion chromatography. *Fresenius J. Anal. Chem.*, 354 (1996) 870-873.
- 3788 El Moll, A., Heimburger, R., Lagarde, F., Leroy, M.J.F. and Maier, E.: Arsenic speciation in marine organisms: from the analytical methodology to the constitution of reference materials. *Fresenius J. Anal. Chem.*, 354 (1996) 550-556.
- 3789 Feldmann, J.: Ion chromatography coupled with inductively-coupled argon plasma mass spectrometry: multielement speciation as well as online matrix separation technique. *Anal. Commun.*, 33 (1996) 11-13; C.A., 124 (1996) 167655h.
- 3790 Haraguchi, H., Itoh, A. and Kimata, C.: Chemical speciation of trace metals in natural water by ultrafiltration/size exclusion chromatography/UV absorption/ICP-MS. *Anal. Sci. Technol.*, 8 (1995) 405-410; C.A., 124 (1996) 211245a.
- 3791 Hi, W., Miyazaki, A. and Haraguchi, H.: New approach to the simultaneous separation of inorganic cations and anions by ion chromatography based on ion-exchange and electrostatic interactions. *Anal. Sci.*, 11 (1995) 999-1000; C.A., 124 (1996) 163454u.
- 3792 Hissner, F., Mattusch, J. and Werner, G.: Determination of metal ions by ion chromatography with precolumn electrochemical preconcentration. *Fresenius J. Anal. Chem.*, 354 (1996) 718-721.
- 3793 Hu, W. and Haraguchi, H.: New approach for controlling the partitioning of analyte ions in ion chromatography with a water mobile phase. *J. Chromatogr. A*, 723 (1996) 251-258.
- 3794 Janos, P., Chroma, H. and Kuban, V.: Ion-chromatographic separation of metal cations in the presence of  $\alpha$ -hydroxyisobutyric acid. *Fresenius J. Anal. Chem.*, 355 (1996) 135-140.
- 3795 Kim, H.-J., Shin, Y.-K. and Kim, S.-J.: (Separation of Hg(II) by using the poly-NTOE (1,12-diaza-3,4:9,10-dibenzo-5,8-dioxacyclopentadeca-1,12-ylene-2,7-dihydroxyoctamethylene in liquid chromatography). *Anal. Sci. Technol.*, 8 (1995) 259-264; C.A., 124 (1996) 134092u.
- 3796 Kitazume, E.: Separation of rare earth and certain inorganic elements by high-speed countercurrent chromatography. *Chem. Anal.*, 132 (1996) 415-443; C.A., 124 (1996) 163313x - a review with 42 refs.
- 3797 Kondoh, A., Oi, T. and Hosoe, M.: Fractionation of barium isotopes in cation-exchange chromatography. *Sep. Sci. Technol.*, 31 (1996) 39-48; C.A., 124 (1996) 157011f.
- 3798 Lagarde, F. and Leroy, M.: (Speciation of trace elements: use and suitability). *Spectra Anal.*, 24 (1995) 32-36; C.A., 124 (1996) 109107v.
- 3799 Louw, I.: Ion chromatographic determination of transition metals in irradiated nuclear reactor surveillance samples. *Fresenius J. Anal. Chem.*, 354 (1996) 432-435.
- 3800 Mitrovic, B., Milacic, R. and Pihlar, B.: Speciation of aluminium in soil extracts by employing cation-exchange fast protein liquid chromatography - inductively coupled plasma atomic emission spectrometry. *Analyst (Cambridge)*, 121 (1996) 627-634.
- 3801 Muralidharan, S., Ma, G. and Freiser, H.: Factors influencing the efficiencies of multistage separations of lanthanides. In: Nash, K.L. and Choppin, G.R. (Editors), *Sep. f Elem. (Proc. Am. Chem. Soc. Symp.)*, 1994, Plenum, New York, 1995, pp. 177-197; C.A., 124 (1996) 127621n.
- 3802 Rodriguez, M., Suarez, J.A. and Espartero, A.G.: Separation of radioactive strontium by extraction using chromatographic resin. *Nucl. Instrum. Methods Phys. Res., Sect. A*, 369 (1996) 348-352; C.A., 124 (1996) 248915v.
- 3803 Saleh, F.Y., Mbamalu, G.E., Jaradat, Q.H. and Brungardt, C.E.: Ion chromatography - photodiode array UV-visible detection of Cr(III) hydrolytic polymerization products in pure and natural waters. *Anal. Chem.*, 68 (1996) 740-745.
- 3804 San Andrés, M.P. and Vera, S.: Chromatographic retention of Ni(II), Co(II) and Cu(II) as diethyldithiocarbamate complexes in presence of surfactant/n-propanol/water systems: determination of micellar binding constants. *J. Liq. Chromatogr.*, 19 (1996) 799-813.

- 3805 Sastry, M.S. and Ghose, A.K.: Column chromatographic separation of cations by anion exchange chromatography using PAR as complexing agent. *J. Indian Chem. Soc.*, 72 (1995) 705-708; C.A., 124 (1996) 134094w.
- 3806 Schulte, L.D., FitzPatrick, J.R., Salazer, R.R., Schake, B.S. and Martinez, B.T.: Removal of plutonium and americium from hydrochloric acid waste streams using extraction chromatography. In: Nash, K.L. and Choppin, G.R. (Editors), *Sep. f Elem., (Proc. Am. Chem. Soc. Symp.) 1994*, Plenum, New York, 1995, pp. 199-208; C.A., 124 (1996) 69470z.
- 3807 Seubert, A.: On-line coupling of ion chromatography and atomic spectrophotometry for ultra trace analysis in high purity molybdenum- and tungsten-silicide. *Fresenius J. Anal. Chem.*, 354 (1996) 788-796.
- 3808 Shijo, Y., Sato, H., Uehara, N. and Aratake, S.: Simultaneous determination of trace amounts of copper, nickel and vanadium in sea-water by high-performance liquid chromatography after extraction and back-extraction. *Analyst (Cambridge)*, 121 (1996) 325-328.
- 3809 Slejkovec, Z., Byrne, A.R., Smidov, B. and Rossbach, M.: Preliminary studies on arsenic species in some environmental samples. *Fresenius J. Anal. Chem.*, 354 (1996) 592-595.
- 3810 Snyder, T.S., Peterson, S.H. and Nayak, U.P.: Continuous, steady-state, chromatographic separation of gadolinium isotopes. U.S. US 5,470,479 (Cl. 210-635; B01D15/08), 28 Nov. 1995, Appl. 264,806, 23 Jun. 1994; 17 p.; C.A., 124 (1996) 69522t.
- 3811 Soylak, M., Sahin, U. and Elci, L.: Spectrophotometric determination of molybdenum in steel samples utilizing selective sorbent extraction on Amberlite XAD-8 resin. *Anal. Chim. Acta*, 322 (1996) 111-115.
- 3812 Srivastava, U., Rupainwar, D.C. and Singh, V.N.: Hydroxylamine hydrochloride as a complexing agent in the separation of cations using cation and anion exchange chromatography. *J. Indian Chem. Soc.*, 72 (1995) 759-761; C.A., 124 (1996) 134095x.
- 3813 Stummeyer, J., Harazin, B. and Wippermann, T.: Speciation of arsenic in water samples by high-performance liquid chromatography-hydride generation - atomic absorption spectrometry at trace levels using a postcolumn reaction system. *Fresenius J. Anal. Chem.*, 354 (1996) 344-351.
- 3814 Sutheimer, S.: Aluminum speciation in natural waters using high performance liquid chromatography with aluminum specific detection. Avail. *Univ. Microfilms Int.*, Order No. DA9604465, 1996, 267 p.; C.A., 124 (1996) 241466r.
- 3815 Tian, S. and Schwedt, G.: Solide-phase extraction of chromium(III)-diphenylcarbazone complex prior to ion-pair chromatography application to geological samples. *Fresenius J. Anal. Chem.*, 354 (1996) 447-450.
- 3816 Tsuji, A., Kotagawa, S., Watanabe, H. and Tsuda, T.: (Chromatographic behaviors by using ion exchange resins in electrochromatography). *Kuromatogurafi*, 16 (1995) 304-305; C.A., 124 (1996) 127883z.
- 3817 Umile, C. and Huber, J.F.K.: Significant reduction of the detection limit in ion chromatography by relative analyte enrichment with column switching. *J. Chromatogr. A*, 723 (1996) 11-17.
- 3818 Van Dy Nguyen: Ion chromatographic determination of trace anions and cations in high purity water. *Fresenius J. Anal. Chem.*, 354 (1996) 738-741.
- 3819 Vinas, P., Campillo, N., Garcis, I.L. and Cordoba, M.H.: Indirect flame atomic absorption detection for liquid chromatographic separation of alkaline metals. *Fresenius J. Anal. Chem.*, 354 (1996) 497-498.
- 3820 Wang, Q., Tsunoda, K.-i. and Akaiwa, H.: Extraction chromatographic behaviors of zirconium and hafnium using sulphoxides as stationary phases. *Anal. Sci.*, 11 (1995) 909-913; C.A., 124 (1996) 163453t.
- 3821 Yan, C.-T., Tan, Y.-A., Jen, J.-F., Hsiech, C. and Shih, T.-S.: Evaluating the stability of As(III) and As(V) in airborne sampling using HPLC. *LC-GC Int.*, 9 (1996) 101-108.
- 3822 Zablotsky, S.V., Rudak, E.A., Stopolyanshaya, L.V. and Wiley, J.R.: Improvements in the method of strontium-90 extraction from environmental samples with dicyclohexyl-18-crown-6. *Kernotechnik*, 60 (1995) 262-264; C.A., 124 (1996) 134161r.
- 3823 Zhang, X., Cornelis, R., de Kimpe, J. and Mees, L.: Arsenic speciation in serum of uraemic patients based on liquid chromatography with hydride generation atomic absorption spectrometry and on-line UV photo-oxidation digestion. *Anal. Chim. Acta*, 319 (1996) 177-185.
- For additional information see C.A.:  
124 (1996) 163508q, 163866e, 248960f.
- See also 2495, 2521, 2619, 2646, 2657, 3309, 3310, 3311, 3312, 3758, 3827, 3832, 3841.
- 38b. Anions*
- 3824 Buckee, G.K.: Determination of anions in beer by ion chromatography. *J. Inst. Brew.*, 101 (1995) 429-430; C.A., 124 (1996) 143858a.
- 3825 Chen, S.-H., Yang, Z.-Y., Wu, H.-L., Kou, H.-S. and Lin, S.-J.: Determination of thiocyanate anion by high-performance liquid chromatography with fluorometric detection. *J. Anal. Toxicol.*, 20 (1996) 38-42; C.A., 124 (1996) 78854e.
- 3826 Doblander, C. and Lackner, R.: Metabolism and detoxification of nitrite by trout hepatocytes. *Biochim. Biophys. Acta*, 1289 (1996) 270-274.
- 3827 Hu, W., Cao, S.-a., Tominaga, M. and Miyazaki, A.: Direct determination of bromide ions in sea water by ion chromatography using water as the mobile phase. *Anal. Chim. Acta*, 322 (1996) 43-47.
- 3828 Jun, X., Lima, J.L.F.C. and Montenegro, M.C.B.S.M.: Simultaneous determination of inorganic anions and carboxylic acids in wine using isocratic separation on a permanently coated reversed-phase column and UV indirect detection. *Anal. Chim. Acta*, 321 (1996) 263-271.
- 3829 Karmarkar, S.V.: Quick ion chromatographic determination of sulphate alone in soil extracts and natural waters. *Commun. Soil Sci. Plant Anal.*, 27 (1996) 843-852; C.A., 124 (1996) 218880n.
- 3830 Karpuk, A.D., Bragina, N.V., Ignatova, Z.N. and Poletaeva, I.L.: Application of monohydric phenols as mobile phase modifiers in the determination of anions by two-column ion chromatography. *J. Anal. Chem.*, 50 (1995) 1149-1153; C.A., 124 (1996) 163387z.
- 3831 Knox, J.H. and Wan, Q.-H.: Surface modification of porous graphite for ion exchange chromatography. *Chromatographia*, 42 (1996) 83-88.

- 3832 Ohta, K., Tanaka, K. and Fritz, J.S.: Non-suppressed ion chromatography of inorganic anions, magnesium and calcium ions using a pyromellitate eluent and its application in evaluating environmental water quality. *J. Chromatogr. A*, 731 (1996) 179-186.
- 3833 Okamoto, T., Isozaki, A. and Nagashima, H.: (Microdetermination of iodine in organic compounds by ion chromatography with a ceramic carbon column). *Bunseki Kagaku*, 45 (1996) 65-70; *C.A.*, 124 (1996) 192603x.
- 3834 Racker, M.: Ion chromatography of cyanide using a miniaturized dual working electrode potentiostatic detector (cyclic voltammetry, step deposition, chronocoulometry, amperometric detection). Avail. *Univ. Microfilms Int.*, Order No.: DA9541649, 1995, 102 p.; *C.A.*, 124 (1996) 134074q.
- 3835 Swiader, J.M. and Freiji, F.G.: Characterizing nitrate uptake in lettuce using very-sensitive ion chromatography. *J. Plant Nutr.*, 19 (1996) 15-27; *C.A.*, 124 (1996) 174561c.
- 3836 Tucker, H.L.: Determination of halogens in a petroleum product by ion chromatography. *Proc.-NOBCChE*, 22 (1995) 213-220; *C.A.*, 124 (1996) 121474v.
- 3837 Yu, B.S., Chen, P., Nie, L.-H. and Yao, S.-Z.: Determination of chloride in human body fluids by ion chromatography with piezoelectric sensor as detector. *Anal. Lett.*, 29 (1996) 43-57.

For additional information see *C.A.*:

124 (1996) 104969t, 115673t, 126772a, 134247y, 192650k, 192686b, 192805q.

See also 2426, 2495, 2507, 2826, 3789, 3791, 3793, 3817, 3818, 3840.

### 39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

- 3838 Aspin, I.P., Hamerton, I., Howlin, B.J., Jones, J.R., Parker, M.J., Russell, J.C. and Vick, T.A.: Development and application of radio-size-exclusion chromatography. *J. Chromatogr. A*, 727 (1996) 61-70.
- 3839 Kruglov, A.V., Andreev, B.M. and Pozhidaev, I.E.: Continuous isotope separation in systems with solid phase. II. Separation of nitrogen isotopes with use of ion-exchange resin. *Sep. Sci. Technol.*, 31 (1996) 471-490; *C.A.*, 124 (1996) 187642e.
- 3840 Toth, G.: Adsorption chromatographic separation of <sup>125</sup>I-labeled low molecular weight substances. In: Allen, J. and Voges, J. (Editors), *Synth. Appl. Isot. Labelled Compd. 1994, Proc. Int. Symp.I, 5th 1994*, Wiley, Chichester, 1995, pp. 807-809; *C.A.*, 124 (1996) 87714m.
- 3841 Yoshida, M., Sumiya, S., Watanabe, H. and Tobita, K.: A rapid separation method for determination of promethium-147 and samarium-151 in environmental samples with high performance liquid chromatography. *J. Radioanal. Nucl. Chem.*, 197 (1995) 219-227; *C.A.*, 124 (1996) 80760q.

For additional information see *C.A.*:

124 (1996) 185507r.

See also 2595, 2917, 3254, 3783, 3797, 3802, 3806, 3810, 3822.