

Electrophoresis

1. REVIEWS AND BOOKS

- 1611 Altria, K.D. (Editor): *Capillary Electrophoresis Guidebook: Principles, Operation, and Applications*. In: *Methods Mol. Biol. (Totowa)*, Humana, Totowa, 1996, 349 pp.; C.A., 124 (1996) 169985q.
- 1612 Buchberger, W.: Electrochemical detectors - tailor made techniques for liquid chromatography and capillary electrophoresis? *Fresenius J. Anal. Chem.*, 354 (1996) 797-802.
- 1613 Chrambach, A.: Quantitative and automated electrophoresis in sieving media. *Electrophoresis (Weinheim)*, 17 (1996) 454-464 - a review with 70 refs.
- 1614 Novotny, M.V.: Capillary electrophoresis. *Curr. Opin. Biotechnol.*, 7 (1996) 29-34; C.A., 124 (1996) 169610p - a review with many refs.
- 1615 Righetti, P.G. (Editor): *Capillary Electrophoresis in Analytical Biotechnology*. CRC, Boca Raton, 1996, 551 pp.; C.A., 124 (1996) 169986r.
- 1616 Timperman, A.T. and Schweedler, J.V.: Capillary electrophoresis with wavelength-resolved fluorescence detection. *Analyst (Cambridge)*, 121 (1996) 45R-52R.

See also 1618, 1637, 1646, 1648, 1666, 1672, 1676, 1679, 1680, 1682, 1686, 1691, 1697, 1698, 1707, 1708, 1712, 1715, 1718, 1720, 1721, 1727, 1745, 1753, 1756, 1759, 1770, 1808, 1819, 1820, 1822, 1874, 1880, 1889, 1906, 1913, 2176, 2231, 2232, 2248, 2289, 2366, 2376, 2377, 2407, 2408.

2. FUNDAMENTALS, THEORY AND GENERAL

2a. General

- 1617 Aalberts, D.P.: Electrophoretic mobility of asymmetric reptating polymers. *Phys. Rev. Lett.*, 75 (1995) 4544-4547; C.A., 124 (1996) 81273v.
- 1618 Altria, K.D.: Fundamentals of capillary electrophoresis theory. *Methods Mol. Biol. (Totowa)*, 52 (1996) 3-13; C.A., 124 (1996) 156714a - a review with 13 refs.
- 1619 Chen, N. and Chrambach, A.: Enhanced field strength and resolution in gel electrophoresis upon substitution of buffer by histidine at its isoelectric point. *Electrophoresis (Weinheim)*, 17 (1996) 699-703.
- 1620 Esteve-Romero, J., Simó-Alfonso, E., Bossi, A., Bresciani, F. and Righetti, P.G.: Sample streaks and smears in immobilized pH gradient gels. *Electrophoresis (Weinheim)*, 17 (1996) 704-708.
- 1621 Hansen, S.H., Tjornelund, J. and Bjornsdottir, I.: Selectivity enhancement in capillary electrophoresis using non-aqueous media. *TrAC*, 15 (1996) 175-180.
- 1622 Harmon, B.J., Leesong, I. and Regnier, F.E.: Moving boundary electrophoretically mediated microanalysis. *J. Chromatogr. A*, 726 (1996) 193-204.

- 1623 Kinawi, A., Hecker-Kia, A. and Groitl, M.M.: Further development of flow-through electrophoresis. *J. Clin. Chem. Clin. Biochem.*, 34 (1996) 49-52; C.A., 124 (1996) 169838u.
- 1624 Masoliver, J. and Weiss, G.H.: Resolution in time of two electrophoretic peaks. *Sep. Sci. Technol.*, 31 (1996) 319-326; C.A., 124 (1996) 197553r.
- 1625 Michalke, B.: Capillary electrophoresis - a useful tool in separation investigation. *Fresenius J. Anal. Chem.*, 354 (1996) 557-565.
- 1626 Semenov, S.N.: (Correlation analysis of fluctuations in continuous observation of electrophoresis). *Zh. Fiz. Khim.*, 69 (1995) 2070-2073; C.A., 124 (1996) 67706g.

See also 1680, 1700, 1717, 2335.

2b. Thermodynamics and theoretical relationships

- 1627 Long, D. and Viovy, J.L.: Theory of gel electrophoresis in high fields: transient regimes at the field onset. *Phys. Rev. E: Stat. Phys. Plasmas, Fluids, Relat. Interdiscip. Top.*, 53 (1996) 803-811; C.A., 124 (1996) 111567p.
- 1628 Patterson, D.H., Harmon, B.J. and Regnier, F.E.: Dynamic modeling of electrophoretically mediated microanalysis. *J. Chromatogr. A*, 732 (1996) 119-132.
- 1629 Wronski, M.: Calculation of pK values and zone lengths of anions in capillary isotachophoresis by the calibration method. *Chem. Anal. (Warsaw)*, 40 (1995) 911-915; C.A., 124 (1996) 127600e.

See also 1697, 1711, 1722, 2254, 2311.

2c. Relationship between structure and electrophoretic behaviour

See 1764.

2d. Measurement of physico-chemical and related values

- 1630 Ishihama, Y., Oda, Y. and Asakawa, N.: A hydrophobicity scale based on the migration index from microemulsion electrokinetic chromatography of anionic solutes. *Anal. Chem.*, 68 (1996) 1028-1032.

3. GENERAL TECHNIQUES

3a. Apparatus and accessories

- 1631 Djordjevic, N.M. and Ryan, K.: An easy way to enhance absorbance detection sensitivity of Waters Quanta-4000 capillary electrophoresis system. *J. Liq. Chromatogr.*, 19 (1996) 201-206.
- 1632 Liang, Z., Chiem, N., Ocvirk, G., Tang, T., Fluri, K. and Harrison, D.J.: Microfabrication of a planar absorbance and fluorescence cell for integrated capillary electrophoresis devices. *Anal. Chem.*, 68 (1996) 1040-1046.

- 1633 Nishigaki, K.: Method and apparatus for electrophoresis. *Jpn. Kokai Tokkyo Koho JP 08 05,609* [96 05,609] (Cl. G01N27/447), 12 Jan. 1996, Appl. 94/159,709, 20 Jun. 1994; 11 p.; C.A., 124 (1996) 192789n.
- 1634 Rebscher, H. and Pyell, U.: Instrumental developments in capillary electrochromatography. *Chromatographia*, 42 (1996) 171-176.
- 1635 Wittwer, H. and Krueger, H.G.: Possibilities and limits of electrophoresis. *CFI, Ceram. Forum Int.*, 72 (1995) 556-560; C.A., 124 (1996) 124414m.
- 1636 Yamada, H. and Kamahori, M.: Microinjector for capillary gel electrophoresis. *Jpn. Kokai Tokkyo Koho JP 08 29,391* [96 29,391] (Cl. G01N27/447), 2 Feb. 1996, Appl. 94/167,666, 20 Jul. 1994; 5 p.; C.A., 124 (1996) 243461j.
- See also 1729.
- 3b. Detectors and detection procedures**
- 1637 Chen, J.-G., Woltman, S.J. and Weber, S.G.: Electrochemical detection of biomolecules in liquid chromatography and capillary electrophoresis. *Adv. Chromatogr.*, 36 (1996) 273-313; C.A., 124 (1996) 140051w - a review with many refs.
- 1638 Clark, B.K. and Sepaniak, M.J.: Evaluation of a spatial scanning laser fluorometric detector for capillary electrophoresis: application to studies of band migration and dispersion. *J. Microcolumn Sep.*, 7 (1995) 593-601; C.A., 124 (1996) 81282x.
- 1639 De Backer, B.L. and Nagels, L.J.: Potentiometric detection of carboxylic aliphatic acids in CZE using ion-selective liquid-membrane electrodes. *Biomed. Chromatogr.*, 9 (1995) 257-258; C.A., 124 (1996) 134357j.
- 1640 Fang, L.: Development of time-of-flight mass spectrometric detector for capillary electrophoresis using electrospray ionization. Avail. *Univ. Microfilms Int.*, Order No.: DA9602874, 1996, 135 p.; C.A., 124 (1996) 249176s.
- 1641 Fishman, H.A., Orwar, O., Allbritton, N.L., Modi, B.P., Shear, J.B., Scheller, R.H. and Zare, N.: Cell-to-cell scanning in capillary electrophoresis. *Anal. Chem.*, 68 (1996) 1181-1186.
- 1642 Kaizu, T. and Donghyun, K.: Detector cell for capillary electrophoresis. *PCT Int. Appl. WO 95 33,196* (Cl. G01N21/05), 7 Dec. 1995, US Appl 250,200, 27 May 1994; 20 pp.; C.A., 124 (1996) 140375e.
- 1643 Karaman, Z.: Towards an automated interpretation of gel electrophoresis. In: Van Ooijen, J.W. and Jansen, J. (Editors), *Biom. Plant Breed.: Appl. Mol. Markers, Proc. Meet. EUCARPIA Sect., 9th 1994*, CPRO-DLO, Wageningen, 1994, pp. 125-131; C.A., 124 (1996) 81286b.
- 1644 LaCourse, W.R. and Owens, G.S.: Pulsed electrochemical detection of nonchromophoric compounds following capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 310-318.
- 1645 Ludwig, M., Pfeifer, P. and Kittler, L.: Integrated optical sensor for the analysis of substances in a liquid phase. *Ger. Offen. DE 4,419,586* (Cl. G01N21/41), 7 Dec. 1995, Appl. 4,419,586, 3 Jun. 1994; 20 pp.; C.A., 124 (1996) 81462f.
- 1646 Luo, G. and Wang, Y.: (Principles and applications of capillary electrophoresis. (II). Detection techniques in capillary electrophoresis). *Sepu*, 13 (1995) 437-440; C.A., 124 (1996) 192413k - a review with 31 refs.
- 1647 Ma, M.S., Han, H.W. and Liu, G.Q.: Laser induced fluorescence detector (LIF) for capillary electrophoresis. *Chin. Chem. Lett.*, 7 (1996) 39-42; C.A., 124 (1996) 197337y.
- 1648 Mank, A.J.G., Lingeman, H. and Gooijer, C.: Diode laser-based detection in liquid chromatography and capillary electrophoresis. *TrAC*, 15 (1996) 1-11 - a review.
- 1649 Ranganathan, V. and De, P.K.: Western blot of proteins from Coomassie-stained polyacrylamide gels. *Anal. Biochem.*, 234 (1996) 102-104.
- 1650 Ren, J., Li, B., Deng, Y. and Cheng, J.: Indirect thermo-optical detection for capillary electrophoresis. *Talanta*, 42 (1995) 1891-1895; C.A., 124 (1996) 156942y.
- 1651 Ren, J.-C., Li, B.-C., Deng, Y.-Z. and Cheng, J.K.: (Studies on indirect thermo-optical detection for capillary electrophoresis). *Gaodeng Xuexiao Huaxue Xuebao*, 16 (1995) 1182-1185; C.A., 124 (1996) 105113w.
- 1652 Roberts, R.E. and Johnson, D.C.: Variation in PED response at a gold microelectrode as a function of waveform parameters when applied to alditols and carbohydrates separated by capillary electrophoresis. *Electroanalysis*, 7 (1995) 1015-1019; C.A., 124 (1996) 105178w.
- 1653 Shojace, N., Patton, W.F., Lim, M.J. and Shepro, D.: Pyrogallol red-molybdate: a reversible, metal chelate stain for detection of proteins immobilized on membrane supports. *Electrophoresis (Weinheim)*, 17 (1996) 687-693.
- 1654 Voegel, P.D. and Baldwin, R.P.: Electrochemical detection with copper electrodes in liquid chromatography and capillary electrophoresis. *Int. Lab.*, 26, March (1996) 16A-16N.
- 1655 Vogt, C., Vogt, J. and Wittrisch, H.: Element-sensitive X-ray detection for capillary electrophoresis. *J. Chromatogr. A*, 727 (1996) 301-310.
- 1656 Wu, X.-Z., Hosaka, A., Kobayashi, E. and Hobo, T.: Sensitivity enhancement effect in off-column detection for capillary electrophoresis. *J. Chromatogr. A*, 726 (1996) 205-210.
- 1657 Xiong, S., Li, J. and Cheng, J.: Multiwavelength fluorescence detection for purity analysis of fluorescence agents using capillary electrophoresis with a charge-coupled device. *Anal. Chim. Acta*, 322 (1996) 187-194.
- See also 1612, 1631, 1632, 1693, 1702, 1716, 1723, 1724, 1725, 1735, 1778, 1782, 1785, 1887, 2010, 2049, 2089, 2116, 2135, 2171, 2334, 2364, 2379, 2427.
- 3c. Stabilization media for electrophoresis**
- 1658 Guo, H., Fang, W., Li, R., Yang, P., Ding, Z. and Shen, E.: (Acryldex cross-linked allyl dextran - a high resolution gel chromatographic media). *Sepu*, 13 (1995) 466-469; C.A., 124 (1996) 11541a.
- 1659 Liu, J.-M., Nakagama, T., Okazawa, H., Wu, X.-Z. and Hobo, T.: Separation and determination of some stereoisomers by capillary gel electrophoresis with cyclodextrin incorporated in polyacryl gel. *Fresenius J. Anal. Chem.*, 354 (1996) 451-454.
- 1660 Liu, P.Z.: Surface-modified polymeric hollow fiber columns for capillary electrophoresis. Avail. *Univ. Microfilms Int.*, Order No.: DA9602114, 1995, 164 p.; C.A., 124 (1996) 192524x.
- 1661 Menchen, S.M., Winnik, M.A. and Johnson, B.F.: Viscous electrophoresis polymer medium and method. *U.S. US 5,468,365* (Cl. 204-299R; C25B7/00), 21 Nov. 1995, US Appl. 950,863, 24 Sep. 1992; 38 pp.; C.A., 124 (1996) 111731n.

- 1662 Miura, R., Yamagata, S. and Yamagata, T.: Gels made from copolymers of acrylamide and glycosaminoglycan derivatives and their use as electrophoretic carriers in identifying enzyme. *Eur. Pat. Appl. EP 693,499* (Cl. C08B37/08), 24 Jan. 1996, JP Appl. 94/191,285, 22 Jul. 1994; 18 p.; C.A., 124 (1996) 179362y.
- 1663 Phang, T., Ji, I. and Ji, T.H.: No need of acetic acid for processing polyacrylamide gels. *Anal. Biochem.*, 234 (1996) 96-97.
- 1664 Simò-Alfonso, E., Gelfi, C., Sebastiano, R., Citterio, A. and Righetti, P.G.: Novel acrylamido monomers with higher hydrophilicity and improved hydrolytic stability: I. Synthetic route and product characterization. *Electrophoresis (Weinheim)*, 17 (1996) 723-731.
- 1665 Simò-Alfonso, E., Gelfi, C., Sebastiano, R., Citterio, A. and Righetti, P.G.: Novel acrylamido monomers with higher hydrophilicity and improved hydrolytic stability: II. Properties of N-acryloylaminopropanol. *Electrophoresis (Weinheim)*, 17 (1996) 732-737.
- See also 1687, 2234, 2276.
- 3d. *Quantitative analysis*
- 1666 Altria, K.D.: Quantitation procedures. *Methods Mol. Biol. (Totowa)*, 52 (1996) 49-60; C.A., 124 (1996) 212169r - a review with 15 refs.
- 1667 Qi, S., Huang, A. and Sun, Y.: Compensation of two kinds of bias in capillary zone electrophoresis and its use in internal normalization quantitation. *Anal. Chem.*, 68 (1996) 1342-1346.
- 3e. *Preparative scale electrophoresis*
- 1668 Corthals, G.L., Margolis, J., Williams, K.L. and Gooley, A.A.: The role of pH and membrane porosity in preparative electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 771-775.
- See also 1946, 2059.
- 3f. *Programmed voltage and buffer gradients*
- See 2154.
4. SPECIAL TECHNIQUES
- 4a. *Automation*
- See 1613, 1643, 1714, 1725, 1849, 2294.
- 4b. *Computerization and modelling*
- 1669 Desruisseaux, C. and Slater, G.W.: Pulsed-field trapping electrophoresis: a computer simulation study. *Electrophoresis (Weinheim)*, 17 (1996) 623-632.
- 1670 Masujima, T. and Tamura, A.: (Imaging of separation dynamics and looking for a new separation base). *Kuromatogurafi*, 16 (1995) 342-343; C.A., 124 (1996) 127885b.
- See also 1628, 1754, 1871, 1884, 1899, 1940, 1948, 1961, 1963, 2174, 2267.
- 4c. *Combination with other physicochemical techniques, (MS, IR etc.)*
- 1671 Hutton, T. and Major, H.J.: Characterizing biomolecules by electrospray ionization-mass spectrometry [ESI-MS] coupled to liquid chromatography and capillary electrophoresis. *Biochem. Soc. Trans.*, 23 (1995) 924-927; C.A., 124 (1996) 197491u.
- 1672 Li, D.T. and Her, G.R.: (Capillary electrophoresis-mass spectrometry). *Huaxue*, 53 (1995) 438-445; C.A., 124 (1996) 169618x - a review with 23 refs.
- 1673 Mordehai, A., Lim, H.K. and Henion, J.D.: Ion spray liquid chromatography/mass spectrometry and capillary electrophoresis/mass spectrometry on a modified benchtop ion trap mass spectrometer. In: March, R.E. and Todd, J.F.J. (Editors), *Pract. Aspects Ion Trap Mass Spectrom.*, CRC, Boca Raton, 1995, pp. 215-237; C.A., 124 (1996) 192914z.
- 1674 Schreiner, M., Strupat, K., Lottspeich, F. and Eckerskorn, C.: Ultraviolet matrix assisted laser desorption ionization-mass spectrometry of electroblotted proteins. *Electrophoresis (Weinheim)*, 17 (1996) 954-961.
- 1675 Tomlinson, A.J., Benson, L.M., Oda, R.P., Braddock, W.D., Riggs, B.L., Katzmann, J.A. and Naylor, S.: Novel modifications and clinical applications of preconcentration-capillary electrophoresis-mass spectrometry. *J. Capillary Electrophor.*, 2 (1995) 97-104; C.A., 124 (1996) 81290y.
- 1676 Wahl, J.H., Udseth, H.R. and Smith, R.D.: Capillary electrophoresis-mass spectrometry in peptide mapping. In: Hancock, W.S. (Editor), *New Methods Pept. Mapp. Charact. Proteins*, CRC, Boca Raton, 1996, pp. 143-179; C.A., 124 (1996) 111287x - a review with 79 refs.
- 1677 Xiong, S., Li, J. and Cheng, J.: (Studies on the coupling technique between high performance capillary electrophoresis and charge-coupled device detector. I. Set-up for multiwavelength fluorescence detection). *Sepu*, 13 (1995) 427-440; C.A., 124 (1996) 163793d.
- 1678 Xiong, S., Li, J. and Cheng, J.: (Studies on the coupling techniques of high performance capillary electrophoresis and charge coupled device detection. IV. Monte Carlo simulation). *Fenxi Huaxue*, 24 (1996) 19-23; C.A., 124 (1996) 163440m.
- See also 1640, 1690, 1701, 1728, 1735, 1738, 1847, 1863, 1866, 1871, 1879, 1881, 1882, 1900, 1907, 1976, 2081, 2167, 2236, 2363, 2380, 2413.
- 4d. *Affinity electrophoresis*
- See 1804, 1886, 2007, 2241, 2385.
- 4e. *Capillary zone electrophoresis and electrokinetic chromatography*
- 1679 Altria, K.D.: Additional application areas of capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 309-344; C.A., 124 (1996) 197272y - a review with 139 refs.
- 1680 Altria, K.D.: Method development/optimization. *Methods Mol. Biol. (Totowa)*, 52 (1996) 29-48; C.A., 124 (1996) 197268b - a review with 30 refs.

- 1681 Altria, K.D.: Method validation. *Methods Mol. Biol. (Totowa)*, 52 (1996) 83-97; C.A., 124 (1996) 197514d.
- 1682 Altria, K.D.: Optimization of precision in quantitative analysis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 61-70; C.A., 124 (1996) 197269c - a review with 21 refs.
- 1683 Altria, K.D.: Optimization of sensitivity. *Methods Mol. Biol. (Totowa)*, 52 (1996) 71-81; C.A., 124 (1996) 197513c.
- 1684 Altria, K.D.: Quick guide to running a successful separation. *Methods Mol. Biol. (Totowa)*, 52 (1996) 119-122; C.A., 124 (1996) 197517g.
- 1685 Altria, K.D.: Troubleshooting [capillary electrophoresis]. *Methods Mol. Biol. (Totowa)*, 52 (1996) 105-118; C.A., 124 (1996) 197516f.
- 1686 Altria, K.D.: Typical operating procedures. *Methods Mol. Biol. (Totowa)*, 52 (1996) 21-28; C.A., 124 (1996) 197267a - a review with 2 refs.
- 1687 Baechmann, K., Bazzanella, A., Goettlicher, B., Haag, I., Han, K.Y., Arnecke, R. and Boehmer, V.: (New pseudostationary phase for electrokinetic chromatography). *GIT Spez. Chromatogr.*, 15 (1995) 96-103; C.A., 124 (1996) 163777b.
- 1688 Boutin, J.A., Henning, P., Lambert, P.-H., Bertin, S., Petit, L., Mahieu, J.-P., Serkiz, B., Volland, J.-P. and Fauchère, J.-L.: Combinatorial peptide libraries: robotic synthesis and analysis of nuclear magnetic resonance, mass spectrometry, tandem mass spectrometry, and high-performance capillary electrophoresis techniques. *Anal. Biochem.*, 234 (1996) 126-141.
- 1689 Burgi, D.S. and Chien, R.-L.: Application and limits of sample stacking in capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 211-226; C.A., 124 (1996) 197520c.
- 1690 Chankvetadze, B., Endresz, G., Schulte, G., Bergenthal, D. and Blaschke, G.: Capillary electrophoresis and ¹H NMR studies on chiral recognition of atropisomeric binaphthyl derivatives by cyclodextrin hosts. *J. Chromatogr. A*, 732 (1996) 143-150.
- 1691 Cunat-Walter, M.A.: (Application of the capillary electrophoresis in the biochemical analysis). *Oesterr. Chem. Z.*, 96 (1995) 195-197; C.A., 124 (1996) 197284d - a review without refs.
- 1692 Cunat-Walter, M.A., Shoikhet, K. and Engelhardt, H.: (pH Change in the buffer vials during a separation in capillary electrophoresis). *GIT Fachz. Lab.*, 39 (1995) 914-921; C.A., 124 (1996) 105145h.
- 1693 Dalhoeven, G.A.M., Joshi, N.V., Rodriguez, N. and Hernandez, L.: Simultaneous measurement of capillary electrophoresis fluorescence peaks and their corresponding spectra. *J. Liq. Chromatogr.*, 18 (1995) 3729-3749.
- 1694 Dasgupta, P.K. and Surowiec, K.: Quantitative injection from a microloop. Reproducible volumetric sample introduction in capillary zone electrophoresis. *Anal. Chem.*, 68 (1996) 1164-1168.
- 1695 Dulay, M.T., Yan, C., Rakestraw, D.J. and Zare, R.N.: Automated capillary electrochromatography: reliability and reproducibility studies. *J. Chromatogr. A*, 725 (1996) 361-366.
- 1696 Dunn, C.D.: Reverse direction capillary electrophoresis: theory and application (sample injection, micellar concentration). Avail. *Univ. Microfilms Int.*, Order No.: DA9533431, 1995, 168 p.; C.A., 124 (1996) 74906a.
- 1697 Guan, F.: Solution chemical equilibrium in high performance capillary electrophoresis separation. *Fenxi Huaxue*, 24 (1996) 109-114; C.A., 124 (1996) 192434t - a review with 17 refs.
- 1698 Guo, X., Xue, J., Zhang, Y. and Lin, B.: (Capillary non-gel sieving electrophoresis). *Shengwu Huaxue Yu Shengwu Wuli Jinzhan*, 22 (1995) 403-407; C.A., 124 (1996) 140032r - a review with 20 refs.
- 1699 Guttman, A.: Capillary gel electrophoresis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 157-169; C.A., 124 (1996) 197518h.
- 1700 Hayashi, Y. and Matsuda, R.: Noise and measurement precision in analytical instruments. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 35-44; C.A., 124 (1996) 75099b.
- 1701 Heegaard, N.H.H. and Roepstorff, P.: Preparative capillary electrophoresis and mass spectrometry for the identification of a putative heparin-binding site in amyloid P component. *J. Capillary Electrophor.*, 2 (1995) 219-223; C.A., 124 (1996) 169799g.
- 1702 Hietpas, P.B. and Ewing, A.G.: On-column and post-column derivatization for capillary electrophoresis with laser-induced fluorescence for the analysis of single cells. *J. Liq. Chromatogr.*, 18 (1995) 3557-3576.
- 1703 Idei, M., Kiss, E. and Kéri, G.: Calculation of modified capacity factor in micellar electrokinetic chromatography. *Electrophoresis (Weinheim)*, 17 (1996) 762-765.
- 1704 Lelièvre, F., Yan, C., Zare, R.N. and Gareil, P.: Capillary electrochromatography: operating characteristics and enantiomeric separations. *J. Chromatogr. A*, 723 (1996) 145-156.
- 1705 Lin, Q.S., Zhang, T.T. and Liu, G.Q.: On-column electro-pumping system for sample introduction in capillary electrophoresis. *Chin. Chem. Lett.*, 7 (1996) 35-38; C.A., 124 (1996) 197557v.
- 1706 Liu, S.: Development and applications of capillary-based analysis systems (electroosmotic flow, microbore column chromatography). Avail. *Univ. Microfilms Int.*, Order No.: DA9541756, 1995, 211 p.; C.A., 124 (1996) 134106b.
- 1707 Majors, R.E.: The role of column diameter in chromatography and capillary electrophoresis. *LC-GC*, 14 (1996) 10-19; C.A., 124 (1996) 98435x - a review with 17 refs.
- 1708 Manabe, T.: (Electrophoresis, as one distinct field of instrumental analysis. New developments in capillary electrophoresis). *Seibutsu Butsuri Kagaku*, 39 (1995) 337-341; C.A., 124 (1996) 111293w - a review with 15 refs.
- 1709 McGuffin, V.L.: Photoactivation: a novel means to mediate electrophoretic separations. *J. Liq. Chromatogr.*, 18 (1995) 3787-3812.
- 1710 Mechref, Y. and El Rassi, Z.: Micellar electrokinetic capillary chromatography with *in-situ* charged micelles. VI. Evaluation of novel chiral micelles consisting of steroidal-glycoside surfactant-borate complexes. *J. Chromatogr. A*, 724 (1996) 285-296.
- 1711 Minarik, M., Gas, B., Rizzi, A. and Kennler, E.: Plate height contribution from wall adsorption in capillary zone electrophoresis of proteins. *J. Capillary Electrophor.*, 2 (1995) 89-96; C.A., 124 (1996) 81289e.
- 1712 Moring, S.E.: Buffers, electrolytes, and additives for capillary electrophoresis. In: Righetti, P.G. (Editor), *Capillary Electrophor. Anal. Biotechnol.*, CRC, Boca Raton, 1996, pp. 37-60; C.A., 124 (1996) 249152f - a review with 43 refs.
- 1713 Nussberger, S., Foret, F., Hebert, S.C., Karger, B.L. and Hediger, M.A.: Nonradioactive monitoring of organic and inorganic solute transport into single *Xenopus* oocytes by capillary zone electrophoresis. *Biophys. J.*, 70 (1996) 998-1005; C.A., 124 (1996) 111563j.

- 1714 Ooyama, M.: (Automated device for capillary electrophoresis). *Jpn. Kokai Tokkyo Koho JP 07,274,935* [95,274,935] (Cl.C12M1/100), 24 Oct. 1995, Appl. 94/74,591, 13 Apr. 1994; 6 pp.; C.A., 124 (1996) 81480k.
- 1715 Otsuka, K. and Terabe, S.: Micellar electrokinetic chromatography. *Methods Mol. Biol. (Totowa)*, 52 (1996) 125-155; C.A., 124 (1996) 139996b - a review with 71 refs.
- 1716 Palm, A., Lindh, C., Hjertén, S. and Pawliszyn, J.: Capillary zone electrophoresis in agarose gels using absorption imaging detection. *Electrophoresis (Weinheim)*, 17 (1996) 766-770.
- 1717 Poppe, H., Cifuentes, A. and Kok, W.T.: Theoretical description of the influence of external radial fields on the electroosmotic flow in capillary zone electrophoresis. *Anal. Chem.*, 68 (1996) 888-893.
- 1718 Pritchett, T., Evangelista, R.A. and Chen, F.-T.A.: Capillary electrophoresis-based immunoassays. *Bio/Technology*, 13 (1995) 1449-1450; C.A., 124 (1996) 49748t - a review with 7 refs.
- 1719 Razeq, S., Tamura, A., Khademizadeh, M. and Masujima, T.: Direct observation of capillary electrophoretic phenomena using a video image analysis system. *Chem. Lett.*, (1996) 93-94; C.A., 124 (1996) 111334k.
- 1720 Righetti, P.G., Gelfi, C. and Chiari, M.: Capillary electrophoresis of DNA. In: Righetti, P.G. (Editor), *Capillary Electrophor. Anal. Biotechnol.*, CRC, Boca Raton, 1996, pp. 431-476; C.A., 124 (1996) 197311k - a review with 237 refs.
- 1721 Shimura, K.: (Affinity probe capillary electrophoresis. Antigen quantification and analysis of specific interaction). *Seibutsu Butsuri Kagaku*, 39 (1995) 349-353; C.A., 124 (1996) 84122f - a review with 9 refs.
- 1722 Slais, K.: Transient electrophoretic processes in capillaries of non-uniform cross-section. *J. Chromatogr. A*, 730 (1996) 247-259.
- 1723 Song, J.M., Kawazumi, H., Inoue, T. and Ogawa, T.: Time-resolved photon-counting fluorimetry with a semiconductor laser for capillary electrophoresis. *J. Chromatogr. A*, 727 (1996) 330-333.
- 1724 Staller, T.D.: Novel detection alternatives for capillary electrokinetic separations (fluorometric photodiode array, peroxyoxalate chemiluminescence). Avail. *Univ. Microfilms Int.*, Order No.: DA9540126, 1995, 173 p.; C.A., 124 (1996) 134339e.
- 1725 Stastna, M., Kahle, V. and Slais, K.: Automated instrumentation for miniaturized displacement electrophoresis with on-column photometric detection. *J. Chromatogr. A*, 730 (1996) 261-272.
- 1726 Swartz, M.E.: Separation and detection of enantiomeric compounds by liquid chromatography and chiral micellar electrokinetic capillary chromatography. Avail. *Univ. Microfilms Int.*, Order No.: DA9601875, 1995, 231 p.; C.A., 124 (1996) 192902u.
- 1727 Takagi, T.: (Capillary electrophoresis. Can we say good-bye SDS-PAGE?). *Seibutsu Butsuri Kagaku*, 39 (1995) 343-348; C.A., 124 (1996) 111294x - a review with 26 refs.
- 1728 Tomlinson, A.J., Guzman, N.A. and Naylor, S.: Enhancement of concentration limits of detection in CE and CE-MS: a review of on-line sample extraction, cleanup, analyte preconcentration, and microreactor technology. *J. Capillary Electrophor.*, 2 (1995) 247-266; C.A., 124 (1996) 197281a.
- 1729 Veraart, J.R., Lingeman, H. and Gooijer, C.: Temperature control in capillary electrophoresis systems. *Biomed. Chromatogr.*, 9 (1995) 271-272; C.A., 124 (1996) 104823r.
- 1730 Wheat, T.E., Chiklis, F.M. and Lilley, K.A.: Methods development in capillary electrophoresis with automated peak tracking by chemometric analysis of diode array detection data. *J. Liq. Chromatogr.*, 18 (1995) 3643-3657.
- 1731 Williams, B.A. and Vigh, G.: Fast, accurate mobility determination method for capillary electrophoresis. *Anal. Chem.*, 68 (1996) 1174-1180.
- 1732 Williams, R.L. and Vigh, G.: Maximization of separation efficiency in capillary electrophoretic chiral separations by means of mobility-matching background electrolytes. *J. Chromatogr. A*, 730 (1996) 273-278.
- 1733 Williams, R.L. and Vigh, G.: The use of alkylmethylmorpholine-based background electrolytes in capillary electrophoresis. *J. Liq. Chromatogr.*, 18 (1995) 3813-3824.
- 1734 Wright, P.B. and Dorsey, J.G.: Silver(I)-mediated separations by capillary zone electrophoresis and micellar electrokinetic chromatography: argentation electrophoresis. *Anal. Chem.*, 68 (1996) 415-424.
- 1735 Xiong, S., Li, J. and Cheng, J.: (Coupling technique of high performance capillary electrophoresis with charge-coupled device detector. VI. Laser-induced fluorescence detection setup and its comparison with common light source setup). *Fenxi Kexue Xuebao*, 11 (1995) 12-15; C.A., 124 (1996) 218948r.
- 1736 Yamaguchi, M., Izumi, T., Isobe, T. and Okuyama, T.: (Resolution of biological macromolecules by capillary electrophoresis using konjac glucomannan as macromolecule-sieving additive). *Nippon Kagaku Kaishi*, (1996) 166-175; C.A., 124 (1996) 197549u.
- 1737 Yang, S.: Characterization of chemical interactions in micellar liquid chromatography and micellar electrokinetic capillary chromatography (phenols, surfactants). Avail. *Univ. Microfilms Int.*, Order No.: DA9542250, 1995, 205 p.; C.A., 124 (1996) 134384r.
- 1738 Zaramella, A., Curcuruto, O., Hamdan, M., Cabrele, C. and Peggion, E.: Investigation of crudes of synthesis of [Leu³¹,Pro³⁴]-neuropeptide Y by capillary zone electrophoresis/mass spectrometry. *Rapid Commun. Mass Spectrom.*, 9 (1995) 1386-1390; C.A., 124 (1996) 146816w.

See also* 1611, 1612, 1614, 1615, 1616, 1618, 1621, 1622, 1625, 1630, 1631, 1632, 1634, 1636, 1637, 1638, 1639, 1640, 1641, 1642, 1644, 1646, 1647, 1648, 1650, 1651, 1652, 1654, 1655, 1656, 1657, 1659, 1660, 1666, 1667, 1671, 1672, 1673, 1675, 1676, 1677, 1678, 1739, 1740, 1741, 1742, 1743, 1744, 1745, 1746, 1747, 1748, 1749, 1750, 1752, 1764, 1765, 1767, 1768, 1769, 1770, 1771, 1772, 1773, 1775, 1776, 1777, 1778, 1779, 1782, 1783, 1784, 1785, 1786, 1787, 1790, 1791, 1793, 1796, 1798, 1806, 1807, 1808, 1813, 1814, 1815, 1816, 1817, 1818, 1819, 1829, 1830, 1832, 1833, 1834, 1835, 1836, 1837, 1838, 1839, 1840, 1841, 1842, 1843, 1844, 1845, 1846, 1847, 1849, 1850, 1851, 1852, 1855, 1856, 1857, 1859, 1860, 1861, 1863, 1865, 1866, 1877, 1880, 1882, 1894, 1895, 1896, 1897, 1902, 1905, 1908, 1914, 1931, 1953, 1973, 1978,

* The number of cross references in this section reflects the exponential growth of capillary electrophoretic applications. This section will be rearranged from January 1st 1997.

1988, 2018, 2069, 2070, 2077, 2092, 2116, 2167, 2168, 2171, 2172, 2173, 2174, 2176, 2177, 2179, 2180, 2181, 2232, 2235, 2236, 2251, 2254, 2280, 2285, 2288, 2289, 2295, 2300, 2301, 2317, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2340, 2341, 2342, 2343, 2344, 2345, 2347, 2349, 2350, 2351, 2352, 2353, 2354, 2356, 2357, 2358, 2359, 2360, 2361, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2393, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2405, 2407, 2408, 2410, 2411, 2412, 2413, 2414, 2421, 2422, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436.

4f. Isotachopheresis

See 1629, 1862, 2340, 2423.

4g. Enantiomers, separation

- 1739 Armstrong, D.W. and Rundlett, K.L.: CE resolution of neutral and anionic racemates with glycopeptide antibiotics and micelles. *J. Liq. Chromatogr.*, 18 (1995) 3659-3674.
- 1740 Chankvetadze, B., Schulte, G. and Blaschke, G.: Reversal of enantiomer elution order in capillary electrophoresis using charged and neutral cyclodextrins. *J. Chromatogr. A*, 732 (1996) 183-187.
- 1741 Copper, C.L., Davis, J.B. and Sepaniak, M.J.: Mechanisms of enantiomeric resolution in cyclodextrin-modified capillary electrophoretic separations of binaphthyl compounds. *Chirality*, 7 (1995) 401-408; *C.A.*, 124 (1996) 75141j.
- 1742 Endresz, G., Chankvetadze, B., Bergenthal, D. and Blaschke, G.: Comparative capillary electrophoretic and nuclear magnetic resonance studies of the chiral recognition of racemic metomidate with cyclodextrin hosts. *J. Chromatogr. A*, 732 (1996) 133-142.
- 1743 Guttman, A., Brunet, S. and Cooke, N.: Capillary electrophoresis separation of enantiomers using cyclodextrin array chiral analysis. *LC-GC Int.*, 9 (1996) 88-100.
- 1744 Kuhn, R.: (Separation of enantiomers by capillary zone electrophoresis using a chiral crown ether). *GIT Fachz. Lab.*, 39 (1995) 1031-1034; *C.A.*, 124 (1996) 192878r.
- 1745 Olechno, J.D. and Nolan, J.A.: Injection methods in capillary electrophoresis. In: Righetti, P.G. (Editor), *Capillary Electrophor. Anal. Biotechnol.*, CRC, Boca Raton, 1996, pp. 61-99; *C.A.*, 124 (1996) 249153g - a review with 83 refs.
- 1746 Rogan, M.M. and Altria, K.D.: Chiral separations by capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 171-196; *C.A.*, 124 (1996) 197519j.
- 1747 Rundlett, K.L., Gasper, M.P., Zhou, E.Y. and Armstrong, D.W.: Capillary electrophoretic enantiomeric separations using the glycopeptide antibiotic, teicoplanin. *Chirality*, 8 (1996) 88-107; *C.A.*, 124 (1996) 242482e.
- 1748 Stalcup, A.M. and Gahm, K.H.: Application of sulfated cyclodextrins to chiral separations by capillary zone electrophoresis. *Anal. Chem.*, 68 (1996) 1360-1368.

- 1749 Szemán, J., Ganzler, K., Salgó, A. and Szejtli, J.: Effect of the degree of substitution of cyclodextrin derivatives on chiral separations by high-performance liquid chromatography and capillary electrophoresis. *J. Chromatogr. A*, 728 (1996) 423-431.
- 1750 Tanaka, Y., Yanagawa, M. and Terabe, S.: Separation of enantiomers by capillary electrophoresis using charged cyclodextrins. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 395-400; *C.A.*, 124 (1996) 75129m.
- 1751 Ward, T.J., Dann, C., III and Brown, A.P.: Separation of enantiomers using vancomycin in a countercurrent process by suppression of electroosmosis. *Chirality*, 8 (1996) 77-83; *C.A.*, 124 (1996) 242480c.
- 1752 Wu, H.-L., Otsuka, K. and Terabe, S.: (Optical resolution of drug metabolites by capillary electrophoresis). *Kuromatogurafi*, 16 (1995) 306-307; *C.A.*, 124 (1996) 164151e.

See also 1659, 1704, 1710, 1726, 1732, 1777, 1835, 1845, 1850, 1851, 1852, 1856, 2330, 2343, 2370, 2372, 2373, 2376, 2381, 2383, 2385, 2386, 2388, 2395, 2399, 2414.

4h. Two dimensional electrophoresis

- 1753 Anderson, N.G. and Anderson, N.L.: Twenty years of two-dimensional electrophoresis: past, present and future. *Electrophoresis (Weinheim)*, 17 (1996) 443-453 - a review with 54 refs.
- 1754 Appel, R.D., Bairoch, A., Sanchez, J.-C., Vargas, J.R., Golaz, O., Pasquali, C. and Hochstrasser, D.F.: Federated two-dimensional electrophoresis database: a simple means of publishing two-dimensional electrophoresis data. *Electrophoresis (Weinheim)*, 17 (1996) 540-546.
- 1755 Jordan, B., Lu, Y.F., Bloodworth, I., Rawson, M. and Maxwell, J.: Analysis of molecular variation using two-dimensional gel electrophoresis. Current applications and prospects for the future. *Proc. Malays. Biochem. Soc. Conf.*, 18th (1993) 11-19; *C.A.*, 124 (1996) 169619y.
- 1756 Jungblut, P., Thiede, B., Zimny-Arndt, U., Müller, E.-C., Scheler, C., Wittmann-Liebold, B. and Otto, A.: Resolution power of two-dimensional electrophoresis and identification of proteins from gels. *Electrophoresis (Weinheim)*, 17 (1996) 839-847 - a review with 76 refs.
- 1757 Tsugita, A., Kamo, M., Kawakami, T. and Ohki, Y.: Two-dimensional electrophoresis of plant proteins and standardization of gel patterns. *Electrophoresis (Weinheim)*, 17 (1996) 855-865.
- 1758 Wheeler, C.H., Berry, S.L., Wilkins, M.R., Corbett, J.M., Ou, K., Gooley, A.A., Humphery-Smith, I., Williams, K.L. and Dunn, M.J.: Characterisation of proteins from two-dimensional electrophoresis gels by matrix-assisted laser desorption mass spectrometry and amino acid compositional analysis. *Electrophoresis (Weinheim)*, 17 (1996) 580-587.
- 1759 Wittmann-Liebold, B., Solbrig-Lebuhn, H. and Jungblut, P.: (High performance 2D-polyacrylamide gel electrophoresis. Application in biochemistry, biotechnology and medicine). *Labor-Praxis*, 19 (1995) 34-40; *C.A.*, 124 (1996) 49707d - a review with 41 refs.

See also 1809, 1820, 1868, 1869, 1872, 1884, 1885, 1899, 1904, 1906, 1911, 1912, 1913, 1925, 1927, 1940, 1951, 1961, 1963, 1971, 1975, 1976, 1984, 1990, 1997, 2007, 2012, 2015, 2033, 2051, 2071, 2073, 2081, 2109, 2145, 2158, 2231, 2248, 2277, 2292, 2309.

4i. Other special techniques

1760 Park, Y.: The separation of polyelectrolyte using an electric field in a packed column. *Sep. Sci. Technol.*, 30 (1995) 3629-3647; *C.A.*, 124 (1996) 111538e.

1761 Perrella, M. and Denisov, I.: Low-temperature electrophoresis methods. *Methods Enzymol.*, 259(Energetics of Biological Macromolecules) (1995) 468-487; *C.A.*, 124 (1996) 169822j.

1762 Ross, G., Dittmann, M., Bek, F. and Rozing, G.: Capillary electrochromatography: enhancement of LC separation in packed capillary columns by means of electrically driven mobile phases. *Int. Lab.*, 26, May (1996) 10A-10F.

1763 Roux-de Balman, H., Sanchez, V. and Changeart, F.J.: Study of the continuous flow electrophoresis process - a sounding rocket program. *Eur. Space Agency [Spec. Publ.]*; ESA SP-370(12th ESA Symposium on European Rocket and Balloon Programmes and Related Research, 1995) (1995) 155-159; *C.A.*, 124 (1996) 169811e.

See also 1843, 2130, 2233.

5. HYDROCARBONS AND HALOGEN DERIVATIVES

5b. Cyclic hydrocarbons, fullerenes

1764 Jinno, K. and Sawada, Y.: Relationships between capacity factors and hydrophobicity of polycyclic aromatic hydrocarbons in cyclodextrin-modified micellar electrokinetic chromatography using surface treated capillaries. *J. Liq. Chromatogr.*, 18 (1995) 3719-3727.

See also 1634, 1690.

5c. Halogen derivatives

See 2414.

5d. Complex hydrocarbon mixtures (incl. analysis of tars, bitumens and mineral oils)

See 1814.

7. PHENOLS

See 1709, 1737, 1741.

8. SUBSTANCES CONTAINING HETEROCYCLIC OXYGEN

8a. Flavonoids

1765 Bridle, P., García-Viguera, C. and Tomás-Barberán, F.A.: Analysis of anthocyanins by capillary zone electrophoresis. *J. Liq. Chromatogr.*, 19 (1996) 537-545.

8b. Aflatoxins and other mycotoxins

1766 Oubrahim, H., Cantin-Esnault, D. and Richard, J.-M.: Novel methods for identification and quantitation of the mushroom nephrotoxin orellanine. *Pharmazie*, 51 (1996) 231-234.

8c. Other compounds with heterocyclic oxygen (incl. tannins)

1767 Bogan, D.P., Killard, A.J. and O'Kennedy, R.: The use of capillary electrophoresis [CE] for studying the *in vitro* glucuronidation of 7-hydroxycoumarin. *J. Capillary Electrophor.*, 2 (1995) 241-245; *C.A.*, 124 (1996) 169800a.

9. OXO COMPOUNDS, ETHERS, EPOXIDES AND QUINONES

1768 Feige, K., Ried, T. and Bächmann, K.: Determination of formaldehyde by capillary electrophoresis in the presence of a dihydroxyacetone matrix. *J. Chromatogr. A*, 730 (1996) 333-336.

10. CARBOHYDRATES

10a. Mono and oligosaccharides. Structural studies

1769 Carney, S.L., Cataerson, B. and Penticost, H.R.: The investigation of glycosaminoglycan mimotope structure using capillary electrophoresis and other complementary electrophoretic techniques. *Electrophoresis (Weinheim)*, 17 (1996) 384-390.

1770 El Rassi, Z. and Mechref, Y.: Recent advances in capillary electrophoresis of carbohydrates. *Electrophoresis (Weinheim)*, 17 (1996) 275-301 - a review with 122 refs.

1771 Evangelista, R.A., Guttman, A. and Chen, F.-T.A.: Acid-catalyzed reductive amination of aldoses with 8-aminopyrene-1,3,6-trisulfonate. *Electrophoresis (Weinheim)*, 17 (1996) 347-351.

1772 Guttman, A. and Herrick, S.: Effect of the quantity and linkage position of mannose(α 1,2) residues in capillary gel electrophoresis of high mannose-type oligosaccharides. *Anal. Biochem.*, 235 (1996) 236-239.

1773 Hoffstetter-Kuhn, S., Alt, G. and Kuhn, R.: Profiling of oligosaccharide-mediated microheterogeneity of a monoclonal antibody by capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 418-422.

1774 Honda, S., Saitoh, Y., Kawakami, S., Suzuki, S. and Takechi, K.: (Analysis of amino sugars by LIF-HPCE as NBD-labeled derivatives). *Kuromatogurafi*, 16 (1995) 310-311; *C.A.*, 124 (1996) 218978a.

- 1775 Honda, S., Togashi, K., Sasaoka, I., Miyoshi, A. and Taga, A.: (Capillary electrophoresis of PMP derivatives of pentose epimers). *Kuromatogurafi*, 16 (1995) 308-309; *C.A.*, 124 (1996) 218985a.
- 1776 Karamanos, N.K., Vanky, P., Tzanakakis, G.N. and Hjerpe, A.: High performance capillary electrophoresis method to characterize heparin and heparan sulfate disaccharides. *Electrophoresis (Weinheim)*, 17 (1996) 391-395.
- 1777 Lee, Y.-H. and Lin, T.-I.: Capillary electrophoretic analysis of cyclodextrins and determination of formation constants for inclusion complexes. *Electrophoresis (Weinheim)*, 17 (1996) 333-340.
- 1778 Lu, B. and Westerlund, D.: Indirect UV detection of carbohydrates in capillary zone electrophoresis by using tryptophan as a marker. *Electrophoresis (Weinheim)*, 17 (1996) 325-332.
- 1779 Mort, A.J. and Chen, E.M.W.: Separation of 8-aminonaphthalene-1,3,6-trisulfonate (ANTS)-labeled oligomers containing galacturonic acid by capillary electrophoresis: application to determining the substrate specificity of endopolygalacturonases. *Electrophoresis (Weinheim)*, 17 (1996) 379-383.
- 1780 O'Neill, R.A., O'Neill, R.A. and Khan, S.H.: Method and apparatus for automated carbohydrate mapping and sequencing. *PCT Int. Appl. WO 95 33,989 (Cl. GO1N27/447)*, 14 Dec. 1995, US Appl. 252,657, 2 Jun. 1994; 63 pp.; *C.A.*, 124 (1996) 197716w.
- 1781 O'Shea, M.G. and Morell, M.K.: High resolution slab gel electrophoresis of 8-amino-1,3,6-pyrenetrisulfonic acid (APTS) tagged oligosaccharides using a DNA sequencer. *Electrophoresis (Weinheim)*, 17 (1996) 681-688.
- 1782 Perez, S.A. and Colón, L.A.: Determination of carbohydrates as their dansylhydrazine derivatives by capillary electrophoresis with laser-induced fluorescence detection. *Electrophoresis (Weinheim)*, 17 (1996) 352-358.
- 1783 Weber, P.L. and Lunte, S.M.: Capillary electrophoresis with pulsed amperometric detection of carbohydrates and glycopeptides. *Electrophoresis (Weinheim)*, 17 (1996) 302-309.
- 1784 Zhang, Z., Pierce, M.L. and Mort, A.J.: Detection and differentiation of pectic enzyme activity *in vitro* and *in vivo* by capillary electrophoresis of products from fluorescent-labeled substrate. *Electrophoresis (Weinheim)*, 17 (1996) 372-378.
- 1785 Zhou, W. and Baldwin, R.P.: Capillary electrophoresis and electrochemical detection of underivatized oligo- and polysaccharides with surfactant-controlled electroosmotic flow. *Electrophoresis (Weinheim)*, 17 (1996) 319-324.
- See also 1644, 1652, 1654, 1787, 1796, 1808.
- 10b. *Polysaccharides, mucopolysaccharides, lipopolysaccharides*
- 1786 Grimshaw, J., Trocha-Grimshaw, J., Fisher, W., Rice, A., Smith, S., Spedding, P., Duffy, J. and Mollan, R.: Quantitative analysis of hyaluronan in human synovial fluid using capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 396-400.
- 1787 Harland, G.B., Okafo, G., Matejtschuk, P., Sellick, I.C., Chapman, G.E. and Camilleri, P.: Fingerprinting of glycans as their 2-aminoacridone derivatives by capillary electrophoresis and laser-induced fluorescence. *Electrophoresis (Weinheim)*, 17 (1996) 406-411.
- 1788 Kouznetsov, D.A., Ivanov, A.A., Veletsky, P.R., Charsky, V.I. and Beklemishev, O.S.: A laboratory model for studying environmentally dependent chemical modifications in textile cellulose. *Text. Res. J.*, 66 (1996) 111-114; *C.A.*, 124 (1996) 119953g.
- 1789 Liu, J., Desai, U.R., Han, X.-J., Toida, T. and Linhardt, R.J.: Strategy for the sequence analysis of heparin. *Glycobiology*, 5 (1995) 765-774; *C.A.*, 124 (1996) 197648a.
- 1790 Malsch, R. and Harenberg, J.: Purity of glycosaminoglycan-related compounds using capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 401-405.
- 1791 Mayer, S. and Schleimer, M.: Quantitative determination of heparinoid mimetics in human and rat plasma by micellar electrokinetic chromatography. *J. Chromatogr. A*, 730 (1996) 297-303.
- 1792 Piraud, M., Froissart, R., Mandon, G., Bernard, A. and Maire, I.: Amniotic fluid for screening of lysosomal storage diseases presenting in utero (mainly as non-immune hydrops fetalis). *Clin. Chim. Acta*, 248 (1996) 143-155.
- 1793 Toida, T. and Linhardt, R.J.: Detection of glycosaminoglycans as a copper(II) complex in capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 341-346.
- See also 1769, 1785, 2079.
- 10c. *Glycoproteins and their constituents*
- 1794 Carpenter, G.H., Proctor, G.B., Pankhurst, C.L., Zhang, X.S. and Shori, D.K.: Lectin probe analysis of the glycosylation of human parotid salivary glycoproteins. *Biochem. Soc. Trans.*, 23 (1995) 538S; *C.A.*, 124 (1996) 197354b.
- 1795 Carron, J.A., Bates, R.C., Smith, A.I., Tetz, T., Arellano, A., Gordon, D.L. and Burns, G.F.: Factor H co-purifies with thrombospondin isolated from platelet secrete. *Biochim. Biophys. Acta*, 1289 (1996) 305-311.
- 1796 Guttman, A., Chen, F.-T.A. and Evangelista, R.A.: Separation of 1-aminopyrene-3,6,8-trisulfonate-labeled asparagine-linked fetuin glycans by capillary gel electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 412-417.
- 1797 Hashimoto, K., Tobe, T., Sumiya, J.-i., Sano, Y., Choi-Miura, N.-H., Ozawa, A., Yasue, H. and Tomita, M.: Primary structure of the pig homologue of human IHRP: inter- α -trypsin inhibitor family heavy chain-related protein. *J. Biochem. (Tokyo)*, 119 (1996) 577-584.
- 1798 Hennebicq-Reig, S., Soudan, B., Tetaert, D., Briand, G., Richet, C., Demeyer, D. and Degand, P.: Capillary electrophoresis separation of mucin motif MUC 5AC glycopeptides from *in vitro* O-glycosylation reactions. *Carbohydr. Res.*, 280 (1996) 121-126; *C.A.*, 124 (1996) 111551d.
- 1799 Hervé, F., Millot, M.-C., Eap, C.B., Duché, J.-C. and Tillement, J.-P.: Two-step chromatographic purification of human plasma α_1 -acid glycoprotein. Its application to the purification of rare phenotype samples of the protein and their study by chromatography on immobilized metal chelate affinity adsorbent. *J. Chromatogr. B*, 678 (1996) 1-14.
- 1800 Jandik, K.A., Kruep, D., Cartier, M. and Linhardt, R.J.: Accelerated stability studies of heparin. *J. Pharm. Sci.*, 85 (1996) 45-51.

- 1801 Kalabay, L., Mathur, S., Bobin, S. and Arnaud, P.: Electrophoretic and isoelectric focusing analysis of human recombinant alpha2-HS glycoprotein produced in insect cells: analysis of the post-translational events. *Electrophoresis (Weinheim)*, 17 (1996) 529-532.
- 1802 Kamemura, K., Furuichi, Y., Umekawa, H. and Takahashi, T.: Purification and characterization of a pod lectin from Great Northern bean, *Phaseolus vulgaris* L. *Biochim. Biophys. Acta*, 1289 (1996) 87-94.
- 1803 Kobayashi, D., Takita, H., Mizuno, M., Totsuka, Y. and Kuboki, Y.: Time-dependent expression of bone sialoprotein fragments in osteogenesis induced by bone morphogenetic protein. *J. Biochem. (Tokyo)*, 119 (1996) 475-481.
- 1804 Lampreaue, F., Alava, M.A. and Pineiro, A.: Developments in lectin affinity immunoelectrophoresis of serum glycoproteins. *TrAC*, 15 (1996) 122-129.
- 1805 Mertz, E., Sareneva, T., Haebel, S., Julkunen, I. and Roepstorff, P.: Mass spectrometric characterization of glycosylated interferon- γ variants separated by gel electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 925-931.
- 1806 Moorhouse, K.G., Rickel, C.A. and Chen, A.B.: Electrophoretic separation of recombinant tissue-type plasminogen activator glycoforms: validation issues for capillary isoelectric focusing methods. *Electrophoresis (Weinheim)*, 17 (1996) 423-430.
- 1807 Oda, R.P. and Landers, J.P.: Effect of cationic buffer additives on the capillary electrophoretic separation of serum transferrin from different species. *Electrophoresis (Weinheim)*, 17 (1996) 431-437.
- 1808 Oda, R.P., Madden, B.J. and Landers, J.P.: Capillary electrophoretic analysis of glycoproteins and glycoprotein-derived oligosaccharides. *Adv. Chromatogr.*, 36 (1996) 163-199; *C.A.*, 124 (1996) 140050v - a review with 99 refs.
- 1809 Packer, N.H., Wilkins, M.R., Golaz, O., Lawson, M.A., Gooley, A.A., Hochstrasser, D.F., Redmond, J.W. and Williams, K.L.: Characterization of human plasma glycoproteins separated by two-dimensional gel electrophoresis. *BioTechnology*, 14 (1996) 66-70; *C.A.*, 124 (1996) 140227h.
- 1810 Toombs, C.F., Degraaf, G.L., Martin, J.P., Geng, J.G., Anderson, D.C. and Shebuski, R.J.: Pretreatment with a blocking monoclonal antibody to P-selectin accelerates pharmacological thrombolysis in a primate model of arterial thrombosis. *J. Pharmacol. Exp. Ther.*, 275 (1995) 941-949.
- 1811 Yasuda, T., Takeshita, H., Tsubota, E., Sawazaki, K., Iida, R., Nadano, D. and Kishi, K.: Alpha-2-HS-glycoprotein (AHSG) polymorphism in semen and saliva. *Electrophoresis (Weinheim)*, 17 (1996) 793-796.
- 1812 Bähre, F. and Maier, H.-G.: Electrophoretic clean-up of organic acids from coffee for GC/MS analysis. *Fresenius J. Anal. Chem.*, 355 (1996) 190-193.
- 1813 Chen, H., Xu, Y., van Lente, F. and Ip, M.P.C.: Indirect ultraviolet detection of biologically relevant organic acids by capillary electrophoresis. *J. Chromatogr. B*, 679 (1996) 49-59.
- 1814 Lin, Q., Liu, G. and Cheng, Z.: (Analysis of organic acids in degradation product of kerogen and oil field water by high performance capillary electrophoresis). *Fenxi Ceshi Xuebao*, 14 (1995) 49-53; *C.A.*, 124 (1996) 206556y.
- 1815 Marsh, D.B. and Nuttall, K.L.: Methylmalonic acid in clinical urine specimens by capillary zone electrophoresis using indirect photometric detection. *J. Capillary Electrophor.*, 2 (1995) 63-67; *C.A.*, 124 (1996) 81288d.
- 1816 Schmitt, P., Kettrup, A., Freitag, D. and Garrison, A.W.: Flocculation of humic substances with metal ions as followed by capillary zone electrophoresis. *Fresenius J. Anal. Chem.*, 354 (1996) 915-920.
- 1817 Simon, P. and Nicot, T.: Capillary electrophoresis and supercritical chromatography, complementary and alternative techniques for the determination of urinary metabolites of styrene. *J. Chromatogr. B*, 679 (1996) 103-112.
- 1818 Weers, M., Balzer, H., Bradbury, A. and Vitzthum, O.G.: Analysis of acids in coffee by capillary electrophoresis. *Colloq. Sci. Int. Cafe, [C.R.]*, 16th(Vol. 1, Seizieme Colloque Scientifique International sur le Cafe, 1995) (1995) 218-223; *C.A.*, 124 (1996) 143904n.
- 1819 Weiss, J. and Reinhard, S.: (Aliphatic carboxylic acids, A difficult chromatographic problem?) *GIT Spez. Chromatogr.*, 15 (1995) 105-111; *C.A.*, 124 (1996) 105084n - a review with 16 refs.

See also 1639, 1741, 1750, 2361, 2405, 2434, 2436.

11d. Lipoproteins and their constituents

- 1820 Barrans, A., Jaspard, B., Barbaras, R., Chap, H., Perret, B. and Collet, X.: Pre- β HDL: structure and metabolism. *Biochim. Biophys. Acta*, 1300 (1996) 73-85 - a review with 150 refs.
- 1821 Bravo, E., Rivabene, R., Bruscalupi, G., Calcabrini, A., Arancia, G. and Cantafora, A.: Age-related changes in lipid secretion of perfused livers from male Wistar rats donors. *J. Biochem. (Tokyo)*, 119 (1996) 240-245.
- 1822 Brouwer, D.A.J., van Doormaal, J.J. and Muskiet, F.A.J.: Clinical chemistry of common apolipoprotein E isoforms. *J. Chromatogr. B*, 678 (1996) 23-41 - a review with 117 refs.
- 1823 Castillo, M., Hortal, J.H., García-Fuentes, E., Zafra, M.F. and Barcia-Peregrín, E.: Coconut oil affects lipoprotein composition and structure of neonatal chicks. *J. Biochem. (Tokyo)*, 119 (1996) 610-616.
- 1824 Furbee, J.W., Jr. and Fless, G.M.: Evaluation of common electrophoretic methods in determining the molecular weight of apolipoprotein(a) polymorphs. *Anal. Biochem.*, 234 (1996) 66-73.
- 1825 Marcovina, S.M., Hobbs, H.H. and Albers, J.J.: Relation between number of apolipoprotein(a) kringle 4 repeats and mobility of isoforms in agarose gel: basis for a standardized isoform nomenclature. *Clin. Chem. (Washington)*, 42 (1996) 436-439.
- 1826 Pearson, J.P. and Keren, D.F.: The effects of heparin on lipoproteins in high-resolution electrophoresis of serum. *Am. J. Clin. Pathol.*, 104 (1995) 468-471; *C.A.*, 124 (1996) 49932y.

11. ORGANIC ACIDS AND LIPIDS

11a. Organic acids and simple esters

- 1812 Bähre, F. and Maier, H.-G.: Electrophoretic clean-up of organic acids from coffee for GC/MS analysis. *Fresenius J. Anal. Chem.*, 355 (1996) 190-193.

- 1827 Tallet, F., Vasson, M.P., Couderc, R., Lefevre, G. and Raichvarg, D.: Characterization of lipoproteins during human cholestasis. *Clin. Chim. Acta*, 244 (1996) 1-15.
- 1828 Zhang, L., Kotite, L. and Havel, R.J.: Identification, characterization, cloning, and expression of apolipoprotein C-IV, a novel sialoglycoprotein of rabbit plasma lipoproteins. *J. Biol. Chem.*, 271 (1996) 1776-1783.

See also 1975.

12. ORGANIC PEROXIDES

See 1831.

13. STEROIDS

13a. General techniques

- 1829 Salo, M., Sirén, H., Volin, P., Wiedmer, S. and Vuorela, H.: Structure-retention relationships of steroid hormones in reversed-phase liquid chromatography and micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 728 (1996) 83-88.

13b. Pregnane and androstane derivatives

- 1830 Fernandez, C., Egginger, G., Wainer, I.W. and Lloyd, D.K.: Separation of testosterone metabolites in microsomal incubates using a new capillary electrophoresis assay. *J. Chromatogr. B*, 677 (1996) 363-368.

13d. Sterols

- 1831 Brown, A.J., Dean, R.T. and Jessup, W.: Free and esterified oxysterol: formation during copper-oxidation of low density lipoprotein and uptake by macrophages. *J. Lipid Res.*, 37 (1996) 320-335.

15. TERPENES AND OTHER VOLATILE AROMATIC COMPOUNDS

15a. Terpenes

- 1832 Mauri, P., Catalano, G., Gardana, C. and Pietta, P.: Analysis of *Stevia* glycosides by capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 367-371.

16. NITRO AND NITROSO COMPOUNDS

- 1833 Kennedy, S., Caddy, B. and Douse, J.M.F.: Micellar electrokinetic capillary chromatography of high explosives utilising indirect fluorescence detection. *J. Chromatogr. A*, 726 (1996) 211-222.

- 1834 Oehrle, S.A.: Analysis of anionic impurities in ammonium dinitrosamide (ADN). Part II. Capillary ion electrophoresis. *J. Energ. Mater.*, 14 (1996) 37-45; *C.A.*, 124 (1996) 121366m.

17. AMINES, AMIDES AND RELATED NITROGEN COMPOUNDS

17a. Amines and polyamines

- 1835 Fillet, M., Bechet, I., Ceccato, A. and Crommen, J.: Optimization of enantiomeric separations for basic compounds in capillary zone electrophoresis using neutral and anionic cyclodextrins as chiral selectors. *Biomed. Chromatogr.*, 9 (1995) 265-266; *C.A.*, 124 (1996) 66731z.

See also 1690.

17b. Catecholamines and their metabolites

- 1836 Bert, L., Robert, F., Denoroy, L. and Renaud, B.: High-speed separation of subnanomolar concentrations of noradrenaline and dopamine using capillary zone electrophoresis with laser-induced fluorescence detection. *Electrophoresis (Weinheim)*, 17 (1996) 523-525.

17c. Urea and guanidine derivatives

- 1837 Geise, R.J., Machnicki, N.I. and Ianniello, R.M.: Determination of diazolidinyl urea in cosmetic formulations by capillary electrophoresis. *J. Soc. Cosmet. Chem.*, 45 (1994) 173-181; *C.A.*, 124 (1996) 97233t.
- 1838 Michalakis, G.M. and Barry, E.F.: Determination of imidurea in cosmetic products by capillary zone electrophoresis and micellar electrokinetic chromatography. *J. Soc. Cosmet. Chem.*, 45 (1994) 193-202; *C.A.*, 124 (1996) 97234u.

17d. Other amine derivatives and amides (excl. peptides)

- 1839 Lin, C.-e., Chiou, W.-c. and Lin, W.-c.: Capillary zone electrophoretic separation of alkylbenzyl quaternary ammonium compounds: Effect of organic modifier. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 419-425; *C.A.*, 124 (1996) 75130e.
- 1840 Lin, C.-e., Chiou, W.-c. and Lin, W.-c.: Separation of alkylbenzyl quaternary ammonium compounds by capillary zone electrophoresis. Effect of organic solvent in sample solution. *J. Chromatogr. A*, 723 (1996) 189-195.

18. AMINO ACIDS AND PEPTIDES; CHEMICAL STRUCTURE OF PROTEINS
- 18a. *Amino acids and their derivatives*
- 1841 Dawson, L.A., Stow, J., Dourish, C.T. and Routledge, C.: The analysis of excitatory amino acids in striatal micro dialyzates using capillary electrophoresis and laser induced fluorescence detection. In: Louilot, A. (Editor), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 6th 1994*, Universite de Bordeaux I, Talence, 1994, pp. 31-32; C.A., 124 (1996) 135876q.
- 1842 De Antonis, K.M.: The analysis of amino acids and peptides by high performance liquid chromatography and capillary electrophoresis using the novel derivatizing agent, 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate. Avail. *Univ. Microfilms Int.*, Order No.: DA9601831, 1995, 349 p.; C.A., 124 (1996) 192909b.
- 1843 Hernandez, L., Paez, X., Tucci, S., Murzi, E., Rodriguez, N. and Baptista, T.: New data on capillary zone electrophoresis and microdialysis coupled to laser induced fluorescence detection. In: Louilot, A. (Editor), *Monit. Mol. Neurosci., Proc. Int. Conf. In Vivo Methods, 6th 1994*, Universite de Bordeaux I, Talence, 1994, pp. 27-28; C.A., 124 (1996) 135874n.
- 1844 Mechref, Y., Smith, J.T. and El Rassi, Z.: Micellar electrokinetic capillary chromatography with *in situ* charged micelles. VII. Expanding the utility of alkylglycoside-borate micelles to acidic and neutral pH for capillary electrophoresis of dansyl amino acids and herbicides. *J. Liq. Chromatogr.*, 18 (1995) 3769-3786.
- 1845 Nouadje, G. and Couderc, F.: (Chiral separation of fluoresceinthiocarbamoyl amino acids by capillary electrophoresis together with fluorescence detection by a laser). *Spectra Anal.*, 24 (1995) 17-21; C.A., 124 (1996) 105141d.
- 1846 Orwar, O., Fishman, H.A., Sundahl, M., Banthia, V., Dadoo, R. and Zare, R.N.: Determination of photodestruction quantum yields using capillary electrophoresis: application to α -phthalaldehyde/ β -mercaptoethanol-labeled amino acids. *J. Liq. Chromatogr.*, 18 (1995) 3833-3846.
- 1847 Salahudeen, I.: Development of electrogenic derivatizing reagents and amino acids analysis using capillary electrophoresis-electrochemistry. Avail. *Univ. Microfilms Int.*, Order No.: DA9544857, 1995, 310 p.; C.A., 124 (1996) 163831q.
- 1848 Suiko, M., Sakakibara, Y., Nakajima, H., Sakaida, H. and Liu, M.-C.: Enzymic sulphation of dopa and tyrosine isomers by HepG2 human hepatoma cells: stereoselectivity and stimulation by Mn^{2+} . *Biochem. J.*, 314 (1996) 151-158.
- 1849 Taga, A., Suginaura, M. and Honda, S.: (Derivatization at the inlet and inside of a capillary in capillary electrophoresis). *Kuro-matogurafi*, 16 (1995) 302-303; C.A., 124 (1996) 127654a.
- 1850 Van Hove, E. and Sandra, P.: Considerations on the enantiomeric separation by MEKC of N-Bz-amino acids with N-dodecylcarboxylvaline as chiral selector. *J. Liq. Chromatogr.*, 18 (1995) 3675-3683.
- 1851 Vogt, C.: (Enantiomeric separation of D/L-carnitine by HPLC and capillary zone electrophoresis (CZE) after derivatization). *GIT Fachz. Lab.*, 39 (1995) 1035-1037; C.A., 124 (1996) 127213f.
- 1852 Wan, H., Engström, A. and Blomberg, L.G.: Direct chiral separation of amino acids derivatized with 2-(9-anthryl)ethyl chloroformate by capillary electrophoresis using cyclodextrins as chiral selectors. Effect of organic modifiers on resolution and enantiomeric elution order. *J. Chromatogr. A*, 731 (1996) 283-292.
- See also 1650, 1651, 1654, 1693, 1702, 1713, 1750, 1751, 1954.
- 18b. *Peptides, peptidic and proteinous hormones, growth factors*
- 1853 Baumann, M., Golabek, A., Lalowski, M. and Wisniewski, T.: Micropreparative gel electrophoresis of low-molecular-weight peptides: purification of highly insoluble amyloid peptide fragments. *Anal. Biochem.*, 236 (1996) 191-198.
- 1854 Borowicz, B., Goluszko, E., Wielinska, L., Jagla, K. and Domaniewski, J.: (Electrophoretic analysis of polypeptides in cells after DNA transfection from mammary carcinoma). *Diagn. Lab.*, 29 (1993) 295-303; C.A., 124 (1996) 81277z.
- 1855 Castagnola, M., Messina, I. and Rossetti, D.V.: Capillary zone electrophoresis for the analysis of peptides. In: Righetti, P.G. (Editor), *Capillary Electrophor. Anal. Biotechnol.*, CRC, Boca Raton, 1996, pp. 239-275; C.A., 124 (1996) 197554s.
- 1856 DeSilva, K., Jiang, Q. and Kuwana, T.: Chiral separation of naphthalene-2,3-dialdehyde labeled peptides by cyclodextrin-modified electrokinetic chromatography. *Biomed. Chromatogr.*, 9 (1995) 295-301; C.A., 124 (1996) 134358k.
- 1857 Hagono, P.A., Baeyens, W.R.G. and van der Weken, G.: Capillary electrophoretic separation of some pharmaceutically important polypeptides. *Biomed. Chromatogr.*, 9 (1995) 291; C.A., 124 (1996) 66713v.
- 1858 Hermansen, L.F., Juul, J. and Sletten, K.: Sodium dodecyl sulfate-polyacrylamide gel electrophoresis and centrifuge blotting: preparation of polypeptides for amino-terminal sequence analysis. *Electrophoresis (Weinheim)*, 17 (1996) 720-722.
- 1859 Herold, M., Ross, G.A., Grimm, R. and Heiger, D.N.: Separation of peptides and protein digests by capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 285-308; C.A., 124 (1996) 197521d.
- 1860 Idei, M., Mezö, I., Vadász, Z., Horváth, A., Seprödi, J., Ércegyi, J., Teplán, I. and Kéri, G.: Comparative analysis of somatostatin analog peptides by capillary electrophoresis and micellar electrokinetic chromatography. *Electrophoresis (Weinheim)*, 17 (1996) 758-761.
- 1861 Janáky, T., Szabó, E., Balásipri, L., Adi, B. and Penke, B.: High-performance separation methods in the analysis of a new peptide family: the galanins. *J. Chromatogr. B*, 676 (1996) 7-12.
- 1862 Takemura, M., Ueno, H. and Kodama, H.: S-(1,2-Dicarboxyethyl)glutathione and glutathione in lens and liver of naphthalene-treated rabbits. *Eur. J. Clin. Chem. Clin. Biochem.*, 34 (1996) 85-90.
- 1863 Tomlinson, A.J. and Naylor, S.: Systematic development of on-line membrane preconcentration-capillary electrophoresis-mass spectrometry for the analysis of peptide mixtures. *J. Capillary Electrophor.*, 2 (1995) 225-233; C.A., 124 (1996) 197634t.
- 1864 Wang, Y., Liu, J., Mamidi, M. and Walker, A.M.: Identification of the major site of rat prolactin phosphorylation as serine 177. *J. Biol. Chem.*, 271 (1996) 2462-2469.

- 1865 Weiler, A., Stoeva, S., Lehmann, R., Grübler, G., Paulus, G. and Voelter, W.: Characterization of natural peptides from bovine tissue using capillary electrophoresis, high performance liquid chromatography, matrix-assisted laser desorption ionization, and Edman degradation. *Electrophoresis (Weinheim)*, 17 (1996) 518-522.
- 1866 Wycherley, D., Rose, M.E., Giles, K., Jarvis, S.A. and McDowall, M.A.: From dipeptides to oligopeptides: analysis by capillary electrophoresis/electrospray mass spectrometry. *Biomed. Chromatogr.*, 9 (1995) 283-284; *C.A.*, 124 (1996) 140289e.
- See also 1688, 1701, 1738, 1750, 1842, 1882, 1933, 1972, 2103, 2398, 2410.
- 18c. Elucidation of structure of proteins and enzymes**
- 1867 Casari, G., Sander, C. and Valencia, A.: Sequence space analysis: identification of functionally or structurally important residues in protein sequence families. In: Bohr, H. and Brunak, S. (Editors), *Protein Folds, [Distance-Based Approaches Protein Struct. Determ., Symp.]*, 2nd 1994, CRC, Boca Raton, 1996, pp. 124-131; *C.A.*, 124 (1996) 50145a.
- 1868 Furui, T., Ikeda, M., Chao-Ming, L., Okita, K. and Nakamura, K.: Protein degradation in human pure pancreatic juice analyzed by two-dimensional gel electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 797-802.
- 1869 Fushitani, K., Higashiyama, K., Asao, M. and Hosokawa, K.: Characterization of the constituent polypeptides of the extracellular hemoglobin from *Lumbricus terrestris*: heterogeneity and discovery of a new linker chain L4. *Biochim. Biophys. Acta*, 1292 (1996) 273-280.
- 1870 Galzie, Z., Qasim, M.A. and Salahuddin, A.: Isolation and characterization of domain I of ovoinhibitor. *Biochim. Biophys. Acta*, 1293 (1996) 113-121.
- 1871 Gevaert, K., Verschelde, J.-L., Puype, M., van Damme, J., Goethals, M., de Boeck, S. and Vandekerckhove, J.: Structural analysis and identification of gel-purified proteins, available in the femtomole range, using a novel computer program for peptide sequence assignment, by matrix-assisted laser desorption ionization - reflectron time-of-flight - mass spectrometry. *Electrophoresis (Weinheim)*, 17 (1996) 918-924.
- 1872 Golaz, O., Wilkins, M.R., Sanchez, J.-C., Appel, R.D., Hochstrasser, D.F. and Williams, K.L.: Identification of proteins by their amino acid composition: an evaluation of the method. *Electrophoresis (Weinheim)*, 17 (1996) 573-579.
- 1873 Houthaevae, T., Gausepohl, H., Mann, M. and Ashman, K.: Automation of micro-preparation and enzymic cleavage of gel electrophoretically separated proteins. *FEBS Lett.*, 376 (1995) 91-94; *C.A.*, 124 (1996) 50114q.
- 1874 Inglis, A.S., Reid, G.E. and Simpson, R.J.: Chemical techniques employed for the primary structural analysis of proteins and peptides. *EXS*, 73 (1995) 141-171; *C.A.*, 124 (1996) 169551v - a review with many refs.
- 1875 Jensen, O.N., Vorm, O. and Mann, M.: Sequence patterns produced by incomplete enzymatic digestion or one-step Edman degradation of peptide mixtures as probes for protein database searches. *Electrophoresis (Weinheim)*, 17 (1996) 938-944.
- 1876 Jensen, P.E.H., Arbelaez, L.F., Shanbhag, V.P. and Stigbrand, T.: Preparation and characterization of a C-terminal fragment of pregnancy zone protein corresponding to the receptor-binding peptide from human α_2 -macroglobulin. *Biochim. Biophys. Acta*, 1293 (1996) 254-258.
- 1877 Kornfelt, T., Vinther, A., Okafo, G.N. and Camilleri, P.: Improved peptide mapping using phytic acid as ion-pairing buffer additive in capillary electrophoresis. *J. Chromatogr. A*, 726 (1996) 223-228.
- 1878 Luquet, G., Testenière, O. and Graf, F.: Characterization and N-terminal sequencing of a calcium binding protein from the calcareous concretion organic matrix of the terrestrial crustacean *Orchestia cavimana*. *Biochim. Biophys. Acta*, 1293 (1996) 272-276.
- 1879 Patterson, S.D., Thomas, D. and Bradshaw, R.A.: Application of combined mass spectrometry and partial amino acid sequence to the identification of gel-separated proteins. *Electrophoresis (Weinheim)*, 17 (1996) 877-891.
- 1880 Rickard, E.C. and Towns, J.K.: The use of capillary electrophoresis for peptide mapping of proteins. In: Hancock, W.S. (Editor), *New Methods Pept. Mapp. Charact. Proteins*, CRC, Boca Raton, 1996, pp. 97-117; *C.A.*, 124 (1996) 80996w - a review with 36 refs.
- 1881 Schnölzer, M., Jedrzejewski, P. and Lehmann, W.D.: Protease-catalyzed incorporation of ^{18}O into peptide fragments and its application for protein sequencing by electrospray and matrix-assisted laser desorption/ionization mass spectrometry. *Electrophoresis (Weinheim)*, 17 (1996) 945-953.
- 1882 Tomlinson, A.J. and Naylor, S.: A strategy for sequencing peptides for dilute mixtures at the low femtomole level using membrane preconcentration-capillary electrophoresis-tandem mass spectrometry (mPC-CE-MS/MS). *J. Liq. Chromatogr.*, 18 (1995) 3591-3615.
- 1883 Wilm, M., Shevchenko, A., Houthaevae, T., Breit, S., Schweigerer, L., Fotsis, T. and Mann, M.: Femtomole sequencing of proteins from polyacrylamide gels by nano-electrospray mass spectrometry. *Nature (London)*, 379 (1996) 466-469; *C.A.*, 124 (1996) 169972h.
- See also 1676, 1758, 1805, 1859, 1891, 1903, 1954, 1992, 2000, 2037, 2046, 2069, 2078, 2090, 2103, 2156.
- 19. PROTEINS**
- 19a. General techniques**
- 1884 Appel, R.D., Sanchez, J.-C., Bairoch, A., Golaz, O., Ravier, F., Pasquali, C., Hughes, G.J. and Hochstrasser, D.F.: The SWISS-2DPAGE database of two-dimensional polyacrylamide gel electrophoresis, its status in 1995. *Nucleic Acids Res.*, 24 (1996) 180-181; *C.A.*, 124 (1996) 139007e.
- 1885 Arnott, D.P., Henzel, W.J. and Stults, J.T.: Identification of proteins from two-dimensional electrophoresis gels by peptide mass fingerprinting. *ACS Symp. Ser.*, 619 (Biochemical and Biotechnological Applications of Electrospray Ionization Mass Spectrometry) (1996) 226-243; *C.A.*, 124 (1996) 197496z.

- 1886 Baek, W.-O., Haupt, K., Colin, C. and Vijayalakshmi, M.A.: Immobilized metal ion affinity gel electrophoresis: quantification of protein affinity to transition metal chelates. *Electrophoresis (Weinheim)*, 17 (1996) 489-492.
- 1887 Burkhart, W.A., Moyer, M.B., Bailey, J.M. and Miller, C.G.: Electroblothing of proteins to teflon tape and membranes for N- and C-terminal sequence analysis. *Anal. Biochem.*, 236 (1996) 364-367.
- 1888 Cade-Treyer, D., Cade, A., Darjo, A. and Jouvion-Moreno, M.: Isoelectric focusing and titration curves in biomedicine and in agrofood industries: a multimedia teaching program. *Electrophoresis (Weinheim)*, 17 (1996) 479-482.
- 1889 Chen, D.: (Protein, gel electrophoresis and the application of gel electrophoresis in protein analysis). *Fenxi Kexue Xuebao*, 11 (1995) 75-86; *C.A.*, 124 (1996) 140031q - a review with 49 refs.
- 1890 Costa, M.A., Duro, G., Porcasi, R., Colombo, P., Locorotondo, G., Geraci, D. and Izzo, V.: Isolation of proteins and nucleic acids by electrophoresis on disposable gel columns. *Electrophoresis (Weinheim)*, 17 (1996) 781-783.
- 1891 Eckerskorn, C. and Grimm, R.: Enhanced *in situ* gel digestion of electrophoretically separated proteins with automated peptide elution onto mini reversed-phase columns. *Electrophoresis (Weinheim)*, 17 (1996) 899-906.
- 1892 Funderburgh, J.L. and Prakash, S.: SDS-polyacrylamide gel electrophoretic analysis of proteins in the presence of guanidinium hydrochloride. *BioTechniques*, 20 (1996) 376-378; *C.A.*, 124 (1996) 197555t.
- 1893 Hartley, J.L.: Gene construction and cloning for a multidomain protein used as a protein size marker ladder in gel electrophoresis. *U.S. US 5,449,758* (Cl. 530-350; CO7K14/00), 12 Sep. 1995, Appl. 160,670, 2 Dec. 1993; 25 pp.; *C.A.*, 124 (1996) 50193q.
- 1894 Haupt, K., Roy, F. and Vijayalakshmi, M.A.: Immobilized metal ion capillary electrophoresis of proteins - a model for affinity capillary electrophoresis using soluble polymer-supported ligands. *Anal. Biochem.*, 234 (1996) 149-154.
- 1895 Huang, M., Mitchell, D. and Bigelow, M.: Highly efficient protein separations in high-performance capillary electrophoresis, using hydrophilic coatings on polysiloxane-bonded columns. *J. Chromatogr. B*, 677 (1996) 77-84.
- 1896 Iki, N. and Yeung, E.S.: Non-bonded poly(ethylene oxide) polymer-coated column for protein separation by capillary electrophoresis. *J. Chromatogr. A*, 731 (1996) 273-282.
- 1897 Janini, G.M., Fisher, R.J., Henderson, L.E. and Issaq, H.J.: Application of capillary zone electrophoresis for the analysis of proteins, protein-small molecules, and protein-DNA interactions. *J. Liq. Chromatogr.*, 18 (1995) 3617-3628.
- 1898 Koegler, W.S. and Ivory, C.F.: Focusing proteins in an electric field gradient. *J. Chromatogr. A*, 726 (1996) 229-236.
- 1899 Lemkin, P.F., Chipperfield, M., Merrill, C. and Zullo, S.: A World Wide Web (WWW) server database engine for an organelle database, MitoDat. *Electrophoresis (Weinheim)*, 17 (1996) 566-572.
- 1900 Liang, X., Bai, J., Liu, Y.-H. and Lubman, D.M.: Characterization of SDS - PAGE-separated proteins by matrix-assisted laser desorption/ionization mass spectrometry. *Anal. Chem.*, 68 (1996) 1012-1018.
- 1901 Lin, B., Dai, Z., Han, X., Jiang, Y., Yang, G. and Lu, Y.: (Separation and analysis of components in tissue culture solution of fetal hypophysis by multi-dimensional analysis system). *Fenxi Ceshi Xuebao*, 14 (1995) 22-27; *C.A.*, 124 (1996) 197701n.
- 1902 Liu, Y., Fu, R. and Gu, J.: Epoxy resin coatings for capillary zone electrophoretic separation of basic proteins. *J. Chromatogr. A*, 723 (1996) 157-167.
- 1903 Moritz, R.L., Eddes, J.S., Reid, G.E. and Simpson, R.J.: S-Pyridylethylation of intact polyacrylamide gels and *in situ* digestion of electrophoretically separated proteins: a rapid mass spectrometric method for identifying cysteine-containing peptides. *Electrophoresis (Weinheim)*, 17 (1996) 907-917.
- 1904 Nakayama, H., Uchida, K., Shinkai, F., Shinoda, T., Okuyama, T., Seta, K. and Isobe, T.: Capillary column high-performance liquid chromatographic-electrospray ionization triple-stage quadrupole mass spectrometric analysis of proteins separated by two-dimensional polyacrylamide gel electrophoresis. Application to cerebellar protein mapping. *J. Chromatogr. A*, 730 (1996) 279-287.
- 1905 Pedron, J., Maldiney, R., Brault, M. and Miginiac, E.: Monitoring of hapten-protein coupling reactions by capillary zone electrophoresis: improvement of abscisic acid-bovine serum albumin coupling and determination of molar coupling ratios. *J. Chromatogr. A*, 723 (1996) 381-388.
- 1906 Rabilloud, T.: Solubilization of proteins for electrophoretic analyses. *Electrophoresis (Weinheim)*, 17 (1996) 813-829 - a review with 147 refs.
- 1907 Schuhmacher, M., Glocker, M.O., Wunderlin, M. and Przybylski, M.: Direct isolation of proteins from sodium dodecyl sulfate-polyacrylamide gel electrophoresis and analysis by electrospray-ionization mass spectrometry. *Electrophoresis (Weinheim)*, 17 (1996) 848-854.
- 1908 Skelsey, K.R. and Bushey, M.M.: Capillary electrophoretic monitoring of the folding and unfolding of β -lactoglobulin B. *J. Chromatogr. Sci.*, 34 (1996) 85-91.
- 1909 Soulié, S., Moller, J.V., Falson, P. and le Maire, M.: Urea reduces the aggregation of membrane proteins on sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Anal. Biochem.*, 236 (1996) 363-364.
- 1910 Teixeira do Amaral Junior, A., Casali, Vicente Wagner, D., Floriano de Moraes, C. and Sediya, Maria Aparecida, N.: (Adjustment of the methodology of protein electrophoresis to obtain isoenzymic phenotypes of pumpkin). *Bragantia*, 53 (1994) 19-24; *C.A.*, 124 (1996) 139330e.
- 1911 Vogt, C.P., Willi, A., Hess, D. and Hunziker, P.E.: Identification of stress proteins in lysates of human cell lines separated by two-dimensional electrophoresis and electroblotted simultaneously onto two different membranes. *Electrophoresis (Weinheim)*, 17 (1996) 892-898.
- 1912 Wilkins, M.R., Pasquali, C., Appel, R.D., Ou, K., Golaz, O., Sanchez, J.-C., Yan, J.X., Gooley, A.A., Hughes, G., et al.: From proteins to proteomes: large scale protein identification by two-dimensional electrophoresis and amino acid analysis. *BioTechnology*, 14 (1996) 61-65; *C.A.*, 124 (1996) 140226g.
- 1913 Wilkins, M.R., Sanchez, J.-C., Williams, K.L. and Hochstrasser, D.F.: Current challenges and future applications for protein maps and post-translational vector maps in proteome projects. *Electrophoresis (Weinheim)*, 17 (1996) 830-838 - a review with 74 refs.

- 1914 Xu, R.J., Vidal-Madjar, C., Sébille, B. and Diez-Masa, J.C.: Separation of basic proteins by capillary zone electrophoresis with coatings of a copolymer of vinylpyrrolidone and vinylimidazole. *J. Chromatogr. A*, 730 (1996) 289-295.
- See also 1620, 1644, 1649, 1653, 1665, 1668, 1674, 1688, 1711, 1716, 1756, 1758, 1872, 2049, 2073.
- 19b. Proteins of cells, viruses and subcellular particles**
- 1915 Astill, D.St.J., Rychkov, G., Clarke, J.D., Hughes, B.P., Roberts, M.L. and Bretag, A.H.: Characteristics of skeletal muscle chloride channel C1C-1 and point mutant R304E expressed in Sf-9 insect cells. *Biochim. Biophys. Acta*, 1280 (1996) 178-186.
- 1916 Dohi, T., Nakasuji, M., Nakanishi, K., Yasugi, E., Yuyama, Y., Sugiyama, N., Tabuchi, Y., Horiuchi, T. and Oshima, M.: Biochemical bases in differentiation of a mouse cell line GSM06 to gastric surface cells. *Biochim. Biophys. Acta*, 1289 (1996) 71-78.
- 1917 Gu, J. and Wada, Y.: Effect of exogenous decorin on cell morphology and attachment of decorin-deficient fibroblasts. *J. Biochem. (Tokyo)*, 119 (1996) 743-748.
- 1918 Kim, B.Y., Ahn, S.C., Kang, D.O., Ko, H.R., Oh, W.K., Lee, H.S., Mheen, T.L., Rho, H.M. and Ahn, J.S.: Inhibition of PDGF-induced phospholipase C activation by herbimycin A. *Biochim. Biophys. Acta*, 1311 (1996) 33-36.
- 1919 Koshikawa, N., Nakamura, T., Tsuchiya, N., Isaji, M., Yasumitsu, H., Umeda, M. and Miyazaki, K.: Purification and identification of a novel and four known serine proteinase inhibitors secreted by human glioblastoma cells. *J. Biochem. (Tokyo)*, 119 (1996) 334-339.
- 1920 Mattson, J.P. and Keeling, D.J.: [³H]Bafilomycin as a probe for the transmembrane proton channel of the osteoclast vacuolar H⁺-ATPase. *Biochim. Biophys. Acta*, 1280 (1996) 98-106.
- 1921 Morré, D.J., Wilkinson, F.E., Kim, C., Cho, N.M., Lawrence, J., Morré, D.M. and McClure, D.: Antitumor sulfonyleurea-inhibited NADH oxidase of cultured HeLa cells shed into media. *Biochim. Biophys. Acta*, 1280 (1996) 197-206.
- 1922 Ohtsuka, T., Kaziro, Y. and Satoh, T.: Analysis of the T-cell activation signaling pathway mediated by tyrosine kinases, protein kinase C, and Ras protein, which is modulated by intracellular cyclic AMP. *Biochim. Biophys. Acta*, 1310 (1996) 223-232.
- 1923 Pei, Z.-M. and Tang, Z.-C.: (A rapid method for isolation and purification of membrane protein). *Zhiwu Shenglixue Tongxun*, 31 (1995) 210-212; *C.A.*, 124 (1996) 169974k.
- 1924 Pitto, M., Chigorno, V., Renlund, M. and Tettamanti, G.: Impairment of ganglioside metabolism in cultured fibroblasts from Salla patients. *Clin. Chim. Acta*, 247 (1996) 143-157.
- 1925 Portig, I., Pankuweit, S., Lottspeich, F. and Maisch, B.: Identification of stress proteins in endothelial cells. *Electrophoresis (Weinheim)*, 17 (1996) 803-808.
- 1926 Pozzi, D., Lisi, A., Lanzilli, G. and Grimaldi, S.: Role of membrane fluidity in Epstein Barr virus (EBV) infectivity on Akata cell line. *Biochim. Biophys. Acta*, 1280 (1996) 161-168.
- 1927 Tanaka, T., Inoue, M., Sakamoto, J. and Sone, N.: Intra- and inter-complex cross-linking of subunits in the quinol oxidase super-complex from thermophilic *Bacillus* PS3. *J. Biochem. (Tokyo)*, 119 (1996) 482-486.
- 1928 Viollet, B., Lefrancois-Martinez, A.-M., Henrion, A., Kahn, A., Raymondjean, M. and Martinez, A.: Immunochemical characterization and transacting properties of upstream stimulatory factor isoforms. *J. Biol. Chem.*, 271 (1996) 1405-1415.
- 1929 Yang, H., Hamada, K., Terashima, H., Izuta, M., Yamaguchi-Sihata, E., Kondoh, O., Satoh, H., Miyazaki, M., Arisawa, M., Miyamoto, C. and Kitada, K.: A point mutation within each of two ATP-binding motifs inactivates the functions of elongation factor 3. *Biochim. Biophys. Acta*, 1310 (1996) 303-308.
- See also 1932, 1934, 1995, 1998, 2047, 2062, 2074, 2099, 2118.
- 19c. Proteins synthesized by genetic manipulation, monoclonal antibodies**
- 1930 Amari, J.V. and Mazsaroff, I.: Analysis of recombinant human interleukin-11 fusion protein derived from *Escherichia coli* lysate by combined size-exclusion and reversed-phase liquid chromatography. *J. Chromatogr. A*, 729 (1996) 113-124.
- 1931 Greve, K.F., Hughes, D.E., Richberg, P., Kats, M. and Karger, B.L.: Liquid chromatographic and capillary electrophoretic examination of intact and degraded fusion protein CTLA4Ig and kinetics of conformational transition. *J. Chromatogr. A*, 723 (1996) 273-284.
- 1932 Porte, F., de Santa Barbara, P., Phalipou, S., Liautard, J.-P. and Widada, J.S.: Change in the N-terminal domain conformation of annexin I that correlates with liposome aggregation is impaired by Ser-27 to Glu mutation that mimics phosphorylation. *Biochim. Biophys. Acta*, 1293 (1996) 177-184.
- 1933 Slominski, A., Ermak, G., Hwang, J., Mazurkiewicz, J., Corliss, D. and Eastman, A.: The expression of proopiomelanocortin (POMC) and of corticotropin releasing hormone receptor (CRH-R) genes in mouse skin. *Biochim. Biophys. Acta*, 1289 (1996) 247-251.
- 1934 Yoshimoto, M., Ushiyama, Y., Sakai, M., Tamaki, S., Hara, H., Takahashi, K., Sawasaki, Y. and Hanada, K.: Characterization of single chain urokinase-type plasminogen activator with a novel amino-acid substitution in the kringle structure. *Biochim. Biophys. Acta*, 1293 (1996) 83-89.
- 1935 Zhang, S.-M., Qi, Z.-B. and Li, H.-M.: (Purification of HCV recombinant protein C₂₇). *Shengwu Huaxue Zazhi*, 11 (1995) 706-710; *C.A.*, 124 (1996) 111561g.
- See also 1797, 1806, 1967, 1985, 2021, 2046, 2053, 2060, 2062, 2075, 2149, 2219.
- 19d. Microbial and plant proteins**
- 1936 Baldan, B., Navazio, L., Friso, L., Mariani, P. and Meggio, F.: Plant calreticulin is specifically and efficiently phosphorylated by protein kinase CK2. *Biochem. Biophys. Res. Commun.*, 221 (1996) 498-502.
- 1937 Beall, B., Cassidy, P.K. and Sanden, G.N.: Analysis of *Bordetella pertussis* isolates from an epidemic by pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 33 (1995) 3083-3086; *C.A.*, 124 (1996) 137280q.

- 1938 Bolte, G., Osman, A., Mothes, T. and Stern, M.: Peptic-tryptic digests of gliadin: contaminating trypsin but not pepsin interferes with gastrointestinal protein binding characteristics. *Clin. Chim. Acta*, 247 (1996) 59-70.
- 1939 Burg, H.C.J. and van den Berg, B.M.: State of the art of large-scale genetic purity testing of hybrid vegetable seeds using isoelectric focusing at PetoSluis. *Electrophoresis (Weinheim)*, 17 (1996) 502-504.
- 1940 Chow, L.-P., Fukaya, N., Miyatake, N., Horimoto, K., Sugiura, Y., Tabuchi, K., Ueno, Y. and Tsugita, A.: Resolution of *Fusarium sporotrichioides* proteins by two-dimensional polyacrylamide gel electrophoresis and identification by sequence homology comparison in protein data base. *J. Biomed. Sci. (Basel)*, 2 (1995) 343-352; *C.A.*, 124 (1996) 136799d.
- 1941 Dwyer, S.C., Legendre, L., Low, P.S. and Leto, T.L.: Plant and human neutrophil oxidative burst complexes contain immunologically related proteins. *Biochim. Biophys. Acta*, 1289 (1996) 231-237.
- 1942 Feist, H., Geschwend, G., Erler, W. and Schimmel, D.: (Proteins from the outer membrane of *Pasteurella*. Polyacrylamide gel electrophoresis of outer membrane proteins as a possibility for differentiation of *Pasteurella haemolytica* strains). *Berl. Muench. Tierarztl. Wochenschr.*, 108 (1995) 249-252; *C.A.*, 124 (1996) 81283y.
- 1943 Frandsen, E.V.G. and Wade, W.G.: Differentiation of human *Capnocytophaga* species by multilocus enzyme electrophoretic analysis and serotyping of immunoglobulin A1 proteases. *Microbiology (Reading)*, 142 (1996) 441-448; *C.A.*, 124 (1996) 198029m.
- 1944 Funk, C., Adamska, I., Green, B.R., Andersson, B. and Renger, G.: The nuclear-encoded chlorophyll-binding photosystem II-S protein is stable in the absence of pigments. *J. Biol. Chem.*, 270 (1995) 30141-30147.
- 1945 Gibson, J.R., Fitzgerald, C. and Owen, R.J.: Comparison of PFGE, ribotyping and phage-typing in the epidemiological analysis of *Campylobacter jejuni* serotype H52 infections. *Epidemiol. Infect.*, 115 (1995) 215-225; *C.A.*, 124 (1996) 140220a.
- 1946 Haq, A., Remo, N. and Al-Sedairy, S.T.: Fractionation of black seed (*Nigella sativa* Linn) proteins by using Rotofor. *J. Liq. Chromatogr.*, 19 (1996) 593-599.
- 1947 Heyworth, M.V. and Vergara, J.A.: Recognition of acidic proteins of *Giardia muris* by anti-trophozoite antibodies. *Adv. Exp. Med. Biol.*, 371B (1995) 939-943; *C.A.*, 124 (1996) 84140k.
- 1948 Jensen, K., Kesmir, C. and Søndergaard, I.: From image processing to classification: IV. Classification of electrophoretic patterns by neural networks and statistical methods enable quality assessment of wheat varieties for breadmaking. *Electrophoresis (Weinheim)*, 17 (1996) 694-698.
- 1949 Kalinin, A.E., Karamyshev, A.L. and Nesmeyanova, M.A.: (The block of alkaline phosphatase processing due to single amino acid substitutions affects phospholipid composition and turnover in *Escherichia coli* cells secreting mutant proteins). *Biokhimiya (Moscow)*, 61 (1996) 100-109.
- 1950 Kumar, D., Srivastava, B.S., Singh, N.B. and Srivastava, R.: Identification of a 25-kilodalton protein of *Mycobacterium bovis* BCG to distinguish BCG strains from *Mycobacterium tuberculosis*. *J. Clin. Microbiol.*, 34 (1996) 224-226; *C.A.*, 124 (1996) 169809k.
- 1951 Laurière, M., Bouchez, I., Doyen, C. and Eynard, L.: Identification of glycosylated forms of wheat storage proteins using two-dimensional electrophoresis and blotting. *Electrophoresis (Weinheim)*, 17 (1996) 497-501.
- 1952 Lee, J.H. and Schoeffl, F.: GUS activity staining in gels: a powerful tool for studying protein interactions in plants. *Plant Mol. Biol. Rep.*, 13 (1995) 346-354; *C.A.*, 124 (1996) 197339a.
- 1953 Lookhart, G.L. and Bean, S.R.: Improvements in cereal protein separations by capillary electrophoresis: resolution and reproducibility. *Cereal Chem.*, 73 (1996) 81-87; *C.A.*, 124 (1996) 173832e.
- 1954 Morel, M.-H. and Bonicel, J.: Determination of the number of cysteine residues in high molecular weight subunits of wheat glutenin. *Electrophoresis (Weinheim)*, 17 (1996) 493-496.
- 1955 Morita, S., Fukase, M., Hoshino, K., Fukuda, Y., Yamaguchi, M. and Morita, Y.: Partial purification and characterization of a novel soybean protease which is inhibited by Kunitz and Bowman-Birk trypsin inhibitors. *J. Biochem. (Tokyo)*, 119 (1996) 711-718.
- 1956 Musienko, M.M., Abu Massalam, Tagel Din M.: (Molecular marking of *Triticum aestivum* L. genotypes by electrophoretic spectra of storage proteins). *Ukr. Bot. Zh.*, 51 (1994) 21-31; *C.A.*, 124 (1996) 81278a.
- 1957 Nagahama, M., Michiue, K. and Sakurai, J.: Membrane-damaging action of *Clostridium perfringens* alpha-toxin on phospholipid liposomes. *Biochim. Biophys. Acta*, 1280 (1996) 120-126.
- 1958 Nesmeyanova, M.A., Krupyaniko, V.I., Kalinin, A.E. and Kadyrova, L.Yu.: (Isolation and some properties of mutant *Escherichia coli* alkaline phosphatases). *Biokhimiya (Moscow)*, 61 (1996) 89-99.
- 1959 Ozolina, N.V., Kuzevanov, V.Ya., Kotova, L.G. and Salyaev, R.K.: (Electrophoretic studies of proteins in morphogenic and non-morphogenic calluses of spring wheat). *Fiziol. Biokhim. Kul't. Rast.*, 27 (1995) 254-258; *C.A.*, 124 (1996) 170790d.
- 1960 Park, T.Y., Kim, J.K., Suh, K.Y. and Yoon, W.M.: (Cultivar identification and seed purity test by electrophoresis of isoenzymes in radish and Chinese cabbage). *Han'guk Wonye Hakhoechi*, 36 (1995) 805-811; *C.A.*, 124 (1996) 169835r.
- 1961 Pasquali, C., Frutiger, S., Wilkins, M.R., Hughes, R.D., Bairoch, A., Schaller, D., Sanchez, J.-C. and Hochstrasser, D.F.: Two-dimensional gel electrophoresis of *Escherichia coli* homogenates: the *Escherichia coli* SWISS-2DPAGE database. *Electrophoresis (Weinheim)*, 17 (1996) 547-555.
- 1962 Pernelle, J.-J.: (Characterization of bacterial strains by electrophoresis of total proteins. Influence of gel calibration on the taxonomic proximity coefficient). *J. Eur. Hydrol.*, (1995) 305-316; *C.A.*, 124 (1996) 140218f.
- 1963 Sanchez, J.-C., Golaz, O., Frutiger, S., Schaller, D., Appel, R.D., Bairoch, A., Hughes, G.J. and Hochstrasser, D.F.: The yeast SWISS-2DPAGE database. *Electrophoresis (Weinheim)*, 17 (1996) 556-565.
- 1964 Shi, J., Sun, L. and Jing, X.: (Protein analysis of 6-crude drugs and their processed products by PAGE). *Zhongguo Zhongyao Zazhi*, 20 (1995) 533-574; *C.A.*, 124 (1996) 212202w.
- 1965 Sugimoto, M. and Suzuki, Y.: Molecular cloning, sequencing, and expression of a cDNA encoding α -glucosidase from *Mucor javanicus*. *J. Biochem. (Tokyo)*, 119 (1996) 500-505.

- 1966 Sugiyama, H. and Natori, S.: Transmethylation reaction is essential for *Sarcophaga* lectin gene activation. *J. Biochem. (Tokyo)*, 119 (1996) 354-359.
- 1967 Taanman, J.-W., Burton, M.D., Marusich, M.F., Kennaway, N.G. and Capaldi, R.A.: Subunit specific monoclonal antibodies show different steady-state levels of various cytochrome-c oxidase subunits in chronic progressive external ophthalmoplegia. *Biochim. Biophys. Acta*, 1315 (1996) 199-207.
- 1968 Trachuk, L.A., Shemyakina, T.M., Chestukhina, G.G. and Stepanov, V.M.: (Chitinases of *Bacillus cereus*: isolation and properties). *Biokhimiya (Moscow)*, 61 (1996) 357-368.
- 1969 Verissimo, A., Morais, P.V., Diogo, A., Gomes, C. and da Costa, M.S.: Characterization of *Legionella* species by numerical analysis of whole-cell protein electrophoresis. *Int. J. Syst. Bacteriol.*, 46 (1996) 41-49; *C.A.*, 124 (1996) 197843k.
- See also 1757, 1910, 2109.
- 19e. *Proteins of blood, serum and blood cells*
- 1970 Benga, G., Banner, M. ad Wrigglesworth, J.M.: Quantitation of the water channel protein aquaporin (CHIP28) from red blood cell membranes by densitometry of silver stained polyacrylamide gels. *Electrophoresis (Weinheim)*, 17 (1996) 715-719.
- 1971 Bini, L., Magi, B., Marzocchi, B., Cellesi, C., Berti, B., Raggiacchi, R., Rossolini, A. and Pallini, V.: Two-dimensional electrophoretic patterns of acute-phase human serum proteins in the course of bacterial and viral diseases. *Electrophoresis (Weinheim)*, 17 (1996) 612-616.
- 1972 Craig, K.L. and Harley, C.B.: Phosphorylation of human pleckstrin on Ser-113 and Ser-117 by protein kinase C. *Biochem. J.*, 314 (1996) 937-942.
- 1973 Craig, S.J., Ashton, D.S., Beddell, C. and Valko, K.: The effect of operating parameters on the analysis of a human alpha-interferon by capillary zone electrophoresis. *J. Liq. Chromatogr.*, 18 (1995) 3629-3641.
- 1974 Daaka, Y., Friedman, H. and Klein, T.W.: Cannabinoid receptor proteins are increased in Jurkat, human T-cell line after mitogen activation. *J. Pharmacol. Exp. Ther.*, 276 (1996) 776-783.
- 1975 Diederichs, J.E.: Plasma protein adsorption patterns on liposomes: establishment of analytical procedure. *Electrophoresis (Weinheim)*, 17 (1996) 607-611.
- 1976 Ducret, A., Bruun, C.F., Bures, E.J., Marhaug, G., Husby, G. and Aebersold, R.: Characterization of human serum amyloid A protein isoforms separated by two-dimensional electrophoresis by liquid chromatography/electrospray ionization tandem mass spectrometry. *Electrophoresis (Weinheim)*, 17 (1996) 866-876.
- 1977 Duncker, B.P., Gauthier, S.Y. and Davies, P.L.: Evidence for a proprotein intermediate during maturation of Type II antifreeze protein in sea raven, *Hemipteris americanus*. *Biochim. Biophys. Acta*, 1292 (1996) 312-316.
- 1978 Guzman, N.A.: Biomedical applications of on-line preconcentration-capillary electrophoresis using an analyte concentrator: investigation of design options. *J. Liq. Chromatogr.*, 18 (1995) 3751-3768.
- 1979 Kötz, K., Hänslers, M., Sauer, H., Kaltenhäuser, S. and Hantschel, H.: Immunoglobulin G galactosylation deficiency determined by isoelectric focusing and lectin affino blotting in differential diagnosis of rheumatoid arthritis. *Electrophoresis (Weinheim)*, 17 (1996) 533-534.
- 1980 Leibl, H., Tomasits, R., Wolf, H.M., Eibl, M.M. and Mannhalter, J.W.: Method for the isolation of biologically active monomeric immunoglobulin A from a plasma fraction. *J. Chromatogr. B*, 678 (1996) 173-180.
- 1981 Luczak, J.B., Zhang, K., Saxon, A. and Morrison, S.L.: Expression of novel secreted isoforms of human immunoglobulin E proteins. *J. Biol. Chem.*, 271 (1996) 3428-3436.
- 1982 Matsumoto, M., Yamaguchi, T., Terada, S. and Kimoto, E.: Effects of intracellular pH on high pressure-induced hemolysis of anion transport inhibitor-treated erythrocytes. *Biochim. Biophys. Acta*, 1280 (1996) 243-250.
- 1983 Nagamine, T., Takagi, H., Sugimoto, H., Takehara, K., Fukui, T. and Mori, M.: Prevalence of biotin-binding immunoglobulin in patients with allergic and autoimmune diseases. *Clin. Chim. Acta*, 245 (1996) 209-217.
- 1984 Schägger, H., Bentlage, H., Ruitenbeek, W., Pfeiffer, K., Rotter, S., Rother, C., Böttcher-Purkl, A. and Lodemann, E.: Electrophoretic separation of multiprotein complexes from blood platelets and cell lines: technique for the analysis of diseases with defects in oxidative phosphorylation. *Electrophoresis (Weinheim)*, 17 (1996) 709-714.
- 1985 Schuster, V.L., Lu, R., Kanai, N., Bao, Y., Rosenberg, S., Prié, D., Ronco, P. and Jennings, M.L.: Cloning of the rabbit homologue of mouse 'basigin' and rat 'OX-47': kidney cell type-specific expression, and regulation in collecting duct cells. *Biochim. Biophys. Acta*, 1311 (1996) 13-19.
- 1986 Taketa, K., Liu, M. and Taga, H.: Interaction of lectins with their ligand carbohydrate of α -fetoprotein: analysis by mixed-lectin affinity electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 483-488.
- 1987 Trotter, P.J., Ho, S.Y. and Storch, J.: Fatty acid uptake by CaCo-2 human intestinal cells. *J. Lipid Res.*, 37 (1996) 336-346.
- 1988 Van de Locht, L.T.F., Kupers, A.W.H.M., Verbruggen, B.W., Linsen, P.C.M., Novakova, I.R.O. and Mensink, E.J.B.M.: Semi-automated detection of the factor V mutation by allele specific amplification and capillary electrophoresis. *Thromb. Haemostasis*, 74 (1995) 1276-1279; *C.A.*, 124 (1996) 77628x.
- 1989 Watanabe, S. and Sato, T.: Effects of free fatty acids on the binding of bovine and human serum albumin with steroid hormones. *Biochim. Biophys. Acta*, 1289 (1996) 385-396.
- 1990 Zhang, X., Teng, X.-Y. and Yan, Y.-L.: (Searching for distinct proteins in cytoplasmic male sterile line of wheat from two-dimensional gel electrophoresis with personal computer). *Shengwu Huaxue Zazhi*, 11 (1995) 669-672; *C.A.*, 124 (1996) 111560f.
- 1991 Zhao, X., Bo, J., Dai, T. and Xia, Z.: (Isolation and purification of prealbumin from human plasma). *Shanghai Dier Yike Daxue Xuebao*, 15 (1995) 198-202; *C.A.*, 124 (1996) 81295d.
- See also 1795, 1801, 1876, 2403, 2406.

19f. *Structural and muscle proteins*

- 1992 Fukami, K., Sawada, N., Endo, T. and Takenawa, T.: Identification of a phosphatidylinositol 4,5-bisphosphate-binding site in chicken skeletal muscle α -actinin. *J. Biol. Chem.*, 271 (1996) 2646-2650.
- 1993 George, J. and Chandrakasan, G.: Molecular characteristics of dimethylnitrosamine induced fibrotic liver collagen. *Biochim. Biophys. Acta*, 1292 (1996) 215-222.
- 1994 Hofmann, K., Fischer, K., Mueller, E. and Kemper, V.: (Control of effectiveness of heat treatment of canned meats and meat-and-bone meal). *Fleischwirtschaft*, 75 (1995) 1227-1231; *C.A.*, 124 (1996) 85251j.
- 1995 Kurasawa, Y., Hanyu, K., Watanabe, Y. and Numata, O.: F-Actin bundling activity of *Tetrahymena* elongation factor 1 α is regulated by Ca²⁺/calmodulin. *J. Biochem. (Tokyo)*, 119 (1996) 791-798.
- 1996 Morimoto, S. and Ohtsuki, I.: Reduced positive feedback regulation between myosin crossbridge and cardiac troponin C in fast skeletal myofibrils. *J. Biochem. (Tokyo)*, 119 (1996) 737-742.
- 1997 Thiede, B., Otto, A., Zimny-Armdt, U., Müller, E.-C. and Jungblut, P.: Identification of human myocardial proteins separated by two-dimensional electrophoresis with matrix-assisted laser desorption/ionization mass spectrometry. *Electrophoresis (Weinheim)*, 17 (1996) 588-599.
- 1998 Thompson, M.G., Mackie, S.C., Thom, A., Hazlerigg, D.G., Morrison, K.S. and Palmer, R.M.: Cyclic AMP stimulates protein synthesis in L6 myoblasts and its effects are additive to those of insulin, vasopressin and 12-O-tetradecanoylphorbol-13-acetate. Possible involvement of mitogen activated protein kinase. *Biochim. Biophys. Acta*, 1311 (1996) 37-44.
- 1999 Wang, Z., Horiuchi, K.Y. and Chacko, S.: Characterization of the functional domains on the C-terminal region of caldesmon using full-length and mutant caldesmon molecules. *J. Biol. Chem.*, 271 (1996) 2234-2242.
- 2000 Xia, T., Akers, K., Eisen, A.Z. and Seltzer, J.L.: Comparison of cleavage site specificity of gelatinases A and B using collagenous peptides. *Biochim. Biophys. Acta*, 1293 (1996) 259-266.

19g. *Protamines, histones and other nuclear proteins*

- 2001 Berthelsen, J., Vandekerhove, J. and Blasi, F.: Purification and characterization of UEF3, a novel factor involved in the regulation of the urokinase and other AP-1 controlled promoters. *J. Biol. Chem.*, 271 (1996) 3822-3830.
- 2002 Deissler, H., Behn-Krappa, A. and Doerfler, W.: Purification of nuclear proteins from human HeLa cells that bind specifically to the unstable tandem repeat (CGG)_n in the human FMR1 gene. *J. Biol. Chem.*, 271 (1996) 4327-4334.
- 2003 Talasz, H., Helliger, W., Puschendorf, B. and Lindner, H.: *In vivo* phosphorylation of histone H1 variants during the cell cycle. *Biochemistry*, 35 (1996) 1761-1767.

19h. *Chromoproteins and metalloproteins*

- 2004 Cartel, N.J.: Purification of human brain metallothionein by organic and reversed-phase high-performance liquid chromatography under acidic conditions. *J. Chromatogr. B*, 676 (1996) 159-164.
- 2005 Delaroche, O., Bordure, P., Lippert, E. and Sagniez, M.: Perilymph detection by β 2-transferrin immunoblotting assay. Application to the diagnosis of perilymphatic fistulae. *Clin. Chim. Acta*, 245 (1996) 93-104.
- 2006 Matsunaga, T., Iwawaki, Y., Watanabe, K., Yamamoto, I., Kegeyama, T. and Yoshimura, H.: Microsomal aldehyde oxygenase (MALDO): purification and characterization of a cytochrome P450 isozyme responsible for oxidation of 9-anthraldehyde to 9-anthracenecarboxylic acid in monkey liver. *J. Biochem. (Tokyo)*, 119 (1996) 617-625.
- 2007 Okazaki, T., Kadohno, N., Katano, M., Yanagisawa, Y., Nagai, T. and Kanno, T.: (Estimation of the affinity between haptoglobin and hemoglobin by two-dimensional affinity electrophoresis). *Seibutsu Butsuri Kagaku*, 40 (1996) 25-29; *C.A.*, 124 (1996) 197523f.
- 2008 Slutter, C.E., Sanders, D., Wittung, P., Malmström, B.G., Asa, R., Richards, J.H., Gray, H.B. and Fee, J.A.: Water-soluble, recombinant Cu α -domain of the cytochrome b $_3$ subunit II from *Thermophilus thermophilus*. *Biochemistry*, 35 (1996) 3387-3395.
- 2009 Vassilyev, V.B., Kachurin, A.M., Rocco, G.-P., Beltrami, M., Salvato, B. and Gaitskhoki, V.S.: (A spectral study of the active center of ceruloplasmin during its copper depletion/repletion). *Biokhimiya (Moscow)*, 61 (1996) 296-307.

See also 1807, 1869, 2081, 2090.

19i. *Proteins of glands, gland products, various zymogens (incl. milk proteins)*

- 2010 Beeley, J.A., Newman, F., Wilson, P.H.R. and Shimmin, I.C.: Sodium dodecyl sulphate-polyacrylamide gel electrophoresis of human parotid salivary proteins: comparison of dansylation, Coomassie Blue R-250 and silver detection methods. *Electrophoresis (Weinheim)*, 17 (1996) 505-506.
- 2011 Bossi, A., Righetti, P.G., Visco, C., Breme, U., Mauriello, M., Valsasina, B., Orsini, G. and Wensch, E.: Investigation on minor degraded derivatives of the recombinant hirudin variant HM2 *Hirudinaria manillensis* isolated by isoelectric focusing in multi-compartment electrolyzers. *Electrophoresis (Weinheim)*, 17 (1996) 932-937.
- 2012 Brodard, V., Bernard, H., Wal, J.M., David, B. and Peltre, G.: Two-dimensional analysis of cow's milk allergens with the IPG-DALT technique. *Adv. Exp. Med. Biol.*, 371B (1995) 875-878; *C.A.*, 124 (1996) 84139s.
- 2013 Cochet, S., Pesliakas, H., Kroviarski, Y., Cartron, J.P. and Bertrand, O.: Chromatography of bovine lactoserum on the immobilized metallized dye Drimarene Rubine R/K 5BL. Modulation of selectivity by metal exchange and application to purification of milk immunoglobulins. *J. Chromatogr. A*, 725 (1996) 237-247.
- 2014 Hashimoto, C., Ayaki, M., Tanaka, K., Yamamoto, R., Fukuda, H., Funai, H., Wada, A. and Inui, Y.: Distribution and subcellular localization of a growth inhibitory factor in hamster liver and its intracellular partner(s). *Biochim. Biophys. Acta*, 1310 (1996) 309-316.

- 2015 He, C. and Jungst, D.: Electrophoretic analysis of biliary proteins: application of high resolution two-dimensional polyacrylamide gel electrophoresis with immobilized pH gradient in the first dimension. *Electrophoresis (Weinheim)*, 17 (1996) 617-619.
- 2016 Hiratsuka, M., Matsuura, T., Watanabe, E., Sato, M. and Suzuki, Y.: Sex and strain differences in constitutive expression of fatty acid ω -hydroxylase (CYP4A-related proteins) in mice. *J. Biochem. (Tokyo)*, 119 (1996) 340-345.
- 2017 Kaminarides, S.E., Kandarakis, J.G. and Moschopoulou, E.: Detection of bovine milk in ovine Halloumi cheese by electrophoresis of α_{s1} -casein. *Aust. J. Dairy Technol.*, 50 (1995) 58-61; C.A., 124 (1996) 85153d.
- 2018 Kinghorn, N.M., Paterson, G.R. and Otter, D.E.: Quantification of the major bovine whey proteins using capillary zone electrophoresis. *J. Chromatogr. A*, 723 (1996) 371-379.
- 2019 Kubo, T., Sasaki, M., Nakamura, J., Sasagawa, H., Ohashi, K., Takeuchi, H. and Natori, S.: Change in the expression of hypopharyngeal-gland proteins of the worker honeybees (*Apis mellifera* L.) with age and/or role. *J. Biochem. (Tokyo)*, 119 (1996) 291-295.
- 2020 Sautin, Yu.Yu., Povorotny, V.Yu. and Mikosha, A.S.: (Prolactin-induced phosphorylation of proteins in adrenal cortex by protein kinase C). *Biochim. Biophys. Acta*, 61 (1996) 34-41.
- 2021 Street, M.D., Donovan, G.R. and Baldo, B.A.: Molecular cloning and characterization of the major allergen Myr p II from the venom of the jumper ant *Myrmecia pilosula*: Myr p I and Myr p II share a common protein leader sequence. *Biochim. Biophys. Acta*, 1305 (1996) 87-97.
- See also 1811, 1901, 2409, 2410.
- 19j. *Proteins of brain, cerebrospinal fluid and eye*
- 2022 Gracz, L.M. and Parsons, S.M.: Purification of active synaptic vesicles from the electric organ of *Torpedo californica* and comparison to reserve vesicles. *Biochim. Biophys. Acta*, 1292 (1996) 293-302.
- 2023 Lichtchenko, K., Nguyen, T. and Südhof, T.C.: Structures, alternative splicing, and neurexin binding of multiple neuroligins. *J. Biol. Chem.*, 271 (1996) 2676-2682.
- 2024 Nye, H.E., Hope, B.T., Kelz, M.B., Iadarola, M. and Nestler, E.J.: Pharmacological studies of the regulation of chronic FOS-related antigen induction by cocaine in the striatum and nucleus accumbens. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1671-1680.
- 2025 Pant, H.C. and Veeranna: Neurofilament phosphorylation. *Biochem. Cell Biol.*, 73 (1995) 575-592.
- 2026 Pedrotti, B., Ulloa, L., Avila, J. and Islam, K.: Characterization of microtubule-associated protein MAP1B: phosphorylation state, light chains, and binding to microtubules. *Biochemistry*, 35 (1996) 3016-3023.
- 2027 Sanchez-Blazquez, P., Garcia-Espana, A. and Garzon, J.: *In vivo* injection of antisense oligodeoxynucleotides to $G\alpha$ subunits and supraspinal analgesia evoked by Mu and Delta opioid agonists. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1590-1596.
- 2028 Sax, C.M., Ilagan, J.G. and Haynes, J.L., II.: Lens-preferred activity of the -1809/+46 mouse α A-crystallin promoter in stably integrated chromatin. *Biochim. Biophys. Acta*, 1305 (1996) 49-53.
- 2029 Yoshimura, Y., Nomura, T. and Yamauchi, T.: Purification and characterization of active fragment of Ca^{2+} /calmodulin-dependent protein kinase II from the post-synaptic density in the rat forebrain. *J. Biochem. (Tokyo)*, 119 (1996) 268-273.
- See also 1904, 1933.
- 19k. *Proteins of neoplastic tissue and transformed cells*
- 2030 Frederick, M.J., Rodriguez, L.V., Johnston, D.A. and Grimm, E.A.: Tumor target binding induces phosphorylation of two Mr 65,000 lymphokine-activated killer proteins. *Cancer Res.*, 56 (1996) 127-137.
- 2031 Frederick, M.J., Rodriguez, L.V., Johnston, D.A., Darnay, B.G. and Grimm, E.A.: Characterization of the Mr 65,000 lymphokine-activated killer proteins phosphorylated after tumor target binding: evidence that pp65a and pp65b are phosphorylated forms of L-plastin. *Cancer Res.*, 56 (1996) 138-144.
- 2032 Itoh, K., Hayashi, A., Nakao, M., Hoshino, T., Seki, N. and Shichijo, S.: Human tumor rejection antigens MAGE. *J. Biochem. (Tokyo)*, 119 (1996) 385-390.
- 2033 Voloudakis-Baltatzis, I.E., Malliopoulos, V., Gaitanarou, H., Kapoula, S., Tselentis, V. and Agnantis, N.J.: Histological, ultrastructural and two-dimensional gel electrophoresis studies of human breast cancer. *Cancer Mol. Biol.*, 2 (1995) 597-611; C.A., 124 (1996) 111547g.
- 2034 Wu, T., Ikezono, T., Angus, C.W. and Shelhamer, J.H.: Tumor necrosis factor- α induces the 85-kDa cytosolic phospholipase A₂ gene expression in human bronchial epithelial cells. *Biochim. Biophys. Acta*, 1310 (1996) 175-184.
- 2035 Yanagisawa, J., Ando, J., Nakayama, J., Kohwi, Y. and Kohwi-Shigematsu, T.: A matrix attachment region (MAR)-binding activity due to a p114 kilodalton protein is found only in human breast carcinomas and not in normal and benign breast disease tissues. *Cancer Res.*, 56 (1996) 457-462.
- 19l. *Specific binding and receptor proteins*
- 2036 Baltensperger, K., Karoor, V., Paul, H., Ruoho, A., Czech, M.P. and Malbon, C.C.: The β -adrenergic receptor is a substrate for the insulin receptor tyrosine kinase. *J. Biol. Chem.*, 271 (1996) 1061-1064.
- 2037 Bisaccia, F., Zara, V., Capobianco, L., Iacobazzi, V., Mazzeo, M. and Palmieri, F.: The formation of a disulfide cross-link between the two subunits demonstrates the dimeric structure of the mitochondrial oxoglutarate carrier. *Biochim. Biophys. Acta*, 1292 (1996) 281-288.
- 2038 Carpenter, L., Poole, R.C. and Halestrap, A.P.: Cloning and sequencing of the monocarboxylate transporter from mouse Ehrlich Lettré tumour cell confirms its identity as MCT1 and demonstrates that glycosylation is not required for MCT1 function. *Biochim. Biophys. Acta*, 1279 (1996) 157-163.
- 2039 Castagna, L., Zarzur, J., Filipetti, M. and Landa, C.: Isolation and partial characterization of N-acetyl-D-galactosamine-binding lectins from *Epiphragmophora trenquellionis* snail. *J. Biochem. (Tokyo)*, 119 (1996) 372-377.
- 2040 Chapin, S.J., Enrich, C., Aroeti, B., Havel, R.J. and Mostov, K.E.: Calmodulin binds to the basolateral targeting signal of the polymeric immunoglobulin receptor. *J. Biol. Chem.*, 271 (1996) 1336-1342.

- 2041 Dahms, N.M., Seetharam, B. and Wick, D.A.: Expression of insulin-like growth factor (IGF)-I receptors, IGF-II/cation-independent mannose 6-phosphate receptors (CI-MPRs), and cation-dependent MPRs in polarized human intestinal Caco-2 cells. *Biochim. Biophys. Acta*, 1279 (1996) 84-92.
- 2042 Fishburn, C.S., Elazar, Z. and Fuchs, S.: Differential glycosylation and intracellular trafficking for the long and short isoforms of the D₂ dopamine receptor. *J. Biol. Chem.*, 270 (1995) 29819-29824.
- 2043 Fukunaga, B.N. and Hankinson, O.: Identification of a novel domain in the aryl hydrocarbon receptor required for DNA binding. *J. Biol. Chem.*, 271 (1996) 3743-3749.
- 2044 Garcia-Higuera, I., Thomas, T.C., Yu, F. and Neer, E.J.: Inter-subunit surfaces in G protein $\alpha\beta\gamma$ heterotrimers. Analysis by cross-linking and mutagenesis of $\beta\gamma$. *J. Biol. Chem.*, 271 (1996) 528-535.
- 2045 Gurdal, H., Tilakaratne, N., Brown, R.D., Fonseca, M., Friedman, E. and Johnson, M.D.: The expression of alpha1 adrenoreceptor subtypes changes with age in the rat aorta. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1656-1662.
- 2046 Hashino, K., Uemori, Y., Kimizuka, F., Kato, I. and Titani, K.: A 31-kDa recombinant fibronectin cell-binding domain fragment: its binding to receptor, cell adhesive activity, and fusion proteins. *J. Biochem. (Tokyo)*, 119 (1996) 604-609.
- 2047 Iwata, E., Nakanishi, T., Ogawa, N., Ohyama, K., Murakami, T. and Takigawa, M.: Neurotrophin-3 increases the DNA-binding activities of several transcription factors in a mouse osteoblastic cell line. *Biochim. Biophys. Acta*, 1311 (1996) 85-92.
- 2048 Kikuchi, Y., Sogawa, K., Watanabe, N., Kobayashi, A. and Fujii-Kuriyama, Y.: Purification and characterization of the DNA-binding domain of BTEB, a GC box-binding transcription factor, expressed in *Escherichia coli*. *J. Biochem. (Tokyo)*, 119 (1996) 309-313.
- 2049 Krajewski, S., Zapata, J.M. and Reed, J.C.: Detection of multiple antigens of western blots. *Anal. Biochem.*, 236 (1996) 221-228.
- 2050 LeMotte, P.K., Keidel, S. and Apfel, C.M.: Characterization of synthetic retinoids with selectivity for retinoic acid or retinoid X nuclear receptors. *Biochim. Biophys. Acta*, 1289 (1996) 298-304.
- 2051 Lutomski, D., Caron, M., Cornillot, J.-D., Bourin, P., Dupuy, C., Pontet, M., Bladier, D. and Joubert-Caron, R.: Identification of different galectins by immunoblotting after two-dimensional polyacrylamide gel electrophoresis with immobilized pH gradients. *Electrophoresis (Weinheim)*, 17 (1996) 600-606.
- 2052 Mariller, C., Allain, F., Kouach, M. and Spik, G.: Evidence that human milk isolated cyclophilin B corresponds to a truncated form. *Biochim. Biophys. Acta*, 1293 (1996) 31-38.
- 2053 Mireskandari, A., Reid, R.L., Kashanchi, F., Dittmer, J., Li, W.-B. and Brady, J.N.: Isolation of a cDNA clone, TRX, encoding a human T-cell lymphotropic virus type-I Tax binding protein. *Biochim. Biophys. Acta*, 1306 (1996) 9-13.
- 2054 Murayama, T. and Ogawa, Y.: Properties of Ryr3 ryanodine receptor isoform in mammalian brain. *J. Biol. Chem.*, 271 (1996) 5079-5084.
- 2055 Ozelebi, F., Rao, R.V., Holicky, E., Madden, B.J., McCormick, D.J. and Miller, L.J.: Phosphorylation of cholecystokinin receptors expressed on chinese hamster ovary cells. Similarities and differences relative to native pancreatic acinar cell receptors. *J. Biol. Chem.*, 271 (1996) 3750-3755.
- 2056 Raffaniello, R.D., Lin, J., Wang, F. and Raufman, J.-P.: Expression of Rab3D in dispersed chief cells from guinea pig stomach. *Biochim. Biophys. Acta*, 1311 (1996) 111-116.
- 2057 Ray, R., Swamy, N., MacDonald, P.N., Ray, S., Haussler, M.R. and Holick, M.F.: Affinity labeling of the $1\alpha,25$ -dihydroxyvitamin D₃ receptor. *J. Biol. Chem.*, 271 (1996) 2012-2017.
- 2058 Rokaw, M.D., Benos, D.J., Palevsky, P.M., Cunningham, S.A., West, M.E. and Johnson, J.P.: Regulation of a sodium channel-associated G-protein by aldosterone. *J. Biol. Chem.*, 271 (1996) 4491-4496.
- 2059 Roth, U. and Schroeder, B.G.: Gel electrophoresis for preparative separation of biologicals such as nicotinic acetylcholine receptors (nAChRs). *Ger. Offen. DE 4,431,309 (Cl. B01D57/02)*, 7 Mar. 1996, Appl. 4,431,309, 2 Sep. 1994; 17 p.; C.A., 124 (1996) 236253q.
- 2060 Schaap, F.G., Specht, B., van der Vusse, G.J., Borchers, T. and Glatz, J.F.C.: One-step purification of rat heart-type fatty acid-binding protein expressed in *Escherichia coli*. *J. Chromatogr. B*, 679 (1996) 61-67.
- 2061 Singh, J.K., Yan, Q., Dawson, G. and Banerjee, P.: Cell-specific regulation of the stably expressed serotonin 5-HT_{1A} receptor and altered ganglioside synthesis. *Biochim. Biophys. Acta*, 1310 (1996) 201-211.
- 2062 Taylor, M.R., Peterson, J.A., Ceriani, R.L. and Couto, J.R.: Cloning and sequence analysis of human butyrophilin reveals a potential receptor function. *Biochim. Biophys. Acta*, 1306 (1996) 1-4.
- 2063 Yaojun, G., Xinhui, L. and Lili, H.: Subtyping of group-specific component and protease inhibitor by rapid isoelectric focusing on PhastSystem. *Electrophoresis (Weinheim)*, 17 (1996) 537-539.
- 2064 Yokote, K., Mori, S., Siegbah, A., Rönstrand, L., Wernstedt, C., Heldin, C.-H. and Claesson-Welsh, L.: Structural determinants in the platelet-derived growth factor α -receptor implicated in modulation of chemotaxis. *J. Biol. Chem.*, 271 (1996) 5101-5111.
- 2065 Zhang, L.X. and Liang, H.G.: Detection of Ca²⁺-binding proteins of photosystem 2 particles by electrophoretic migration combined with terbium fluorescence. *Photosynthetica*, 31 (1995) 621-624; C.A., 124 (1996) 812247q.

See also 1878, 1929, 1932.

19m. Urinary proteins

- 2066 Hobbs, G.A. and Levinson, S.S.: Hemoglobin interference with urinary Bence Jones protein analysis on electrophoresis. *Ann. Clin. Lab. Sci.*, 26 (1996) 71-75; C.A., 124 (1996) 199816c.
- 2067 Lindqvist, A. and Åkerström, B.: Bovine α -microglobulin/bikunin. Isolation and characterization of liver cDNA and urinary α 1-microglobulin. *Biochim. Biophys. Acta*, 1306 (1996) 98-106.
- 2068 Servais, M.D., Ritzerfeld, M., Stefanidis, I., Melzer, H. and Mann, H.: (Evaluation and documentation of electrophoretic urinary protein analysis using laser densitometry). *Nieren-Hochdruckkrankh.*, 24 (1995) 566-569; C.A., 124 (1996) 140192t.

- 19n. *Other proteins (incl. proteinous inhibitors of enzymic activity)*
- 2069 Forest, E., Pétillot, Y., Gagnon, J., Cotton, J. and Vita, C.: Determination of the C-terminal form of an anemone toxin using capillary electrophoresis and mass spectrometry. *Electrophoresis (Weinheim)*, 17 (1996) 962-964.
- 2070 Frokiaer, H., Mortensen, K., Sorensen, H. and Sorensen, S.: Characterization of proteintype proteinase inhibitors by high performance capillary electrophoresis. *J. Liq. Chromatogr.*, 19 (1996) 57-67.
- 2071 Ishimura, R., Noda, K., Hattori, N., Shiota, K. and Ogawa, T.: Analysis of rat placental plasma membrane proteins by two-dimensional gel electrophoresis. *Mol. Cell. Endocrinol.*, 115 (1995) 149-159; *C.A.*, 124 (1996) 81262r.
- 2072 Miskin, R. and Abramovitz, R.: One-phase reverse zymography after denaturing gel electrophoresis: high sensitivity detection of activity of plasminogen activator inhibitor 2 and other protease inhibitors. *Fibrinolysis*, 9 (1995) 331-342; *C.A.*, 124 (1996) 110511d.
- 2073 Nakamura, A. and Goto, S.: Analysis of protein carbonyls with 2,4-dinitrophenyl hydrazine and its antibodies by immunoblot in two-dimensional gel electrophoresis. *J. Biochem. (Tokyo)*, 119 (1996) 768-774.
- 2074 Schaeffer, H.: Native electrophoresis for isolation of mitochondrial oxidative phosphorylation protein complexes. *Methods Enzymol.*, 260(Mitochondrial Biogenesis and Genetics, Pt. A) (1995) 190-202; *C.A.*, 124 (1996) 81008n.
- 2075 Schnizer, R., van Heeke, G., Amatore, D. and Schuster, S.M.: Histidine-49 is necessary for the pH-dependent transition between active and inactive states of the bovine F₁-ATPase inhibitor protein. *Biochim. Biophys. Acta*, 1292 (1996) 241-248.
- See also 1870, 1909, 1911, 1919, 1955, 1977.
20. ENZYMES AND ENZYME ACTIVITY ESTIMATION
- 2076 Ballve, R.M.L., Medina Filho, H.P., Bordigno, R. and Lima, M.M.A.: Methodology for starch gel electrophoresis and protocols for isoenzymes of 32 plant genera. *Braz. J. Genet.*, 18 (1995) 491-502; *C.A.*, 124 (1996) 81250k.
- 2077 Salerno, C. and Crifo, C.: Application of capillary zone electrophoresis for rapid screening of enzymic defects in purine metabolism. *Adv. Exp. Med. Biol.*, 370(Purine and Pyrimidine Metabolism in Man VIII) (1994) 811-814; *C.A.*, 124 (1996) 48914g.
- 20a. *Oxidoreductases*
- 2078 Bulitta, C., Ganea, C., Fahimi, H.D. and Völkl, A.: Cytoplasmic and peroxisomal catalases of the guinea pig liver: evidence for two distinct proteins. *Biochim. Biophys. Acta*, 1293 (1996) 55-62.
- 2079 Fernandes, A.I. and Gregoriadis, G.: Synthesis, characterization and properties of sialylated catalase. *Biochim. Biophys. Acta*, 1293 (1996) 90-96.
- 2080 Heinova, D., Blahovec, J. and Rosival, I.: Lactate dehydrogenase isozyme patterns in bird, carp and mammalian sera. *Eur. J. Clin. Chem. Clin. Biochem.*, 34 (1996) 91-95.
- 2081 Homma-Takeda, S., Shinyashiki, M., Nakai, I., Tohyama, C., Kumagai, Y. and Shimajo, N.: Direct detection of mercury-bound metalloproteins (metallothionein and Cu, Zn-superoxide dismutase) using a combination of gel electrophoresis and one dimensional synchrotron radiation X-ray fluorescence analysis. *Anal. Lett.*, 29 (1996) 601-611.
- 2082 Imamura, Y., Ryu, A., Koga, T., Higuchi, T., Otagiri, M., Nozawa, M. and Akita, H.: Purification and catalytic properties of a novel acetohexamide-reducing enzyme from rabbit heart. *J. Biochem. (Tokyo)*, 119 (1996) 648-652.
- 2083 Iwakura, M. and Honda, S.: Stability and reversibility of thermal denaturation are greatly improved by limiting terminal flexibility of *Escherichia coli* dihydrofolate reductase. *J. Biochem. (Tokyo)*, 119 (1996) 414-420.
- 2084 Kobayashi, A., Jiang, L.L. and Hashimoto, T.: Two mitochondrial 3-hydroxyacyl-CoA dehydrogenases in bovine liver. *J. Biochem. (Tokyo)*, 119 (1996) 775-782.
- 2085 Liu, J., Liu, Y., Parkinson, A. and Klaassen, C.D.: Effect of oleanolic acid on hepatic toxicant-activating and detoxifying systems in mice. *J. Pharmacol. Exp. Ther.*, 275 (1995) 768-774.
- 2086 Loseva, O.I., Osipov, V.V., Vanyakin, E.N. and Gavriushkin, A.V.: (Isolation of superoxide dismutase with the use of a method of free flow electrophoresis). *Biotechnologiya*, (1995) 25-29; *C.A.*, 124 (1996) 139378b.
- 2087 Morgunov, I.G., Sharyshev, A.A., Peskova, E.B. and Finogenova, T.V.: (Mitochondrial aldehyde dehydrogenase from the ethanol-assimilating yeast *Candida maltosa*). *Biokhimiya (Moscow)*, 61 (1996) 131-141.
- 2088 Nakayama, N., Takemae, A. and Shoun, H.: Cytochrome P450foxy, a catalytically self-sufficient fatty acid hydroxylase of the fungus *Fusarium oxysporum*. *J. Biochem. (Tokyo)*, 119 (1996) 435-440.
- 2089 Obinger, C., Pfeiffer, S., Hofstetter, W., Wutka, R. and Ebermann, R.: Fast and sensitive staining technique for glucose oxidase in polyacrylamide gel. *J. Chromatogr. A*, 731 (1996) 293-298.
- 2090 Ohgiya, N., Yokota, H., Takahashi, M., Komoro, S. and Yuasa, A.: Purification and properties of a new β -naphthoflavone inducible cytochrome P-450, aryl hydrocarbon hydroxylase from rat kidney. *Biochim. Biophys. Acta*, 1289 (1996) 122-130.
- 2091 Rawat, U.B. and Rao, M.B.: Purification, kinetic characterization and involvement of tryptophan residue at the NADPH binding site of xylose reductase from *Neurospora crassa*. *Biochim. Biophys. Acta*, 1293 (1996) 222-230.
- 2092 Xue, Q. and Yeung, E.S.: Determination of lactate dehydrogenase isoenzymes in single lymphocytes from normal and leukemia cell lines. *J. Chromatogr. B*, 677 (1996) 233-240.
- See also 1967, 2006, 2128.
- 20b. *Transferases (excl. E.C. 2.7.-.-)*
- 2093 Bohets, H.H., Nouwen, E.J., de Broe, M.E. and Dierickx, P.J.: The cytosolic glutathione S-transferase isoenzymes in the dog kidney cortex as compared with the corresponding MDCK renal cell line. *Biochim. Biophys. Acta*, 1311 (1996) 93-101.

- 2094 Chang, C.C.Y., Chen, J., Thomas, M.A., Cheng, D., del Priore, V.A., Newton, R.S., Pape, M.E. and Chang, T.: Regulation and immunolocalization of acyl-coenzyme A:cholesterol acyltransferase in mammalian cells as studied with specific antibodies. *J. Biol. Chem.*, 270 (1995) 29532-29540.
- 2095 Dean, R.A., Zhang, J., Brzezinski, M.R. and Bosron, W.F.: Tissue distribution of cocaine methyl esterase and ethyl transferase activities: correlation with carboxylesterase protein. *J. Pharmacol. Exp. Ther.*, 275 (1995) 965-971.
- 2096 Honke, K., Yamane, M., Ishii, A., Kobayashi, T. and Makita, A.: Purification and characterization of 3'-phosphoadenosine-5'-phosphosulfate:GalCer sulfotransferase from human renal cancer cells. *J. Biochem. (Tokyo)*, 119 (1996) 421-427.
- 2097 Kura, T., Takahashi, Y., Takayama, T., Ban, N., Saito, T., Kuga, T. and Niitsu, Y.: Glutathione S-transferase- π is secreted as a monomer into human plasma by platelets and tumor cells. *Biochim. Biophys. Acta*, 1292 (1996) 317-323.
- 2098 Lavallée, B., Provost, P. and Bélanger, A.: Formation of pregnenolone- and dehydroepiandrosterone-fatty acid esters by lecithin-cholesterol acyltransferase in human plasma high density lipoproteins. *Biochim. Biophys. Acta*, 1299 (1996) 306-312.
- 2099 MacNulty, E.E. and Ryder, N.S.: Characterization of prenyl protein transferase enzymes in a human keratinocyte cell line. *Biochim. Biophys. Acta*, 1289 (1996) 41-50.
- 2100 Radomska, A., Drake, R.R., Zhu, X., Veronese, M.E., Little, J.M., Nowell, S., McManus, M.E., Lester, R. and Falany, C.N.: Photoaffinity labeling of human recombinant sulfotransferases with 2-azidoadenosine 3',5'-[5'-³²P]bisphosphate. *J. Biol. Chem.*, 271 (1996) 3195-3199.
- 2101 Tsai, S.-P., Wang, L.-Y., Yeh, H.-I. and Tam, M.F.: The 30 kDa protein co-purified with chick liver glutathione S-transferases is a carbonyl reductase. *Biochim. Biophys. Acta*, 1292 (1996) 289-292.
- 210c. *Transferases transferring phosphorus containing groups (E.C. 2.7.-.-)*
- 2102 Bachelot, C., Rameh, L., Parsons, T. and Cantley, L.C.: Association of phosphatidylinositol 3-kinase, via the SH2 domains of p85, with focal adhesion kinase in polyoma middle t-transformed fibroblasts. *Biochim. Biophys. Acta*, 1311 (1996) 45-52.
- 2103 Bensen, E.S., Umphress, J.L., Traugh, J.A., Pinna, L.A. and Tuazon, P.T.: A membrane-bound protein kinase from rabbit reticulocytes is an active form of multipotential S6 kinase. *Biochim. Biophys. Acta*, 1292 (1996) 249-258.
- 2104 Croquet, F., Bréhier, A., Gil, S., Davy, J. and Féger, J.: Five isoenzymes of protein kinase C are expressed in normal and STZ-diabetic rat hepatocytes: effect of phorbol 12-myristate 13-acetate. *Biochim. Biophys. Acta*, 1315 (1996) 163-168.
- 2105 Fukui, Y. and Morita, F.: Two phosphorylations specific to the tail region of the 204-kDa heavy chain isoform of porcine aorta smooth muscle myosin. *J. Biochem. (Tokyo)*, 119 (1996) 783-790.
- 2106 Gallagher, P.J., Garcia, J.G.N. and Herring, B.P.: Expression of a novel myosin light chain kinase in embryonic tissues and cultured cells. *J. Biol. Chem.*, 270 (1995) 29090-29095.
- 2107 Gehrman, T., Vereb, G., Schmidt, M., Klix, D., Meyer, H.E., Varsányi, M. and Heilmeyer, L.M.G., Jr.: Identification of a 200 kDa polypeptide as type 3 phosphatidylinositol 4-kinase from bovine brain by partial protein and cDNA sequencing. *Biochim. Biophys. Acta*, 1311 (1996) 53-63.
- 2108 He, Q., Skog, S., Wu, C., Johansson, A. and Tribukait, B.: Existence of phosphorylated and dephosphorylated forms of cytosolic thymidine kinase (TK1). *Biochim. Biophys. Acta*, 1289 (1996) 25-30.
- 2109 Klier, H., Magdolen, V., Schrickler, R., Strobel, G., Lottspeich, F. and Bandlow, W.: Cytoplasmic and mitochondrial forms of yeast adenylate kinase 2 are N-acetylated. *Biochim. Biophys. Acta*, 1280 (1996) 251-256.
- 2110 Meyer, R., Höfelein, C., Lüthy, J. and Candrian, U.: Polymerase chain reaction-restriction fragment length polymorphism analysis: a simple method for species identification in food. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1542-1551.
- 2111 San Agustín, J.T. and Witman, G.B.: Detection of flagellar protein kinases on polyvinylidene difluoride membranes following sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Methods Cell Biol.*, 47(Cilia and Flagella) (1995) 135-140; C.A., 124 (1996) 49947g.
- 2112 Skalter, R., Bergstein, M. and Livneh, Z.: β^* , A UV-inducible shorter form of the β subunit of DNA polymerase II of *Escherichia coli*. II. Overproduction, purification, and activity as a polymerase processivity clamp. *J. Biol. Chem.*, 271 (1996) 2491-2496.
- 2113 Szyszka, R., Bou, G. and Ballesta, J.P.G.: RAP kinase, a new enzyme phosphorylating the acidic P proteins from *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta*, 1293 (1996) 213-221.
- 2114 Tobin, A.B., Keys, B. and Nahorski, S.R.: Identification of a novel receptor kinase that phosphorylates a phospholipase C-linked muscarinic receptor. *J. Biol. Chem.*, 271 (1996) 39-7-3916.
- 2115 Walter, J., Schnölzer, M., Pyerin, W., Kinzel, V. and Kübler, D.: Induced release of cell surface protein kinase yields CK1- and CK2-like enzymes in tandem. *J. Biol. Chem.*, 271 (1996) 111-119.
- 2116 Williams, S.J., Schwer, C., Krishnarao, A.S.M., Heid, C., Karger, B.L. and Williams, P.M.: Quantitative competitive polymerase chain reaction: analysis of amplified products of the HIV-1 gag gene by capillary electrophoresis with laser-induced fluorescence detection. *Anal. Biochem.*, 236 (1996) 146-152.
- 2117 Yi, X.B., Sweitzer, N.M. and de S. Otero, A.: Neutralizing antibodies to nucleoside diphosphate kinase inhibit the enzyme *in vitro* and *in vivo*: evidence for two distinct mechanisms of activation of atrial currents by ATP₂S. *Biochim. Biophys. Acta*, 1310 (1996) 334-342.
- 2118 Zimmerman, U.-J., Wang, M., Nelson, J.B., Ekwunife, F.S. and Liu, L.: Secretagogue-induced proteolysis of cAMP-dependent protein kinase in intact rat alveolar epithelial type II cells. *Biochim. Biophys. Acta*, 1311 (1996) 117-123.

See also 2029.

20d. *Hydrolases, acting on ester bonds (E.C. 3.1.-.-)*

- 2119 Atkins Latham, K., Rajendran, S., Carmical, J.R., Lee, J.C. and Lloyd, R.S.: T4 endonuclease V exists in solution as a monomer and binds to target sites as a monomer. *Biochim. Biophys. Acta*, 1292 (1996) 324-334.
- 2120 Bernal, C. and Pidgeon, C.: Affinity purification of phospholipase A₂ on immobilized artificial membranes containing and lacking the glycerol backbone. *J. Chromatogr. A*, 731 (1996) 139-151.
- 2121 Boschetti, N., Brodbeck, U., Jensen, S.P., Koch, C. and Norgaard-Pedersen, B.: Monoclonal antibodies against a C-terminal peptide of human brain acetylcholinesterase distinguishing between erythrocyte and brain acetylcholinesterases. *Clin. Chem. (Washington)*, 42 (1996) 19-23.
- 2122 Bucht, G. and Hjalmarsson, K.: Residues in *Torpedo californica* acetylcholinesterase necessary for processing to a glycosyl phosphatidylinositol-anchored form. *Biochim. Biophys. Acta*, 1292 (1996) 223-232.
- 2123 Chetoui, H., Delhalle, E., Osterrieth, P. and Rousseaux, D.: Combination of biotyping and electrophoretic patterns of esterases for differentiation of nosocomial *Serratia marcescens* strains. *Res. Microbiol.*, 146 (1995) 579-586; *C.A.*, 124 (1996) 81285a.
- 2124 Donnelly, J.G., Ooi, D.S., Burns, B.F. and Goel, R.: Chronic increased serum lipase without evidence of pancreatitis: tumor-derived lipase? *Clin. Chem. (Washington)*, 42 (1996) 462-464.
- 2125 Hata, K., Tokuhira, H. and Miura, M.: New identification of bone specific alkaline phosphatase in rat serum by an electrophoretic methods. *Rinsho Kagaku (Nippon Rinsho Kagakkai)*, 24 (1995) 219-222; *C.A.*, 124 (1996) 196838a.
- 2126 Hattori, M., Adachi, H., Aoki, J., Tsujimoto, M., Arai, H. and Inoue, K.: Cloning and expression of a cDNA encoding the β -subunit (30-kDa subunit) of bovine brain platelet-activating factor acetylhydrolase. *J. Biol. Chem.*, 270 (1995) 31345-31352.
- 2127 Huijbregts, R.P.H., de Kroon, A.I.P. and de Kruijff, B.: Rapid transmembrane movement of C₆-NBD-labeled phospholipids across the inner membrane of *Escherichia coli*. *Biochim. Biophys. Acta*, 1280 (1996) 41-50.
- 2128 Jing, Y.: (Gel electrophoresis of rhizobial esterase, superoxide dismutase and glucose-6-phosphate dehydrogenase). *Huanan Shifan Daxue Xuebao, Ziran Kexueban*, (1995) 36-40; *C.A.*, 124 (1996) 168908e.
- 2129 Nakshatri, H. and Currie, R.A.: Differential whole-cell extract preparation and electrophoretic mobility shift assay to evaluate the effect of tyrosine phosphatases on DNA binding activity of transcription factors. *Anal. Biochem.*, 236 (1996) 178-181.
- 2130 Reitz, M., Köllner, D., Preiss, U. and Gutjahr, P.: Reduced neutral DNase activity in the urine of children after cytostatic therapy. *Arzneim.-Forsch.*, 46 (1996) 438-441.
- 2131 Varier, A., Dadlani, M. and Sharma, S.P.: Phenol test and electrophoresis of seed esterases for testing genetic purity in seed lots of pearl millet (*Pennisetum glaucum*) inbreds. *Indian J. Agric. Sci.*, 65 (1995) 789-792; *C.A.*, 124 (1996) 111546f.
- 2132 Yaojun, G., Lili, H., Xuegang, Y. and Zhongyi, Y.: Phenotyping of esterase D and acid phosphatase by rapid isoelectric focusing on PhastSystem. *Electrophoresis (Weinheim)*, 17 (1996) 535-536.

See also 1918, 2095.

20e. *Hydrolases, acting on glycosyl compounds (E.C. 3.2.-.-)*

- 2133 Cueva, R., Muñoz, M.D., Andaluz, E., Basco, R.D. and Larriba, G.: Preferential transfer of truncated oligosaccharides to the first sequon of yeast exoglucanase in *Saccharomyces cerevisiae* alg3 cells. *Biochim. Biophys. Acta*, 1289 (1996) 336-342.
- 2134 Guadalupi, R., Bernard, M., Orlicchio, A., Foglietti, M.J. and Emiliani, C.: Purification and properties of human urinary β -D-mannosidase. *Biochim. Biophys. Acta*, 1293 (1996) 9-16.
- 2135 Qi, S. and Su, T.: (Gel electrophoresis active dyeing method for determination of α -galactosidase). *Weishengwuxue Tongbao*, 22 (1995) 253-254; *C.A.*, 124 (1996) 168898b.
- 2136 Srisomsap, C., Svasti, J., Surarit, R., Champattanachai, V., Sawangareetrakul, P., Boonpuan, K., Subhasitanont, P. and Chokchaichamnankit, D.: Isolation and characterization of an enzyme with β -glucosidase and β -fucosidase activities from *Dalbergia cochinchinensis* Pierre. *J. Biochem. (Tokyo)*, 119 (1996) 585-590.

See also 1784, 1968.

20f. *Other hydrolases*

- 2137 Arthur, J.S.C. and Crawford, C.: Investigation of the interaction of m-calpain with phospholipids: calpain-phospholipid interactions. *Biochim. Biophys. Acta*, 1293 (1996) 201-206.
- 2138 Azarkan, M., Wintjens, R.T., Smolders, N., Nijs, M. and Looze, Y.: S-Pegylthiopapain, a versatile intermediate for the preparation of the fully active form of the cysteine proteinase archetype. *J. Chromatogr. A*, 724 (1996) 185-192.
- 2139 Bandell, M., Hall, M.J., Wang, G., Seto-Young, D. and Perlin, D.S.: Probing the cytoplasmic LOOP1 domain of the yeast plasma membrane H⁺-ATPase by targeted factor Xa proteolysis. *Biochim. Biophys. Acta*, 1280 (1996) 81-90.
- 2140 Fabunmi, R.P., Baker, A.H., Murray, E.J., Booth, R.F.G. and Newby, A.C.: Divergent regulation of growth factors and cytokines of 95 kDa and 72 kDa gelatinase s and tissue inhibitors of metalloproteinases 1, -2 and -3 in rabbit aortic smooth muscle cells. *Biochem. J.*, 315 (1996) 335-342.
- 2141 Ferrington, D.A., Reijneveld, J.C., Bär, P.R. and Bigelow, D.J.: Activation of the sarcoplasmic reticulum Ca²⁺-ATPase induced by exercise. *Biochim. Biophys. Acta*, 1279 (1996) 203-213.
- 2142 Fukusen, N. and Aoki, Y.: Purification and characterization of novel trypsin-like serine proteases from mouse spleen. *J. Biochem. (Tokyo)*, 119 (1996) 633-638.
- 2143 Goodwin, S.B., Schneider, R.E. and Fry, W.E.: Use of cellulose-acetate electrophoresis for rapid identification of allozyme genotypes of *Phytophthora infestans*. *Plant Dis.*, 79 (1995) 1181-1185; *C.A.*, 124 (1996) 81248r.
- 2144 Hansell, E.J., Frisch, S.M., Trenble, P., Murnane, J.P. and Werb, Z.: Simian virus 40 transformation alters the actin cytoskeleton, expression of matrix metalloproteinases and inhibitors of metalloproteinases, and invasive behavior of normal and ataxia-telangiectasia human skin fibroblasts. *Biochem. Cell Biol.*, 73 (1995) 373-389.

- 2145 Hoijer, M.A., Melief, M.-J., Keck, W. and Hazenberg, M.P.: Purification and characterization of N-acetylmuramyl-L-alanine amidase from human plasma using monoclonal antibodies. *Biochim. Biophys. Acta*, 1289 (1996) 57-64.
- 2146 Houde, M., Tremblay, P., Masure, S., Opendakker, G., Oth, D. and Mandeville, R.: Synergistic and selective stimulation of gelatinase B production in macrophages by lipopolysaccharide, *trans*-retinoic acid and CGP 41251, a protein kinase C regulator. *Biochim. Biophys. Acta*, 1310 (1996) 193-200.
- 2147 Itoh, M., Masuda, K., Ito, Y., Akizawa, T., Yoshioka, M., Imai, K., Okada, Y., Sato, H. and Seiki, M.: Purification and refolding of recombinant human proMMP-7 (pro-matrilysin) expressed in *Escherichia coli* and its characterization. *J. Biochem. (Tokyo)*, 119 (1996) 667-673.
- 2148 Kitazono, A., Kitano, A., Kabashima, T., Ito, K. and Yoshimoto, T.: Prolyl aminopeptidase is also present in *Enterobacteriaceae*: cloning and sequencing of the *Hafnia alvei* enzyme-gene and characterization of the expressed enzyme. *J. Biochem. (Tokyo)*, 119 (1996) 468-474.
- 2149 Maeda, Y., Ueda, H., Kazami, J., Kawano, G., Suzuki, E. and Nagamune, T.: Truncation of Vargula luciferase still results in retention of luminescence. *J. Biochem. (Tokyo)*, 119 (1996) 601-603.
- 2150 Makowski, G.S. and Ramsby, M.L.: Calibrating gelatin zymograms with human gelatinase standards. *Anal. Biochem.*, 236 (1996) 353-356.
- 2151 Matsuki, H., Fujimoto, N., Iwata, K., Knäuper, V., Okada, Y. and Hayakawa, T.: A one-step sandwich enzyme immunoassay for human matrix metalloproteinase 8 (neutrophil collagenase) using monoclonal antibodies. *Clin. Chim. Acta*, 244 (1996) 129-143.
- 2152 Motley, A.M., Tabak, H.F., Smeitink, J.A.M., Poss-The, B.T., Barth, P.G. and Wanders, R.J.A.: Non-rhizomelic and rhizomelic chondrodysplasia punctata within a single complementation group. *Biochim. Biophys. Acta*, 1315 (1996) 153-158.
- 2153 Ohuchi, E., Azumano, I., Yoshida, S., Iwata, K. and Okada, Y.: A one-step sandwich immunoassay for human matrix metalloproteinase 7 (matrilysin) using monoclonal antibodies. *Clin. Chim. Acta*, 244 (1996) 181-198.
- 2154 Sättler, A., Kanka, S., Maurer, K.-H. and Riesner, D.: Thermostable variants of subtilisin selected by temperature-gradient gel electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 784-792.
- 2155 Tyulkova, N.A. and Sandalova, T.P.: (The comparative study of the temperature effect on the bacterial luciferase). *Biokhimiya (Moscow)*, 61 (1996) 275-287.
- 2156 Ueshima, S., Okada, K. and Matsuo, O.: Stabilization of plasmin by lysine derivatives. *Clin. Chim. Acta*, 245 (1996) 7-18.
- 2158 Kakei, M. and Nakahara, H.: Aspects of carbonic anhydrase and carbonate content during mineralization of the rat enamel. *Biochim. Biophys. Acta*, 1289 (1996) 226-230.
- 2159 Kromer, S., Gardeström, P. and Samuelsson, G.: Regulation of the supply of cytosolic oxaloacetate for mitochondrial metabolism via phosphoenolpyruvate carboxylase in barley leaf protoplasts. I. The effect of covalent modification on PEPC activity, pH response, and kinetic properties. *Biochim. Biophys. Acta*, 1289 (1996) 343-350.
- 2160 Santha, R., Rao, N.A. and Vaidyanathan, C.S.: Identification of the active-site peptide of 2,3-dihydroxybenzoic acid decarboxylase from *Aspergillus oryzae*. *Biochim. Biophys. Acta*, 1293 (1996) 191-200.

20h. Isomerases

- 2161 Giacomelli, S., Leone, M.-G., Grima, J., Silvestrini, B. and Cheng, C.Y.: Astrocytes synthesize and secrete prostaglandin D synthetase *in vitro*. *Biochim. Biophys. Acta*, 1310 (1996) 269-276.

20i. Ligases

- 2162 Gonen, H., Stancovski, I., Shkedy, D., Hadari, T., Bercovich, B., Bengal, E., Mesilati, S., Abu-Hatoum, O., Schwartz, A.L. and Ciechanover, A.: Isolation, characterization, and partial purification of a novel ubiquitin-protein ligase, E3. Targeting of protein substrates via multiple and distinct recognition signals and conjugating enzymes. *J. Biol. Chem.*, 271 (1996) 302-310.

20j. Complex mixtures and incompletely identified enzymes

- 2163 Andrade, M.D. and Skalka, A.M.: Multimerization determinants reside in both the catalytic core and C terminus of avian sarcoma virus intergrase. *J. Biol. Chem.*, 270 (1995) 29299-29306.
- 2164 Flint, S.H., Hartley, N.J., Avery, S.M. and Hudson, J.A.: A comparison between starch and polyacrylamide gels for the analysis of *Listeria monocytogenes* using multilocus enzyme electrophoresis. *Lett. Appl. Microbiol.*, 22 (1996) 16-17; *C.A.*, 124 (1996) 197383k.
- 2165 Uchida, Y., Shimmi, O., Sudoh, M., Arisawa, M. and Yamada-Okabe, H.: Characterization of chitin synthase 2 of *Saccharomyces cerevisiae* II: both full size and processed enzymes are active for chitin synthesis. *J. Biochem. (Tokyo)*, 119 (1996) 659-666.
- 2166 Young, K., Shi, G., Chang, Y., Chang, B., Chang, L., Lai, M., Chuang, W. and Wu, H.: Interaction of streptokinase and plasminogen. Studied with truncated streptokinase peptides. *J. Biol. Chem.*, 270 (1995) 29601-29606.

See also 1622, 1628, 1920, 2166.

See also 1662.

20g. Lyases

- 2157 Kaihara, R., Matsushashi, S., Kusakabe, T., Kondo, T., Iwanaga, A. and Hori, K.: Monoclonal anti-human aldolase C antibodies that react to the isozyme group-specific sequences and generally conserved sequences of human aldolase C. *J. Biochem. (Tokyo)*, 119 (1996) 281-290.

21. PURINES, PYRIMIDINES, NUCLEIC ACIDS AND THEIR CONSTITUENTS
- 21a. *Purines, pyrimidines, nucleosides, nucleotides*
- 2167 Barry, J.P., Muth, J., Law, S.-J., Karger, B.L. and Vouros, P.: Analysis of modified oligonucleotides by capillary electrophoresis in a polyvinylpyrrolidone matrix coupled with electrospray mass spectrometry. *J. Chromatogr. A*, 732 (1996) 159-166.
- 2168 Bory, C., Chantin, C. and Bouliou, R.: Comparison of capillary electrophoretic and liquid chromatographic determination of hypoxanthine and xanthine for the diagnosis of xanthinuria. *J. Chromatogr. A*, 730 (1996) 329-331.
- 2169 Chen, T.-Z., Lin, S.-B., Wu, J.-C., Choo, K.-B. and Au, L.-C.: A method for screening antisense oligodeoxyribonucleotides effective for mRNA translation-arrest. *J. Biochem. (Tokyo)*, 119 (1996) 252-255.
- 2170 Cibelli, G., Jüngling, S., Schoch, S., Gerdes, H.-H. and Thiel, G.: Identification of a functional cAMP response element in the secretogranin II gene. *Eur. J. Biochem.*, 236 (1996) 171-179.
- 2171 Evangelista, R.A., Liu, M.-S., Rampal, S. and Chen, F.-T.A.: Characterization of fluorescent nucleoside triphosphates by capillary electrophoresis with laser-induced fluorescence detection: active of alkaline phosphatase and DNA polymerase. *Anal. Biochem.*, 235 (1996) 89-97.
- 2172 Grune, T. and Perrett, D.: Rapid simultaneous measurement of nucleotides, nucleosides and bases in tissues by capillary electrophoresis. *Adv. Exp. Med. Biol.*, 370 (Purine and Pyrimidine Metabolism in Man VIII) (1994) 805-810; *C.A.*, 124 (1996) 49912s.
- 2173 Leeds, J.M., Graham, M.J., Truong, L.A. and Cummins, L.L.: Quantitation of phosphorothioate oligonucleotides in human plasma. *Anal. Biochem.*, 235 (1996) 36-43.
- 2174 Rabel, S.R., Riley, C.M., Trueworthy, R. and Stobaugh, J.F.: Aspects involving selectivity manipulation for the separation of 6-mercaptapurine metabolites by capillary electrophoresis. *J. Capillary Electrophor.*, 2 (1995) 57-61; *C.A.*, 124 (1996) 156203h.
- 2175 Saneyoshi, M., Kohsaka-Ichikawa, M., Yahata, A., Kimura, S., Izuta, S. and Yamaguchi, T.: Synthetic nucleosides and nucleotides. XXXV. Synthesis and biological evaluations of 5-fluoropyrimidine nucleosides and nucleotides of 3-deoxy- β -D-ribofuranose and related compounds. *Chem. Pharm. Bull.*, 43 (1995) 2005-2009.
- 2176 Schwartz, H.E. and Ulfelder, K.J.: Analysis of bases, nucleosides, and (oligo) nucleotides by capillary electrophoresis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 227-264; *C.A.*, 124 (1996) 197271x - a review with 83 refs.
- 2177 Sevcik, J., Adam, T. and Mazacova, H.: A fast and simple screening method for detection of 2,8-dihydroxyadenine urolithiasis by capillary zone electrophoresis. *Clin. Chim. Acta*, 245 (1996) 85-92.
- 2178 Shida, T., Iwasaki, H., Shinagawa, H. and Kyogoku, Y.: Characterization and comparison of synthetic immobile and mobile Holliday junctions. *J. Biochem. (Tokyo)*, 119 (1996) 653-658.
- 2179 Siren, H., Jumppanen, J.H., Peltonen, K. and Riekkola, M.-L.: Identification of nucleic acids and oligonucleotides by capillary zone electrophoresis with the four marker technique. *J. Liq. Chromatogr.*, 18 (1995) 3577-3589.
- 2180 Wang, C.-C., McCann, W.P. and Beale, S.C.: Measurement of adenosine by capillary zone electrophoresis with on-column isotachophoretic preconcentration. *J. Chromatogr. B*, 676 (1996) 19-28.
- 2181 Zhao, T., Liu, Q. and Cheng, J.: (Separation of determination of pyrimidine bases by capillary zone electrophoresis). *Fenxi Huaxue*, 24 (1996) 180-183; *C.A.*, 124 (1996) 169837t.
- See also 2240.
- 21b. *Nucleic acids, RNA*
- 2182 Aizawa, A., Kazahari, K., Yoneyama, T. and Ono, M.: Significance of the chromatin structure in expression of the rat prolactin gene. *J. Biochem. (Tokyo)*, 119 (1996) 296-301.
- 2183 Asiedu, D.K., Froyland, L., Vaagenes, H., Lie, O., Demoz, A. and Berge, R.K.: Long-term effect of tetradecylthioacetic acid: a study on plasma lipid profile and fatty acid composition and oxidation in different rat organs. *Biochim. Biophys. Acta*, 1300 (1996) 86-96.
- 2184 Bachvarov, D.R., Saint-Jacques, E., Larrivée, J.F., Levesque, L., Rioux, F., Drapeau, G. and Marceau, F.: Cloning and pharmacological characterization of the rabbit bradykinin B2 receptor. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1623-1630.
- 2185 Bencherif, M., Fowler, K., Lukas, R.J. and Lippello, P.M.: Mechanisms of up-regulation of neural nicotinic acetylcholine receptors in clonal cell lines and primary cultures of fetal rat brain. *J. Pharmacol. Exp. Ther.*, 275 (1995) 987-994.
- 2186 Brown, V.D., Wang, Z.-F., Williams, A.S. and Marzluff, W.F.: Structure of a cluster of mouse histone genes. *Biochim. Biophys. Acta*, 1306 (1996) 17-22.
- 2187 Cheung, H.S., Sallis, J.D. and Struve, J.A.: Specific inhibition of basic calcium phosphate and calcium pyrophosphate crystal-induction of metalloproteinase synthesis by phosphocitrate. *Biochim. Biophys. Acta*, 1315 (1996) 105-111.
- 2188 Costeas, P.A., Tonelli, L.A. and Chinsky, J.M.: Molecular cloning of the murine branched chain α -ketoacid dehydrogenase E2 subunit: presence of 3' B1 repeat elements. *Biochim. Biophys. Acta*, 1305 (1996) 25-28.
- 2189 Covic, L. and Lew, R.R.: *Arabidopsis thaliana* cDNA isolated by functional complementation shows homology to serine/threonine protein kinases. *Biochim. Biophys. Acta*, 1305 (1996) 125-129.
- 2190 Dabis-Cox, M.I., Fletcher, T.L., Turner, J.N., Szarowski, D. and Shain, W.: Three-day exposure to low-dose ethanol alters guanine nucleotide binding protein expression in the developing rat hippocampus. *J. Pharmacol. Exp. Ther.*, 276 (1996) 758-764.
- 2191 Ferris, M.J., Muyzer, G. and Ward, D.M.: Denaturing gradient gel electrophoresis profiles of 16S rRNA-defined populations inhibiting a hot spring microbial mat community. *Appl. Environ. Microbiol.*, 62 (1996) 340-346; *C.A.*, 124 (1996) 195085r.
- 2192 Fraji, B.M. and Gonzales, R.A.: A third human tissue transglutaminase homologue as a result of alternative gene transcripts. *Biochim. Biophys. Acta*, 1306 (1996) 63-74.
- 2193 Gehris, A.L., Brandli, D.W., Lewis, S.D. and Bennett, V.D.: The exon encoding the fibronectin type III-9 repeat is constitutively included in the mRNA from chick limb mesenchyme and cartilage. *Biochim. Biophys. Acta*, 1311 (1996) 5-12.

- 2194 Han, Y.-C. and Pruetz, S.B.: Mechanism of ethanol-induced suppression of a primary antibody response in a mouse model for binge drinking. *J. Pharmacol. Exp. Ther.*, 275 (1995) 950-957.
- 2195 Hatch, C.L. and Bonner, W.M.: An upstream region of the H2AZ gene promoter modulates promoter activity in different cell types. *Biochim. Biophys. Acta*, 1305 (1996) 59-62.
- 2196 Hattori, Y., Yamasaki, M. and Itoh, N.: The rat FGF-5 mRNA variant generated by alternative splicing encodes a novel truncated form of FGF-5. *Biochim. Biophys. Acta*, 1306 (1996) 31-33.
- 2197 Honda, S.-i., Harada, N. and Takagi, Y.: The alternative exons 1 of the mouse aromatase cytochrome P-450 gene. *Biochim. Biophys. Acta*, 1305 (1996) 145-150.
- 2198 Hotta, K., Kuwajima, M., Ono, A., Nakajima, H., Shingu, R., Miyagawa, J.-i., Namba, M., Hanafusa, T., Noguchi, T., Kono, N. and Matsuzawa, Y.: Disordered expression of hepatic glycolytic and gluconeogenic enzymes in Otsuka Long-Evans Tokushima fatty rats with spontaneous long-term hyperglycemia. *Biochim. Biophys. Acta*, 1289 (1996) 145-149.
- 2199 Hotta, K., Kuwajima, M., Ono, A., Uenaka, R., Nakajima, H., Miyagawa, J.-i., Namba, M., Hanafusa, T., Horiuchi, M., Nikaido, H., Hayakawa, J.-i., Kono, N., Saheki, T. and Matsuzawa, Y.: Altered expression of carnitine palmitoyltransferase II in liver, muscle, and heart of mouse strain with juvenile visceral steatosis. *Biochim. Biophys. Acta*, 1289 (1996) 131-135.
- 2200 Kalachnyuk, L.G.: (Individual isoacceptor tyrosine tRNAs from bovine liver. Isolation and characterization). *Biopolim. Kletka*, 10 (1994) 88-91; C.A., 124 (1996) 50154c.
- 2201 Ketchner, S.L. and Malkin, R.: Nucleotide sequence of the PetM gene encoding a 4 kDa subunit of the cytochrome *b₅f* complex from *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, 1273 (1996) 195-197.
- 2202 Komatsu, N., Hiraoka, Y., Shiozawa, M., Ogawa, M. and Aiso, S.: Cloning and expression of *Xenopus laevis* xSox12 cDNA. *Biochim. Biophys. Acta*, 1305 (1996) 117-119.
- 2203 Komine, Y., Kitabatake, M. and Inokuchi, H.: 10Sa RNA is associated with 70S ribosome particles in *Escherichia coli*. *J. Biochem. (Tokyo)*, 119 (1996) 463-467.
- 2204 Kuzumaki, T., Matsuda, A., Ito, K. and Ishikawa, K.: Cell adhesion to substratum and activation of tyrosine kinases are essentially required for G1/S phase transition in BALB/c 3T3 fibroblasts. *Biochim. Biophys. Acta*, 1310 (1996) 185-192.
- 2205 Langerak, A.W., van der Linden-van Beurden, C.A.J. and Versnel, M.A.: Regulation of differential expression of platelet-derived growth factor α - and β -receptor mRNA in normal and malignant human mesothelial cell lines. *Biochim. Biophys. Acta*, 1305 (1996) 63-70.
- 2206 Mahnke-Zizelman, D.K., Eddy, R., Shows, T.B. and Sabina, R.L.: Characterization of the human AMPD3 gene reveals that 5' exon usage is subject to transcriptional control by three tandem promoters and alternative splicing. *Biochim. Biophys. Acta*, 1306 (1996) 75-92.
- 2207 Mann, E.A., Jump, M.L. and Giannella, R.A.: Cell line-specific transcriptional activation of the promoter of the human guanylyl cyclase C/heat-stable enterotoxin receptor gene. *Biochim. Biophys. Acta*, 1305 (1996) 7-10.
- 2208 Matsuno, F., Chowdhury, S., Gotoh, T., Iwase, K., Matsuzaki, H., Takatsuki, K., Mori, M. and Takiguchi, M.: Induction of the C/EBP β gene by dexamethasone and glucagon in primary-cultured rat hepatocytes. *J. Biochem. (Tokyo)*, 119 (1996) 524-532.
- 2209 Mineura, K., Watanabe, K., Yanagisawa, T. and Kowada, M.: Quantification of O⁶-methylguanine-DNA methyltransferase mRNA in human brain tumors. *Biochim. Biophys. Acta*, 1289 (1996) 105-109.
- 2210 Miyamoto, K.-i., Shiraga, T., Morita, K., Yamamoto, H., Haga, H., Taketani, Y., Tamai, I., Sai, Y., Tsuji, A. and Takeda, E.: Sequence, tissue distribution and developmental changes in rat intestinal oligopeptide transporter. *Biochim. Biophys. Acta*, 1305 (1996) 34-38.
- 2211 Nielsen, M.S., Petersen, C.M., Gliemann, J. and Madsen, P.: Cloning and sequencing of a human cDNA encoding a putative transcription factor containing a bromodomain. *Biochim. Biophys. Acta*, 1306 (1996) 14-16.
- 2212 Noti, J.D. and Reinemann, B.C.: Simplified probe preparation facilitates S1 nuclease analysis. *BioTechniques*, 20 (1996) 174-178; C.A., 124 (1996) 166531c.
- 2213 Ogurusu, T. and Shingai, R.: Cloning of a putative γ -aminobutyric acid (GABA) receptor subunit p3 cDNA. *Biochim. Biophys. Acta*, 1305 (1996) 15-18.
- 2214 Park, S.-K., Lee, H.-S. and Lee, S.-T.: Characterization of the human full-length PTK7 cDNA encoding a receptor protein tyrosine kinase-like molecule closely related to chick KLG. *J. Biochem. (Tokyo)*, 119 (1996) 235-239.
- 2215 Plath, A., Peters, F. and Einspanier, R.: Detection and quantitation of specific mRNAs by ribonuclease protection assay using denaturing horizontal polyacrylamide gel electrophoresis: a radioactive and nonradioactive approach. *Electrophoresis (Weinheim)*, 17 (1996) 471-472.
- 2216 Roglic, A., Prossnitz, E.R., Cavanagh, S.L., Pan, Z., Zou, A. and Ye, R.D.: cDNA cloning of a novel G protein-coupled receptor with a large extracellular loop structure. *Biochim. Biophys. Acta*, 1305 (1996) 39-43.
- 2217 Saito, H., Okuda, M., Terada, T., Sasaki, S. and Inui, K.-I.: Cloning and characterization of a rat H⁺/peptide cotransporter mediating absorption of β -lactam antibiotics in the intestine and kidney. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1631-1637.
- 2218 Saito, H., Terada, T., Okuda, M., Sasaki, S. and Inui, K.-I.: Molecular cloning and tissue distribution of rat peptide transporter PEPT2. *Biochim. Biophys. Acta*, 1280 (1996) 173-177.
- 2219 Satoh, A., Takayama, E., Kojima, K., Ogawa, H., Yamori, T., Sato, S., Kawaguchi, T., Tsuroo, T., Katsura, Y., Kina, T. and Matsumoto, I.: Expression of carbohydrate-binding protein p33/41 in human tumor cell lines. *J. Biochem. (Tokyo)*, 119 (1996) 346-353.
- 2220 Schuetz, E.G., Furuya, K.N. and Schuetz, J.D.: Interindividual variation in expression of P-glycoprotein in normal human liver and secondary hepatic neoplasms. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1011-1018.
- 2221 Scofield, M.A., Liu, F., Abel, P.W. and Jeffries, W.B.: Quantification of steady state expression of mRNA for alpha-1 adrenergic receptor subtypes using reverse transcription and a competitive polymerase chain reaction. *J. Pharmacol. Exp. Ther.*, 275 (1995) 1035-1042.

- 2222 Smetanina, N.S., Leonova, J.Y., Levy, N. and Huisman, T.H.J.: The $\alpha\beta$ - and $\alpha2\alpha1$ -globin mRNA ratios in different forms of α -thalassaemia. *Biochim. Biophys. Acta*, 1315 (1996) 188-192.
- 2223 Tamarappoo, B.K., McDonald, K.K. and Kilberg, M.S.: Expressed human hippocampal ASCT1 amino acid transporter exhibits a pH-dependent change in substrate specificity. *Biochim. Biophys. Acta*, 1279 (1996) 131-136.
- 2224 Tao, J. and Frankel, A.D.: Arginine-binding RNAs resembling TAR identified by *in vitro* selection. *Biochemistry*, 35 (1996) 2229-2238.
- 2225 Tetsu, O., Kanno, R., Isono, K., Taniguchi, M. and Kanno, M.: Cloning and characterization of two transcripts generated from the mel-13 gene positioned adjacent to the mammalian Polycomb group-related gene mel-18. *Biochim. Biophys. Acta*, 1305 (1996) 109-112.
- 2226 Webb, J.C., Sun, X.-M., McCarthy, S.N., Neuwirth, C., Thompson, G.R., Knight, B.L. and Soutar, A.K.: Characterization of mutations in the low density lipoprotein (LDL) - receptor gene in patients with homozygous familial hypercholesterolemia and frequency of these mutations in FH patients in the United Kingdom. *J. Lipid Res.*, 37 (1996) 368-381; *C.A.*, 124 (1996) 368-381.
- 2227 Yang, W., Dyck, J.R.B. and Fliegel, L.: Regulation of NHE1 expression in L6 muscle cells. *Biochim. Biophys. Acta*, 1306 (1996) 107-113.
- 2228 Yang, Y., Hoefflich, A., Butenandt, O. and Kiess, W.: Opposite regulation of IGF-I and IGF-I receptor mRNA and concomitant changes of GH receptor and IGF-II/M6P receptor mRNA in human IM-9 lymphoblasts. *Biochim. Biophys. Acta*, 1310 (1996) 317-324.
- 2229 Yuan, H.T., Bingle, C.D. and Kelly, F.J.: Differential patterns of antioxidant enzyme mRNA expression in guinea pig lung and liver during development. *Biochim. Biophys. Acta*, 1305 (1996) 163-171.
- 2230 Zou, Z., Nomura, M., Takihara, Y., Yasunaga, T. and Shimada, K.: Isolation and characterization of retinoic acid-inducible cDNA clones in F9 cells: a novel cDNA family encodes cell surface proteins sharing partial homology with MHC class I molecules. *J. Biochem. (Tokyo)*, 119 (1996) 319-328.
- See also 1617, 1890, 1974, 1985, 2034, 2045, 2053, 2056, 2061, 2078, 2146, 2243, 2262, 2283, 2284, 2302.
- 21c. *Nucleic acids, DNA*
- 2231 Asakawa, J.: (High resolutional two-dimension electrophoresis of DNA). *Tanpakushitsu Kakusan Koso*, 41 (1996) 170-177; *C.A.*, 124 (1996) 107722t - a review with 8 refs.
- 2232 Baba, Y.: (Human genome analysis and gene diagnosis by capillary electrophoresis). *Bunseki Kagaku*, 44 (1995) 883-893; *C.A.*, 124 (1996) 46628t - a review with 31 refs.
- 2233 Baez-Camargo, M., Lopez-Canovas, L., Riveron, A.M., Chavez, P. and Orozco, E.: Fast procedure to distinguish circular and linear DNA molecules in pulsed field gel electrophoresis. *Anal. Lett.*, 29 (1996) 745-753.
- 2234 Barbieri, R., Duro, G., Morgante, A. and Izzo, V.: Rapid DNA elution procedure from agarose gels. *Anal. Biochem.*, 235 (1996) 106-107.
- 2235 Barron, A.E., Sunada, W.M. and Blanch, H.W.: The effects of polymer properties on DNA separations by capillary electrophoresis in uncross-linked polymer solutions. *Electrophoresis (Weinheim)*, 17 (1996) 744-757.
- 2236 Barry, J.P., Norwood, C. and Vouros, P.: Detection and identification of benzo[a]pyrene diol epoxide adducts to DNA utilizing capillary electrophoresis - electrospray mass spectrometry. *Anal. Chem.*, 68 (1996) 1432-1438.
- 2237 Bendich, A.J.: Structural analysis of mitochondrial DNA molecules from fungi and plants using moving pictures and pulsed-field gel electrophoresis. *J. Mol. Biol.*, 255 (1996) 564-588; *C.A.*, 124 (1996) 168525c.
- 2238 Beyazova, M.L., Brodsky, B.C., Shearer, M.C. and Horan, A.C.: Preparation of actinomycete DNA for pulsed-field gel electrophoresis. *Int. J. Syst. Bacteriol.*, 45 (1995) 852-854; *C.A.*, 124 (1996) 81284z.
- 2239 Boerglum, A.D., Mullaart, E., Kvistgaard, A.B., Uitterlinden, A.G., Vijg, J. and Kruse, T.A.: Two-dimensional DNA typing as a genetic marker system in humans. *Cytogenet. Cell Genet.*, 71 (1995) 260-265; *C.A.*, 124 (1996) 107882v.
- 2240 Burkhalter, P.W., Müller, C., Lüthy, J. and Candrian, U.: Detection of Salmonella spp. in eggs: DNA analyses, culture techniques, and serology. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1531-1537.
- 2241 Cantor, C.R., Ito, T. and Smith, C.L.: DNA purification by triplex-affinity capture and affinity capture electrophoresis. *U.S. Pat.* 5,482,836 (Cl. 435-6; C12Q1/68), 9 Jan. 1996, Appl. 4,552, 14 Jan. 1993; 20 pp.; *C.A.*, 124 (1996) 137812q.
- 2242 Carlsson, C., Larsson, A. and Jonsson, M.: Influence of optical probing with YOYO on the electrophoretic behavior of the DNA molecule. *Electrophoresis (Weinheim)*, 17 (1996) 642-651.
- 2243 Chiari, M., Campoleoni, A., Conti, P., Felli, C., Patrosso, M.C. and Brogren, C.H.: Electrophoretic separation of biopolymers in a matrix of polyacrylamide covalently linked to agarose. *Electrophoresis (Weinheim)*, 17 (1996) 473-478.
- 2244 Coque, T.M. and Murray, B.E.: Identification of *Enterococcus faecalis* strains by DNA hybridization and pulsed-field gel electrophoresis. *Comments. J. Clin. Microbiol.*, 33 (1995) 3368-3369; *C.A.*, 124 (1996) 136875a.
- 2245 Darini, A.L. da Costa, Zucchi, O.L. Domingues, Carbonell, G.V., Furlanetto, A. and Levy, C.E.: (Plasmid size determination by three methods). *Rev. Latinoam. Microbiol.*, 37 (1995) 217-225; *C.A.*, 124 (1996) 195112x.
- 2246 Destro, M.T., Leitao, M.F.F. and Farber, J.M.: Use of molecular typing methods to trace the dissemination of *Listeria monocytogenes* in a shrimp processing plant. *Appl. Environ. Microbiol.*, 62 (1996) 705-711; *C.A.*, 124 (1996) 173966b.
- 2247 Dey, R., Naik, S. and Patel, M.S.: Tissue-specific expression of the human pyruvate dehydrogenase α (Pdha-1)/chloramphenicol acetyltransferase fusion gene in transgenic mice. *Biochim. Biophys. Acta*, 1305 (1996) 189-195.
- 2248 Friedman, K.L. and Brewer, B.: Analysis of replication intermediates by two-dimensional agarose gel electrophoresis. *Methods Enzymol.*, 262(DNA Replication) (1995) 613-627; *C.A.*, 124 (1996) 77510c - a review with 15 refs.
- 2249 Fueki, T., Sugiura, S. and Yamaguchi, K.: Open strands adjacent to iterons promote the binding of the replication initiator protein (Rep) of pSC101 to the unit sequence of the iterons *in vitro*. *Biochim. Biophys. Acta*, 1305 (1996) 181-188.

- 2250 Fujiwaka, H.: Gel electrophoresis and reagents therefor. *Jpn Kokai Tokkyo Koho JP 07,270,378 [95,270,378]*, (Cl. G01N27/447), 20 Oct. 1995, Appl. 94/85,670, 30 Mar. 1994; 6 p.; C.A., 124 (1996) 75075r.
- 2251 Gelfi, C., Simó-Alfonso, E., Sebastiano, R., Citterio, A. and Righetti, P.G.: Novel acrylamido monomers with higher hydrophilicity and improved hydrolytic stability: III. DNA separations by capillary electrophoresis in poly(N-acryloylaminopropanol). *Electrophoresis (Weinheim)*, 17 (1996) 738-743.
- 2252 Green, M., Wald, E.R., Dashesky, B., Barbadora, K. and Wadowsky, R.M.: Field inversion gel electrophoretic analysis of *Legionella pneumophila* strains associated with nosocomial legionellosis in children. *J. Clin. Microbiol.*, 34 (1996) 175-176; C.A., 124 (1996) 169808j.
- 2253 Hamano, K., Takasawa, T., Kurazono, T., Okuyama, Y. and Nishigaki, K.: (Genome profiling - establishment and practical evaluation of its methodology). *Nippon Kagaku Kaishi*, (1996) 54-61; C.A., 124 (1996) 136924r.
- 2254 Hubert, S.J., Slater, G.W. and Viovy, J.-L.: Theory of capillary electrophoretic separation of DNA using ultradilute polymer solutions. *Macromolecules*, 29 (1996) 1006-1009; C.A., 124 (1996) 81264t.
- 2255 Ishizuka, T., Kita, K., Sonoda, T., Ishijima, S., Sawa, K., Suzuki, N. and Tatibana, M.: Cloning and sequencing of human complementary DNA for the phosphoribosylpyrophosphate synthetase-associated protein 39. *Biochim. Biophys. Acta*, 1306 (1996) 27-30.
- 2256 Jacobs, D. and Neilan, B.A.: Long-term preservation of DNA in agarose gels using 70% ethanol. *BioTechniques*, 19 (1995) 892-894; C.A., 124 (1996) 49931x.
- 2257 Kozłowski, P., Trzcinska-Danielewicz, J. and Toczko, K.: Identification and sequence analysis of a rap gene from the true slime mold *Physarum polycephalum*. *Biochim. Biophys. Acta*, 1305 (1996) 29-33.
- 2258 Kuryama, A.: (Gel electrophoresis methods for isolation of microorganism DNA from sample containing free DNA of unknown origin). *Jpn. Kokai Tokkyo Koho JP 07,322,899 [95,322,899]* (Cl. C12Q1/68), 12 Dec. 1995, Appl. 94/116,266, 30 May 1994, 8 pp.; C.A., 124 (1996) 167498j.
- 2259 Larsson, A., Aakerman, B. and Jonsson, M.: DAPI staining of DNA: effect of change in charge, flexibility, and contour length on orientational dynamics and mobility of the DNA during agarose gel electrophoresis. *J. Phys. Chem.*, 100 (1996) 3252-3263; C.A., 124 (1996) 140117x.
- 2260 Liu, P.Y.-F., Shi, Z.-Y., Lau, Y.-J., Hu, B.-S., Shyr, J.-M., Tsai, W.-S., Lin, Y.-H. and Tseng, C.Y.: Epidemiological typing of *Flavimonas oryzihabitans* by PCR and pulsed-field gel electrophoresis. *J. Clin. Microbiol.*, 34 (1996) 68-70; C.A., 124 (1996) 108888p.
- 2261 Matsuda, M., Tsukada, M., Fukuyama, M., Kato, Y., Ishida, Y., Honda, M. and Kaneuchi, C.: Detection of genomic variability among isolates of *Campylobacter jejuni* from chickens by crossed-field gel electrophoresis. *Cytobios*, 82 (1995) 73-79; C.A., 124 (1996) 169798f.
- 2262 Mazuruk, K., Schoen, T.J., Chader, G.J., Iwata, T. and Rodriguez, I.R.: Structural organization and chromosomal localization of the human ribosomal protein L9 gene. *Biochim. Biophys. Acta*, 1305 (1996) 151-162.
- 2263 Namsaraev, E.A., Lanzov, V.A. and Akhmedov, A.T.: Partial purification and characterization of two types of homologous DNA pairing activity from rat testis nuclei. *Biochim. Biophys. Acta*, 1305 (1996) 172-180.
- 2264 Ozawa, M. and Muramatsu, T.: Molecular cloning and expression of a mouse α -1,3 fucosyltransferase gene that shows homology with the human α -1,3 fucosyltransferase IV gene. *J. Biochem. (Tokyo)*, 119 (1996) 302-308.
- 2265 Paesen, G.C., Feng, X. and Happ, G.M.: Structure of a D-protein gene and amino-acid sequences of the highly repetitive D-proteins secreted by the accessory glands of the mealworm beetle. *Biochim. Biophys. Acta*, 1293 (1996) 171-176.
- 2266 Peskin, A.V.: (The role of oxygen radicals formed during the membrane-redox functioning in nuclear DNA damage). *Biochimiya (Moscow)*, 61 (1996) 65-72.
- 2267 Platt, D.J., Browning, L.M. and Candlish, D.: Molecular analysis of *Salmonella enterica* serotype Dublin: building bridges between population genetic and molecular epidemiological studies. *Electrophoresis (Weinheim)*, 17 (1996) 667-671.
- 2268 Porter, J.F., Connor, K. and Donachie, W.: Differentiation between human and ovine isolates of *Bordetella parapertussis* using pulsed-field gel electrophoresis. *FEMS Microbiol. Lett.*, 135 (1996) 131-135; C.A., 124 (1996) 169810d.
- 2269 Rodríguez, M.A., Báez-Camargo, M., Delgadillo, D.M. and Orozco, E.: Cloning and expression of an *Entamoeba histolytica* NADP⁺-dependent alcohol dehydrogenase gene. *Biochim. Biophys. Acta*, 1306 (1996) 23-26.
- 2270 Sakai, A., Kato, M., Fukasawa, M., Ishiguro, M., Furuya, E. and Sakakibara, R.: Cloning of cDNA encoding for a novel isozyme of fructose 6-phosphate,2-kinase/fructose 2,6-bisphosphatase from human placenta. *J. Biochem. (Tokyo)*, 119 (1996) 506-511.
- 2271 Sasagawa, N., Saitoh, N., Shimokawa, M., Sorimachi, H., Maruyama, K., Arahata, K., Ishiura, S. and Suzuki, K.: Effect of artificial (CTG) repeat expansion on the expression of myotonin protein kinase (MtPK) in COS-1 cells. *Biochim. Biophys. Acta*, 1315 (1996) 112-116.
- 2272 Schmitz, F.J., Geisel, R., Wagner, S., Heinz, H.P., Idel, H. and Hadding, U.: (Standardized pulsed-field gel electrophoresis by the GenePath Strain Typing system for DNA fingerprinting in nosocomial infections). *Laboratoriumsmedizin*, 19 (1995) 512-522; C.A., 124 (1996) 169817m.
- 2273 Sigmon, Melody Jane Moore: The effect of etidium bromide on mobility of DNA and quantitation of UV-induced lesions in DNA using agarose gel electrophoresis. Avail. *Univ. Microfilms Int.*, Order No.: DA9600688, 1994, 83 pp.; C.A., 124 (1996) 166393j.
- 2274 Starchev, K., Sturm, J. and Weill, G.: Dynamics of DNA molecules in agarose gel under sinusoidal electric fields. *Electrophoresis (Weinheim)*, 17 (1996) 465-470.
- 2275 Suhara, W., Yoneyama, M., Yonekawa, H. and Fujita, T.: Structure of mouse interferon stimulated gene factor 3γ (ISGF3 γ /p48) cDNA and chromosomal localization of the gene. *J. Biochem. (Tokyo)*, 119 (1996) 231-234.
- 2276 Suttorp, M., von Neuhoff, N., Tiemann, M., Dreger, P., Schaub, J., Löffler, H., Parwaresch, R. and Schmitz, N.: Precast commercial polyacrylamide gels for separation of DNA amplicates by temperature gradient gel electrophoresis: application to clonality analysis of lymphomas. *Electrophoresis (Weinheim)*, 17 (1996) 672-677.

- 2277 Suzuki, H., Yaoi, T., Kawai, J., Hara, A., Kuwajima, G. and Watanabe, S.: Restriction landmark cDNA scannig (RLCS): a novel cDNA display system using two-dimensional gel electrophoresis. *Nucleic Acids Res.*, 24 (1996) 289-294; *C.A.*, 124 (1996) 166553m.
- 2278 Tamai, Y., Yokono, T., Sunagawa, M., Kojima, Y. and Miura, K.: Separation of large DNA molecules from *Pleurotus cornucopiae* by pulsed-field gel electrophoresis. *Mokuzai Gakkaishi*, 41 (1995) 952-955; *C.A.*, 124 (1996) 77661c.
- 2279 Tarese, R.V., Jr., Cases, S., Ruland, S.L., Kaydem, M.J., Wong, J.S., Young, S.G. and Hamilton, R.L.: A novel function for apolipoprotein B: lipoprotein synthesis in the yolk sac is critical for material-fetal lipid transport in mice. *J. Lipid Res.*, 37 (1996) 347-360.
- 2280 Vincent, U., Patra, G., Therasse, J. and Gareil, P.: Quantitation of polymerase chain reaction-amplified DNA fragments by capillary electrophoresis and laser-induced fluorescence detection. *Electrophoresis (Weinheim)*, 17 (1996) 512-517.
- See also 1617, 1665, 1720, 1822, 1890, 1974, 2021, 2192, 2206, 2287, 2290, 2300, 2302, 2321, 2323.
- 21d. *Structural studies on RNA and RNA mapping*
- 2281 Ishizuka, T., Sawa, K., Kita, K., Ino, H., Sonoda, T., Suzuki, N. and Tatibana, M.: Promoter region of the rat phosphoribosylpyrophosphate synthetase-associated protein 39. *Biochim. Biophys. Acta*, 1306 (1996) 34-37.
- 2282 Laughrea, M. and Jetté, L.: Kissing-Loop model of HIV-1 genome dimerization: HIV-1 RNAs can assume alternative dimeric forms, and all sequences upstream or downstream of hairpin 248-271 are dispensable for dimer formation. *Biochemistry*, 35 (1996) 1589-1598.
- 2283 Smallman, D.S., Schnare, M.N. and Gray, M.W.: RNA:RNA interactions in the large subunit ribosomal RNA of *Euglena gracilis*. *Biochim. Biophys. Acta*, 1305 (1996) 1-6.
- 2284 Zarghami, N. and Diamandis, E.P.: Detection of prostate-specific antigen mRNA and protein in breast tumors. *Clin. Chem. (Washington)*, 42 (1996) 361-366.
- 21e. *Structural studies on DNA and DNA mapping*
- 2285 Baba, Y., Nishimura, A., Sumita, C., Tshako, M.T. and Ogihara, T.: (Analysis of repeated sequence (VNTR and microsatellite) in human genome using capillary electrophoresis). *Kuromatogurafi*, 16 (1995) 312-313; *C.A.*, 124 (1996) 166487t.
- 2286 Benore-Parsons, M. and Anderson, L.: Purification of DNA fragments from lyophilized agarose gels. *Nucleic Acids Res.*, 23 (1995) 4926-4927; *C.A.*, 124 (1996) 111534a.
- 2287 Blank, R.D., Sklar, C.A. and Martin, M.L.: Denaturing gradient gel electrophoresis to diagnose multiple endocrine neoplasia type 2. *Clin. Chem. (Washington)*, 42 (1996) 598-603.
- 2288 Cheng, J., Shoffner, M.A., Mitchelson, K.R., Kricka, L.J. and Wilding, P.: Analysis of ligase chain reaction products amplified in a silicon-glass chip using capillary electrophoresis. *J. Chromatogr. A*, 732 (1996) 151-158.
- 2289 Cohen, A.S., Smisek, D.L. and Wang, B.H.: Emerging technologies for sequencing antisense oligonucleotides: capillary electrophoresis and mass spectrometry. *Adv. Chromatogr.*, 36 (1996) 127-162; *C.A.*, 124 (1996) 136764p - a review with 115 refs.
- 2290 Guillemm, C., Lellouche, F., Darnige, L., Schandelong, A., Gefroy, F. and Dorval, I.: Rapid detection of the G1691A mutation of coagulation factor V by PCR-mediated site-directed mutagenesis. *Clin. Chem. (Washington)*, 42 (1996) 329.
- 2291 Haapalainen, M., Karp, M. and Metzler, M.C.: Isolation of strong promoters from *Clavibacter xyli* subsp. *cynodontis* using a promoter probe plasmid. *Biochim. Biophys. Acta*, 1305 (1996) 130-134.
- 2292 Hampe, J., Marczinek, K., Preuss, A. and Nürnberg, P.: Marker pattern instabilities as a major cause of reproducibility problems in two-dimensional DNA fingerprinting. *Electrophoresis (Weinheim)*, 17 (1996) 659-666.
- 2293 Holz, O., Joeres, R., Kaestner, A., Krause, T. and Magnussen, H.: Reproducibility of basal and induced DNA single-strand breaks detected by the single-cell gel electrophoresis assay in human peripheral mononuclear leukocytes. *Int. Arch. Occup. Environ. Health*, 67 (1995) 305-310; *C.A.*, 124 (1996) 81249s.
- 2294 Jacobson, S.C. and Ramsey, J.M.: Integrated microdevice for DNA restriction fragment analysis. *Anal. Chem.*, 68 (1996) 720-723.
- 2295 Kamahori, M. and Kambara, H.: (Analysis of DNA fragments by capillary electrophoresis). *Seibutsu Butsuri Kagaku*, 39 (1995) 363-370; *C.A.*, 124 (1996) 111548h.
- 2296 Kayanoki, Y., Fujii, J., Islam, K.N., Suzuki, K., Kawata, S., Matsuzawa, Y. and Taniguchi, N.: The protective role of glutathione peroxidase in apoptosis induced by reactive oxygen species. *J. Biochem. (Tokyo)*, 119 (1996) 817-822.
- 2297 Kim, S., Yoo, H.J. and Hahn, J.H.: Postelectrophoresis capillary scanning method for DNA sequencing. *Anal. Chem.*, 68 (1996) 936-939.
- 2298 Kim, W.S.: A simple and quick method of isolating DNA fragments from agarose gel suitable for molecular procedures. *Methods Mol. Cell. Biol.*, 5 (1994; Pub. 1995) 242-244; *C.A.*, 124 (1996) 166576w.
- 2299 Kiryanov, G.I., Kintsurashvili, L.N., Manamshyan, T.A., Noskov, V.A. and Smirnova, T.A.: (Chromatin structure and enzymatic methylation of α -amylase gene DNA in barley aleurone layer cells during gene expression induction). *Biokhimiya (Moscow)*, 61 (1996) 55-64.
- 2300 Koch, M., Lehmann, R., Pfohl, M., Voelter, W. and Liebich, H.: Analysis of the deletion/insertion polymorphism of the angiotensin I-converting enzyme gene by capillary electrophoresis. *Clin. Chim. Acta*, 248 (1996) 197-203.
- 2301 Kolesar, J.M., Rizzo, J.D. and Kuhn, J.G.: Quantitative analysis of NQO1 gene expression by RT-PCR and CE-LIF. *J. Capillary Electrophor.*, 2 (1995) 287-290; *C.A.*, 124 (1996) 195067m.
- 2302 Law, P.T.W., Tsui, S.K.W., Lam, W.Y., Luk, S.C.W., Hwang, D.M., Liew, C.C., Lee, C.Y., Fung, K.P. and Waye, M.M.Y.: A novel cDNA encoding a human homologue of ribosomal protein L29. *Biochim. Biophys. Acta*, 1305 (1996) 105-108.
- 2303 Lee, J.-G.: The assessment of oxidant-induced DNA single-strand breaks in human and animal lung cells: use of the single-cell gel electrophoresis assay. *Avail. Univ. Microfilms Int.*, Order No.: DA9542235, 1995, 144 pp.; *C.A.*, 124 (1996) 109444c.
- 2304 Lopez-Solache, I., Luu-The, v., Seralini, G.-E. and Labrie, F.: Heterogeneity of rat type I 5 α -reductase cDNA: cloning, expression and regulation by pituitary implants and dihydrotestosterone. *Biochim. Biophys. Acta*, 1305 (1996) 139-144.

- 2305 McGraw, D.W., Jacobi, S.E., Hiller, F.C. and Cornett, L.E.: Structural and functional analysis of the 5'-flanking region of the rat β_2 -adrenergic receptor gene. *Biochim. Biophys. Acta*, 1305 (1996) 135-138.
- 2306 McIndoe, R.A., Hood, L. and Bumgarner, R.E.: An analysis of the dynamic range and linearity of an infrared DNA sequencer. *Electrophoresis (Weinheim)*, 17 (1996) 652-658.
- 2307 Mishima, Y., Suda, T. and Kominami, R.: Formation of a triple-stranded DNA between d(GGA:TCC) repeats and d(GGA) repeat oligonucleotides. *J. Biochem. (Tokyo)*, 119 (1996) 805-810.
- 2308 Monteith, D.K. and Vanstone, J.: Comparison of the microgel electrophoresis assay and other assays for genotoxicity in the detection of DNA damage. *Mutat. Res.*, 345 (1995) 97-103; *C.A.*, 124 (1996) 109117y.
- 2309 Mullaart, E., Vijg, J. and de Vos, G.J.: Method for detecting mutations by two-dimensional electrophoresis of amplified labeled genome fragments. *PCT Int. Appl. WO 96 00,796 (Cl. C12Q1/68)*, 11 Jan. 1996, NL Appl. 94/1,082, 28 Jun. 1994; 23 pp.; *C.A.*, 124 (1996) 137817v.
- 2310 Murray, V., Matias, C., McFadyen, W.D. and Wickham, G.: The influence of linker chain length on the sequence specificity of DNA damage by *n*-bromoalkylphenanthridinium bromides in plasmid DNA and in intact human cells. *Biochim. Biophys. Acta*, 1305 (1996) 79-86.
- 2311 Nair, T.M., Madhusudan, K., Nagaraja, V., Kulkarni, B.D., Majumdar, H.K. and Singh, R.: Theoretical permutation gel electrophoretic analysis of a curved DNA fragment located in circular permutation. *Electrophoresis (Weinheim)*, 17 (1996) 633-641.
- 2312 Oliveira, M.A.A., Caballero, O.L., Neto, E.D., Koury, M.C., Romanha, A.J., Carvalho, J., Hartskeerl, R.A. and Simpson, A.J.G.: Use of nondenaturing silver-stained polyacrylamide gel analysis of polymerase chain reaction amplification products for the differential diagnosis of *Leptospira interrogans* infection. *Diagn. Microbiol. Infect. Dis.*, 22 (1995) 343-348; *C.A.*, 124 (1996) 77658g.
- 2313 Paesen, G.C., de A. Zanotto, P.M. and Nuttall, P.A.: A tick homologue of the human DNA helicase II 70-kDa subunit. *Biochim. Biophys. Acta*, 1305 (1996) 120-124.
- 2314 Pandrangi, R., Petras, M., Ralph, S. and Vrzc, M.: Alkaline single cell gel (comet) assay and genotoxicity monitoring using bullheads and carp. *Environ. Mol. Mutagen.*, 26 (1995) 345-356; *C.A.*, 124 (1996) 78851b.
- 2315 Petersen, N.E., Nissen, H., Hansen, T.S., Rasmussen, K., Brock, A. and Horder, M.: R325X mutation in exon 15 of the hydroxymethylbilane synthase gene identified in two Danish families with acute intermittent porphyria. *Clin. Chem. (Washington)*, 42 (1996) 106-107.
- 2316 Qin, C., Shen, J., Huang, S. and Wang, G.: (Single cell gel electrophoresis assay: a DNA breakage detection technique). *Shengwu Huaxue Yu Shengwu Wuli Jinzha*, 22 (1995) 517-520; *C.A.*, 124 (1996) 169597q.
- 2317 Schlenck, A., Visvikis, S., O'Kane, M. and Siest, G.: High-resolution separation of PCR product and gene diagnosis by capillary gel electrophoresis. *Biomed. Chromatogr.*, 10 (1996) 48-50; *C.A.*, 124 (1996) 107940n.
- 2318 Seo, H.G., Fujiwara, N., Kaneto, H., Asahi, M., Fujii, J. and Taniguchi, N.: Effect of a nitric oxide synthase inhibitor, S-ethylisothiourea, on cultured cells and cardiovascular functions of normal and lipopolysaccharide-treated rabbits. *J. Biochem. (Tokyo)*, 119 (1996) 553-558.
- 2319 Stoerker, J., Hurwitz, C., Rose, N.C., Silberstein, L.E. and Highsmith, W.E.: Heteroduplex generator in analysis of Rh blood group alleles. *Clin. Chem. (Washington)*, 42 (1996) 356-360.
- 2320 Tubiello, G., Carrera, P., Soriani, N., Morandi, L. and Ferrari, M.: Mutational analysis of muscle and brain specific promoter regions of dystrophin gene in DMD/BMD Italian patients by denaturing gradient gel electrophoresis (DGGE). *Mol. Cell. Probes*, 9 (1995) 441-446; *C.A.*, 124 (1996) 142708q.
- 2321 Tuo, J., Loft, S., Thomsen, M.S. and Poulsen, H.E.: *Ex vivo* time-dependent cell DNA-degradation shown by single cell gel electrophoresis. *Pharmacol. Toxicol. (Copenhagen)*, 78 (1996) 55-57; *C.A.*, 124 (1996) 140240g.
- 2322 Vandenbroeck, K., Consiglio, A., Martino, G. and Grimaldi, L.M.E.: Improved conditions for the analysis of large variable number of tandemly repeated (VNTR) unit polymorphisms. *Electrophoresis (Weinheim)*, 17 (1996) 678-680.
- 2323 Wang, G., Godinger, D., Aronovitch, J. and Samuni, A.: Opposing effects of nitroxide free radicals in *Escherichia coli* mutants deficient in DNA repair. *Biochim. Biophys. Acta*, 1305 (1996) 71-78.
- 2324 Wu, C.-G., Hakvoort, T.B.M., Lamers, W.H. and Chamuleau, R.A.F.M.: Isolation of up- and down-regulated cDNAs associated with hepatocellular carcinoma by a subtraction-enhanced display technique. *Biochim. Biophys. Acta*, 1315 (1996) 169-175.
- 2325 Zaporozhets, D.A. and Mironov, A.S.: Physical mapping of *Escherichia coli* genome rearrangements by means of pulse-field gel electrophoresis and PCR method. *Russ. J. Genet. (Transl. of Genetika (Moscow))*, 32 (1996) 17-27; *C.A.*, 124 (1996) 195586e.
- See also 2179, 2184, 2217, 2239, 2394.
- 21f. *Complex mixtures of nucleic acids and their fragments*
- 2326 Xian, J., Harrington, M.G. and Davidson, E.H.: DNA-protein binding assays from a single sea urchin egg: a high-sensitivity capillary electrophoresis method. *Proc. Natl. Acad. Sci. U.S.A.*, 93 (1996) 86-90; *C.A.*, 124 (1996) 111552e.
- See also 2047.
22. ALKALOIDS
- 2327 Hufschmid, E., Theurillat, R., Wilder-Smith, C.H. and Thormann, W.: Characterization of the genetic polymorphism of dihydrocodeine O-demethylation in man via analysis of urinary dihydrocodeine and dihydromorphine by micellar electrokinetic capillary chromatography. *J. Chromatogr. B*, 678 (1996) 43-51.

23. OTHER SUBSTANCES CONTAINING HETEROCYCLIC NITROGEN

23a. Porphyrins and other pyrroles

2328 Yao, Y.J. and Li, S.F.Y.: Determination of erythrocyte porphyrins by epi-illumination fluorescence microscope with capillary electrophoresis. *J. Liq. Chromatogr.*, 19 (1996) 1-15.

23c. Indole derivatives and plant hormones (gibberelins)

2329 Lancas, F.M. and Galhiane, M.S.: Separation of plant growth regulators in citrus crops by capillary electrophoresis. *J. Capillary Electrophor.*, 2 (1995) 77-79; *C.A.*, 124 (1996) 111545e.

2330 Zhang, L., Jin, S., Zhou, X., Gu, J.-L. and Fu, R.-N.: Separation of diastereomers of 1,3-disubstituted tetrahydro-1-carboline by capillary electrophoresis with β -cyclodextrin. *Chromatographia*, 42 (1996) 385-388.

23d. Pyridine derivatives

2331 McKillop, A.G., Smith, R.M., Rowe, R.C. and Wren, S.A.C.: Optimisation of the separation of the dimethylpyridines by capillary electrophoresis. *J. Chromatogr. A*, 730 (1996) 321-328.

23e. Other N-heterocyclic compounds

2332 Chen, C.-C. and Lin, C.-E.: Analysis of copper corrosion inhibitors by capillary zone electrophoresis. *Anal. Chim. Acta*, 321 (1996) 215-218.

2333 Lin, C.-E. and Chen, C.-C.: Migration behavior and separation of aromatic triazole and thiazole compounds by capillary zone electrophoresis. *J. Chromatogr. A*, 731 (1996) 299-303.

2334 Milofsky, R. and Spaeth, S.: Determination of orotic acid in aqueous solutions by micellar electrokinetic capillary chromatography using sensitized lanthanide-ion luminescence detection. *Chromatographia*, 42 (1996) 12-16.

2335 Wu, J., Wong, M.K., Lee, H.K. and Ong, C.N.: Orthogonal array design for optimizing the capillary zone electrophoretic analysis of heterocyclic amines. *J. Chromatogr. Sci.*, 34 (1996) 139-145.

See also 2327.

24. ORGANIC SULPHUR COMPOUNDS (INCL. GLUCOSINOLATES)

2336 Abo, M., Tachibana, M., Okubo, A. and Yamazaki, S.: Application of micellar electrokinetic chromatography for monitoring the kinetic resolution of sulfoxides by dimethyl sulfoxide reductase. *Bunseki Kagaku*, 44 (1995) 835-838; *C.A.*, 124 (1996) 86489e.

2337 Ercal, N., Le, K., Treeratphan, P. and Matthews, R.: Analysis of thiol-containing compounds in biological samples by capillary zone electrophoresis. *Biomed. Chromatogr.*, 10 (1996) 15-18; *C.A.*, 124 (1996) 111566n.

2338 Kok, S.J., Koster, E.H.M., Gooijer, C., Velthorst, N.H. and Brinkman, U.A.T.: Separation of twenty-one naphthalene sulfonates by means of capillary electrophoresis. *J. High Resolut. Chromatogr.*, 19 (1996) 99-104.

2339 Okada, T.: Design of chromatographic and electrophoretic separation based on ion-pair formation of aromatic disulfonates with polyammonium ions. *Anal. Chem.*, 68 (1996) 1158-1163.

2340 Wronski, M.: Separation of urinary thiols as tributyltinmercaptides and determination using capillary isotachopheresis. *J. Chromatogr. B*, 676 (1996) 29-34.

2341 Zenki, M., Yukutake, H. and Irizawa, M.: (Separation of naphthalenesulphonates by capillary zone electrophoresis). *Bunseki Kagaku*, 45 (1996) 181-184; *C.A.*, 124 (1996) 192884q.

See also 2332, 2333, 2412.

25. ORGANIC PHOSPHORUS COMPOUNDS (INCL. SUGAR PHOSPHATES)

2342 Jara, D. and Borros, S.: (Preparation of a method for the determination of monofluorophosphate in toothpastes and similar products by capillary electrophoresis). *Afinidad*, 52 (1995) 393-398; *C.A.*, 124 (1996) 97238y.

2343 Shaw, C.J. and Silverman, C.E.: Enantiomeric separation of aminophosphonic acids by capillary electrophoresis. *Chirality*, 8 (1996) 84-87; *C.A.*, 124 (1996) 242481d.

See also 2103, 2346.

26. ORGANOMETALLIC AND RELATED COMPOUNDS

26a. Organometallic compounds

2344 Carro-Diaz, A.M., Lorenzo-Ferreira, R.A. and Cela-Torrijos, R.: Capillary electrophoresis of methylmercury with injection by sample stacking. *J. Chromatogr. A*, 730 (1996) 345-351.

2345 Li, S.F.Y. and Li, K.: Determination of organometallic compounds in environmental samples by capillary electrophoresis after on-column or off-column preconcentration. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 459-466; *C.A.*, 124 (1996) 75172v.

See also 1625, 2340.

26c. Coordination compounds

2346 Pandey, A.K. and Singh, R.K.P.: A novel method for the study of simple and mixed metal-complexes of some bio-active ligands. *Proc. Natl. Acad. Sci., India, Sect. A 1995*, 65 (1995) 35-41; *C.A.*, 124 (1996) 186755g.

2347 Semenova, O.P., Orinak, A., Matisova, E. and Timerbaev, A.R.: Stability study of novel zinc complexes of pharmaceutical relevance by CE. *J. Capillary Electrophor.*, 2 (1995) 203-207; *C.A.*, 124 (1996) 156242v.

2348 Tewari, B.B.: Ionophoretic technique study of binary complexes of lead(II) and uranyl(II) with sulphur containing organic ligands. *Trans SAEST*, 30 (1995) 100-103; *C.A.*, 124 (1996) 163451r.

28. ANTIBIOTICS

- 2349 Castaneda Penalvo, G., Julien, E. and Fabre, H.: Cross validation of capillary electrophoresis and high-performance liquid chromatography for cefotaxime and related impurities. *Chromatographia*, 42 (1996) 159-164.
- 2350 Chen, C.-L. and Gu, X.: Determination of tetracycline residues in bovine milk, serum, and urine by capillary electrophoresis. *J. Assoc. Off. Anal. Chem.*, 78 (1995) 1369-1377.
- 2351 Fabre, H. and Castaneda Penalvo, G.: Capillary electrophoresis as an alternative method for the determination of cefotaxime. *J. Liq. Chromatogr.*, 18 (1995) 3877-3887.
- 2352 Flurer, C.L.: Analysis of macrolide antibiotics by capillary electrophoresis. *Electrophoresis (Weinheim)*, 17 (1996) 359-366.
- 2353 Van Schepdael, A., van den Bergh, I., Roets, E. and Hoogmartens, J.: Purity control of oxytetracycline by capillary electrophoresis. *J. Chromatogr. A*, 730 (1996) 305-311.

See also 1747, 2392.

29. INSECTICIDES, PESTICIDES AND OTHER AGROCHEMICALS

29a. General techniques

- 2354 Süsse, H. and Müller, H.: Pesticide analysis by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 730 (1996) 337-343.

29b. Chlorinated insecticides

- 2355 Grainger, J., McClure, P.C., Liu, Z., Botero, B., Sirimanne, S., Patterson, D.G., Jr., Sewer, M., Gillyard, C., Kimata, K. et al.: Isomer identification of chlorinated dibenzo-*p*-dioxins by orthogonal spectroscopic and chromatographic techniques. *Chemosphere*, 32 (1996) 13-23; *C.A.*, 124 (1996) 163838x.

See also 1739.

29d. Carbamates

- 2356 Lancas, F.M., Rissato, S.R. and Galhiane, M.S.: Off-line SFE-CZE analysis of carbamates residues in Tobacco samples. *Chromatographia*, 42 (1996) 323-327.

29e. Herbicides

- 2357 Schmitt, P., Garrison, A.W., Freitag, D. and Kettrup, A.: Separation of *s*-triazine herbicides and their metabolites by capillary zone electrophoresis as a function of pH. *J. Chromatogr. A*, 723 (1996) 169-177.

See also 1739, 1844.

30. SYNTHETIC AND NATURAL DYES

30a. Synthetic dyes

- 2358 Masar, M., Kaniansky, D. and Madajova, V.: Separation of synthetic food colourants by capillary zone electrophoresis in a hydrodynamically closed separation compartment. *J. Chromatogr. A*, 724 (1996) 327-336.
- 2359 Xiong, S., Li, J. and Cheng, J.: (Studies on the coupling technique of high performance capillary electrophoresis and charge-coupled device detection. II. Separation of fluorescent dye mixtures). *Fenxi Huaxue*, 24 (1996) 54-56; *C.A.*, 124 (1996) 120100h.

See also 1657.

30b. Chloroplast and other natural pigments

- 2360 Ciavatta, C., Govi, M., Sitti, L. and Gessa, C.: Capillary electrophoresis of humic acid fractions. *Commun. Soil Sci. Plant Anal.*, 26 (1995) 3305-3313; *C.A.*, 124 (1996) 54674v.
- 2361 Pompe, S., Heise, K.-H. and Nitsche, H.: Capillary electrophoresis for a "finger-print" characterization of fulvic and humic acids. *J. Chromatogr. A*, 723 (1996) 215-218.

See also 1816.

31. PLASTICS AND THEIR INTERMEDIATES

- 2362 Bastos-Gonzales, D., Hidalgo-Alvarez, R. and de las Nieves, F.J.: Electrokinetic behavior of polystyrenes latexes with different surface groups: effect of heat treatment. *J. Colloid Interface Sci.*, 177 (1996) 372-379; *C.A.*, 124 (1996) 156983n.
- 2363 Nielen, M.W.F. and van de Ven, H.J.F.M.: Characterization of (methoxymethyl)melamine resin by combined chromatographic/mass spectrometric techniques. *Rapid Commun. Mass Spectrom.*, 10 (1996) 74-81; *C.A.*, 124 (1996) 88812d.
- 2364 Xia, J., Dubin, P.L., Izumi, T., Hirata, M. and Kokofuta, E.: Dynamic and electrophoretic light scattering of poly(dimethyldiallylammonium chloride) in salt-free solutions. *J. Polym. Sci., Part B: Polym. Phys.*, 34 (1996) 497-503; *C.A.*, 124 (1996) 118632q.
- 2365 Zimmerman, S.B. and Murphy, L.D.: Electrophoresis of polyethylene glycols and related materials as sodium dodecyl sulfate complexes. *Anal. Biochem.*, 234 (1996) 190-193.

See also 2414.

32. DRUG ANALYSIS

32a. Drug analysis, general techniques

- 2366 Altria, K.D.: Application of capillary electrophoresis to pharmaceutical analysis. *Methods Mol. Biol. (Totowa)*, 52 (1996) 265-284; *C.A.*, 124 (1996) 156160s - a review with 60 refs.

- 2367 Altria, K.D., Elgey, J., Lockwood, P. and Moore, D.: An overview of the applications of capillary electrophoresis to the analysis of pharmaceutical raw materials and excipients. *Chromatographia*, 42 (1996) 332-342.
- 2368 Bingcheng, L., Yibing, J., Yuang, C., Epperlein, U. and Koppenhoefer, B.: Separation of study enantiomers by capillary electrophoresis: α -cyclodextrin as chiral solvation agent. *Chromatographia*, 42 (1996) 106-110.
- 2369 Carducci, C.N., Lucangioli, S.E., Rodríguez, V.G. and Fernández Otero, G.C.: Application of extraction disks in dissolution tests of clenbuterol and levothyroxine tablets by capillary electrophoresis. *J. Chromatogr. A*, 730 (1996) 313-319.
- 2370 Desiderio, C. and Fanali, S.: (Stereoselective analysis of pharmaceutical compounds, in biological fluids using capillary electrophoresis). *Boll. Chim. Farm.*, 134 (1995) 541-546; *C.A.*, 124 (1996) 193174v.
- 2371 Gomez, F.A., Mirkovich, J.N., Dominguez, V.M., Liu, K.W. and Macias, D.M.: Multiple-plug binding assays using affinity capillary electrophoresis. *J. Chromatogr. A*, 727 (1996) 291-299.
- 2372 Guttman, A., Aumatell, A., Brunet, S. and Cooke, N.: Cyclodextrin array chiral analysis. *Am. Lab. (Shelton)*, 27 (1995) 18-22; *C.A.*, 124 (1996) 156208p.
- 2373 Guttman, A., Brunet, S. and Cooke, N.: Capillary electrophoresis separation of enantiomers by cyclodextrin array chiral analysis. *LC-GC*, 14 (1996) 32-42; *C.A.*, 124 (1996) 156213m.
- 2374 Herbert, B.J.: Octanol-water partition coefficient estimation by capillary electrophoresis and subsequent relevance to bioavailability prediction. *Avail. Univ. Microfilms Int.*, Order No.: DA9541640, 1995, 177 pp.; *C.A.*, 124 (1996) 111549j.
- 2375 Lurie, I.S.: Application of capillary electrophoresis to the analysis of seized drugs. *Int. Lab.*, 26, March (1996) 21-29.
- 2376 Marzo, A. and Balant, L.P.: Investigation of xenobiotic metabolism by CYP2D6 and CYP2C19: importance of enantioselective analytical methods. *J. Chromatogr. B*, 678 (1996) 73-92 - a review with 105 refs.
- 2377 Oravcova, J., Böhs, B. and Lindner, W.: Drug-protein binding studies. New trends in analytical and experimental methodology. *J. Chromatogr. B*, 677 (1996) 1-28 - a review with 241 refs.
- 2378 Rickard, E.C., Bopp, R.J., Skanchy, D.J., Chetwyn, K.L., Pahlen, B. and Stobaugh, J.F.: Role of capillary electrophoresis method in the drug development process. *Chirality*, 8 (1996) 108-121; *C.A.*, 124 (1996) 212179u.
- 2379 Swartz, M.E. and Brown, P.R.: Use of mathematically enhanced spectral analysis and spectral contrast techniques for the liquid chromatographic and capillary electrophoretic detection and identification of pharmaceutical compounds. *Chirality*, 8 (1996) 67-76; *C.A.*, 124 (1996) 242479j.
- See also 1666, 1730, 1742, 1743, 1748, 1749, 1752, 1835, 1857, 1964, 2401.
- 32b. Antirheumatics and antiinflammatory drugs**
- 2380 Ashcroft, A.E., Major, H.J., Lowes, S. and Wilson, I.D.: Identification of non-steroidal anti-inflammatory drugs and their metabolites in solid phase extracts of human urine using capillary electrophoresis-mass spectrometry. *Anal. Proc.*, 32 (1995) 459-462; *C.A.*, 124 (1996) 44586d.
- 2381 Aturki, Z., Camera, E., La Torre, F. and Fanali, S.: Direct chiral resolution of tiaprofenic acid in pharmaceutical formulations by capillary zone electrophoresis using cyclodextrins as chiral selectors. *J. Capillary Electrophor.* 1995, 2 (1995) 213-217; *C.A.*, 124 (1996) 127241p.
- 2382 Bechet, I., Fillet, M., Hubert, P. and Crommen, J.: Improvement of achiral resolution in capillary zone electrophoresis by use of cyclodextrin additives. *Biomed. Chromatogr.*, 9 (1995) 267-268; *C.A.*, 124 (1996) 66732a.
- 2383 Kataoka, M., Imamura, M., Nishijima, K. and Nishi, K.: (Simultaneous determination method for the active ingredients in cold medicines by capillary electrophoresis). *Iyakuhin Kenkyu*, 27 (1996) 49-56; *C.A.*, 124 (1996) 242501k.
- See also 1751.
- 32c. Autonomic and cardiovascular drugs**
- 2384 Clothier, J.G., Jr. and Tomellini, S.A.: Chiral separation of verapamil and related compounds using micellar electrokinetic capillary chromatography with mixed micelles of bile salt and polyoxyethylene ethers. *J. Chromatogr. A*, 723 (1996) 179-187.
- 2385 Ljungberg, H. and Nilsson, S.: Protein-based capillary affinity gel electrophoresis for chiral separation of β -adrenergic blockers. *J. Liq. Chromatogr.*, 18 (1995) 3685-3698.
- 2386 Swartz, M.E., Mazzeo, J.R., Grover, E.R., Brown, P.R. and Aboul-Enein, H.Y.: Separation of piperidine-2,6-dione drug enantiomers by micellar electrokinetic capillary chromatography using synthetic chiral surfactants. *J. Chromatogr. A*, 724 (1996) 307-316.
- See also 1739, 2369.
- 32d. Central nervous system drugs**
- 2387 Liu, J., Zhou, W., Wu, M. and Wang, E.: (Determination of acetaminophen and its hydrolyzate by capillary electrophoresis-electrochemical detection). *Fenxi Huaxue*, 23 (1995) 1256-1260; *C.A.*, 124 (1996) 66735d.
- 2388 Sängers-van de Griend, C.E., Grönningsson, K. and Westerlund, D.: Chiral separation of local anaesthetics with capillary electrophoresis. Evaluation of the inclusion complex of the enantiomers with heptakis(2,6-di-O-methyl)- β -cyclodextrin. *Chromatographia*, 42 (1996) 263-268.
- 2389 Tomita, M. and Okuyama, T.: Application of capillary electrophoresis to the simultaneous screening and quantitation of benzodiazepines. *J. Chromatogr. B*, 678 (1996) 331-337.
- See also 2391.
- 32e. Chemotherapeutics (exc. cytostatics and antibiotics)**
- 2390 Chen, Y., Han, F. and Yuan, Z.: (Application of capillary zone electrophoresis in analysis of antagonists as *p*-hydroxybenzoic acid and sulphanilamide). *Fenxi Huaxue*, 23 (1995) 1281-1283; *C.A.*, 124 (1996) 66736e.
- 2391 Nickerson, B., Scypinski, S., Sokoloff, H. and Sahota, S.: Separation of sulfisoxazole, phenazopyridine, and their related impurities by micellar electrokinetic chromatography. *J. Liq. Chromatogr.*, 18 (1995) 3847-3875.

See also 1734.

32f. Cytostatics

2392 Fletcher, M.C. and Fox, K.R.: Dissociation kinetics of echinomycin from CpG binding sites in different sequence environments. *Biochemistry*, 35 (1996) 1064-1075.

2393 Wenclawiak, B.W. and Wollmann, M.: Separation of platinum(II) anti-tumour drugs by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 724 (1996) 317-326.

32g. Other drug categories

2394 Bailly, C., Hamy, F. and Waring, M.J.: Cooperativity in the binding of echinomycin to DNA fragments containing closely spaced CpG sites. *Biochemistry*, 35 (1996) 1150-1161.

2395 Esquisabel, A., Goyenaga, P.F., Vicente, A.S., Calvo, M.B. and Pedraz, J.L.: Capillary electrophoresis for the chiral separation of salbutamol. In: *World Meet. Pharm., Biopharm Pharm. Technol., 1st, APgl, Chatenay Malabry*, 1995, pp. 121-122; *C.A.*, 124 (1996) 242472b.

2396 Geise, R.J. and Machnicki, N.I.: A study of parabens as model hydrophobic compounds by capillary electrophoresis and their determination in cosmetic formulations. *J. Capillary Electrophor.*, 2 (1995) 69-75; *C.A.*, 124 (1996) 97218s.

2397 Mälkki-Laine, L. and Hartikainen, E.: Electrokinetic behaviour of salbutamol and its decomposition products and determination of salbutamol by micellar electrokinetic capillary chromatography. *J. Chromatogr. A*, 724 (1996) 297-306.

2398 Oylar, A.R., Armstrong, B.L., Cha, J.Y., Zhou, M.X., Yang, Q., Robinson, R.I., Dunphy, R. and Burinsky, D.J.: Hydrophilic interaction chromatography on amino-silica phases complements reversed-phase high-performance liquid chromatography and capillary electrophoresis for peptide analysis. *J. Chromatogr. A*, 724 (1996) 378-383.

2399 Ren, J. and Liu, H.: Chiral separation of dioxypromethazine enantiomers by capillary electrophoresis using β -cyclodextrin as a chiral selector. *J. Chromatogr. A*, 732 (1996) 175-181.

2400 Shibabi, Z.K., Rocco, M.V. and Hinsdale, M.E.: Analysis of the contrast agent iopamidol in serum by capillary electrophoresis. *J. Liq. Chromatogr.*, 18 (1995) 3825-3831.

See also 1837, 1838, 1851, 2334, 2342, 2367, 2396.

32h. Toxicological and forensic applications

2401 Steinmann, L. and Thormann, W.: Toxicological drug screening and confirmation by electrokinetic capillary techniques: concept of an automated system. *J. Capillary Electrophor.*, 2 (1995) 81-88; *C.A.*, 124 (1996) 109093n.

33. CLINICO-CHEMICAL APPLICATIONS

33a. General papers and reviews

2402 Ishioka, N.: (A new approach for clinical chemistry. Genetic variant and chemical structure). *Kuromatogurafi*, 16 (1995) 284-285; *C.A.*, 124 (1996) 140236k.

33b. *Complex mixtures and profiling (single compounds by cross-reference only)*

2403 Alkmin, M.G.A., Landgraf, I.M., Vieira, M.F.P., Camargo, M.C.C. and Goncalves, M.I.C.: Diagnosis of bacterial meningitis and septicemia by serum counterimmunoelectrophoresis. *Braz. J. Med. Biol. Res.*, 28 (1995) 1065-1068; *C.A.*, 124 (1996) 81251m.

2404 Gao, C., Zhu, Y., Zhang, J., Liu, J., Wang, M., Li, R. and Han, Y.: (Establishment and application of diagnostic method of hepatoma using AFP-ferritin enzyme-linked-counter immunoelectrophoresis). *Shandong Yike Daxue Xuebao*, 33 (1995) 111-114; *C.A.*, 124 (1996) 140239p.

2405 Jariego, C.M. and Hernanz, A.: Determination of organic acids by capillary electrophoresis in screening of organic acidurias. *Clin. Chem. (Washington)*, 42 (1996) 477-478.

2406 Shah, R., Green, M., Barbadora, K.A., Wagener, W.C., Schwartz, B., Facklam, R.R. and Wald, E.R.: Comparison of M and T type antigen testing to field inversion gel electrophoresis in the differentiation of strains of group A Streptococcus. *Pediatr. Res.*, 38 (1995) 988-992; *C.A.*, 124 (1996) 142967y.

See also 1792, 1815, 1825, 1924, 1979, 1983, 1990, 2005, 2121, 2124, 2168, 2173, 2177, 2226, 2284, 2287, 2290, 2300, 2315, 2319, 2328.

34. FOOD ANALYSIS

34a. General papers and reviews

2407 Lindeberg, J.: Addendum to "Capillary electrophoresis in food analysis". *Food Chem.*, 55 (1995) 95-101; *C.A.*, 124 (1996) 53928u - a review with 41 refs.

2408 Lindeberg, J.: Capillary electrophoresis in food analysis. *Food Chem.*, 55 (1995) 73-94; *C.A.*, 124 (1996) 53927t - a review with 170 refs.

34b. *Complex mixtures (single compounds by cross-reference only)*

2409 Molina, E., Romos, M. and Martin-Alvarez, P.J.: Prediction of the percentages of cows', goats' and ewes' milk in "Iberico" cheese by electrophoretic analysis of whey proteins. *Z. Lebensm.-Unters. Forsch.*, 201 (1995) 331-335; *C.A.*, 124 (1996) 173970y.

2410 Strickland, M., Weimer, B.C. and Broadbent, J.R.: Capillary electrophoresis of Cheddar cheese. *J. Chromatogr. A*, 731 (1996) 305-313.

See also 1818, 1994, 2017, 2358.

35. ENVIRONMENTAL ANALYSIS

35a. General papers and reviews

See 2345, 2355, 2435.

35c. *Water pollution (complex mixtures; single compounds by cross-reference only)*

See 2338, 2422, 2428, 2429.

36. SOME TECHNICAL PRODUCTS AND COMPLEX MIXTURES

36a. *Surfactants*

2411 Monclus, A., Rodriguez, M. and Masana, I.: The analysis of anionic surfactants in toiletries using CE. *LC-GC Int.*, 9 (1996) 79-87.

See also 1737, 2341.

36c. *Complex mixtures, technical products and unidentified compounds*

2412 Dahlman, O. and Maansson, K.: Analysis of low molecular weight lignin-derived sulphonates by capillary zone electrophoresis. *J. Wood Chem. Technol.*, 16 (1996) 47-60; *C.A.*, 124 (1996) 205327n.

2413 Harakuwe, A.H., Haddad, P.R. and Thomas, R.: Capillary zone electrophoresis method for the determination of sulphate in concrete. *Anal. Proc.*, 32 (1995) 493-494; *C.A.*, 124 (1996) 64499f.

2414 Marina, M.L., Benito, I., Díez-Masa, J.C. and González, M.J.: Chiral separation of polychlorinated biphenyls by micellar electrokinetic chromatography with γ -cyclodextrin as modifier in the separation buffer. *Chromatographia*, 42 (1996) 269-272.

See also 1833.

37. CELLS, CELLULAR PARTICLES AND SUPRAMOLECULAR STRUCTURES

2415 Bauer, J. and Weber, G.: Sodium chloride in preparative free-flow cell electrophoresis: recent developments. *Electrophoresis (Weinheim)*, 17 (1996) 526-528.

2416 Budde, A., Knippel, E., Grümmer, G., Treichler, J., Brockmann, H., Donath, E. and Bäuml, H.: Electrophoretic fingerprinting and multiparameter analysis of cells and particles. *Electrophoresis (Weinheim)*, 17 (1996) 507-511.

2417 Chang, H.-T., Zakharov, S.F. and Chrambach, A.: Separation and isolation of subcellular-sized particles by electrophoresis in polymer solution using the commercial scanning apparatus. *Electrophoresis (Weinheim)*, 17 (1996) 776-780.

2418 Janzen, J., Song, X. and Brooks, D.E.: Interfacial thickness of liposomes containing poly(ethylene glycol)-cholesterol from electrophoresis. *Biophys. J.*, 70 (1996) 313-320; *C.A.*, 124 (1996) 140229k.

2419 Jy, W., Horstman, L.L., Homolak, D. and Ahn, Y.S.: Electrophoretic properties of platelets from normal, thrombotic and ITP patients by doppler electrophoretic light scattering analysis. *Platelets*, 6 (1995) 354-358; *C.A.*, 124 (1996) 81272u.

2420 Toll, E., Castillo, F.J., Crespi, P., Crevecoeur, M. and Greppin, H.: Purification of plasma membranes from leaves of conifer and deciduous tree species by phase partitioning and free-flow electrophoresis. *Physiol. Plant.*, 95 (1995) 399-408; *C.A.*, 124 (1996) 111640g.

See also 1937.

38. INORGANIC COMPOUNDS

38a. *Cations*

2421 Baraj, B., Sastre, A., Martínez, M. and Spahiu, K.: Simultaneous determination of chloride complexes of Pt(IV) and Pd(II) by capillary zone electrophoresis with direct UV absorbance detection. *Anal. Chim. Acta*, 319 (1996) 191-197.

2422 Colburn, B.A., Sepaniak, M.J. and Hinton, E.R.: A screening protocol for the direct determination of low ppb levels of uranyl cation using arsenazo III and capillary electrophoresis. *J. Liq. Chromatogr.*, 18 (1995) 3699-3718.

2423 Hirokawa, T., Nishimoto, K. and Nishiyama, F.: Isotachophoretic separation of Fe(II) and Fe(III) by using 1,10-phenanthroline as a complex-forming agent. *J. Chromatogr. A*, 723 (1996) 389-394.

2424 Prella, S., Todt, F., Langner, K. and Ebeling, R.: (Migration of ions in gelatin). *Prax. Naturwiss., Chem.*, 44 (1995) 32-34; *C.A.*, 124 (1996) 201170h.

2425 Schlegel, D., Mattusch, J. and Wennrich, R.: Speciation analysis of arsenic and selenium compounds by capillary electrophoresis. *Fresenius J. Anal. Chem.*, 354 (1996) 535-539.

2426 Timberbaev, A.R., Semenova, O.P., Buchberger, W. and Bonn, G.K.: Speciation studies by capillary electrophoresis. Simultaneous determination of chromium(III) and chromium(VI). *Fresenius J. Anal. Chem.*, 354 (1996) 414-419.

2427 Wen, J. and Cassidy, R.M.: Anodic and cathodic pulse amperometric detection of metal ions separated by capillary electrophoresis. *Anal. Chem.*, 68 (1996) 1047-1053.

2428 Xu, J. and Ma, Y.: Determination of iron in rainwater, lakewater, and tapwater by high performance capillary electrophoresis after precolumn complexation with 1,10-phenanthroline. *J. Microcolumn Sep.*, 8 (1996) 137-143; *C.A.*, 124 (1996) 241492w.

See also 1625, 1655, 1713, 2433.

38b. *Anions*

2429 Choi, O.-K. and Cho, J.-S.: Analysis of inorganic anions in various drinking waters by capillary electrophoresis. *Anal. Sci. Technol.*, 8 (1995) 835-841; *C.A.*, 124 (1996) 211249e.

2430 Francois, C., Morin, P. and Dreux, M.: Adjusting the selectivity of inorganic anion separation by capillary electrophoresis. *J. High Resolut. Chromatogr.*, 19 (1996) 5-16.

2431 Galceran, M.T., Puignou, L. and Díez, M.: Comparison of different electroosmotic flow modifiers in the analysis of inorganic anions by capillary electrophoresis. *J. Chromatogr. A*, 732 (1996) 167-174.

- 2432 Hauser, P.C., Hong, A.P.C. and Renner, N.D.: Surface charge reversal for inorganic anion determination in capillary electrophoresis with an ion-selective microelectrode as detector. *J. Capillary Electrophor.*, 2 (1995) 209-212; *C.A.*, 124 (1996) 192580n.
- 2433 Jia, L., Zhang, H. and Hu, Z.: Separation and determination of chromium(VI), molybdenum(VI) and vanadium(V) in steel using capillary ion electrophoresis with direct UV detection. *J. Liq. Chromatogr.*, 19 (1996) 565-581.
- 2434 Soga, T. and Inoue, Y.: Simultaneous analysis of halides, oxoanions and metal oxoacids using capillary electrophoresis. In: Hatano, H. and Hanai, T. (Editors), *Int. Symp. Chromatogr., 35th Anniv. Res. Group Liq. Chromatogr. Jpn.*, World Scientific, Singapore, 1995, pp. 443-448; *C.A.*, 124 (1996) 74895w.
- 2435 Takayanagi, T., Wada, E., Oshima, M. and Motomizu, S.: (Determination of inorganic anions in environmental samples by capillary electrophoresis). *Kankyo Seigyo*, 17 (1995) 49-55; *C.A.*, 124 (1996) 218856j.
- 2436 Wang, T. and Li, S.F.Y.: Separation of polyphosphates and polycarboxylates by capillary electrophoresis in a carrier electrolyte containing adenosine 5'-triphosphate and cetyltrimethylammonium bromide with indirect UV detection. *J. Chromatogr. A*, 723 (1996) 197-205.

See also 1629.

39. RADIOACTIVE AND OTHER ISOTOPE COMPOUNDS

See 1655.