

Author Index Vol. 768

- Ahmed, A., see Govindaraju, K. 768(1997)3
- Altria, K.D., see Kelly, M.A. 768(1997)73
- Åman, P., see Fredriksson, H. 768(1997)325
- Amigo, L., see Recio, I. 768(1997)47
- Andersson, R., see Fredriksson, H. 768(1997)325
- Angioni, A., see Pirisi, F.M. 768(1997)207
- Bächmann, K., see Göttlicher, B. 768(1997)320
- Bandino, G., see Pirisi, F.M. 768(1997)207
- Belchior, A.P., see Bronze, M.R. 768(1997)143
- Berloni, A., see Cappiello, A. 768(1997)215
- Brinkman, U.A.Th., see Slobodník, J. 768(1997)239
- Bronze, M.R., Vilas Boas, L.F. and Belchior, A.P.
Analysis of old brandy and oak extracts by capillary electrophoresis 768(1997)143
- Cabras, P., see Pirisi, F.M. 768(1997)207
- Cappiello, A., Famiglini, G. and Berloni, A.
Large volume injection of acidic pesticides by reversed-phase micro high-performance liquid chromatography 768(1997)215
- Cetta, G., see Grimm, R. 768(1997)57
- Chang, C.-C., see Lin, C.-E. 768(1997)105
- Chen, Y.
Critical conditions for separating the microheterogeneous components of glycoproteins by capillary electrophoresis 768(1997)39
- Clark, B.J., see Kelly, M.A. 768(1997)73
- Degenhardt, M., see Kunkel, A. 768(1997)17
- Degenhardt, M. and Wätzig, H.
Purity control of carbamazepine by micellar electrokinetic chromatography 768(1997)113
- Dyne, K., see Grimm, R. 768(1997)57
- Erim, F.B.
Effect of cationic polymer on the separation of phenols by capillary electrophoresis 768(1997)161
- Fall, J., see Voelkel, A. 768(1997)271
- Famiglini, G., see Cappiello, A. 768(1997)215
- Ferguson, P.D., Goodall, D.M. and Loran, J.S.
Systematic approach to links between separations in capillary electrophoresis and liquid chromatography. IV. Application of binding constant-retention factor relationship to the separation of 2-, 3- and 4-methylbenzoate anions using β -cyclodextrin as selector 768(1997)29
- Fish, B.J.
Separation and quantification of geometric isomers in a leukotriene antagonist using cyclodextrins in capillary electrophoresis. Comparison to high-performance liquid chromatographic methods 768(1997)81
- Fredriksson, H., Andersson, R., Koch, K. and Åman, P.
Calibration of a size-exclusion chromatography system using fractions with defined amylopectin unit chains 768(1997)325
- Garau, V.L., see Pirisi, F.M. 768(1997)207
- Goodall, D.M., see Ferguson, P.D. 768(1997)29
- Gooijer, C., see Veraart, J.R. 768(1997)307
- Gordon, D.B., see Lord, G.A. 768(1997)9
- Göttlicher, B. and Bächmann, K.
Investigation of the separation efficiency of hydrophobic compounds in suspension electrokinetic chromatography 768(1997)320
- Govindaraju, K., Ahmed, A. and Lloyd, D.K.
Electroosmotic flow mobilities in open and packed capillaries with spermine and other amine flow modifiers 768(1997)3
- Grasserbauer, M., see Silgoner, I. 768(1997)259
- Grimm, R., Zanaboni, G., Viglio, S., Dyne, K., Valli, M., Cetta, G. and Iadarola, P.
Complete resolution of imidodipeptide mixtures in urine of prolidase-deficient patients using micellar electrokinetic chromatography 768(1997)57
- Günter, S., see Kunkel, A. 768(1997)125
- Hayase, S., Oda, Y., Honda, S. and Kakehi, K.
High-performance capillary electrophoresis of hyaluronic acid: determination of its amount and molecular mass 768(1997)295
- Helfferich, F.G.
Non-linear waves in chromatography. III. Multicomponent Langmuir and Langmuir-like systems 768(1997)169
- Honda, S., see Hayase, S. 768(1997)295
- Hows, M.E.P., Perrett, D. and Kay, J.
Optimisation of a simultaneous separation of sulphonamides, dihydrofolate reductase inhibitors and β -lactam antibiotics by capillary electrophoresis 768(1997)97
- Hsieh, Y.-Z., see Kuo, K.-L. 768(1997)334
- Iadarola, P., see Grimm, R. 768(1997)57
- Kakehi, K., see Hayase, S. 768(1997)295
- Karim dos Santos, M., see Pirisi, F.M. 768(1997)207
- Kay, J., see Hows, M.E.P. 768(1997)97
- Kelly, M.A., Altria, K.D. and Clark, B.J.
Approaches used in the reduction of buffer electrolysis effects for routine capillary electrophoresis procedures in pharmaceutical analysis 768(1997)73
- King, B.W., see Lord, G.A. 768(1997)9
- Koch, K., see Fredriksson, H. 768(1997)325
- Kollroser, M., see Trathnigg, B. 768(1997)223
- Kunkel, A., Degenhardt, M., Schirm, B. and Wätzig, H.
Performance of instruments and aspects of methodology and validation in quantitative capillary electrophoresis. An update 768(1997)17
- Kunkel, A., Günter, S. and Wätzig, H.
Quantitation of acetaminophen and salicylic acid in plasma using capillary electrophoresis without sample pretreatment. Improvement of precision 768(1997)125
- Kuo, K.-L. and Hsieh, Y.-Z.
Determination of preservatives in food products by cyclodextrin-modified capillary electrophoresis with multiwavelength detection 768(1997)334
- Lin, C.-E., Chang, C.-C. and Lin, W.-C.
Migration behavior and separation of sulfonamides in capillary zone electrophoresis. III. Citrate buffer as a background electrolyte 768(1997)105
- Lin, W.-C., see Lin, C.-E. 768(1997)105

- Lingeman, H., see Veraart, J.R. 768(1997)307
 Liška, I., see Slobodník, J. 768(1997)239
 Lloyd, D.K., see Govindaraju, K. 768(1997)3
 Loran, J.S., see Ferguson, P.D. 768(1997)29
 Lord, G.A., Gordon, D.B., Myers, P. and King, B.W.
 Tapers and restrictors for capillary electrochromatography and capillary electrochromatography-mass spectrometry 768(1997)9
 Louter, A.J.H., see Slobodník, J. 768(1997)239
 Meyer, V.R.
 Example of gradient elution in normal-phase liquid chromatography 768(1997)315
 Myers, P., see Lord, G.A. 768(1997)9
 Oda, Y., see Hayase, S. 768(1997)295
 Pérez-Rodríguez, M.-L., see Recio, I. 768(1997)47
 Perrett, D., see Hows, M.E.P. 768(1997)97
 Perrett, D., see Taylor, M.R. 768(1997)67
 Pirisi, F.M., Angioni, A., Cabras, P., Garau, V.L., Sanjust di Teulada, M.T., Karim dos Santos, M. and Bandino, G.
 Phenolic compounds in virgin olive oils. I. Low-wavelength quantitative determination of complex phenols by high-performance liquid chromatography under isocratic elution 768(1997)207
 Prošek, M., see Vovk, I. 768(1997)329
 Ramos, M., see Recio, I. 768(1997)47
 Recio, I., Pérez-Rodríguez, M.-L., Ramos, M. and Amigo, L.
 Capillary electrophoretic analysis of genetic variants of milk proteins from different species 768(1997)47
 Rosenberg, E., see Silgoner, I. 768(1997)259
 Sanjust di Teulada, M.T., see Pirisi, F.M. 768(1997)207
 Schirm, B., see Kunkel, A. 768(1997)17
 Schouten, Y., see Veraart, J.R. 768(1997)307
 Silgoner, I., Rosenberg, E. and Grasserbauer, M.
 Determination of volatile organic compounds in water by purge-and-trap gas chromatography coupled to atomic emission detection 768(1997)259
 Šlais, K., see Šťastná, M. 768(1997)283
 Slobodník, J., Louter, A.J.H., Vreuls, J.J., Liška, I. and Brinkman, U.A.Th.
 Monitoring of organic micropollutants in surface water by automated on-line trace-enrichment liquid and gas chromatographic systems with ultraviolet diode-array and mass spectrometric detection 768(1997)239
 Šťastná, M. and Šlais, K.
 Displacement electrophoresis of ampholytes in a continuous pH gradient moving in a capillary with a non-constant cross-section 768(1997)283
 Taylor, M.R. and Teale, P.
 Gradient capillary electrochromatography of drug mixtures with UV and electrospray ionisation mass spectrometric detection 768(1997)89
 Taylor, M.R., Westwood, S.A. and Perrett, D.
 Direct monitoring of enzyme reactions using micellar electrokinetic capillary chromatography. Optimisation of drug glucuronide and sulfate conjugate hydrolysis 768(1997)67
 Teale, P., see Taylor, M.R. 768(1997)89
 Trathnigg, B. and Kollroser, M.
 Liquid chromatography of polyethers using universal detectors. V. Quantitative aspects in the analysis of low-molecular-mass poly(ethylene glycol)s and their derivatives by reversed-phase high-performance liquid chromatography with an evaporative light scattering detector 768(1997)223
 Valli, M., see Grimm, R. 768(1997)57
 Veraart, J.R., Schouten, Y., Gooijer, C. and Lingeman, H.
 Evaluation of phytic acid as a buffer additive for the separation of proteins in capillary electrophoresis 768(1997)307
 Viglio, S., see Grimm, R. 768(1997)57
 Vilas Boas, L.F., see Bronze, M.R. 768(1997)143
 Voelkel, A. and Fall, J.
 Inverse gas chromatography for the examination of fractions separated from oil vacuum distillation residues 768(1997)271
 Vovk, I. and Prošek, M.
 Quantitative evaluation of chromatograms from totally illuminated thin-layer chromatographic plates 768(1997)329
 Vreuls, J.J., see Slobodník, J. 768(1997)239
 Wätzig, H., see Degenhardt, M. 768(1997)113
 Wätzig, H., see Kunkel, A. 768(1997)17
 Wätzig, H., see Kunkel, A. 768(1997)125
 Westwood, S.A., see Taylor, M.R. 768(1997)67
 Wren, S.A.C.
 Mobility measurements on dansylated amino acids 768(1997)153
 Yeung, E.S., see Zhang, N. 768(1997)135
 Zanaboni, G., see Grimm, R. 768(1997)57
 Zhang, N. and Yeung, E.S.
 Simultaneous separation and genetic typing of four short tandem repeat loci by capillary electrophoresis 768(1997)135